

- Inorg. Chem.*, **7**, 847 (1968).
 (16) J. P. Collman, N. Kim, J. L. Hoard, G. Lang, L. J. Radonovich, and C. A. Reed, Abstracts, 167th National Meeting of the American Chemical Society, Los Angeles, Calif., April 1974, No. INOR 29; full paper to be submitted to *J. Am. Chem. Soc.*
 (17) J. L. Hoard, *Science*, **174**, 1295 (1971).
 (18) M. Zerner, M. Gouterman, and H. Kobayashi, *Theor. Chim. Acta*, **6**, 363 (1966).
 (19) K. Moffat, R. S. Loe, and B. M. Hoffman, *J. Am. Chem. Soc.*, **96**, 5259 (1974).
 (20) J. P. Collman, R. R. Gagne, C. A. Reed, T. R. Halbert, G. Lang, and W. T. Robinson, *J. Am. Chem. Soc.*, **97**, 1427 (1975).

Barbara Gonzalez, Jay Kouba
 Sharon Yee, Christopher A. Reed*

Department of Chemistry, University of Southern California
 Los Angeles, California 90007

John F. Kirner, W. Robert Scheidt*

Department of Chemistry, University of Notre Dame
 Notre Dame, Indiana 46556

Received March 7, 1975

Stereospecific Acid-Catalyzed Rearrangement of 1,6-Dimethylpentacyclo[6.4.0.0^{2,7}.0^{3,10}.0^{6,9}]dodecane- 5,12-dione to a Bisnordiadamantane

Sir:

On treatment with trifluoroacetic acid at room temperature, pentacyclo[6.4.0.0^{2,7}.0^{3,10}.0^{6,9}]dodecane-5,12-diones having a methyl group at C-2 (**1**), synthesized photochemically from Diels-Alder dimers (**2**) of cyclohexa-2,4-dienones, reverted easily to **2** in high yield.¹ Stabilization of a carbonium ion at C-2 by a methyl group was assumed to be the most important requirement for this acid-catalyzed reversal.^{1b}

A cage compound, with a methyl group at C-1, on acid




Figure 1. Stereodrawing of compound **4** executed with the experimentally determined coordinates from a crystal structure analysis. The shaded ellipses represent oxygen atoms.


treatment would be expected to release most of the strain of the bicyclo[2.2.0]hexane system in a different way as shown here by the stereospecific rearrangement of the representative **3**.

When **3** was heated under reflux in trifluoroacetic acid for 15 min, or in benzene with *p*-TsOH for 45 min, the isomeric rearrangement product **4** was isolated in almost quantitative yield: mp 182–184°, from *n*-hexane; ν_{Nujol} 1735 cm^{-1} ; m/e 272 (M^+); ^1H nmr δ_{CDCl_3} 0.84 (s, 6 H), 0.96 (s, 6 H), 0.96 (s, 6 H), 1.03 (s, 6 H), 2.38 (m, 4 H), 2.48 (m, 2 H); ^{13}C nmr δ_{CDCl_3} 11.6 (CH_3), 20.6 (CH_3), 26.8 (CH_3), 48.4 (C), 52.2 (CH), 53.8 (CH), 57.2 (C), 57.5 (CH).

Protonation of one of the carbonyl groups in **3** causes the formation of the methyl-stabilized carbonium cation at C-1 through rearrangement of either bond a or b, followed by another set of *twofold Wagner-Meerwein rearrangements* to yield a less strained cage compound, such as **4**, **5**, or **6**. A few precedents of such rearrangements have been reported in simpler cases, propellanes² and spiranone.³ Both **4** and **5** have a twofold axis of symmetry whereas **6** has not. Since ^1H and ^{13}C NMR spectra clearly indicate that the product is symmetric, **6** is excluded. Although it is impossible to distinguish between **4** and **5** by the usual spectral data, there is a marked difference in their dipole moments: estimated value for **4** ca. 4.0 D; and for **5** ca. 0.0 D. The observed value (ca. 4.2 D) shows that **4** is the correct structure, a conclusion which is confirmed by Roentgen-ray analysis.

Compound **4** crystallizes in the monoclinic space group $P2_1/c$ with $a = 6.818$ (4) Å, $b = 12.586$ (5) Å, $c = 17.966$ (8) Å, and $\beta = 106.8$ (1)°. There is one molecule per asymmetric unit corresponding to a calculated crystal density of 1.22 g/cm^3 . The structure was solved by the symbolic addition procedure for centrosymmetric crystals⁴ and refined by full-matrix least-squares methods⁵ to an R factor of 5.8%. The stereodrawing⁶ in Figure 1 which was constructed with the experimentally determined atomic coordinates displays the results of the X-ray analysis. Within experimental error, the molecule has twofold rotation symmetry.⁷ So far cedrone⁸ seems to be the only other representative of the interesting bisnordiadamantane type.

The most important reason for the favored rearrangement of bond a must lie in the stability difference between the rearranged cations **3a** and **3b** both of which arise by conversion of the strained four-, six-, six-membered ring system in **3** (thick line) to the more stable five, five, six system with release of strain energy. Inspection of models clearly indicates that the six-membered cationic structures, **3a** and **3b** (shaded parts), are quite different, though the strain in the remainder of the molecules may be the same. The six-membered ring in **3a** is present in a normal chair conforma-



tion, whereas in **3b** the chair conformation is strongly distorted by a directly fused four-membered ring, destabilizing the structure of **3b**.

References and Notes

- (1) (a) T. Iwakuma, O. Yonemitsu, N. Kanamaru, K. Kimura, and B. Witkop, *Angew. Chem., Int. Ed. Engl.*, **12**, 72 (1973); (b) T. Iwakuma, K. Hirao, and O. Yonemitsu, *J. Am. Chem. Soc.*, **96**, 2570 (1974).
- (2) N. P. Peet, R. L. Cargill, and D. F. Bushey, *J. Org. Chem.*, **38**, 1218 (1973); for a review see R. L. Cargill, T. E. Jackson, N. P. Peet, and D. M. Pond, *Acc. Chem. Res.*, **7**, 106 (1974).
- (3) B. P. Mundy and R. D. Otzenberger, *J. Org. Chem.*, **38**, 2109 (1973).
- (4) J. Karle and I. L. Karle, *Acta Crystallogr.*, **21**, 849 (1966).
- (5) W. R. Busing, K. O. Martin, H. A. Levy, R. D. Ellison, W. C. Hamilton, J. A. Ibers, C. K. Johnson, and W. E. Thiessen, ORXFLS3, Oak Ridge National Laboratory, Oak Ridge, Tenn. 1971.
- (6) Both stereodrawings were drawn by computer using program ORTEP. C. K. Johnson, ORTEP, Report ORNL-3794, Oak Ridge National Laboratory, Oak Ridge, Tenn.

(7) Complete structural data will be submitted for publication by J. L. Flippen, *Acta Crystallogr.*

(8) J. A. Beisler, J. V. Silverton, A. Penttila, D. H. S. Horn, and H. M. Fales, *J. Am. Chem. Soc.*, **93**, 4850 (1971).

Ken-ichi Hirao, Mikio Taniguchi
Takeo Iwakuma, Osamu Yonemitsu*

Faculty of Pharmaceutical Sciences, Hokkaido University
Sapporo, Japan


Judith L. Flippen, Isabella L. Karle
Laboratory for the Structure of Matter
Naval Research Laboratory
Washington, D.C. 20375

Bernhard Witkop*
Laboratory of Chemistry, NIAMDD,
National Institutes of Health
Bethesda, Maryland 20014
Received December 4, 1974

Organoselenium Chemistry. α -Lithio Selenoxides and Selenides. Preparation and Further Transformation to Olefins, Dienes, and Allylic Alcohols

Sir:

Functionalized organolithium reagents have assumed an important role in synthetic organic chemistry. We report here the preparation and some reactions of several selenium stabilized organolithium reagents,¹ and synthetic applications based on the facile syn elimination of the selenoxide function.²⁻⁴ The reaction products of selenium stabilized organolithium compounds with electrophiles could in general undergo selenoxide elimination in two directions.⁵ Elimina-



tion away from the newly formed C-C bond (path a) results in an overall transformation synthetically equivalent to the operation of a vinyl anion; whereas, elimination across the new C-C bond (path b) results in a coupling of halides to form an olefin (alkyl selenides are usually prepared by nucleophilic displacements of halides or sulfonates by PhSe-Na). To be synthetically useful, reactions of this type must have one pathway predominant, either because one elimination is blocked (no H_a or H_b) or by the operation of factors favoring one pathway over the other. We have found systems in which the exclusive operation of either path a or path b can be achieved.

Methyl phenyl and benzyl phenyl⁷ selenoxides cannot undergo selenoxide elimination, and so preparation and handling pose no special problems. They are rapidly deprotonated at -78° by lithium diisopropylamide (LDA) giving the anions **1** and **2**. Longer chain alkyl selenoxides having β -hydrogens must be handled below 0° to avoid the elimination reaction. We have developed procedures for the low temperature in situ oxidation of phenyl alkyl selenides (by ozonization in ether at -78°,⁸ or reaction with *m*-chloroperbenzoic acid in THF at -10°⁹), and for deprotonation