

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/261915593>

ChemInform Abstract: Ti(III)–Catalyzed Cyclizations of Ketoepoxypolyprenes: Control over the Number of Rings and Unexpected Stereoselectivities.

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · APRIL 2014

Impact Factor: 12.11 · DOI: 10.1021/ja411942h · Source: PubMed

CITATIONS

8

READS

66

10 AUTHORS, INCLUDING:



Alba Millán

University of Bristol

15 PUBLICATIONS 133 CITATIONS

SEE PROFILE



Duane Choquesillo-Lazarte

Spanish National Research Council

138 PUBLICATIONS 1,159 CITATIONS

SEE PROFILE



Juan M. Garcia-Ruiz

University of Granada

279 PUBLICATIONS 4,420 CITATIONS

SEE PROFILE



Juan M Cuerva

University of Granada

115 PUBLICATIONS 2,110 CITATIONS

SEE PROFILE

Article

Ti(III)-Catalyzed Cyclizations of Ketoepoxypolyprenes: Control over the Number of Rings and Unexpected Stereoselectivities

Sara P. Morcillo, Delia Miguel, Sandra Resa, Ana Martin-Lasanta, Alba Millan, Duane Choquesillo-Lazarte, Juan Manuel Garcia-Ruiz, Antonio J. Mota, Jose Justicia, and Juan M. Cuerva

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/ja411942h • Publication Date (Web): 26 Apr 2014

Downloaded from <http://pubs.acs.org> on May 1, 2014

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications
High quality. High impact.

Ti(III)-Catalyzed Cyclizations of Ketoepoxypolyprenes: Control over the Number of Rings and Unexpected Stereoselectivities

Sara P. Morcillo,[†] Delia Miguel,[†] Sandra Resa,[†] Ana Martín-Lasanta,[†] Alba Millán,[†] Duane Choquesi-llo-Lazarte,[#] Juan M. García-Ruiz,[#] Antonio J. Mota,^{‡*} José Justicia,^{†*} and Juan M. Cuerva^{†*}


[†]Department of Organic Chemistry, Faculty of Sciences, University of Granada, C. U. Fuentenueva s/n, 18071 Granada, Spain. [#]Laboratorio de Estudios Cristalográficos, IACT, CSIC-University of Granada, P. T. Ciencias de la Salud, 18100, Granada, Spain. [‡]Department of Inorganic Chemistry, Faculty of Sciences, University of Granada, C. U. Fuentenueva s/n, 18071 Granada, Spain.

ABSTRACT: We describe a new strategy to control the number of cyclization steps in bioinspired radical (poly)cyclizations involving epoxypolyenes containing keto units positioned along the polyene chain. This approach provides an unprecedentedly straightforward access to natural terpenoids with pendant unsaturated side chains. Additionally, in the case of bi- and tricyclizations, decalins with *cis* stereochemistry have been obtained as a consequence of the presence of the ketone. The preferential formation of *cis*-fused adducts was rationalized using DFT calculations. This result is completely unprecedented in biomimetic cyclizations and permits the access to natural terpenoids with this stereochemistry, as well as to non-natural analogues

Introduction

In recent years, bioinspired radical cyclizations have emerged as a powerful tool for the efficient synthesis of different terpenic structures, constituting one of the best ways for the preparation of structures such as these.¹ It is also worth noting that, in many cases, these cyclizations represent a complementary approach to more developed biomimetic cationic cyclizations.² Moreover, it is expected that new chemo- and stereoselectivities could be obtained taking into account the significant differences between their corresponding reactive intermediates and the catalysts used. One of the main differences is the nature of the alkenes present in the polyprenic starting material. Electron deficient alkenes are not usually suitable functionalities for cationic cyclizations of simple polyprenic precursors.^{2,3} On the other hand, these electron deficient alkenes can efficiently react with carbon centered radicals.⁴ Thus, the expected final products using conventional cationic cyclizations and radical ones might not be the same. Within this context, we focused our attention on Ti(III)-catalyzed radical cyclizations^{5–7} of epoxypolyprenes (**A**) presenting an internal keto functionality in their structures (Scheme 1). We hypothesized that enol radical **B**, obtained during the bioinspired radical cyclization, could be readily trapped by highly oxophilic radical species Cp₂TiCl.^{8,9}


Scheme 1. Working hypothesis



After an acidic quenching, the final (poly)cyclic product would provide a saturated ketone, which could be transformed in other functions present in natural terpenes. Notably, the

number of rings in the final carbocycle would depend on the position of the keto group, and not on the number of prenyl subunits in the starting polyprene, as is the case in previous cationic or radical cyclizations.^{1,2} Thus, the cyclization might be terminated even in the presence of additional unsaturations. This result would mimic the incomplete polycyclizations, which are responsible for the presence of terpenes with (poly)preenic side chains in nature.

Figure 1. Synthesized compounds 1–6.



These compounds are widespread and are present in several natural organisms, such as plants, insects, fungus, etc.¹⁰ Moreover, they exhibit diverse and interesting biological properties such as anti-inflammatory, cytotoxic, antifeedant, and antimicrobial effects.¹⁰ All these facts contribute to making them attractive targets in organic synthesis. Nevertheless, efficient methods for their synthesis have remained elusive


until now owing to the fact that the efficient control of the number of cyclization steps is challenging. A remarkable example is the use of allyl silanes by Corey's group to orientate the termination in cationic biomimetic cyclizations.¹¹ Nevertheless, the preparation of the starting materials containing such functionality is not trivial,¹¹ increasing the complexity and the number of steps in the synthetic sequence. Interestingly, we have also observed that in some substrates tested in this work, the keto group can dramatically affect the stereochemistry of the reaction, resulting in unprecedented bioinspired cyclizations yielding exclusively *cis*-fused decalins.¹² *Cis*-decalins¹³ are not very common in terpenes, although it has been very recently suggested that they may possess enhanced biological activities compared with their *trans* analogs.¹⁴

In this study we conclude that α,β -unsaturated ketones located in suitable positions of an epoxypolyprene chain can be used to control two key features in the synthesis of natural terpenes: i) the number of carbocycles, and ii) the stereochemistry of the cyclization. As an example of the usefulness of this new protocol, we have used some of the resulting cyclization products in the synthesis of terpenic structures **1-6** (see Figure 1). Thus, functional group interconversions from original saturated ketone to α,β -unsaturated ketone or alkene groups allow the total synthesis of two natural terpenes **1** and **2**,^{15,16} the unnatural *cis*-fused terpenes **3-5**, and an advanced intermediate **6** in the synthesis of fregenedadiol.¹⁷ Interestingly, more common *trans*-decalines are also available from α,β -unsaturated ketones using known protocols.¹⁸


Results and discussion

Cyclization reaction. The required starting epoxypolyprenes were straightforwardly synthesized following the method depicted in Scheme 2. First, we prepared the corresponding hydroxypolyprenes using a modification of our previously described α -prenylation protocol,^{7j,19} yielding the corresponding hydroxypolyprenes **7-19** (see Scheme 3).²⁰

Scheme 2. General protocol for the preparation of starting ketoepoxypolyprenes **33-45**.



Scheme 3. Hydroxypolyprenes **7-19** from Ti(III)-mediated Barbier-type reactions.




























Subsequent hydroxyl group oxidation of allylic alcohols **7-19** by Dess-Martin periodinane to yield the corresponding ketones **20-32**,^{20,21} and regioselective epoxidation using described methods⁷ allowed us to prepare a set of starting ketoepoxypolyprenes **33-45** with different lengths and functionalities.²⁰

With these starting ketoepoxypolyprenes **33-45** in hand, we submitted them to our titanocene(III)-bioinspired cyclization procedure using substoichiometric amounts of Cp_2TiCl .⁷ In this case, we selected the usual aprotic combination of $\text{Me}_3\text{SiCl}/2,4,6\text{-collidine}$ and Mn as a regenerating agent for Cp_2TiCl species since it is compatible with the presence of the oxirane. The results are summarized in Table 1.

Table 1. Ti(III)-catalyzed cyclization of epoxypolyprenes **33-45**.^a

Entry	Starting epoxide	Product ^b	Yield (%)
-------	------------------	----------------------	-----------

1			99 ^c
2			64 ^c
3			80
4			67 ^d
5			59
6			45
7			67
8			60
9			56 ^e
10			59 ^e
11			50
12			41




^a Conditions: Cp₂TiCl₂ (0.2 mmol), Mn (8 mmol), 2,4,6-collidine (7 mmol), and TMSCl (4 mmol). ^b Products containing a methyl group in the α position of the ketone (C-6 or C-8) were usually isolated as a mixture of α:β epimers. See SI for details. ^c An additional minor isomer is also observed. See SI for details. ^d 1:1 mixture of epimers at C-9. ^e 1:1 mixture of epimers at C-13. ^f 9:1 mixture of *trans-transoid-cis* and *trans-transoid-trans* isomers.

The results depicted in Table 1 show that our initial hypothesis was correct, allowing us to prepare different mono-, bi-, and even tricyclic compounds, controlling very efficiently the cyclization sequence. In all cases, we did not detect subsequent additions of the enol radical to other alkenes present in the molecule, even in trace amounts (entries 2-4, 9, 11 and 13), independently of the substitution pattern and functionality of the final side chain. Nevertheless, such intermediate radical or the final titanocene(IV) enolate is able to react with the carbonyl group present in the side chain of substrate **37**, giving aldol **50** (entry 5). We also tried regioisomeric ketone **38**. In this case, a Michael-type addition exclusively yielded cyclopentane **51** (entry 6). As expected, the relative position of carbonyl group is essential for the regioselectivity of the radical addition.

Furthermore, the yields observed were comparable and even higher to those obtained in the parent bioinspired protocol.^{1,7} The cyclization process is also highly stereoselective with the exception of the methyl group in vicinal position to the ketone group, which was usually obtained as a mixture of epimers. This presents a minor drawback, however, since this position is usually modified in the subsequent synthetic steps. In monocycles **46-49** we mainly observed a *cis*-relationship between the hydroxyl group at C-3 and the side chain at C-5.⁷ Remarkably, bicyclic products **52-56** presented a *cis*-fused decalin despite the fact that radical additions to double bonds with *E* configuration usually yield *trans* decalines.¹ This unprecedented stereochemistry was confirmed by X-ray analysis of *p*-nitrobenzoate **59**, a simple derivative of **52**. Thus, we could also assign the relative configuration of the side chain at C-9.

Figure 2. Crystal structure for compound **59**.




Tricyclic products **57-58** mainly presented a *trans* fusion between the two first cyclohexanes, and a *cis* fusion between the second and third cyclohexane rings,²⁰ suggesting that the *cis* stereochemistry is intrinsic to the radical addition reaction

when the ketone group is present. The reason for this surprising stereoselectivity was not clear. The use of different titanocene(III) catalysts with different steric or electronic characteristics always yielded the *cis*-fused compounds. However, in the presence of other functional group in the same position, such as acetate, *trans*-fused structures were obtained.^{7j}

Theoretical calculations. To shed light on this unexpected chemo- and stereoselectivity, we made use of theoretical calculations²⁰ carried out at the DFT-B3LYP level²²⁻²⁴ with the Gaussian 09 program.²⁵ The geometries were fully optimized by the gradient technique using polarized 6-31G* basis set for all the atoms.²⁶ This theoretical level has been extensively used to rationalize titanocene(III) chemistry.^{7a,d,f,k,9a,b,d} The nature of the optimized structures, either transition states or intermediates, was assessed through a frequency calculation, and the changes of Gibbs free reaction energies (ΔG values) were obtained by taking into account zero-point energies, thermal motion, and entropy contribution at standard conditions (temperature of 298.15 K, pressure of 1 atm). We have concentrated our discussion on the corresponding enthalpic values.

A key point of this theoretical study is the suitable selection of the titanocene(III) active specie in the catalytic cycle. Using epoxypolyprene **I** as model, we explored the coordination capabilities of epoxide and ketone groups toward different oxophilic titanocene(III) complexes. Taking into account that THF is used as solvent, the presence of a THF molecule as ligand has also been considered to maintain all the potential intermediates coordinatively saturated, thus avoiding any artificial bias during the calculations.^{27,28} Cationic complex $\text{Cp}_2\text{Ti}^+(\text{THF})_2$ resulted to be the best ligand for compound **I**.^{29,30} Moreover, Gansäuer's group has experimentally demonstrated that coordination of epoxides to Cp_2TiCl is accompanied by dissociation of the chloride anion to form the true cationic active specie.³¹ Therefore, we selected titanocene(III) cationic complexes as the active species in our catalytic cycle. In this case, we also guaranty a homogeneous treatment of the energies, avoiding artificial charge separations. After an exhaustive sampling of structures presenting different conformations and binding geometries, we could conclude that complex **II** is slightly favored (3.6 kcal mol⁻¹) in free energy terms (Scheme 4). As expected, the complex possessing both groups coordinated simultaneously to Cp_2Ti^+ (**IV**) is the most favorable one by 11 kcal mol⁻¹ and can be considered the resting state of the system.^{8,32}


UB3LYP/6-31G(d) (C,H,O,Ti) E + ZPE (kcal mol⁻¹) (G)




Scheme 4. Calculated minimum energy structures **II-IV** of epoxypolyprene **I** and $\text{Cp}_2\text{Ti}^+(\text{THF})_2$

Homolytic epoxide opening of such complexes (**II-IV**) leads to a tertiary radical able to undergo 6-*endo*-trig radical cyclizations to yield monocyclic radicals. Taking into account the thermodynamic stability of **IV** we initially focused our attention in such intermediate. Homolytic epoxide opening of **IV** to yield radical **V** is slightly endothermic and takes place with a calculated activation energy of 16.6 kcal mol⁻¹ (Scheme 5a). Then, we explored different cyclization processes from radical **V** to monocyclic radical type **VI**. For this rigid structure, we could only find one chair-like and one boat-like transition states. The lowest energy for the transition state was found for templated structure **VI_{boat}** (17.8 kcal mol⁻¹). This cyclization is slightly endothermic (3.5 kcal mol⁻¹) in clear contrast with the thermodynamically unfavorable monocycle **VI_{chair}** (17.1 kcal mol⁻¹). These calculated values can be explained based on the geometrical constraints of intermediates **V** (Scheme 5c). In the transition state from **V_{boat}** to **VI_{boat}** the geometry of the system allows an efficient conjugation between the alkene and the titanocene(IV)-activated carbonyl group and, therefore, the radical addition is highly favoured. On the other hand, in the transition state from **V_{chair}** to **VI_{chair}** the alkene is rotated and the conjugation is disrupted, disfavoring the radical addition.

Taking into account the modest binding energies calculated before (Scheme 4) a dynamic situation is expected at room temperature. Within this context, a Curtin-Hammett scenario could be operative. In this sense, we also studied the behavior of energetically unfavorable non-templated radical **VII**, which derived from complex **II** (Scheme 5b) to rule out such possibility. The lowest activation energy (16.2 kcal mol⁻¹ referred to **VII**) corresponds to a monocyclization process yielding monocyclic radical **VIII_{trans}**, which possesses the opposite stereochemistry to that experimentally observed. However, the difference between the lowest activation energies for both reaction pathways is less than 2 kcal mol⁻¹. Consequently the experimental results, in which the final monocyclic radical present a *cis* relationship between the hydroxyl group at C-3 and the side chain at C-5, can be explained based on a template effect derived from the coordination capabilities of the titanium center. Such reaction pathway is globally favored by 9 kcal mol⁻¹. Spin density of key monocyclic **VI_{boat}** was then calculated showing that the electron is mainly located at the carbon atom and could be able to undergo subsequent radical additions (Scheme 6).




Scheme 5. Activation and reaction energies calculated for cyclization reactions of the model radicals a) V, b) VII, and c) geometrical details of transition states V_{chair} to VI_{chair} and V_{boat} to VI_{boat} (some atoms have been omitted for clarity). Energies are referenced to resting state IV from Scheme 4. TS=transition state, ZPE=zeropoint energy.




Scheme 6. Calculated spin densities for (a) V, (b) TS-V_{boat}-VI_{boat} and (c) VI. The isodensity surfaces represented correspond to a cut-off value of 0.018 e·bohr⁻³.

At these point we explored three subsequent reactions of this enol radical (Scheme 7): (a) a second radical addition, (b) a hydrogen-atom transfer from THF, and (c) a reaction with another titanocene(III) radical center. In the first case, the templated structure VI_{boat} can only cyclize by the β-face to yield the bicyclic radical IX. This reaction is slightly endothermic (3.9 kcal mol⁻¹) and presents a moderate activation energy (11.7 kcal mol⁻¹). With this calculated data, we can consider that the cyclization reaction is in fact reversible³³ and the monocyclic radical VI_{boat} dominates the equilibrium. On the other hand, the solvent (THF) can be considered as a hydrogen-atom source and the enol radical could be directly deactivated by a hydrogen-atom transfer reaction. We explored such possibility to explain the incapacity of intermediate VI_{boat} to continue the cyclization process. We found a viable transition state in which the hydrogen atom is transferred to the carbon atom.³⁴ Such reaction resulted energetically disfavored (7.4 kcal mol⁻¹) and only could explain the experimental results if we assume that the resulting THF-derived radical diffuses in the solvent and are irreversibly trapped. As we proposed in our working hypothesis, the direct reaction of templated enol radical VI_{boat} with Cp₂TiCl, present in the reaction media, is barrierless and highly exothermic (-43.9 kcal mol⁻¹). This favorable process joined to the stabilization of the radical by the carbonyl group could also explain why subsequent cyclizations were never observed. Finally, the cycle can proceed by reaction of the titanocene(IV) alkoxides XI by other oxophilic species such as TMSCl. The final Ti(IV) chlorides can be reduced by manganese dust thus closing the catalytic cycle.




Scheme 7. Activation and reaction energies calculated for reactions of the model radical VI_{boat} . TS=transition state, ZPE=zeropoint energy.

All the previous findings can be extended to the rest of the described cyclizations. Nevertheless, the reason of the observed *cis* stereoselectivity in the bi- and tricyclization processes is not clear. In this case, we used model monocyclic radical **XII** in our calculations. This specie is in equilibrium with two Ti-templates **XIII**. Such species are energetically favored in free energy terms, specially the *exo* form **XIII_{exo}** (-12.2 kcal mol⁻¹). Interestingly, the penalty cost to dissociate the carbonyl group is low, allowing interconversion from **XIII_{exo}** to **XIII_{endo}** at room temperature, via radical **XII** or even through a direct rotation (Scheme 8). In Scheme 9, we summarized the activation energies for the formation of bicyclic products **XIV** and **XV** from both potential intermediates **XII** and **XIII_{endo}**. As it can be seen, cyclization of **XII** mainly would lead to *trans*-decalins in contrast with the observed stereochemistry. Intermediate **XIII_{endo}** can only evolve to the final *cis* product. Interestingly, in this case we could find two transition states leading to two diastereoisomers. The computed energy for the transition state to yield **XV_b**, which present the experimentally observed stereochemistry, is only 6.9 kcal mol⁻¹.




Scheme 8. Calculated energies for the proposed Ti-templated radical intermediates, **XIII_{exo}** and **XIII_{endo}**, on the stereoselective *cis* bicycle formation pathway using intermediate **XII** as reference, ZPE=zeropoint energy.



Scheme 9. Activation and reaction energies calculated for reactions of the model radicals **XII** and **XIII**. TS=transition state, ZPE=zeropoint energy.

On the other hand, the observed *cis* ring closing in the formation of the tricycle cannot be due to any template effect, which is not plausible from a structural point of view. To study this situation we used model bicyclic radical compound **XVI**. Remarkably, the side chain presenting the ketone group adopts less easily the required *trans* 6-membered transition state once the bicycle (*trans*) is formed. We calculated both reaction mechanisms as it is shown in Scheme 10. Once again the *cis* pathway was favored (**XVI_{cis}** to **XVII_{cis}**) due to a low-energy transition state (12.5 kcal mol⁻¹), even if a pre-arrangement (of about 4.1 kcal mol⁻¹) is required.³⁵




Scheme 10. Activation and reaction energies calculated for reactions of the model radicals **XVI**. TS=transition state, ZPE=zeropoint energy.

To sum up, theoretical calculations revealed that the presence of the ketone group is relevant for two processes. Firstly, it is able to stabilize transient radicals and also to react with oxophilic radical species in the reaction media, thus avoiding subsequent cyclization reactions. Secondly, it forces the corresponding side chain to follow a reaction pathway through the β -face, promoting the *cis* stereochemistry in the final cyclization reaction. The reason for this preferred stereochemistry is a combination of template (for the bicyclic structures, see above) and/or conformational effects (for the tricyclic structures).

Synthetic applications. Monocyclic structures **46-49** are versatile synthons and they can be easily transformed into different terpenic substructures. Thus, the synthesis of sesquiterpene **1**, isolated from the plant *Atermisia chamaemelifolia*,¹⁵ started from monocycle **49** (Scheme 11). It was transformed in the diacetate **60** in two steps with excellent global yield. Subsequently, the hydroxyl group was mesylated and the mesyl group eliminated by basic treatment, yielding the corresponding trisubstituted alkene in good yield (61 %, two steps). Finally, selective acetylation of secondary hydroxyl group gave **1**, in only 5 steps from **49** and in a 48 % overall yield. Sesquiterpene **2**, obtained from the plant *Celastopholis glauca*,¹⁶ was prepared from cyclization product **48** using a similar procedure in only 5 steps (57 % overall yield). Gratifyingly, this new strategy allows access to elusive natural terpenoids with excellent yields and a reasonable number of synthetic steps, thus demonstrating the usefulness of our methodology.


Scheme 11. Synthesis of sesquiterpenes **1** and **2**.



A similar synthetic sequence allowed us to prepare the *cis*-fused terpenic structures **3-4**, which are structurally related to existing natural labdanes (Scheme 12).^{36,37} As expected, comparison between their ¹H and ¹³C NMR spectra with biblio-


graphic *trans* stereoisomers showed differences between the natural compounds and our synthetic analogues.

Scheme 12. Synthesis of *cis*-fused compounds **3-4**.



Beyond elimination processes, the dehydrogenation of cyclohexanones present in compounds **46-58** might also yield interesting terpenic building blocks. Thus for example, we explored the possibility with model substrate **63** (Scheme 13). Two conjugated double bonds can be introduced sequentially in the structure using Br₂ in AcOH and SeO₂.²⁰ Compounds **64** and **65** are related with dri-8-en-7-one³⁸ and dri-5,8-dien-7-one,³⁹ interesting building blocks for drimane synthesis. The same sequence was carried out with compound **66**. α,β -Unsaturated ketone **5** was obtained in good yield and is structurally related with known labdanes, as rhinocerotinoic acid.⁴⁰ The second oxidation step yielded compound **6**, an advanced precursor for the synthesis of fregenedadiol,¹⁷ a bicyclic diterpene containing an aromatic ring.

Scheme 13. Transformation of **63** and **66** into natural related structures.



Conclusions

To sum up, in this paper we have described a new strategy to control Ti(III)-catalyzed bioinspired radical cyclizations using α,β -unsaturated ketones placed in suitable positions in the starting ketoepoxypolyenes. These starting materials are easily prepared using previously described titanocene(III)-mediated Barbier-type prenylations. This method allowed us to control the number of cyclizations, yielding the corresponding cyclic products with complete selectivity and high yields. Preparing such structures is challenging using other classical biomimetic cyclization processes. The presence of ketone groups in the final cyclization products facilitates the installation of tri- or tetrasubstituted alkenes usually present in natural terpenes. This alkene regioselectivity is also complementary to the exocyclic regioselectivity observed in the parent Cp₂TiCl-catalyzed reaction. This procedure has allowed us to synthesize the natural sesquiterpenoids **1** and **2**, in a few steps and with acceptable yields, meeting the demand for selectivity and atom- and step economy required for any modern synthesis.⁴¹ Additionally, the presence of ketone groups in the polyene leads to the formation of *cis*-fused decalins from *E*-alkenes, which is unprecedented in the field of biomimetic radical cyclizations. A possible explanation has been explored by means of DFT calculations, which showed that either template

or conformational effects could operate to give this preferred stereochemistry. This fact is especially relevant for the exploration of new compounds with important biological activities.

Experimental Section

General Details. Deoxygenated solvents and reagents were used for all reactions involving Cp_2TiCl_2 . THF was freshly distilled from Na. CH_2Cl_2 was freshly distilled from P_2O_5 . Products were purified by flash chromatography on Merck silica gel 50. Yields refer to analytically pure samples. NMR spectra were recorded in NMR 300, 400 and 500 MHz spectrometers.

General procedure for Ti^{III} -catalyzed bioinspired cyclizations: Strictly deoxygenated THF (20 mL) was added to a mixture of Cp_2TiCl_2 (0.2 mmol) and Mn dust (8 mmol) under an Ar atmosphere and the suspension was stirred at room temperature until it turned lime green (after about 15 min). Then, a solution of epoxypolyprene (1 mmol), 2,4,6-collidine (7 mmol) in THF (2 mL), and Me_3SiCl (4 mmol) were added and the mixture was stirred for 16 h. The reaction was then quenched with 2N HCl and extracted with EtOAc. The organic layer was washed with brine, dried (anhyd Na_2SO_4) and the solvent removed. Products **46–58** were isolated by flash chromatography of the residue (hexane/EtOAc) and characterized by spectroscopic techniques. Results are depicted in Table 1. See SI for more experimental details.

ASSOCIATED CONTENT

General experimental details. Synthesis of all new substrates and compounds **1–6**. ^1H NMR and ^{13}C NMR spectra of all new compounds. Computational data. This material is available free of charge via the internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

* mota.avila@gmail.com; jjusti@ugr.es; jmcuerva@ugr.es

Funding Sources

This research was funded by the Regional Government of Andalucía (project P09-FQM-4571), the Spanish MICINN (project CTQ-2011.22455), the University of Granada (CEI-BIOTIC), and “Factoría de Cristalización, CONSOLIDER INGENIO-2010”.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENT

SPM and DM thank the Regional Government of Andalucía for their contracts. AML is grateful to the MICINN for her FPU fellowship. Authors thank the Centro de Servicios de Informática y Redes de Comunicaciones (CSIRC), Universidad de Granada, for providing the computing time.

REFERENCES

- (1) Justicia, J.; Álvarez de Cienfuegos, L.; Campaña, A. G.; Miguel, D.; Jakoby, V.; Gansäuer, A.; Cuerva, J. M. *Chem. Soc. Rev.* **2011**, *40*, 3525–3537.
- (2) (a) Wendt, K. U.; Schulz, G. E.; Corey, E. J.; Liu, D. R. *Angew. Chem. Int. Ed.* **2000**, *39*, 2812–2833. (b) Yoder, R. A.; Johnston, J. N. *Chem. Rev.* **2005**, *105*, 4730–4756. (c) Yamamoto, H.; Futatsugi, K. *Angew. Chem. Int. Ed.* **2005**, *44*, 1924–1942.

- (3) Li, S.; Chiu, P. *Tetrahedron Lett.* **2008**, *49*, 1741–1744.
- (4) (a) *Radicals in Organic Synthesis*; Renaud, P., Sibi, M. P., Eds.; Wiley-VCH: Weinheim, Germany, 2001; Vol. 2. (b) Curran, D. P. In *Comprehensive Organic Synthesis*; Trost, B. M.; Fleming, I. Eds.; Pergamon Press: Oxford, UK, 1991; Vol. 4, pp 715–830.
- (5) For seminal paper, see: (a) RajanBabu, T. V.; Nugent, W. A. *J. Am. Chem. Soc.* **1989**, *111*, 4525–4527. (b) Gansäuer, A.; Bluhm, H.; Rinker, B.; Narayan, S.; Schick, M.; Lauterbach, T.; Pierobon, M. *Chem. Eur. J.* **2003**, *9*, 531–542. (c) Gansäuer, A.; Pierobon, M.; Bluhm, H. *Angew. Chem. Int. Ed.* **1998**, *37*, 101–103. (d) Gansäuer, A.; Bluhm, H.; Pierobon, M. *J. Am. Chem. Soc.* **1998**, *120*, 12849–12859. (e) Gansäuer, A.; Bluhm, H. *Chem. Commun.* **1998**, 2143–2144. (f) Gansäuer, A.; Lauterbach, T.; Bluhm, H.; Noltemeyer, M. *Angew. Chem. Int. Ed.* **1999**, *38*, 2909–2910.
- (6) For recent reviews, see: (a) Cuerva, J. M.; Justicia, J.; Oller-López, J. L.; Oltra, J. E. *Top. Curr. Chem.* **2006**, *264*, 63–91. (b) Gansäuer, A.; Justicia, J.; Fan, C. -A.; Worgull, D.; Piester, F. *Top. Curr. Chem.* **2007**, *279*, 25–52.
- (7) (a) Justicia, J.; Rosales, A.; Buñuel, E.; Oller-López, J. L.; Valdivia, M.; Haidour, A.; Oltra, J. E.; Barrero, A. F.; Cárdenas, D. J.; Cuerva, J. M. *Chem. Eur. J.* **2004**, *10*, 1778–1788. (b) Justicia, J.; Oltra, J. E.; Cuerva, J. M. *Tetrahedron Lett.* **2004**, *45*, 4293–4296. (c) Justicia, J.; Oltra, J. E.; Cuerva, J. M. *J. Org. Chem.* **2004**, *69*, 5803–5806. (d) Justicia, J.; Oller-López, J. L.; Campaña, A. G.; Oltra, J. E.; Cuerva, J. M.; Buñuel, E.; Cárdenas, D. J. *J. Am. Chem. Soc.* **2005**, *127*, 14911–14921. (e) Justicia, J.; Oltra, J. E.; Barrero, A. F.; Guadafío, A.; González-Coloma, A.; Cuerva, J. M. *Eur. J. Org. Chem.* **2005**, 712–718. (f) Justicia, J.; Campaña, A. G.; Bazdi, B.; Robles, R.; Cuerva, J. M.; Oltra, J. E. *Adv. Synth. Catal.* **2008**, *350*, 571–576. (g) Gansäuer, A.; Worgull, D.; Justicia, J. *Synthesis* **2006**, 2151–2154. (h) Gansäuer, A.; Rosales, A.; Justicia, J. *Synlett* **2006**, 927–929. (i) Gansäuer, A.; Justicia, J.; Rosales, A.; Worgull, D.; Rinker, B.; Cuerva, J. M.; Oltra, J. E. *Eur. J. Org. Chem.* **2006**, 4115–4127. (j) Jiménez, T.; Morcillo, S. P.; Martín-Lasanta, A.; Collado-Sanz, D.; Cárdenas, D. J.; Gansäuer, A.; Justicia, J.; Cuerva, J. M. *Chem. Eur. J.* **2012**, *18*, 12825–12833. (k) Justicia, J.; Jiménez, T.; Miguel, D.; Contreras-Montoya, R.; Chahboun, R.; Álvarez-Manzaneda, E.; Collado-Sanz, D.; Cárdenas, D. J.; Cuerva, J. M. *Chem. Eur. J.* **2013**, *19*, 12825–12833.
- (8) (a) Gansäuer, A.; Lauterbach, Geich-Gimbel, D. *Chem. Eur. J.* **2004**, *10*, 4983–4990. (b) Friedrich, J.; Dolg, M.; Gansäuer, A.; Geich-Gimbel, D.; Lauterbach, T. *J. Am. Chem. Soc.* **2005**, *127*, 7071–7077. (c) Friedrich, J.; Walczak, K.; Dolg, M.; Piester, F.; Lauterbach, T.; Worgull, D.; Gansäuer, A. *J. Am. Chem. Soc.* **2008**, *130*, 1788–1796. (d) Gansäuer, A. *Chem. Commun.* **1997**, 457–458. (e) Gansäuer, A.; Moschioni, M.; Bauer, D. *Eur. J. Org. Chem.* **1998**, 1923–1927. (f) Gansäuer, A.; Bauer, D. *Eur. J. Org. Chem.* **1998**, 2673–2676. (g) Gansäuer, A.; Bauer, D. *J. Org. Chem.* **1998**, *63*, 2070–2071. (h) Yamamoto, Y.; Hattori, R.; Miwa, T.; Nakagai, Y.; Kubota, T.; Yamamoto, C.; Okamoto, Y.; Itoh, K. *J. Org. Chem.* **2001**, *66*, 3865–3870. (i) Paradas, M.; Campaña, A. G.; Estévez, R. E.; Álvarez de Cienfuegos, L.; Jiménez, T.; Robles, R.; Cuerva, J. M.; Oltra, J. E. *J. Org. Chem.* **2009**, *74*, 3616–3619. (j) Feurer, M.; Frey, G.; Luu, H.-T.; Kratzert, D.; Streuff, J. *Chem. Commun.* **2014**, ASAP DOI: 10.1039/C3CC46894A. (k) Frey, G.; Luu, H. -T.; Bichovski, P.; Feurer, M.; Streuff, J. *Angew. Chem. Int. Ed.* **2013**, *52*, 7131–7134. (l) Streuff, J.; Feurer, M.; Bichovski, P.; Frey, G.; Gellrich, U. *Angew. Chem. Int. Ed.* **2012**, *51*, 8661–8664. (m) Streuff, J. *Chem. Eur. J.* **2011**, *17*, 5507–5510.
- (9) (a) Cuerva, J. M.; Campaña, A. G.; Justicia, J.; Rosales, A.; Oller-López, J. L.; Robles, R.; Cárdenas, D. J.; Buñuel, E.; Oltra, J. E. *Angew. Chem. Int. Ed.* **2006**, *45*, 5522–5526. (b) Paradas, M.; Campaña, A. G.; Marcos, M. L.; Justicia, J.; Haidour, A.; Robles, R.; Cárdenas, D. J.; Oltra, J. E.; Cuerva, J. M. *Dalton Trans.* **2010**, *39*, 8796–8800. (c) Jiménez, T.; Campaña, A. G.; Bazdi, B.; Paradas, M.; Arráez-Román, D.; Segura-Carretero, A.; Fernández-Gutiérrez, A.; Oltra, J. E.; Robles, R.; Justicia, J.; Cuerva, J. M. *Eur. J. Org. Chem.* **2010**, 4288–4295. (d) Paradas, M.; Campaña, A. G.; Jiménez, T.; Robles, R.; Oltra, J. E.; Buñuel, E.; Justicia, J.; Cárdenas, D. J.; Cuerva, J. M. *J. Am. Chem. Soc.* **2010**, *132*, 12748–12756. (e) Gansäuer, A.; Behlendorf, M.; Cangönül, A.; Kube, C.; Cuerva, J. M.; Friedrich,

- J.; van Gastel, M. *Angew. Chem. Int. Ed.* **2012**, *51*, 3266–3270. (f) Gansäuer, A.; Klatte, M.; Braendle, G. M.; Friedrich, J. *Angew. Chem. Int. Ed.* **2012**, *51*, 8891–8894. (g) Millán, A.; Álvarez de Cienfuegos, L.; Miguel, D.; Campaña, A. G.; Cuerva, J. M. *Org. Lett.* **2012**, *14*, 5984–5987. (h) Kessler, M.; Hansen, S.; Godemann, C.; Spannenberg, A.; Beweries, T. *Chem. Eur. J.* **2013**, *19*, 6350–6357.
- (10) (a) Frijia, L. M. T.; Frade, R. F. M.; Alfonso, C. A. M. *Chem. Rev.* **2011**, *111*, 4418–4452 and references cited therein. For other examples, see: (b) Akihisa, T.; Arai, K.; Kimura, Y.; Koike, K.; Kokke, W. C. M.; Shibata, C. T.; Nikaido, T. *J. Nat. Prod.* **1999**, *62*, 265–268. (c) Kimura, I.; Yoshikawa, M.; Kobayashi, S.; Sugihara, Y.; Suzuki, M.; Oominami, H.; Murakami, T.; Matsuda, H.; Doiphode, V. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 985–989. (d) Pale-Grosdemange, C.; Merkofer, T.; Rohmer, M.; Poralla, K. *Tetrahedron Lett.* **1999**, *40*, 6009–6012. (e) Matsuda, H.; Morikawa, T.; Ando, S.; Oominami, H.; Murakami, T.; Kimura, I.; Yoshikawa, M. *Bioorg. Med. Chem.* **2004**, *12*, 3037–3046.
- (11) For examples of controlled cyclizations by using allyl silane moieties, see: (a) Corey, E. J.; Burk, R. M. *Tetrahedron Lett.* **1987**, *28*, 6413–6416. (b) Corey, E. J.; Luo, G.; Lin, L. S. *J. Am. Chem. Soc.* **1997**, *119*, 9927–9928. (c) Mi, Y.; Schreiber, J. V.; Corey, E. J. *J. Am. Chem. Soc.* **2002**, *124*, 11290–11291.
- (12) Suzuki, E.; Ueda, M.; Ohba, S.; Sugai, T.; Shoji, M. *Tetrahedron Lett.* **2013**, *54*, 1589–1592.
- (13) For recent synthetic approaches to *cis* decalins see: (a) Petrovic, D.; Brückner, R. *Org. Lett.* **2011**, *13*, 6524–6527. (b) Jung, M. E. Guzaev, M. *Org. Lett.* **2012**, *14*, 5169–5171. (c) Jung, M. E. Guzaev, M. *J. Org. Chem.* **2013**, *78*, 7518–7526.
- (14) Maschek, J. A.; Mevers, E.; Diyabalanage, T.; Chen, L.; Ren, Y.; McClintock, J. B.; Amsler, C. D.; Wu, J.; Baker, B. J. *Tetrahedron* **2012**, *68*, 9095–9104.
- (15) (a) Marco, J. A.; Sanz-Cervera, J. F.; Morante, M. D.; García-Lliso, V.; Vallés-Xirau, J.; Jakupovic, J. *Phytochemistry* **1996**, *41*, 837–844. For previous synthesis, see: (b) Barrero, A. F.; Álvarez-Manzaneda, E. J.; Chahboun, R.; Rodríguez Rivas, A.; Linares Palomino, P. *Tetrahedron* **2000**, *56*, 6099–6113. (c) Uttaro, J. -P.; Audran, G.; Palombo, E.; Monti, H. *J. Org. Chem.* **2003**, *68*, 5407–5410.
- (16) Etse, J. T.; Gray, A. I.; Waterman, P. G. *J. Nat. Prod.* **1988**, *51*, 314–318.
- (17) Marcos, I. S.; Basabe, P.; Laderas, M.; Diez, D.; Jorge, A.; Rodilla, J. M.; Moro, R. F.; Lighgow, A. M.; Barata, I. G.; Urones, J. G. *Tetrahedron* **2003**, *59*, 2333–2343.
- (18) Singh, V.; Iyer, S. R.; Pal, S. *Tetrahedron* **2005**, *61*, 9197–9231.
- (19) For related regioselective Barbier-type additions see: (a) Rosales, A.; Oller-López, J. L.; Justicia, J.; Gansäuer, A.; Oltra, J. E.; Cuerva, J. M. *Chem. Comm.* **2004**, 2628–2629. (b) Estévez, R. E.; Justicia, J.; Bazzi, B.; Fuentes, N.; Paradas, M.; Choquesillo-Lazarte, D.; García-Ruiz, J. M.; Robles, R.; Gansäuer, A.; Cuerva, J. M.; Oltra, J. E. *Chem. Eur. J.* **2009**, *15*, 2774–2791. (c) Justicia, J.; Sancho-Sanz, I.; Álvarez-Manzaneda, E.; Oltra, J. E.; Cuerva, J. M. *Adv. Synth. Catal.* **2009**, *351*, 2295–2300. (d) Sancho-Sanz, I.; Miguel, D.; Millán, A.; Estévez, R. E.; Oller-López, J. L.; Álvarez-Manzaneda, E.; Robles, R.; Cuerva, J. M.; Justicia, J. *J. Org. Chem.* **2011**, *76*, 732–735.
- (20) See Supporting Information for experimental details.
- (21) Dess, D. B.; Martin, J. C. *J. Am. Chem. Soc.* **1991**, *113*, 7277–7287.
- (22) Becke, A. D. *Phys. Rev. A* **1988**, *38*, 3098–3100.
- (23) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785–789.
- (24) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648–5652.
- (25) Gaussian 09, Revision B.01, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, Jr., J. A.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.;
- Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, N. J.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian, Inc., Wallingford CT, **2010**.
- (26) Hariharan, P. C.; Pople, J. A. *Mol. Phys.* **1974**, *27*, 209–214.
- (27) (a) Gansäuer, A.; Kube, C.; Daasbjerg, K.; Sure, R.; Grimme, S.; Fianu, G. D.; Sadasivam, D. V.; Flowers, R. A. *J. Am. Chem. Soc.* **2014**, *136*, 1663–1671. (b) Morcillo, S. P.; Martínez-Peragón, A.; Jakoby, V.; Mota, A. J.; Kube, C.; Justicia, J.; Cuerva, J. M.; Gansäuer, A. *Chem. Commun.* **2014**, *50*, 2211–2213.
- (28) 2,4,6-Collidine presents in the reaction media is a potential ligand for titanocene(III). Nevertheless, control experiments using overstoichiometric amounts of Cp₂TiCl (2.5 equiv) in the absence of 2,4,6-collidine yielded the same stereoselections (but lower yields) that those observed using the catalytic protocol with added 2,4,6-collidine. These results rule out any potential interference of 2,4,6-collidine with active species in the catalytic cycle and, therefore, was not considered during the calculations.
- (29) Neutral complex Cp₂TiCl(THF) presented somewhat worse initial coordination capabilities than cationic Cp₂Ti⁺(THF)₂ by 3.3 kcal mol⁻¹. See Supporting Information.
- (30) Theoretical studies support the existence of such cationic species when Cp₂TiCl is generated from the mixture of Cp₂TiCl₂ and Zn dust: ref 9e. Although in our case, we usually use manganese dust as coreductant, control experiments showed that the same stereoselection is obtained using Zn dust as coreductant. This fact also supports similar active species in both reactions.
- (31) The described experiments were carried out at 10 mM, similar to that used in our experiments: Cangönül, A.; Behlendorf, M.; Gansäuer, A.; van Gastel, M. *Inorg. Chem.* **2013**, *52*, 11859–11866.
- (32) In this case we considered the THF molecule in the proximity of the complex. When the THF molecule is removed from the calculated structure **IV** the system resulted to be more favourable by 3.7 kcal mol⁻¹ in free energy terms.
- (33) Radical additions of enol radicals have been described as reversible reactions: Snider, B. B. *Chem. Rev.* **1996**, *96*, 339–364.
- (34) We also found the transition state for the hydrogen-atom transfer to the oxygen atom but the calculated energy is much higher (19.0 kcal mol⁻¹).
- (35) The reason of this low-energy transition state seems to be derived from a minimization of the unfavourable 1,3-diaxial interactions generated by the new cycle. In the case of transition state from **XVI_{cis}** to **XVII_{cis}** the intermediate adopts a distorted structure in which such interactions are slightly minimized. In the case of transition state from **XVI_{trans}** to **XVII_{trans}** the unfavourable 1,3-diaxial interaction between the axial methyl groups is unavoidable.
- (36) Bohlmann, F.; Zdero, C.; Hoffmann, E.; Mahanta, P. K.; Dorner, W. *Phytochem.* **1978**, *17*, 1917–1922.
- (37) (a) Urones, J. G.; Sánchez Marcos, I.; Basabe Barcala, P.; Martín Garrido, N. *Phytochem.* **1988**, *27*, 501–504. (b) De Pascual Teresa, J.; Urones, J. G.; Basabe, P.; Muñoz, M. A.; Marcos, I. S. *Phytochem.* **1985**, *24*, 791–793.
- (38) Banerjee, A. K.; Correa, J. A.; Laya-Mino, M. *J. Chem. Res.* **1998**, 710–711.
- (39) Koltsa, M. N.; Mironov, G. N.; Malinovskii, S. T.; Vlad, P. F. *Russ. Chem. Bull.* **1996**, *1*, 216–222.
- (40) Dekker, T. G.; Fourie, T. G.; Elmaré, M.; Snyckers, F. O.; van der Schyf, C. J. S. *Afr. J. Chem.* **1988**, *41*, 33–35.
- (41) (a) Trost, B. M. *Science* **1991**, *254*, 1471–1477. (b) Trost, B. M. *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 259–281. (c) Fürstner, A. *Synlett* **1999**, 1523–1533. (d) Fürstner, A.; Leitner, A. *Angew. Chem. Int. Ed.* **2003**, *42*, 308–311. (e) Baran, P. S.; Maimone, T. J.; Richter, J. M. *Nature*, **2007**, *446*, 404–408.

TOC

