

## Two-Directional Chain Synthesis: An Application to the Synthesis of (+)-Mycoticin A

Christopher S. Poss, Scott D. Rychnovsky,<sup>1</sup> and Stuart L. Schreiber\*

Department of Chemistry  
Harvard University  
Cambridge, Massachusetts 02138

Received January 25, 1993

Two-directional synthesis involves the simultaneous, stereoselective homologation of the termini of a nascent chain.<sup>2</sup> The potential for efficiency associated with double processing is best realized when the problem of terminus differentiation is solved effectively. This is required for synthetic targets that lack overall symmetry. The strategies for achieving differentiation of chain termini are linked to the symmetry properties of the two-directionally synthesized intermediates. Enantiotopic termini were selectively differentiated in intermediates used in the syntheses of (+)-KDO<sup>3</sup> and (-)-riboflavin<sup>3</sup> and in the stereochemical assignment of (+)-mycoticins A and B.<sup>4</sup> Diastereotopic termini were selectively differentiated in intermediates used in the syntheses of (-)-FK506<sup>5</sup> and the ansa chain of streptovaricin A,<sup>6</sup> whereas homotopic termini were differentiated in a synthesis of (-)-hikizimycin.<sup>7</sup> We now report the two-directional synthesis of a C<sub>2</sub>-symmetric fragment of the skipped polyol chain in the oxopolyene<sup>8,9</sup> macrolide antibiotics mycoticins A and B.<sup>10</sup> Differentiation and sequential homologation of the chain's homotopic termini have resulted in the first synthesis of a member of the oxopolyene macrolide family,<sup>9,11</sup> (+)-mycoticin A (1) (Figure 1).

The C<sub>17</sub>–C<sub>27</sub> fragment of mycoticin A was prepared in enantiomerically pure form using the class B two-directional chain synthesis strategy<sup>2</sup> (Scheme I). Thus, acylation of the sodium anion of  $\alpha$ -keto sulfone 3<sup>12,13</sup> with anhydride 4<sup>12</sup> followed by zinc metal reduction<sup>14</sup> gave the diketone 5 in 64% overall yield. Two-directional catalytic asymmetric reduction using the Noyori–Akutagawa catalyst<sup>15</sup> and protection followed by dissolving metal




Figure 1




Figure 2

reduction and ozonolysis<sup>16</sup> afforded the bis- $\beta$ -keto ester 6. The conversion of 6 to the tris-acetonide 7 was achieved in four steps (30% overall yield), including a second double catalytic asymmetric reduction<sup>17</sup> and a one-pot sequence involving a double reduction and a two-directional Grignard addition.<sup>18</sup> Double ozonolysis of 7 followed by base-catalyzed epimerization<sup>19</sup> and sodium borohydride reduction furnished a C<sub>2</sub>-symmetric diol. Terminus differentiation was accomplished via monoprotection of the homotopic alcohols with 1 equiv of TBSCl to give a statistical mixture of products from which alcohol 8 could be isolated directly in 49% yield and in 67% yield with one round of recycling. A Sharpless oxidation<sup>20</sup> of 8 followed by an amidation using benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate<sup>21</sup> afforded the amide 9.

Coupling of 9 (Scheme II) with the vinyl lithium reagent derived from 10<sup>12</sup> (Figure 2) followed by a Luche reduction,<sup>22</sup> ozonolysis, and protection provided two diastereomeric  $\alpha$ -mesyloxy ketones 11 (ca. 1:1). Although attempts to reductively cleave the  $\alpha$ -keto mesylates using SmI<sub>2</sub> failed,<sup>23</sup> a novel dissolving metal reduction sequence was developed that accomplished several goals. Treatment of 11 with lithium in buffered ammonia resulted in the cleavage of the mesyloxy and *p*-methoxy benzyl groups and the stereoselective (>15:1 syn:anti)<sup>24</sup> reduction of the ketone. Acetalization of the resultant diol yielded tetrakis-acetonide 12, which was identical in all respects to material obtained by degradation of 1.<sup>25</sup>

Compound 12 was converted to 13 in three steps: deprotection

(1) Present address: Department of Chemistry, University of Minnesota, Minneapolis, Minnesota 55455.

(2) For a description of the two-directional chain synthesis strategy, see: Schreiber, S. L. *Chem. Scr.* **1987**, *27*, 563.

(3) Smith, D. B.; Wang, Z.; Schreiber, S. L. *Tetrahedron* **1990**, *46*, 4793.

(4) (a) Schreiber, S. L.; Goulet, M. T. *Tetrahedron Lett.* **1987**, *28*, 6001.

(b) Schreiber, S. L.; Goulet, M. T.; Sammakia, T. *Tetrahedron Lett.* **1987**, *28*, 6005. (c) Schreiber, S. L.; Goulet, M. T.; Shulte, G. J. *Am. Chem. Soc.* **1987**, *109*, 4718. (d) Schreiber, S. L.; Goulet, M. T. *J. Am. Chem. Soc.* **1987**, *109*, 8120.

(5) Nakatsuka, M.; Ragan, J. A.; Sammakia, T.; Smith, D. B.; Uehling, D. E.; Schreiber, S. L. *J. Am. Chem. Soc.* **1990**, *112*, 5583.

(6) (a) Schreiber, S. L.; Wang, Z.; Schulte, G. *Tetrahedron Lett.* **1988**, *29*, 4085. (b) Wang, Z.; Schreiber, S. L. *Tetrahedron Lett.* **1990**, *31*, 31.

(7) (a) Ikemoto, N.; Schreiber, S. L. *J. Am. Chem. Soc.* **1990**, *112*, 9657.

(b) Ikemoto, N.; Schreiber, S. L. *J. Am. Chem. Soc.* **1992**, *114*, 2524.

(8) Oxopolyene refers to polyene macrolides in which the lactone carbonyl is conjugated to the polyene.

(9) For the first synthesis of a polyene macrolide (amphotericin B), see: Nicolaou, K. C.; Daines, R. A.; Ogawa, Y.; Chackraborty, T. K. *J. Am. Chem. Soc.* **1988**, *110*, 4696 and references cited therein.

(10) Isolation: (a) Burke, R. C.; Swartz, J. H.; Chapman, S. S.; Huang, W. J. *Invest. Dermatol.* **1954**, *23*, 163. Constitutional assignment: (b) McCaustland, D. J. Ph.D. Dissertation, Yale University, 1960. (c) Wasserman, H. H.; Van Verth, J. E.; McCaustland, D. J.; Borowitz, I. J.; Kamber, B. J. *Am. Chem. Soc.* **1967**, *89*, 1535.

(11) Roflamycin: (a) Lipshutz, B. H.; Moretti, R.; Crow, R. *Tetrahedron Lett.* **1989**, *30*, 15. Roxaticin: (b) Rychnovsky, S. D.; Rodriguez, C. J. *Org. Chem.* **1992**, *57*, 4793. (c) Mori, Y.; Asai, M.; Furukawa, H. *Heterocycles* **1992**, *34*, 1281.

(12) Preparation details for compounds 3, 4, 10, 14, and 17 will be provided in a full account of this research.

(13) White, J. D.; Avery, M. A.; Choudhry, S. C.; Dhingra, O. P.; Kang, M.-C.; Whittle, A. J. *J. Am. Chem. Soc.* **1983**, *105*, 6517.

(14) Russell, G. A.; Mikol, G. J. *J. Am. Chem. Soc.* **1966**, *88*, 5498.

(15) Kawano, H.; Ishii, Y.; Saburi, M.; Uchida, Y. *J. Chem. Soc., Chem. Commun.* **1988**, 87.

(16) For prior uses of an anisole as a latent  $\beta$ -keto ester, see: (a) Birch, A. J. *J. Chem. Soc.* **1944**, 430. (b) Evans, D. A.; Sjogren, E. B. *Tetrahedron Lett.* **1986**, *27*, 3119. (c) Evans, D. A.; Gauchet-Prunet, J. A.; Carreira, E. M.; Charette, A. B. *J. Org. Chem.* **1991**, *56*, 741.

(17) Noyori, R.; Ohkuma, T.; Kitamura, M.; Takaya, H.; Sayo, N.; Kumabayashi, H.; Akutagawa, S. *J. Am. Chem. Soc.* **1987**, *109*, 5856.

(18) (a) Burke, S. D.; Deaton, D. N.; Olsen, R. J.; Armistead, D. M.; Blough, B. E. *Tetrahedron Lett.* **1987**, *28*, 3905. (b) Schreiber, S. L.; Kelly, S. E.; Porco, J. A., Jr.; Sammakia, T.; Suh, E. M. *J. Am. Chem. Soc.* **1988**, *110*, 6210.

(19) (a) Muxfeldt, H.; Haas, G.; Hardtmann, G.; Kathawala, F.; Mooberry, J. B.; Vedejs, E. *J. Am. Chem. Soc.* **1979**, *101*, 689. (b) Stork, G.; Paterson, I.; Lee, F. K. C. *J. Am. Chem. Soc.* **1982**, *104*, 4686. (c) Schreiber, S. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1984**, *106*, 7200.

(20) (a) Carlsen, P. H. J.; Katsuki, T.; Martin, V. S.; Sharpless, K. B. *J. Org. Chem.* **1981**, *46*, 3936. (b) Denis, J.-N.; Greene, A. E.; Serra, A. A.; Luche, M.-J. *J. Org. Chem.* **1986**, *51*, 46.

(21) Castro, B.; Evin, G.; Selve, C.; Seyer, R. *Synthesis* **1977**, 413.


(22) (a) Luche, J. L. *J. Am. Chem. Soc.* **1978**, *100*, 2226. (b) Luche, J. L.; Gemal, A. L. *J. Am. Chem. Soc.* **1981**, *103*, 5454.

(23) Molander, G.; Hahn, G. *J. Org. Chem.* **1986**, *51*, 1135.


(24) Dissolving metal reduction of 5-hydroxy-2,6-dimethyl-3-heptanone produced an 11–12:1 syn:anti ratio of 1,3-diols. Unpublished results of C.S.P.

(25) Compound 12 was derived from natural mycoticins by the following sequence: isolation from *Streptomyces ruber*,<sup>10b</sup> protection of the mycoticins as their tetraacetates, ozonolysis followed by NaBH<sub>4</sub> workup and bisprotection of the diol with excess TBSCl.

## Scheme I



## Scheme II



of the two TBS groups, monoprotection, and Swern oxidation.<sup>26</sup> Aldehyde **13**<sup>27</sup> was coupled with sulfone **14**<sup>12</sup> under Julia's conditions<sup>28</sup> to furnish the differentially protected diol **15**. After a deprotection, Dess–Martin oxidation,<sup>29</sup> deprotection sequence that provided aldehyde **16**, the polyene was introduced by condensation with the lithium salt of phosphonate **17**.<sup>12,30</sup> Hydrolysis of the ethyl ester and macrolactonization using Yamaguchi's protocol<sup>31</sup> gave the tetrakis-acetonide of (+)-mycotycin A, (**18**),<sup>32,33</sup> which was identical to material derived from natural mycotycin A. Final deprotection of **18** produced (+)-mycotycin A (**1**) in 75% yield.

The synthesis of (+)-mycotycin A confirms the earlier stereochemical assignment that was based upon synthetic and

spectroscopic analyses of degradation products from the oxypolyene.<sup>4</sup> These and other synthetic studies<sup>3–7</sup> and the work described herein relied heavily upon the efficient material processing that can be achieved using two-directional chain synthesis strategies.<sup>2</sup>

**Acknowledgment.** Support for this work was provided by the NIGMS (GM-38627). We also gratefully acknowledge the support of the NIH for a postdoctoral fellowship to S.D.R. Mass spectra were obtained by Dr. A. Tyler, Ms. L. Romo, and Mr. R. Valentekovich at the Harvard University Mass Spectrometry Facility. NMR data were obtained on a Bruker AM-500 for which we acknowledge the NIH (1-S10-RR04870-01) and NSF (CHE88-14019).

**Supplementary Material Available:** Complete spectral data for compounds **1**, **3**, **5**–**10**, **12**–**18**, and (+)-mycotycin A;  $^1\text{H}$  NMR and  $^{13}\text{C}$  NMR spectra of natural and synthetic compounds **12** and **18** are included (8 pages). Ordering information is given on any current masthead page.

(33) Compound **18** was isolated as a mixture of olefin isomers. The major isomer ( $[\alpha]_D^{25} = +182.3^\circ$  ( $c = 0.32$ ,  $\text{CHCl}_3$ )) was identical to **18** derived from natural material ( $[\alpha]_D^{25} = +261.9^\circ$  ( $c = 1.09$ ,  $\text{CHCl}_3$ )) in all respects except optical rotation. This difference is attributed to a small amount of olefin isomers present in the synthetic material. Mycotycin A itself has been reported to yield a range of optical rotations from  $+63.4^\circ$  to  $-41.3^\circ$ , depending upon the duration of light exposure.<sup>10b</sup>

(26) Mancuso, A. J.; Swern, D. *Synthesis* **1981**, 165.

(27) The remainder of the synthesis was completed with material derived from degradation.

(28) (a) Julia, M.; Paris, J. M. *Tetrahedron Lett.* **1973**, 4833. (b) Kocienski, P. J.; Lythgoe, B. J. *Chem. Soc., Perkin Trans.* **1980**, 1400.

(29) Dess, D. B.; Martin, J. C. *J. Org. Chem.* **1983**, *48*, 4155.

(30) Boschelli, D.; Takemasa, T.; Yasuhiro, N.; Masamune, S. *Tetrahedron Lett.* **1985**, *26*, 5239.

(31) Inanaga, J.; Kuniko, H.; Hiroko, S.; Katsuki, T.; Yamaguchi, M. *Bull. Chem. Soc. Jpn.* **1979**, *52*, 1989.

(32) Natural and synthetic (+)-mycotycin A, tetraacetonide, and (+)-mycotycin A produce five olefinic isomers upon light exposure as studied by HPLC (Rainin Microsorb column,  $10.0 \times 250.0 \text{ mm}^2$ , 15%  $\text{EtOAc}$ /hexane, 4 mL/min) and  $^1\text{H}$  NMR. The major (all-trans) isomer reconverts into the five-component mixture after a few hours of light exposure; in addition, the four minor isomers individually convert to the five-component mixture under the same conditions.