

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/225288904>

# Combinatorial Solid-Phase Synthesis of 4,6-Diaryl and 4-Aryl, 6-Alkyl-1,3,5-triazines and Their Application to Efficient Biofuel Production

ARTICLE · JUNE 2012

DOI: 10.1021/co300007a · Source: PubMed

CITATIONS

6

READS

21

8 AUTHORS, INCLUDING:



**Hyung-Ho Ha**

Sunchon National University

33 PUBLICATIONS 367 CITATIONS

SEE PROFILE



**Marc Vendrell**

The University of Edinburgh

38 PUBLICATIONS 1,143 CITATIONS

SEE PROFILE



**Jacqueline T Bork**

University of Maryland Medical Center

10 PUBLICATIONS 126 CITATIONS

SEE PROFILE



**Youngsook Lee**

Pohang University of Science and Technology

95 PUBLICATIONS 4,057 CITATIONS


SEE PROFILE

## Combinatorial Solid-Phase Synthesis of 4,6-Diaryl and 4-Aryl, 6-Alkyl-1,3,5-triazines and Their Application to Efficient Biofuel Production

Jaon Y. H. Kim,<sup>#,†</sup> Jae Wook Lee,<sup>#,‡,⊗</sup> Woo Sirl Lee,<sup>§</sup> Hyung-Ho Ha,<sup>§,||</sup> Marc Vendrell,<sup>⊥</sup> Jacqueline T. Bork,<sup>‡</sup> Youngsook Lee,<sup>\*,†</sup> and Young-Tae Chang<sup>\*,§,⊥</sup><sup>†</sup>Department of Life Science, Pohang University of Science and Technology, Pohang 790-784, Korea<sup>‡</sup>Department of Chemistry, New York University, New York 10003, United States<sup>§</sup>Department of Chemistry and MedChem Program of Life Sciences Institute, National University of Singapore, 117543 Singapore<sup>||</sup>College of Pharmacy, Suncheon National University, Suncheon 570-742, Korea<sup>⊥</sup>Laboratory of Bioimaging Probe Development, Singapore Bioimaging Consortium, Agency for Science, Technology and Research (A\*STAR), 11 Biopolis Way, #02-02 Helios Building, 138667 Singapore

## S Supporting Information

**ABSTRACT:** Herein we report the solid-phase synthesis of a combinatorial aryl, alkyl-triazine library and its application to biofuel production. The combination of Grignard reactions and solid supported Suzuki coupling reactions afforded unique 120 triazine compounds with high purities and minimum purification steps. Through an unbiased phenotypic screening for improved biofuel generation in oleaginous yeast, we found one diaryl triazine derivative (E4) which increased the biolipid production up to 86%.



**KEYWORDS:** triazines, Grignard reaction, Suzuki coupling, biolipids, high-throughput screening, chemical genetics

The ease of manipulation and wide range of biological properties of the triazine scaffold have attracted the interest of combinatorial and medicinal chemists to explore its derivatization.<sup>1–5</sup> As a result, a number of synthetic approaches reported by our group and others have converted the triazine structure into a key scaffold in drug discovery programs.<sup>6–11</sup> 1,3,5-Triazine derivatives can be readily prepared from cyanuric chloride using amine, thiol, and alcohol building blocks that involve the formation of C–heteroatom bonds.<sup>12–16</sup> These syntheses have been adapted to solid-phase methodologies to facilitate the construction of large collections.<sup>17–21</sup> In contrast, the derivatization of the triazine scaffold forming C–C bonds has been much lesser explored. For instance, although aryl–aryl motifs are well-known pharmacophores and widely employed in the derivatization of privileged scaffolds,<sup>22,23</sup> the synthesis of diaryl or aryl and alkyl-substituted triazines are limited to a few examples.<sup>24</sup> Our group recently reported efficient synthetic schemes for triazine libraries with single C–C bonds using either Grignard reactions<sup>25</sup> or Suzuki couplings.<sup>26</sup> We envisioned that a triazine library with double C–C bond derivatization may open a new chemical space for the discovery of novel bioactivities.

Lipids are one of the most important topics of biological research because numerous diseases, including diabetes, obesity, and cancer, are caused by their abnormal uptake, synthesis, metabolism, storage, or mobilization.<sup>27,28</sup> In addition, the global energy crisis has made lipid research even more important, since biolipids produced by microorganisms and plants hold a promising potential to provide sustainable fuel for

the human society. Chemical genetics approaches have been used to find chemicals that alter the neutral lipid accumulation in mammalian cells and simple animal models such as *Caenorhabditis elegans* and zebrafish. These experiments led to identification of new factors that regulate lipid metabolism<sup>28–31</sup> and opened new ways to treat lipid-related diseases. We envisioned that similar approaches might be useful to discover new chemicals that can increase biofuel production in microorganisms and plants, since neutral lipids can be readily converted to biofuel. We screened our new library of 4,6-diaryl and 4-aryl,6-alkyl triazines in a forward chemical genetics approach to identify compounds that increased neutral lipid accumulation in *Yarrowia lipolytica*, an oleaginous yeast well-known for its versatile lipid-related metabolic pathways.<sup>32</sup>


Herein we report the solid-phase synthesis of 4,6-diaryl and 4-aryl, 6-alkyl 1,3,5-triazines by combining Grignard reactions and Suzuki couplings and their evaluation for efficient biofuel generation, which yielded to the discovery of E4 as a unique chemical regulator of the biolipid production. We designed the orthogonal synthesis of a two-dimensional library by combining a solution phase Grignard reaction followed by a solid-phase Suzuki coupling (Scheme 1). Cyanuric chloride was initially derivatized by 12 different Grignard reagents under reported conditions<sup>25</sup> including alkyl and aryl groups on the 6-position of the triazine scaffold (e.g., linear and  $\beta$ -branched alkyl groups,

Received: January 20, 2012


Revised: May 14, 2012

Published: June 11, 2012



Scheme 1. Synthesis of 4,6-Diaryl and 4-Aryl, 6-Alkyl-1,3,5-triazine Compounds<sup>a</sup>

<sup>a</sup>Reagents and conditions: (a)  $R_1$ -MgX, THF, 0 °C to r.t., 8 h; (b) PAL-PS amine-derivatized resin, THF, DIEA, 60 °C, 3 h; (c)  $Pd(PPh_3)_4$ , dioxane,  $R_2$ -B(OH)<sub>2</sub>, Cs<sub>2</sub>CO<sub>3</sub>, 90 °C, 15 h; (d) TFA-DCM (1:9), r.t., 0.5 h.




**Figure 1.** Building blocks for library synthesis: (a) Grignard reagents, (b) boronic acids.

differently substituted aromatic rings, Figure 1a). The resulting 2,4-dichloro-6-alkyl-1,3,5-triazines were loaded onto an amine-derivatized PAL-polystyrene (PAL-PS) resin to accelerate their further combinatorial derivatization.<sup>33</sup> PAL-PS resins are compatible with palladium-catalyzed cross-coupling conditions,<sup>34–36</sup> and have been used for the preparation of tagged-triazine libraries.<sup>6–8</sup>

Polymer-supported triazines were then derivatized using optimized Suzuki coupling conditions<sup>26</sup> with 10 different boronic acids that included both electron-withdrawing and electron-donating groups (Figure 1b). A final acidic cleavage with a trifluoroacetic acid-dichloromethane solution (1:9) rendered a collection of 4,6-diaryl and 4-aryl, 6-alkyl 1,3,5-triazines (120 compounds) in average purities around 90% without running any purification step (Supporting Information, Table S1).

Next we examined the library in a chemical genetics approach for efficient biofuel generation. To identify triazine compounds that promoted lipid accumulation in *Y. lipolytica*, we cultured the cells to midlog phase, treated them with the individual compounds of the library, and analyzed their neutral lipid content by staining the cells with a lipid sensing fluorescent BODIPY dye (D3922, Invitrogen).<sup>37</sup> In this



**Figure 2.** (a) Chemical structures of E4 and the analogues E4OH and E4Bz; (b) Fluorescent microscope images of BODIPY-stained *Y. lipolytica* cells treated with DMSO (left) and 10  $\mu$ M of E4 (right). Cells were grown for 12 h at 28 °C in an air-shaking incubator (180 rpm) after being treated with DMSO or E4; (c) BODIPY fluorescence average intensity of cells treated with E4, E4OH, E4Bz and DMSO (10  $\mu$ M) ( $n = 3$ , bars as SD). The same number of cells ( $OD_{600} = 0.5$ ) was used for all the experiments.

screening we found that only one triazine compound (E4, Figure 2a) induced a significant increase of the lipid production, with a fluorescence intensity enhancement up to 86% at 10  $\mu$ M (Figure 2b). Furthermore, we observed that that none of the single C–C bond triazine derivatives from previously synthesized libraries (588 Grignard<sup>25</sup> and 160 Suzuki derivatives<sup>26</sup>) showed any significant activity, asserting the unique biochemical properties of the newly synthesized triazines. To examine the role that the ethyleneglycol group played in the activity of E4, we synthesized and tested two analogues with shorter and aromatic versions of the linker (i.e., respectively E4OH and E4Bz, Figure 2a). All three compounds showed a similar dose-dependent increase of the lipid accumulation (Supporting Information, Figure S4), which indicated the substituted diaryl triazine core as the main one responsible for the biological activity. Moreover, none of these compounds did show substantial toxicity or inhibition of the cell growth, even at high concentration. Finally, we determined the amount of triacylglycerol (TAG) extracted from the same number of cells treated with E4, E4OH, and E4Bz and compared it to DMSO-treated control cells. This experiment confirmed the capacity of E4 and its analogues to increase the content of TAG (Supporting Information, Figure S4a). The

amount of biochemically extracted TAG was quantified by densitometry on iodine-stained TLC plates, and corroborated the results obtained by fluorescence staining. These results proved the application of 4,6-diaryl 1,3,5-triazines as candidate molecules to increase lipid accumulation in *Y. lipolytica*.

In conclusion, we prepared in solid-phase a collection of 4,6-diaryl and 4-aryl, 6-alkyl 1,3,5-triazines using Grignard reactions and Suzuki couplings and performed an unbiased phenotypic screening to discover E4 as a unique small molecule with efficient biolipid generation activity. Further biochemical studies toward target identification and mechanism elucidation will be reported in due course.

## ■ ASSOCIATED CONTENT

### Supporting Information

Synthetic procedures for the library synthesis, characterization data for the hit compounds and procedures for the bioassays. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: ylee@postech.ac.kr (Y.L.), chmcyt@nus.edu.sg (Y.-T.C.). fax: +82 542792199 (Y.L.), +65 67791691 (Y.-T.C.). Phone: +82 542792296 (Y.L.), +65 65166774 (Y.-T.C.).

### Present Address

⊗The Scripps Research Institute, SR207 10550 North Torrey Pines Road, SR202 La Jolla, CA 92037.

### Author Contributions

#These authors contributed equally to this work.

### Author Contributions

Y.S.L. and Y.T.C. conceived and designed the research and wrote the manuscript. J.Y.H.K. performed the biological experiments and wrote the manuscript. J.W.L., W. S.L., H.-H.H., and M.V. performed organic synthesis and wrote the manuscript. J.T.B. carried out organic synthesis. All authors have given approval to the final version of the manuscript.

### Funding

The authors are grateful to A\*STAR SSCC Grant (SSCC10/024) and the Global Frontier Program (2010-0029723) of the Ministry of Education, Science and Technology of Korea for the financial support.

### Notes

The authors declare no competing financial interest.

## ■ REFERENCES

- (1) Lee, H. K.; Chui, W. K. Combinatorial Mixture Synthesis and Biological Evaluation of Dihydrophenyl Triazine Antifolates. *Bioorg. Med. Chem.* **1999**, *7*, 1255–1262.
- (2) Zhou, C.; Min, J.; Liu, Z.; Young, A.; Deshazer, H.; Gao, T.; Chang, Y. T.; Kallenbach, N. R. Synthesis and biological evaluation of novel 1,3,5-triazine derivatives as antimicrobial agents. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 1308–1311.
- (3) Vu, C. B.; Pan, D.; Peng, B.; Kumaravel, G.; Smits, G.; Jin, X.; Phadke, D.; Engber, T.; Huang, C.; Reilly, J.; Tam, S.; Grant, D.; Hetu, G.; Petter, R. C. Novel Diamino Derivatives of [1,2,4]-triazolo[1,5-a][1,3,5]triazine as Potent and Selective Adenosine A<sub>2a</sub> Receptor Antagonists. *J. Med. Chem.* **2005**, *48*, 2009–2018.
- (4) Lee, H. K.; Rana, T. M. Microwave-Assisted Parallel Synthesis of a 4,6-Diamino-2,2-dimethyl-1,2-dihydro-1-phenyl-s-triazine Library. *J. Comb. Chem.* **2004**, *6*, 504–508.
- (5) Henke, B. R.; Consler, T. G.; Go, N.; Hale, R. L.; Hohman, D. R.; Jones, S. A.; Lu, A. T.; Moore, L. B.; Moore, J. T.; Orband-Miller, L. A.; Robinett, R. G.; Shearin, J.; Spearing, P. K.; Stewart, E. L.; Turnbull, P. S.; Weaver, S. L.; Williams, S. P.; Wisely, G. B.; Lambert, M. H. A New Series of Estrogen Receptor Modulators That Display Selectivity for Estrogen Receptor  $\beta$ . *J. Med. Chem.* **2002**, *45*, 5492–5505.
- (6) Moon, H. S.; Jacobson, E. M.; Khersonsky, S. M.; Luzung, M. R.; Walsh, D. P.; Xiong, W.; Lee, J. W.; Parikh, P. B.; Lam, J. C.; Kang, T. W.; Rosania, G. R.; Schier, A. F.; Chang, Y. T. A Novel Microtubule Destabilizing Entity from Orthogonal Synthesis of Triazine Library and Zebrafish Embryo Screening. *J. Am. Chem. Soc.* **2002**, *124*, 11608–11609.
- (7) Williams, D.; Jung, D. W.; Khersonsky, S. M.; Heidary, N.; Chang, Y. T.; Orlow, S. J. Identification of Compounds that Bind Mitochondrial F1F0 ATPase by Screening a Triazine Library for Correction of Albinism. *Chem. Biol.* **2004**, *11*, 1251–1259.
- (8) Khersonsky, S. M.; Jung, D. W.; Kang, T. W.; Walsh, D. P.; Moon, H. S.; Jo, H.; Jacobson, E. M.; Shetty, V.; Neubert, T. A.; Chang, Y. T. Facilitated Forward Chemical Genetics Using Tagged Triazine Library and Zebrafish Embryo Screening. *J. Am. Chem. Soc.* **2003**, *125*, 11804–11805.
- (9) Guo, Z.; Wu, D.; Zhu, Y. F.; Tucci, F. C.; Pontillo, J.; Saunders, J.; Xie, Q.; Struthers, R. S.; Chen, C. A. Convenient One-pot Synthesis of Asymmetric 1,3,5-Triazine-2,4,6-triones and its Application Towards a Novel Class of Gonadotropin-releasing Hormone Receptor Antagonists. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 693–698.
- (10) Balboni, G.; Lazzari, L.; Trapella, C.; Negri, L.; Lattanzi, R.; Giannini, E.; Nicotra, A.; Melchiorri, P.; Visentin, S.; De Nuccio, C.; Salvadori, S. Triazine Compounds as Antagonists at Bv8-Prokineticin Receptors. *J. Med. Chem.* **2008**, *51*, 7635–7639.
- (11) Krecmerová, M.; Holý, A.; Piskala, A.; Masojdková, M.; Andrei, G.; Naesens, L.; Neyts, J.; Balzarini, J.; Clercq, E.; De; Snoeck, R. Antiviral Activity of Triazine Analogues of 1-(S)-[3-Hydroxy-2-(phosphonomethoxy)propyl]cytosine (Cidofovir) and Related Compounds. *J. Med. Chem.* **2007**, *50*, 1069–1077.
- (12) Kido, H.; Goodrow, M. H.; Griffith, V.; Lucas, A. D.; Gee, S. J.; Hammock, B. D. Development of an Enzyme-linked Immunosorbent Assay for the Detection of Hydroxytriazines. *J. Agric. Food Chem.* **1997**, *45*, 414–424.
- (13) Lippa, K.; Roberts, A. Nucleophilic Aromatic Substitution Reactions of Chloroazines with Bisulfide (HS<sup>-</sup>) and Polysulfides (Sn<sup>2-</sup>). *Environ. Sci. Technol.* **2002**, *36*, 2008–2018.
- (14) Chen, Y.; Wang, D. X.; Huang, Z. T.; Wang, M. X. Synthesis, Structure, and Functionalization of Homo Heterocalix[2]arene[2]-triazines: Versatile Conformation and Cavity Structures Regulated by the Bridging Elements. *J. Org. Chem.* **2010**, *75*, 3786–3796.
- (15) Umali, A. P.; Crampton, H. L.; Simanek, E. E. Triazine Dendrimers with Orthogonally Protected Amines on the Periphery. Masking Amines with Dde and BOC Groups Provides an Alternative to Carrying Protected Alcohols and Disulfides through an Iterative Synthesis. *J. Org. Chem.* **2007**, *72*, 9866–9874.
- (16) Lim, J.; Mintzer, M. A.; Perez, L. M.; Simanek, E. E. Synthesis of Odd Generation Triazine Dendrimers Using a Divergent, Macromonomer Approach. *Org. Lett.* **2010**, *12*, 1148–1151.
- (17) Gopalsamy, A.; Yang, H. Combinatorial Synthesis of Heterocycles: Solid-Phase Synthesis of 6-Amino-2,4-dioxo-3,4-dihydro-1,3,5-triazine Derivatives. *J. Comb. Chem.* **2001**, *3*, 278–283.
- (18) Bork, J. T.; Lee, J. W.; Khersonsky, S. M.; Moon, H. S.; Chang, Y. T. Novel Orthogonal Strategy toward Solid Phase Synthesis of 1,3,5-Substituted Triazines. *Org. Lett.* **2003**, *5*, 117–120.
- (19) Khersonsky, S. M.; Chang, Y. T. Safety-Catch Approach to Orthogonal Synthesis of Triazine Library. *J. Comb. Chem.* **2004**, *6*, 474–477.
- (20) Hu, Z.; Ma, T.; Chen, Z.; Ye, Z.; Zhang, G.; Lou, Y.; Yu, Y. Solid-Phase Synthesis and Antitumor Evaluation of 2,4-Diamino-6-aryl-1,3,5-triazines. *J. Comb. Chem.* **2009**, *11*, 267–273.
- (21) Kong, K. H.; Tan, C. K.; Lam, Y. Traceless Solid-Phase Synthesis of 6-Amino- and 6-Hydroxyimino-1,3,5-triazine-2,4-diones and 1,3,5-Triazine-2,4,6-triones. *J. Comb. Chem.* **2009**, *11*, 1050–1060.



- (22) Tounge, B. A.; Reynolds, C. H. Defining Privileged Reagents Using Subsimilarity Comparison. *J. Chem. Inf. Comput. Sci.* **2004**, *44*, 1810–1815.
- (23) Tejedor, D.; Garcia-Tellado, F. Chemo-differentiating ABB' Multicomponent Reactions. Privileged Building Blocks. *Chem. Soc. Rev.* **2007**, *36*, 484–491.
- (24) Konno, S.; Osawa, N.; Yamanaka, H. Synthesis of 6-Substituted 3,5-Diaryl-1,2,4-triazines as Potential Herbicidal Agents. *J. Agric. Food Chem.* **1995**, *43*, 838–842.
- (25) Lee, J. W.; Bork, J. T.; Ha, H. H.; Samanta, A.; Chang, Y. T. Novel Orthogonal Synthesis of Tagged Combinatorial Triazine Library via Grignard Reaction. *Aus. J. Chem.* **2009**, *62*, 1000–1006.
- (26) Lee, J. W.; Ha, H. H.; Vendrell, M.; Bork, J. T.; Chang, Y. T. Combinatorial Solid-Phase Synthesis of 6-aryl-1,3,5-triazines via Suzuki Coupling. *Aus. J. Chem.* **2011**, *64*, 540–544.
- (27) Kawasumi, M.; Nghiem, P. Chemical Genetics: Elucidating Biological Systems with Small Molecules. *J. Invest. Dermatol.* **2007**, *127*, 1577–1584.
- (28) Lemieux, G. A.; Liu, J.; Mayer, N.; Bainton, R. J.; Ashrafi, K.; Werb, Z. A Whole-organism Screen Identifies New Regulators of Fat Storage. *Nat. Chem. Biol.* **2011**, *7*, 206–213.
- (29) Ishitsuka, R.; Saito, T.; Osada, H.; Ohno-Iwashita, Y.; Kobayashi, T. Fluorescence Image Screening for Chemical Compounds Modifying Cholesterol Metabolism and Distribution. *J. Lipid Res.* **2011**, *52*, 2084–2094.
- (30) Murphy, S.; Martin, S.; Parton, R. G. Quantitative Analysis of Lipid Droplet Fusion: Inefficient Steady State Fusion but Rapid Stimulation by Chemical Fusogens. *PLoS One* **2010**, *5*, e15030.
- (31) Clifton, J. D.; Lucumi, E.; Myers, M. C.; Napper, A.; Hama, K.; Farber, S. A.; Smith, A. B.; Huryn, D. M.; Diamond, S. L.; Pack, M. Identification of Novel Inhibitors of Dietary Lipid Absorption Using Zebrafish. *PLoS One* **2010**, *5*, e12386.
- (32) Beopoulos, A.; Cescut, J.; Haddouche, R.; Uribelarrea, J. L.; Molina-Jouve, C.; Nicaud, J. M. *Yarrowia lipolytica* as a Model for Bio-oil Production. *Prog. Lipid Res.* **2009**, *48*, 375–387.
- (33) Albericio, F.; Barany, G. Acidolytic Cleavage of Tris(alkoxy)-benzylamide (PAL) “internal reference” Amino Acyl (IRAA) Anchoring Linkages: Validation of Accepted Procedures in Solid-Phase Peptide Synthesis (SPPS). *Int. J. Pept. Protein Res.* **1993**, *41*, 307–312.
- (34) Gauthier, D. R., Jr.; Limanto, J.; Devine, P. N.; Desmond, R. A.; Szumigala, R. H., Jr.; Foster, B. S.; Volante, R. P. Palladium-Catalyzed Regioselective Arylation of Imidazo[1,2-b][1,2,4]triazine: Synthesis of an  $\alpha_{2/3}$ -Selective GABA Agonist. *J. Org. Chem.* **2005**, *70*, 5938–5945.
- (35) Kuo, G. H.; DeAngelis, A.; Emanuel, S.; Wang, A.; Zhang, Y.; Connolly, P. J.; Chen, X.; Gruninger, R. H.; Rugg, C.; Fuentes-Pesquera, A.; Middleton, S. A.; Jolliffe, L.; Murray, W. V. Synthesis and Identification of [1,3,5]Triazine-pyridine Biheteroaryl as a Novel Series of Potent Cyclin-Dependent Kinase Inhibitors. *J. Med. Chem.* **2005**, *48*, 4535–4546.
- (36) Garnier, E.; Audoux, J.; Pasquinet, E.; Suzenet, F.; Poullain, D.; Lebre, B.; Guillaumet, G. Easy Access to 3- or 5-Heteroaryl-amino-1,2,4-triazines by  $S_NAr$ ,  $S_NH$ , and Palladium-Catalyzed N-Hetero-arylations. *J. Org. Chem.* **2004**, *69*, 7809–7815.
- (37) Bozaquel-Morais, B. L.; Madeira, J. B.; Maya-Monteiro, C. M.; Masuda, C. A.; Montero-Lomeli, M. A New Fluorescence-based Method Identifies Protein Phosphatases Regulating Lipid Droplet Metabolism. *PLoS One* **2010**, *5*, e13692.