

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/10660985>

De Novo Asymmetric Bio- and Chemocatalytic Synthesis of Saccharides – Stereoselective Formal O –Glycoside Bond Formation Using Palladium Catalysis

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · AUGUST 2003

Impact Factor: 12.11 · DOI: 10.1021/ja0347538 · Source: PubMed

CITATIONS

48

READS

18

4 AUTHORS, INCLUDING:



Rienk Eelkema

Delft University of Technology

51 PUBLICATIONS 1,404 CITATIONS

SEE PROFILE



Adriaan J Minnaard

University of Groningen

224 PUBLICATIONS 6,951 CITATIONS

SEE PROFILE

De Novo Asymmetric Bio- and Chemocatalytic Synthesis of Saccharides – Stereoselective Formal *O*-Glycoside Bond Formation Using Palladium Catalysis

Alex C. Comely, Rienk Eelkema, Adriaan J. Minnaard, and Ben L. Feringa*


Department of Organic and Molecular Inorganic Chemistry, Stratingh Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

Received February 19, 2003; E-mail: b.l.feringa@chem.rug.nl

The chemical synthesis of carbohydrate domains in saccharides and glycoconjugates such as antibiotics, antitumor agents, glycoproteins, and glycolipids is now recognized as a major frontier for organic chemistry.¹ Fundamental to the synthesis of such carbohydrates and their derivatives is the selectivity of α - or β -*O*-glycoside bond formation which typically entails the coupling of one nucleophilic (O-donating) glycoside to another electrophilic glycosyl donor;^{2,3} anomeric stereoselectivity is a complex issue usually dependent on the nature of the donor C2 substituent.¹

Catalytic stereoselective formation of the acetal linkage onto pyranones⁴ of type **1** (Scheme 1) presents a conceptually different solution to this stereochemical problem by providing a stereodefined platform whose chiral information can be relayed around the ring. Such an acetal can be a formal α - or β -glycoside bond depending on the enantiomer of **1**, the stereocontrol in the Pd-catalyzed step, and the chemistry used to elaborate the ring.⁵

Scheme 1. Iterative Saccharide Synthesis: Stereoselective Acetal Bond Formation Using Pd Catalysis





Here, we present a novel integrated approach to the de novo catalytic asymmetric synthesis of saccharides uniting two protocols: the enzymatic resolution of racemic acetoxypyranones **1**⁶ with a highly stereoselective palladium-catalyzed acetal bond formation onto this embryonic sugar (Scheme 1). Resulting from subsequent steps to elaborate the ring into a diversity of natural and unnatural sugars, a free hydroxyl group can be stereoselectively coupled again to **1**, giving rise to an iterative catalytic asymmetric saccharide synthesis. A blank slate for saccharide synthesis, the versatility of this cyclic enone platform has been appreciated for some time.⁷

Despite the widespread use of phenols as nucleophiles in the palladium-catalyzed allylic substitution reaction,⁸ aliphatic alcohols have received scant attention.^{9,10} During early investigations, however, we found that the substitution reaction of enantiomerically pure 6-acetoxy-2*H*-pyran-3(6*H*)-one (–)-**1**⁶ with simple primary and secondary aliphatic alcohols as solvent proceeded with nearly complete retention of stereochemistry.¹¹

Efforts to improve the viability of this methodology resulted in the coupling depicted in Table 1. The use of 10 mol % Pd(OAc)₂ and triphenyl phosphite in DCM at –30 °C¹² was found to convert pyranone (–)-**1** into the benzyl alcohol adduct **2A** in high yield

Table 1. Stereoselective Acetal Bond Formation Using Pd Catalysis



adduct	donor	% yield	% ee/de	adduct	donor	% yield	% de ^e
2A	(–)- 1	83	94 ^b	2H	(–)- 1	65 ^a	91
2B	(–)- 1	87	98 ^{b,c}	2I	(–)- 1	70 ^a	97
2C	(–)- 1	98	99 ^b	2J	(–)- 1	60	<i>f</i>
2D	(–)- 1	84	98 ^b				
2E	(±)- 1	69	nd ^{b,d}				
2F	(–)- 1	78 ^a	97 ^b				
2G	(–)- 1	77 ^a	94 ^e				
	(–)- 3	88 ^a	94 ^e				
	(+)- 3	96 ^a	98 ^e				

^a Isolated yield of unique stereoisomer. ^b 10% Pd(OAc)₂, P(OPh)₃, DCM, –30 °C; stereoselectivities were determined by chiral HPLC analysis. ^c Enantiomeric excess before chromatography. ^d Coupled to racemic **1** only. ^e 5% Pd₂(dba)₃, PPh₃, DCM, –10 °C; diastereoselectivities were determined from ¹H NMR. ^f Mixture of isomers.


and 94% ee. Particularly rewarding were the still higher yields and ee's for anisyl nucleophiles **B** and **C** and the *ortho*-nitrobenzyl alcohol **D**, useful mimics of benzyl linkers^{1b,13} applied to the solid-phase synthesis of saccharides.¹⁴ In preliminary experiments to apply the protocol to the solid-phase, photocleavable **E**, immobilized onto phenolic polystyrene, was also coupled efficiently to racemic **1**. Representative of Mucin-type glycosylation found in the glycopeptides of mammals and other eukaryotes,¹⁵ adduct **2F** was also prepared with excellent stereoselectivity.

Key to the feasibility of the protocol is the success of a first iteration: a stereoselective coupling reaction of enantiopure glycosyl donor with a sugar derivative. The results are illustrated in Table 1. Initial attempts using the Pd(OAc)₂/P(OPh)₃ catalyst system failed, but, to our relief, use of Pd₂(dba)₃/PPh₃ successfully mediated formation of the desired adducts **2G–2J**. Primary alcohol **G**, a 6-deprotected glucopyranose, underwent coupling with (–)-**1** and both (R)-(–)-**3** and (S)-(+)-**3**¹⁶ to afford the stereoisomers of the

products with excellent yield (77–96%) and diastereoselectivity (94–98%). Crucially, similar success was found with the more sterically demanding substrates 4-deprotected glucopyranose **H** and 3-deprotected glucofuranose **I** bearing a secondary alcohol moiety, and good yields (57–76%) and excellent stereoselectivities (82–97%) were obtained during both *R*- and *S*-acetal bond formation. All adducts were isolated as unique diastereomers by simple column chromatography with the exception of that with **J**, deprotected at the anomeric center.

A preliminary application of our iterative approach is depicted in Scheme 2. Diastereoselective catalytic *cis*-dihydroxylation of enone adduct **2C** was effected by $\text{RuCl}_3/\text{NaIO}_4$,¹⁷ and the resulting diol was protected to the dioxolane **4C** under standard conditions. Subsequent reduction using $\text{Zn}(\text{BH}_4)_2$ ¹⁸ gave **5C**, a β -L-ribose.¹⁹ Coupling of this sugar under the catalytic conditions previously described successfully afforded the disaccharide precursor **6C** with 96% de.²⁰


Scheme 2. Preliminary Application of Iterative Saccharide Synthesis^a



^a (i) $\text{RuCl}_3 \cdot 3\text{H}_2\text{O}$ (20 mol %), NaIO_4 ; (ii) 2,2-DMP, acetone, PTSA; (iii) $\text{Zn}(\text{BH}_4)_2$; (iv) $(-)\text{-1}$, $\text{Pd}_2(\text{dba})_3$ (5 mol %), PPh_3 , CH_2Cl_2 .

Unsuccessful endeavors to alkylate the methylene position of **4C** led to an appraisal of prefunctionalized pyranone substrate **7** in the palladium-catalyzed allylic substitution reaction (Scheme 3). Prepared enantiopure employing a Sharpless dihydroxylation protocol,⁷ⁱ **7** indeed underwent substitution with complete retention of stereochemistry, giving **8**. 4,4-Dimethyl-substituted pyranone **9**,^{7i,16} applicable to the asymmetric synthesis of L-noviose,²¹ a constituent of the antibiotic novobiocin, also participated with high stereoselectivity to afford **10**.

Scheme 3. C4-Substituted Glycosyl Donors




Efforts to elaborate on this chemistry by providing a view of an iterative catalytic solid-phase protocol are ongoing.

Acknowledgment. We thank Dr. Richard van Delden for the preparative HPLC separation. Financial support from NRSC-C Catalysis and the Dutch Ministry of Economic Affairs under the EET scheme (grant nos. EETK97107 and EETK99104) is gratefully acknowledged.

Supporting Information Available: Experimental procedures and spectral data for all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (a) Collins, P.; Ferrier, R. *Monosaccharides. Their Chemistry and Their Roles in Natural Products*; Wiley: U.K., 1995. (b) Nicolaou, K. C.; Mitchell, H. J. *Angew. Chem., Int. Ed.* **2001**, *40*, 1576.
- (a) Toshima, K.; Tatsuta, K. *Chem. Rev.* **1993**, *93*, 1503. (b) Wulff, G.; Röhle, G. *Angew. Chem., Int. Ed. Engl.* **1974**, *13*, 157.
- Danishefsky, S. J.; Bilodeau, M. T. *Angew. Chem., Int. Ed. Engl.* **1996**, *35*, 1380.
- (a) Gryniewicz, G.; Barszczak, B.; Zamojski, A. *Synthesis* **1979**, 364. (b) Mucha, B.; Hoffmann, H. M. R. *Tetrahedron Lett.* **1989**, *30*, 4489. (c) Gryniewicz, G. *Carbohydr. Res.* **1980**, *80*, 53. (d) Gryniewicz, G.; Zamojski, A. Z. *Naturforsch.* **1980**, *35b*, 1024. (e) Knol, J.; Jansen, J. F. G. A.; Van Bolhuis, F.; Feringa, B. L. *Tetrahedron Lett.* **1991**, *32*, 7465.
- The authors recognize that to adopt α - and β -nomenclature for **2** is not strictly appropriate until a second chiral center is introduced to the ring.
- Van den Heuvel, M.; Cuiper, A. D.; Van der Deen, H.; Kellogg, R. M.; Feringa, B. L. *Tetrahedron Lett.* **1997**, *38*, 1655.
- For a review, see: Holder, N. L. *Chem. Rev.* **1982**, *82*, 287. For selected examples, see: (a) Plaumann, D. E.; Fitzsimmons, B. J.; Ritchie, B. M.; Fraser-Reid, B. J. *Org. Chem.* **1982**, *47*, 941. (b) Achmatowicz, O., Jr.; Burzyska, M. H. *Tetrahedron* **1982**, *38*, 3507. (c) Panfil, I.; Chmielewski, M. *Tetrahedron* **1985**, *41*, 4713. (d) Georgiadis, M. P.; Couladouros, E. A. *J. Heterocycl. Chem.* **1991**, *28*, 1325. (e) López Tudanca, P. L.; Jones, K.; Brownbridge, P. J. *Chem. Soc., Perkin Trans. I* **1992**, 533. (f) Taniguchi, T.; Nakamura, K.; Ogasawara, K. *Synlett* **1996**, 971. (g) Taniguchi, T.; Ohnishi, H.; Ogasawara, K. *Chem. Commun.* **1996**, 1477. (h) Caddick, S.; Khan, S.; Frost, L. M.; Smith, N. J.; Cheung, S.; Paireau, G. *Tetrahedron* **2000**, *56*, 8953. (i) Hoffmann, H. M. R.; Krumwiede, D.; Mucha, B.; Oehlerking, H. H.; Prahst, G. W. *Tetrahedron* **1993**, *49*, 8999. (j) Harris, J. M.; Keränen, M. D.; Nguyen, H.; Young, V. G.; O'Doherty, G. A. *Carbohydr. Res.* **2000**, *328*, 17.
- For a review on Pd-catalyzed allylic substitution, see: Trost, B. M.; Van Vranken, D. L. *Chem. Rev.* **1996**, *96*, 395. For examples using phenols, see: Trost, B. M.; Tsui, H.-C.; Toste, F. D. *J. Am. Chem. Soc.* **2000**, *122*, 3534 and references therein.
- (a) Cuiper, A. D.; Kellogg, R. M.; Feringa, B. L. *Chem. Commun.* **1998**, 655. (b) Trost, B. M.; Schroeder, G. M. *J. Am. Chem. Soc.* **2000**, *122*, 3785. (c) Kim, H.; Lee, C. *Org. Lett.* **2002**, *4*, 4369.
- Glycosides have been allyl protected using a Pd-catalyzed allylic substitution reaction: Lakhmiri, R.; Lhoste, P.; Sinou, D. *Tetrahedron Lett.* **1989**, *30*, 4669.
- Van der Deen, H.; Van Oeveren, A.; Kellogg, R. M.; Feringa, B. L. *Tetrahedron Lett.* **1999**, *40*, 1755.
- Racemates of compounds **2A–2F** were prepared using $(\pm)\text{-1}$ and 5 mol % $\text{Pd}(\text{OAc})_2$ at ambient temperature. Higher catalyst loading and reduced temperature were found necessary for high stereoselectivities.
- Roussel, F.; Takhi, M.; Schmidt, R. R. *J. Org. Chem.* **2001**, *66*, 8540.
- (a) *Solid Support Oligosaccharide Synthesis and Combinatorial Carbohydrate Libraries*; Seeberger, P. H., Ed.; Wiley: New York, 2001. (b) Osborn, H. M. I.; Khan, T. H. *Tetrahedron* **1999**, *55*, 1807.
- For a review on the synthesis of these biomolecules, see: Herzner, H.; Reipen, T.; Schultz, M.; Kunz, H. *Chem. Rev.* **2000**, *100*, 4495.
- $(\pm)\text{-3}$ and $(\pm)\text{-9}$ were separated by preparative chiral HPLC.
- Murphy, P. V.; O'Brien, J. L.; Smith, A. B., III. *Carbohydr. Res.* **2001**, *334*, 327.
- Gensler, W. J.; Johnson, F. A.; Sloan, A. D. B. *J. Am. Chem. Soc.* **1960**, *82*, 6074.
- Achmatowicz, O.; Gryniewicz, G. *Carbohydr. Res.* **1977**, *54*, 193.
- Preliminary steps toward a disaccharide from glucofuranose adduct **2I** proceeded also with complete diastereoselectivity to give **4I**, albeit so far with low yield.



- (a) Achmatowicz, O., Jr.; Gryniewicz, G.; Szechner, B. *Tetrahedron* **1976**, *32*, 1051. (b) Ferroud, D.; Collard, J.; Klich, M.; Dupuis-Hamelin, C.; Mauvais, P.; Lassaigue, P.; Bonnefoy, A.; Musicki, B. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 2881.

JA0347538