See discussions, stats, and author profiles for this publication at: https://www.researchgate.net/publication/7642290

# Chimeric $(\alpha/\beta + \alpha)$ -peptide ligands for the BH3-recognition cleft of Bcl-xL: Critical role of the molecular scaffold in protein surface recognition

ARTICLE in JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · SEPTEMBER 2005		
Impact Factor: 12.11 · DOI: 10.1021/ja053678t · Source: PubMed		
CITATIONS	READS	
126	32	

### 7 AUTHORS, INCLUDING:



### Hee-Seung Lee

Korea Advanced Institute of Science and Tec...



SEE PROFILE



### Naoki Umezawa

Nagoya City University

44 PUBLICATIONS 1,844 CITATIONS

SEE PROFILE



## **Shaomeng Wang**

University of Michigan

291 PUBLICATIONS 14,343 CITATIONS

SEE PROFILE



Published on Web 08/06/2005

# Chimeric $(\alpha/\beta + \alpha)$ -Peptide Ligands for the BH3-Recognition Cleft of Bcl-x<sub>L</sub>: Critical Role of the Molecular Scaffold in Protein Surface Recognition

Jack D. Sadowsky,† Margaret A. Schmitt,† Hee-Seung Lee,†.‡ Naoki Umezawa,†.§ Shaomeng Wang,II York Tomita,\*,<sup>⊥</sup> and Samuel H. Gellman\*,<sup>†</sup>

Department of Chemistry, University of Wisconsin, Madison, Wisconsin 53706, Departments of Internal Medicine and Medicinal Chemistry, University of Michigan, Ann Arbor, Michigan 48109, and Lombardi Comprehensive Cancer Center, Georgetown University Medical Center, Washington, D.C. 20007

Received June 4, 2005; E-mail: gellman@chem.wisc.edu; yat@georgetown.edu

Molecules that bind to specific protein surface sites are of fundamental interest, from the perpective of molecular recognition, and practical interest, from the perspective of medicine. Such molecules should disrupt specific protein-protein interactions, which are frequently associated with human diseases. Traditional "small molecule" approaches, very successful for enzyme inhibition, have been less productive for generation of protein-protein interaction antagonists, 1 although some recent achievements are very impressive.<sup>2</sup> We and others are interested in the prospect that unnatural oligomers with discrete folding propensities ("foldamers") might provide a rational and general basis for development of molecules that block protein-protein interactions.<sup>3</sup> Here we explore this possibility in the context of Bcl-x<sub>L</sub>/BH3 domain interactions, a system that is attractive for fundamental studies because considerable structural information is available.4

Interactions within the Bcl-2 protein family control the fate of a cell in response to cytotoxic stimuli. In many cancers, anti-apoptotic Bcl-2 proteins such as Bcl-x<sub>L</sub> are overexpressed and protect malignant cells from death (apoptosis) by direct interaction with pro-apoptotic proteins such as Bak and Bad.<sup>5</sup> Thus, inhibitors of the Bcl-x<sub>L</sub>/Bak interaction could be therapeutically useful. A 16residue peptide from the BH3 domain of Bak binds as an α-helix to a hydrophobic groove on Bcl-x<sub>L</sub>, burying four hydrophobic Bak side chains (Val-74, Leu-78, Ile-81, and Ile-84).6 Many small molecule ligands for the BH3-recognition domain have been described. Most have only modest activity ( $K_i$  values in competition binding assays typically  $> 1 \mu M$ ), perhaps because of the large surfaces buried in the Bcl-x<sub>I</sub>/Bak 16-mer complex; however, a potent small molecule has very recently been reported.<sup>2c</sup> Numerous medium-length α-peptides (16-32 residues) have shown high affinity for Bcl-x<sub>L</sub>.8 Foldamers that mimic the α-helical display of Bak side chains might be a good source of Bcl-x<sub>L</sub>/Bak interaction antagonists.<sup>3c</sup> Foldamers can be proteolytically stable,<sup>9</sup> an advantage relative to α-peptide inhibitors. 9 We describe foldamers containing both  $\alpha$ - and  $\beta$ -amino acid residues that compete effectively with the Bak 16-mer for binding to Bcl-xL; the development of these antagonists illustrates principles that may be general for foldamerbased approaches to inhibitors of protein-protein interactions.

Our initial efforts focused on  $\beta$ -peptides as potential ligands for the BH3-recognition cleft of Bcl-x<sub>L</sub>, because it is possible to design  $\beta$ -peptides that will adopt specific helical conformations, the 12or the 14-helix, and display side chains in predictable arrangements.  $^{10}$  Of these two  $\beta$ -peptide scaffolds, the 12-helix appears to be a better structural match to the  $\alpha$ -helix. <sup>3b,10</sup> After evaluating > 100

We wondered whether the failure of **1** and related  $\alpha/\beta$ -peptides to show greater efficacy in the FP assay reflects a local mismatch between some portion of the helical scaffold and the BH3recognition cleft of Bcl-x<sub>L</sub>. To test this hypothesis, we examined chimeric oligomers in which either the N-terminal portion or the C-terminal portion of 1 is replaced by an  $\alpha$ -amino acid segment based on an  $\alpha$ -peptide known to bind tightly to Bcl-x<sub>L</sub>. For example,  $(\alpha/\beta + \alpha)$  oligomer 2 contains the first nine residues of  $\alpha/\beta$ -peptide 1, but the last six  $\alpha$ -residues are related to the C-terminal segment of the Bak 16-mer, with Phe-13 of 2 intended to correspond to

Ile-84 of Bak. In  $(\alpha + \alpha/\beta)$  oligomer 3, the first nine residues correspond to positions 72-81 of Bak with Val-74 replaced by

Leu; the final seven residues correspond to the C-terminal portion

of  $\alpha/\beta$ -peptide 1. These complementary chimeric analogues of 1

show very different activities in the FP assay: for  $(\alpha/\beta + \alpha)$ 

University of Wisconsin.

12-helical  $\beta$ -peptide sequences, however, we could obtain only very

weak Bcl-x<sub>L</sub> ligands (IC<sub>50</sub>  $> 500 \mu$ M in a competition fluorescence

helix and 14/15-helix formed by  $\alpha/\beta$ -peptides (oligomers with a 1:1 alternation of  $\alpha$ - and  $\beta$ -amino acid residues along the backbone),<sup>12</sup> and we designed new Bcl-x<sub>L</sub> ligand candidates based on these secondary structures. Both helices appear to be promoted by  $\beta$ -residues with a five-membered ring constraint. <sup>12</sup>  $\alpha/\beta$ -Peptide designs based on the 11-helix were no more successful than the pure  $\beta$ -peptide designs (IC<sub>50</sub> > 500  $\mu$ M). In contrast, designs based on the 14/15-helix displayed significant activity in the FP assay, although these  $\alpha/\beta$ -peptides were not as effective as  $\alpha$ -peptides corresponding to natural BH3 domain sequences. For example,  $\alpha/\beta$ peptide 15-mer 1 displayed IC<sub>50</sub> = 40  $\mu$ M ( $K_i = 1.5 \mu$ M), <sup>13</sup> while  $IC_{50} = 0.67 \ \mu M \ (K_i = 0.025 \ \mu M)$  for the unlabeled Bak 16-mer (Figure 1).<sup>14</sup> In **1** and related  $\alpha/\beta$ -peptide designs, Leu-6 is intended to play the role of the Leu residue conserved in all BH3 domains reported to date (e.g., Leu-78 of Bak). We speculate that ACPC-3,  $\beta^3$ -homonorleucine-9 ( $\beta^3$ -hNle-9), and  $\beta^3$ -hPhe-13 of 1 also contribute to the hydrophobic surface required for binding to the BH3-recognition cleft of Bcl-x<sub>L</sub>. Arg-4 and Asp-11 of 1 may be involved in electrostatic interactions with residues on the edge of the Bcl-x<sub>L</sub> cleft, as proposed for analogous residues in the Bak 16mer.6

Current address: Korea Advanced Institute of Science and Technology (Korea).

<sup>§</sup> Current address: Nagoya City University (Japan).

University of Michigan.

<sup>&</sup>lt;sup>⊥</sup> Georgetown University.





Figure 1. Competition FP data for binding to Bcl- $x_L$  of 1–3, Bak 16-mer, and control hexa-α-peptide Ac-GDAFNR-NH<sub>2</sub>. A fluorophore-labeled Bak 16-mer peptide was used as the displaced fluorescent probe (see Supporting Information for details).

oligomer **2** IC<sub>50</sub> = 0.059  $\mu$ M ( $K_i$  = 0.0019  $\mu$ M), while for ( $\alpha$  +  $\alpha/\beta$ ) oligomer **3** IC<sub>50</sub> > 700  $\mu$ M (Figure 1). Thus, **2** is 10-fold more potent than the Bak 16-mer. This result suggests that the 14/15-helical  $\alpha/\beta$ -peptide scaffold is well-suited to occupy a portion of the BH3-recognition cleft on Bcl-x<sub>L</sub>, but that some other foldamer backbone, distinct from the  $\beta$ - and  $\alpha/\beta$ -peptides we have examined thus far, will be necessary to replace the proteolytically susceptible  $\alpha$ -peptide segment<sup>15</sup> for optimal fit to the remainder of this cleft.


Several control studies were conducted with  $(\alpha/\beta+\alpha)$  oligomer 2 and related compounds. The hexa- $\alpha$ -peptide corresponding to the C-terminal segment of 2 (Ac-GDAFNR-NH2) at 500  $\mu M$  displayed no interaction with Bcl-xL in the FP assay (Figure 1). Thus, the  $\alpha$ -peptide segment of chimeric oligomer 2 is probably not the dominant contributor to Bcl-xL binding affinity.  $^{16}$ 

Binding of 2 to  $^{15}$ N-labeled Bcl- $x_L$  was examined via  $[^{1}$ H, $^{15}$ N]-HSQC NMR measurements (Figure 2). Most of the Bcl- $x_L$  amide N-H cross-peaks were significantly shifted upon addition of 50  $\mu$ M 2 to 100  $\mu$ M Bcl- $x_L$  (Figure 2A). The pattern of shifts and resonance broadening induced by addition of 2 are comparable to the effects induced by addition of the Bak 16-mer  $\alpha$ -peptide (Figure 2B).

A fluorescein-labeled derivative of **2** (**Flu-2**) was prepared to compare binding to Bcl- $x_L$  with binding to unrelated proteins. Direct FP titration of 50 nM **Flu-2** with protein indicates  $K_d = 0.014 \, \mu \text{M}$  for **Flu-2** binding to Bcl- $x_L$  (Figure 3). In contrast, no binding to bovine  $\gamma$ -globulin (BGG) could be detected at 500  $\mu \text{M}$  BGG, and the onset of binding to bovine serum albumin (BSA) occurred above 10  $\mu \text{M}$  BSA (Figure 3). Thus, binding of **Flu-2** to BGG or BSA is at least  $10^3$ -fold weaker than binding to Bcl- $x_L$ . Both BGG and BSA are promiscuous receptors for hydrophobic ligands; <sup>17</sup> the failure of **Flu-2** to bind tightly to either of these proteins suggests




**Figure 2.** HSQC NMR binding assays. (A) Overlay of  $^{15}$ N-Bcl- $^{1$ 



**Figure 3.** Direct FP titration of fluorescein-labeled ( $\alpha/\beta + \alpha$ )-peptide **Flu-2** with Bcl-x<sub>L</sub>, BSA, and BGG.

that the affinity of **2** for Bcl-x<sub>L</sub> is not simply the result of its hydrophobicity, but instead reflects complementarity to the BH3-recognition cleft. As a further test of such complementarity, we compared ( $\alpha/\beta + \alpha$ ) oligomer **4** and its enantiomer in the competition FP assay (Figure 4). Oligomer **4** is an isomer of **2** in which  $\beta^3$ -hNle-9 has been replaced by  $\beta^3$ -hLeu; this small change leads to slightly improved affinity for Bcl-x<sub>L</sub> (IC<sub>50</sub> = 0.029  $\mu$ M,  $K_i = 0.0007 \ \mu$ M for **4**). The enantiomer of **4**, however, displays very low affinity for Bcl-x<sub>L</sub> (IC<sub>50</sub> > 1000  $\mu$ M).

The folding of  $(\alpha/\beta + \alpha)$  oligomer **5** in CD<sub>3</sub>OH was examined by 2D NMR; <sup>18</sup> **5** has two modifications relative to **2** (Ala-2  $\rightarrow$  Lys and Lys-8  $\rightarrow$  Ile), which moderately diminish binding to the BH3-recognition cleft of Bcl-x<sub>L</sub> (IC<sub>50</sub> = 0.40  $\mu$ M). Good dispersion of <sup>1</sup>H resonances was observed for **5**, which allowed assignment of many NOEs between backbone protons. Numerous i,i+3 NOEs were observed along the entire length of **5** (Figure 5). Of particular importance are the three  $\alpha$ -residue H<sub> $\alpha$ </sub>(i)  $\rightarrow$   $\beta$ -residue H<sub> $\alpha$ </sub>(i+3) NOEs in the  $\alpha/\beta$ -peptide segment of **5**. This NOE pattern is predicted for the 14/15-helix but not for the 11-helix. <sup>12</sup> In contrast,  $\alpha$ -residue H<sub> $\alpha$ </sub>(i)  $\rightarrow$   $\alpha$ -residue NH(i+2) NOEs are predicted for the 11-helix but not for the 14/15-helix, <sup>12</sup> and none of these NOEs



**Figure 4.** Competition FP binding assay for binding to Bcl-x<sub>L</sub> of  $(\alpha/\beta+\alpha)$ -peptide 4 and its enantiomer.



Figure 5. NOEs observed for oligomer 5 in CD<sub>3</sub>OH. NOEs consistent with the 14/15-helix only (blue arcs), both the 11- and 14/15-helix (black solid arcs), and α-helix (red arcs). Ambiguously assigned NOEs are represented by dotted arcs.

is observed for **5**. Thus, the NMR data suggest that **5** has a substantial propensity to adopt the 14/15-helical secondary structure in its N-terminal region, a propensity that is likely to be manifested also by closely related molecules such as **2**. Interestingly, the i,i+3 NOEs involving the C-terminal  $\alpha$ -peptide region of **5** suggest that the 14/15-helical  $\alpha/\beta$ -segment can nucleate helix formation in the short  $\alpha$ -peptide segment.

Our results show that foldamer-based designs can provide tight-binding ligands for a large protein recognition site ( $K_i$  for  $\mathbf{4}=0.7$  nM). The path from  $\beta$ - to  $\alpha/\beta$ - to  $(\alpha/\beta+\alpha)$ -peptide ligands such as 2 leads us to conclude that foldamer-based strategies for disrupting protein—protein interactions will grow in scope and efficacy as the number of foldamer scaffolds with distinct shapes is increased. The tight binding of chimeric  $(\alpha/\beta+\alpha)$ -peptides to Bcl-x<sub>L</sub> suggests that combining different foldamer scaffolds will be an effective and perhaps general strategy for protein ligand design.

**Acknowledgment.** This research was funded by the NIH (GM56414). We thank Z. Nikolovska-Coleska (U. Mich.) for assistance with FP assays and Yumi Udea (Georgetown) for assistance with protein purification. We are grateful to G. Wagner (Harvard), A. Lugovskoy (Biogen Idec), and P. Hajduk, E. Olejniczak, and colleagues (Abbott) for providing Bcl-x<sub>L</sub> NMR assignments. The fluorescence plate reader is in the UW-Madison W. M. Keck Center for Chemical Genomics. An NSF predoctoral fellowship and an NIH Molecular Biophysics grant (GM08293-14) partially supported J.D.S. and M.A.S., respectively.

**Supporting Information Available:** Experimental details, peptide characterization data (HPLC, MS), and other results. This material is available free of charge via the Internet at http://pubs.acs.org.

### References

- (1) (a) Arkin, M. R.; Wells, J. A. Nat. Rev. Drug Discovery 2004, 3, 301–317.
   (b) Cochran, A. G. Chem. Biol. 2000, 7, R85-R94.
   (c) Gadek, T. R.; Nicholas, J. B. Biochem. Pharmacol. 2003, 65, 1-8.
   (d) Berg, T. Angew. Chem., Int. Ed. 2003, 42, 2462-2481.
   (e) Peczuh, M. W.; Hamilton, A. D. Chem. Rev. 2000, 100, 2479-2494.
- (2) (a) Arkin, M. R.; Randal, M.; DeLano, W. L.; Hyde, J.; Luong, T. N.; Oslob, J. D.; Raphael, D. R.; Taylor, L.; Wang, J.; McDowell, R. S.; Wells, J. A.; Braisted, A. C. *Proc. Natl. Acad. Sci. U.S.A.* 2003, 100, 1603–1608. (b) Vassilev, L. T.; Vu, B. T.; Graves, B.; Carvajal, D.; Podlaski, F.; Filipovic, Z.; Kong, N.; Kammlott, U.; Lukacs, C.; Klein, C.; Fotouhi, N.; Liu, E. A. *Science* 2004, 303, 844–848. (c) Oltersdorf, T., et al. *Nature* 2005, 435, 677–681.
- (3) (a) Werder, M.; Hauser, H.; Abele, S.; Seebach, D. Helv. Chim. Acta 1999, 82, 1774–1783. (b) Lee, H.-S.; Syud, F. A.; Wang, X.; Gellman, S. H. J. Am. Chem. Soc. 2001, 123, 7721–7722. (c) Ernst, J. T.; Becerril, J.; Park, H. S.; Yin, H.; Hamilton, A. D. Angew. Chem., Int. Ed. 2003, 42, 535–539. (d) Kritzer, J. A.; Lear, J. D.; Hodsdon, M. E.; Schepartz, A. J. Am. Chem. Soc. 2004, 126, 9468–9469.
- (4) Petros, A. M.; Olejniczak, E. T.; Fesik, S. W. Biochim. Biophys. Acta 2004, 1644, 83–94.
- (a) Adams, J. M.; Cory, S. Science 1998, 281, 1322-1326.
   (b) Coultas, L.; Strasser, A. Semin. Cancer Biol. 2003, 13, 115-123.
   (c) Letai, A.; Bassik, M. C.; Walensky, L. D.; Sorcinelli, M. D.; Weiler, S.; Korsmeyer, S. J. Cancer Cell 2002, 2, 183-192.
- (6) Sattler, M.; Liang, H.; Nettesheim, D.; Meadows, R. P.; Harlan, J. E.; Eberstadt, M.; Yoon, H. S.; Shuker, S. B.; Chang, B. S.; Minn, A. J.; Thompson, C. B.; Fesik, S. W. Science 1997, 275, 983–986.
- (7) Reviews of small molecule-based approaches to Bcl-x<sub>L</sub> inhibition: (a) Baell, J. B.; Huang, D. C. S. *Biochem. Pharmacol.* 2002, 64, 851–863.
  (b) O'Neill, J.; Manion, M.; Schwartz, P.; Hockenbery, D. M. *Biochim. Biophys. Acta* 2004, 1705, 43–51. (c) For a designed Bak mimic with K<sub>i</sub> = 0.1 μM, see: Kutzki, O.; Park, H. S.; Ernst, J. T.; Orner, B. P.; Yin, H.; Hamilton, A. D. *J. Am. Chem. Soc.* 2002, 124, 11838–11839.
- (8) (a) Petros, A. M.; Nettesheim, D. G.; Wang, Y.; Olejniczak, E. T.; Meadows, R. P.; Mack, J.; Swift, K.; Matayoshi, E. D.; Zhang, H.; Thompson, C. B.; Fesik, S. W. Protein Sci. 2000, 9, 2528–2534. (b) Walensky, L. D.; Kung, A. L.; Escher, I.; Malia, T. J.; Barbuto, S.; Wright, R. D.; Wagner, G.; Verdine, G. L.; Korsmeyer, S. J. Science 2004, 305, 1466–1470. (c) Gemperli, A. C.; Rutledge, S. E.; Maranda, A.; Schepartz, A. J. Am. Chem. Soc. 2005, 127, 1596–1597.
- (9) Frackenpohl, J.; Arvidsson, P. I.; Schreiber, J. V.; Seebach, D. Chem-BioChem 2001, 2, 445–455.
- (10) Cheng, R. P.; Gellman, S. H.; DeGrado, W. F. *Chem. Rev.* **2001**, *101*, 3219—3232.
- (11) Competition FP assay involved displacement of BODIPY-TMR-labeled Bak 16-mer from Bcl-x<sub>L</sub>. See Supporting Information for details.
- (12) (a) Hayen, A.; Schmitt, M. A.; Ngassa, F. N.; Thomasson, K. A.; Gellman, S. H. Angew. Chem., Int. Ed. 2004, 43, 505-510. See also: (b) De Pol, S.; Zorn, C.; Klein, C. D.; Zerbe, O.; Reiser, O. Angew. Chem., Int. Ed. 2004, 43, 511-514.
- (13)  $K_i$  calculated from IC<sub>50</sub>: Wang, R.; Nikolovska-Coleska, Z.; Fang, X.; Wang, S. The Ki Calculator. http://sw16.im.med.umich.edu/software/calc\_ki/ (accessed June 2005). Parameters:  $K_d$  for BODIPY-Bak tracer = 1 nM, [BODIPY-Bak] = 33 nM, [Bcl- $x_L$ ] = 20 nM.
- (14) This K<sub>i</sub> for unlabeled Bak 16-mer is similar to the reported value: Zhang, H.; Nimmer, P.; Rosenberg, S. H.; Ng, S.; Joseph, M. Anal. Biochem. 2002, 307, 70–75.
- (15) Rapid proteolysis of the  $\alpha$ -peptide segment of 2 is observed upon treatment with Pronase, while the  $\alpha/\beta$  segment remains intact for at least 24 h.
- (16) The  $\alpha/\beta$ -peptide oligomer corresponding to residues 1–9 of chimeric oligomer **4**, an isomer of **2** (see text), has an IC<sub>50</sub> of 120  $\mu$ M in the FP assay, suggesting that the  $\alpha/\beta$  portions of **2** and **4** target the BH3 binding site and contribute substantially to binding.
- (17) (a) Ali, M.; Smith, L. M.; Feeney, R. E. Biochim. Biophys. Acta 1969, 194, 256–264. (b) Church, W. R.; Rawitch, A. B.; Ebner, K. E. Arch. Biochem. Biophys. 1981, 206, 285–290. (c) Alizadeh-Pasdar, N.; Li-Chan, E. C. Y. J. Agric. Food Chem. 2000, 48, 328–334.
- (18) Among α- and β-peptides, alcohol solvents are known to promote secondary structure relative to water; the folding observed in such organic solvents generally reflects the conformational propensities in water. (a) Luo, P.; Baldwin, R. L. Biochemistry 1997, 36, 8413–8421. (b) Abele, S.; Guichard, G.; Seebach, D. Helv. Chim. Acta 1998, 81, 2141–2156. (c) Arvidsson, P. I.; Rueping, M.; Seebach, D. J. Chem. Soc., Chem. Commun. 2001, 649–650.

JA053678T