

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6270173>

Cyclic Oxonitriles: Stereodivergent Grignard Addition—Alkylations

ARTICLE *in* THE JOURNAL OF ORGANIC CHEMISTRY · AUGUST 2007

Impact Factor: 4.72 · DOI: 10.1021/jo070678y · Source: PubMed

CITATIONS

8

READS

31

4 AUTHORS, INCLUDING:



Fraser Fleming

Drexel University

130 PUBLICATIONS 1,575 CITATIONS

SEE PROFILE



Omar W Steward

Duquesne University

80 PUBLICATIONS 846 CITATIONS

SEE PROFILE

Published in final edited form as:

J Org Chem. 2007 July 6; 72(14): 5270–5275. doi:10.1021/jo070678y.

Cyclic Oxonitriles: Stereodivergent Grignard Addition-Alkylations

Fraser F. Fleming*, Guoqing Wei, Zhiyu Zhang, and Omar W. Steward

Department of Chemistry and Biochemistry, Duquesne University, Pittsburgh, Pennsylvania 15282-1530.

Abstract

Sequential carbonyl addition-conjugate addition of Grignard reagents to cyclic 5–7-membered oxoalkenenitriles efficiently generates cyclic magnesiated nitriles. Alkylations of these magnesiated nitriles exhibit diastereoselectivities that depend intimately on the size of the carbocyclic ring: 5-membered oxonitriles generate magnesiated nitriles whose alkylations are controlled by steric constraints whereas 6- and 7-membered oxonitriles generate internally coordinated, *C*-magnesiated nitriles whose alkylations are controlled by stereoelectronic effects. Reversing the alkylation selectivity of 6-membered *C*-magnesiated nitriles is achieved by conversion to an *N*-metalated nitrile in which steric, rather than electronic, effects direct the electrophile trajectory. Collectively, the conjugate addition-alkylation generates highly substituted, cyclic 5–7-membered nitriles containing three new stereocenters with selective access to diastereomers at the quaternary nitrile-bearing carbon.

Introduction

Conjugate additions to alkenenitriles are notoriously difficult.¹ Highly nucleophilic organometallics are predisposed to add to the polarized nitrile group whereas more modest nucleophiles, such as cuprates, often fail to react.² Three general strategies have emerged to address the difficulty of conjugate addition to alkenenitriles:¹ activating the alkenenitrile by conjugation with an additional electron-withdrawing group,³ using radicaloid species for the conjugate addition,⁴ and employing highly nucleophilic organometallic reagents in intramolecular conjugate additions to *E*-alkenenitriles where addition to the nitrile group is geometrically prevented.⁵ The latter strategy is particularly effective with cyclic nitriles in which highly nucleophilic alkylmagnesium alkoxides are temporarily tethered to the proximal alkenenitrile, effectively using intramolecular delivery to achieve a formal intermolecular conjugate addition.⁶

Cyclic γ -oxoalkenenitriles⁷ are excellent electrophiles for chelation-controlled conjugate addition reactions. Addition of excess MeMgCl to 3-oxo-1-cyclohexene-1-carbonitrile (**1b**)⁸ triggers a complex cascade (Scheme 1) in which carbonyl addition initially affords a halomagnesium alkoxide that suffers halogen-methyl exchange with a second equivalent of MeMgCl to generate the methylmagnesium alkoxide **2**. Tethering the highly nucleophilic methylmagnesium alkoxide of **2** in close proximity to the alkenenitrile triggers a stereoelectronically controlled axial conjugate addition.^{6b} Equilibration of the resulting *N*-magnesiated nitrile **3** to the more stable *C*-magnesiated nitrile **4** is favored by internal coordination and a conformation in which the two methyl groups adopt an equatorial orientation.

flemingf@duq.edu.

Alkylations of the *C*-magnesiased nitrile **4** are highly unusual. Alkylating **4** with alkyl halides and sulfonates occurs with retention of the *C*-Mg configuration (**4**→**5a**), whereas aldehyde and acyl cyanide acylations proceed with inversion of stereochemistry (**4**→**5b**).⁹ The origin of this electrophile-dependent stereoselectivity stems from the size and accessibility of the electrophile's anti-bonding orbital. Highly reactive, planar electrophiles with large π^* orbitals can achieve a favorable collinear overlap with the small orbital of the *C*-Mg bond in an invertive S_E2 alkylation, whereas sterically demanding electrophiles with smaller anti-bonding orbitals can only access the *C*-Mg bond through a side-on overlap resulting in a retentive S_E2 alkylation.¹⁰

The electrophile-dependent alkylations of the internally coordinated *C*-magnesiased nitrile **4** directly contrast with alkylations of comparable *N*-¹¹ and *C*-metalated nitriles.¹² In general, deprotonating cyclic nitriles with lithium amides generate planar *N*-lithiated nitriles (**7**; Scheme 2) whose alkylations are controlled by steric effects.¹³ For example, deprotonating the conformationally constrained nitrile **6** exposes two potentially nucleophilic faces with preferential interception of electrophiles occurring from the more sterically accessible equatorial direction (**7**→**9**). Alkylating the corresponding *C*-magnesiased nitrile **11**, generated by bromine-magnesium exchange (**10**→**11**), generates the equatorially substituted nitrile **9** regardless of the nature of the electrophile.¹²

The disparate alkylation stereoselectivities of typical *N*- and *C*-metalated nitriles such as **7** and **11**, when compared with the internally coordinated *C*-magnesiased nitrile **4**, stimulated a series of alkylations with analogous 5- and 7-membered magnesiased nitriles. Surprisingly, sequential Grignard additions to 5-membered oxonitriles generate magnesiased nitriles whose alkylations are controlled by steric constraints whereas 6- and 7-membered oxonitriles generate internally coordinated *C*-magnesiased nitriles whose alkylations are controlled by stereoelectronic effects. Surveying an array of addition-alkylations with cyclic oxonitriles of varying ring size generates highly substituted 5–7-membered nitriles, establishes fundamental stereoselectivity preferences for alkylations of the intermediate magnesiased nitriles, and provides a strategy for changing the stereochemistry at the quaternary nitrile-bearing carbon.

Results and Discussion

Grignard addition-alkylations to oxonitrile **1b** rapidly assembles highly substituted nitriles (Scheme 1). The unusual alkylation stereoselectivity of the intermediate magnesiased nitrile was investigated by an extensive series of alkylations with the magnesiased nitrile **4** derived by sequential carbonyl and conjugate additions with MeMgCl.⁹ Probing the alkylation stereoselectivity for the analogous 5-membered nitrile (Scheme 3) therefore employed MeMgCl for the carbonyl and conjugate additions to oxonitrile **1a**¹⁴ to provide a direct point of comparison. Adding MeMgCl to the 5-membered oxonitrile **1a** causes rapid carbonyl addition, with the subsequent conjugate addition via the methylmagnesium alkoxide **12** being significantly more challenging than for the 6-membered oxonitrile **1b**.¹⁵ The difficulty arises, in part, because the rehybridization required during the intramolecular conjugate addition requires a pyramidalization of the β -carbon that is less readily accommodated in a 5-membered ring than a 6-membered ring.^{6b} The reactivity difference is apparent in the efficiency differences for the addition-alkylations of the 5- and 6-membered oxonitriles **1a** and **1b** (49% and 86% yield, respectively; Table 1, entries 1 and 7).

The chelation-controlled conjugate addition to oxonitrile **1a** affords the 5-membered magnesiased nitrile **13** whose alkylations reveal an immediate point of departure with comparable alkylations of the 6-membered nitrile **4**. The methylation of **4** occurs from the axial direction and is completely selective (Scheme 1) whereas intercepting **13** with MeI is only modestly selective (Scheme 3), installing the new methyl group predominantly from the

pseudo-equatorial orientation (Table 1, compare the configuration of the nitrile-bearing carbon in entries 1 and 7). The stereoselectivity differences suggest that the initially formed bis-magnesiates nitrile **13a** does not equilibrate to an internally coordinated nitrile analogous to **4**, but exists as a diastereomeric mixture of *C*-magnesiates nitriles **13b** and **13c** (Scheme 3).¹⁶ Presumably internal coordination between the proximal oxygen and magnesium atoms is prevented in **13b** because of strain within the ensuing bicyclic scaffold. The *C*-magnesiates nitrile **13c** is likely to be the more favored diastereomer because the small nitrile¹⁷ is positioned in a sterically compressed environment while the large solvated magnesium occupies the less demanding pseudo equatorial orientation. Retentive alkylation from **13c** leads to the major diastereomer **14a**.

Alkylations of **13** with electrophiles larger than MeI are completely selective (Table 1, entries 2–5). Intercepting **13** with cyclohexanone is an exception that likely stems from a facile retro-aldol-like fragmentation¹⁸ causing rapid equilibration to a mixture of diastereomers (Table 1, entry 6). The otherwise consistent preference of the 5-membered nitrile **13** for alkylation opposite to the hydroxyl group, contrasts with alkylations of the 6-membered, *C*-magnesiates nitrile **4** where alkylation with MeI proceeds with stereochemical retention (Table 1, entry 7), inversion for acylation with MeOCOCN (Table 1, entry 8), and essentially no stereocontrol with CH₂=CHCH₂Br (Table 1, entry 9).⁹ Collectively, the stereoselectivity differences underscore the structural differences between the magnesiates 5- and 6-membered nitriles **4** and **13**.

Grignard addition-alkylations with the 7-membered oxonitrile **1c**¹⁴ bear a striking similarity to those of the 6-membered oxonitrile **1b** (Table 1, entries 10–12). Addition of MeMgCl to oxonitrile **1c** and alkylating the magnesiates nitrile intermediate with MeI installs a methyl substituent in an axial orientation, although the efficiency of the carbonyl- and conjugate additions are diminished relative to the 6-membered oxonitrile (Table 1, compare entries 7 and 10).²¹ Intercepting the magnesiates nitrile derived from **1c** with methyl cyanoformate causes acylation from the equatorial direction, consistent with stereochemical inversion akin to that of the 6-membered oxonitrile (Table 1, compare entries 8 and 11). Collectively these alkylation preferences suggest formation of the internally coordinated, *C*-magnesiates nitrile **16** which intercepts methyl cyanoformate by an initial invertive acylation leading to **17** (Scheme 4). Subsequent internal attack of the magnesium alkoxide on the proximal nitrile is facilitated in the flexible 7-membered ring²² with the resulting imino lactone reacting with excess methyl cyanoformate to afford **15b**. The overall conversion installs 5 new bonds in a remarkable cascade.

Intercepting the *C*-magnesiates nitrile **16** with pivaloyl chloride and MeMgCl results in the formation of the enamide **15c** (Table 1, entry 12). The unusual conversion to an enamide upon alkylation with pivaloyl chloride is a signature of internally coordinated, *C*-magnesiates nitriles.²³ Mechanistically, *N*-acylation of the magnesiates nitrile **16** (Scheme 4) is favored with the large, highly polarized electrophile pivaloyl chloride which affords the reactive acyl ketenimine **18**. Rapid addition to the acyl ketenimine **18** is facilitated by performing the acylation in the presence of the Grignard reagent which attacks from the more sterically accessible face opposite the neighboring secondary methyl group. Protonation of the resulting bis-metalated enamide intermediate ultimately affords the enamide **15c**.

Comparative alkylations and acylations with 6- and 7-membered magnesiates nitriles are consistent with forming internally coordinated *C*-magnesiates nitriles. Analogous alkylations with the corresponding 5-membered magnesiates nitrile **13** implicate a structurally different metalated nitrile. The reactivity difference is evident on intercepting **13** with pivaloyl chloride which affords only the pseudo-equatorially oriented ketone **14d** (Table 1, entry 4) as does acylation with the sterically more demanding triethylacetyl chloride²⁴ (Table 1, entry 5). The

complete absence of enamides in the acylations of the 5-membered magnesiated nitrile **13**²⁵ is consistent with the intermediacy of a *C*-magnesiated nitrile devoid of internal coordination.

Divergent stereoselectivities in multi-component addition-alkylations of 5–7-membered oxonitriles reveals the key role of internal chelation in alkylations of *C*-magnesiated nitriles. The stereoselectivity differences suggested an intriguing strategy for reversing the alkylation stereoselectivity of the *C*-magnesiated nitrile **4** by conversion to the corresponding *N*-metalated nitrile **19** (Scheme 5). Brief optimization identified a procedure in which the sequential addition of 6 equivalents of BuLi and MeI, preferentially affords the equatorially methylated nitrile **5d** and diastereomer **5a** in a 12:1 ratio (Scheme 5).²⁶ The stereoselectivity is consistent with an initial conversion of **4** to the magnesiate **19**,²⁷ followed by alkylation from the more accessible equatorial face. Extending the protocol to alkylations with allyl bromide and propyl iodide similarly favored equatorial alkylation, although to a more modest extent (Table 2, compare entry 1 with entries 2 and 3).

Assuming that an equilibrium exists between the *C*- and *N*-magnesiated nitriles **4** and **19**, several additives were screened with the aim of generating a more stable magnesiate analogous to **19**. Although LiF was ineffective (Table 1, entry 4), lithium butoxide led to equal amounts of the two diastereomers (Table 1, entry 5). Formation of a strong Mg–O bond²⁸ and the entropic advantage of forming an ate complex with a bifunctional ligand, provided the impetus to use lithium alkoxides derived from 1,2-diols. Adding the lithium alkoxide obtained by treating ethylene glycol with BuLi improved the preference for equatorial alkylation although the reagent derived from 1,2-cyclohexanediol is significantly more effective (Table 2, entries 6 and 7, respectively). The latter protocol provides an effective method for equatorial alkylations. Addition of the alkoxide and alkylation with allyl bromide favors equatorial alkylation by 7.9:1 whereas alkylating **4** with allyl bromide is essentially non-selective (compare Table 1, entry 9 with Table 2, entries 2 and 8). The stereodivergent cyclohexanediol alkylation strategy is also effective for the sequential addition of two different Grignard reagents followed by alkylation (Table 2, entry 11).

Employing the lithium alkoxide derived from cyclohexanediol and alkylating with cyclopropylmethyl iodide affords the direct alkylation product **5i** and the ring-opened nitrile **5g** in a 1.8:1 ratio (Table 2, entry 12). This deployment of cyclopropylmethyl iodide as a single electron transfer probe²⁹ suggests that S_N2 and single electron transfer mechanisms are of roughly equal efficiency.⁹ In both cases the preferred diastereomer results from an equatorial alkylation, consistent with the steric influence of a bulky axial magnesiate. Effectively, the conversion of the *C*-magnesiated nitrile **4** to the *N*-metalated nitrile **20** (Scheme 5) extends the carbonyl addition-conjugate addition-alkylation strategy by providing selective access to diastereomers at the quaternary, nitrile-bearing stereocenter.

The Grignard addition-alkylation of 5–7 membered oxonitriles efficiently installs contiguous quaternary-tertiary-quaternary stereocenters. Unfortunately, introduction of an additional quaternary center through conjugate addition to a tetra-substituted alkenenitrile appears not to be possible. The strategy was pursued with the bromooxonitrile **21**, prepared by bromination of **1b** (Scheme 6). Addition of excess MeMgCl to **21** triggers sequential carbonyl- and conjugate additions to afford a metalated nitrile postulated to be **22**,³⁰ which ejects bromide to ultimately generate **24**. Despite the use of excess MeMgCl, to promote a chelation-controlled conjugate addition from the magnesiate **23**,³¹ no conjugate addition is observed. The inability to coax the conjugate addition is consistent with the deactivating effect of β-alkyl substituents in alkenenitriles.¹

Conclusion

Grignard addition-alkylations to cyclic 5–7-membered oxonitriles efficiently generate highly substituted cyclic nitriles. The strategy installs contiguous quaternary-tertiary-quaternary centers imbedded within diverse cyclic nitriles. Stereoselective alkylations of the intermediate magnesiated nitriles depend intimately on the size of the carbocyclic ring: 5-membered oxonitriles generate magnesiated nitriles whose alkylations are controlled by steric constraints whereas 6- and 7-membered oxonitriles generate internally coordinated, *C*-magnesiated nitriles whose alkylations are controlled by stereoelectronic effects. Conversion of 6-membered *C*-magnesiated nitriles to the corresponding *N*-metalated nitrile is readily achieved by adding lithium alkoxides which allows steric, rather than electronic, effects to control the alkylation stereoselectivity. Synthetically, this interconversion between *C*-magnesiated and *N*-lithiated nitriles allows stereodivergent alkylations at the quaternary nitrile-bearing carbon. Collectively, the conjugate addition-alkylation installs three new stereocenters in cyclic 5–7-membered nitriles, establishes fundamental stereoselectivity preferences for alkylations of the intermediate magnesiated nitriles, and provides a strategy for changing the stereochemistry at the quaternary nitrile-bearing carbon.

Experimental

General Procedure for Oxonitrile Synthesis

Two individual watchglasses having a thin (< 3mm) layer of powdered chromium trioxide (12 equiv) or 2,3-dimethylpyrazole (12 equiv) were dried overnight (> 16 h) in a vacuum desiccator (< 5mm/Hg) containing P₂O₅. Dry 2,3-dimethylpyrazole was added to a magnetically stirred, –20 °C, CH₂Cl₂ solution (1.2 M) of dry CrO₃. After stirring at –20 °C for 15 min, neat nitrile (1 equiv) was added. The mixture was stirred between –10 °C to –12 °C for 3 h and then the temperature raised to 0 °C. Aqueous NaOH (5M) was then added and stirring continued at 0 °C for 1 h. The crude reaction mixture was then refrigerated (–10 °C) overnight causing the aqueous phase to freeze and the organic phase to be removed by decanting. The aqueous phase was further extracted twice with CH₂Cl₂, the organic extracts were then combined, and the resulting organic material was then washed sequentially with HCl, H₂O and brine. The organic extract was dried (Na₂SO₄) and then concentrated to give a dark yellow oil that was purified by radial chromatography to afford pure oxonitrile.

General Grignard Addition-Alkylation Procedure with Oxonitrile 1a

A THF solution of MeMgCl (2.5 equiv) was added to a –78 °C THF solution (0.1M) of oxonitrile **1a**, and the resulting solution was stirred at –78 °C for 1 h and then warmed to room temperature. After 0.5 h, the solution was cooled to 0 °C and the electrophile (2–10 equiv) was then added neat. After 1 h, saturated aqueous NH₄Cl was added and the resulting mixture was extracted with EtOAc. The extracts were combined, washed with brine and dried (Na₂SO₄). Concentration and purification of the crude product by radial chromatography afforded the pure nitrile.

General Grignard Addition-Alkylation Procedure with Oxonitrile 1c

A THF solution of MeMgCl (2.5–6 equiv) was added to a –78 °C THF solution (0.1 M) of oxonitrile **1c**, and the resulting solution was stirred at –78 °C for 1 h and then warmed to room temperature. After 1.5 h, the electrophile (2.5–6equiv) was added neat either at room temperature or with prior cooling to –78 °C. Subsequent addition of saturated NH₄Cl and extraction with EtOAc afforded a crude product that was washed with brine and dried (MgSO₄), concentrated, and purified by radial chromatography to afford the pure product.

General Procedure for the Addition-Alkylation with Stereochemical Inversion

A THF solution of MeMgCl (1.05–1.1 equiv) was added to a –15 °C THF solution (0.1 M) of the oxonitrile **1b**. After 2 h, a second Grignard reagent (1.2–1.5 equiv), if different from MeMgCl, was added and the solution was then allowed to warm to room temperature. After 2 h, the solution was transferred by syringe to a 0 °C THF solution (0.3 M) of cis-cyclohexane-1,2-diol (3 equiv) to which BuLi (3.3 equiv) had been added. The resulting solution was cooled to –78 °C and after 0.5 h, the electrophile (3.0–10 equiv) was added neat. After 1 h, the solution was allowed to warm to room temperature, and after 16 h, saturated aqueous NH₄Cl was added, and the resulting mixture was extracted with EtOAc. The extracts were combined, washed with brine and dried (NaSO₄). Concentration and purification of the crude product by radial chromatography afforded the pure nitrile.

2-Bromo-3-oxocyclohex-1-ene-1-carbonitrile (**21**)

A 1 mL CH₂Cl₂ solution of the bromine (240 mg, 1.5 mmol) was added to a 0 °C, 4 mL CH₂Cl₂ solution of **1b** (181 mg, 1.5 mmol). After 1.5 h at 0 °C, Et₃N (0.32 mL, 2.25 mmol) was added dropwise. The solution was stirred at room temperature for 1.5 h and was washed with 3% HCl twice and saturated brine once and dried (MgSO₄). Concentration and radial chromatography (stepped gradient 1:9, 3:17, 1:4, 1:3 EtOAc/Hexanes) to afford **21** (210 mg, 70%) as a light yellow oil. IR (film) 2221, 1706, 1692 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 2.10–2.18 (m, 2H), 2.62–2.70 (m, 4H); ¹³C NMR (300 MHz, CDCl₃): δ 21.9, 30.6, 37.4, 116.2, 131.1, 134.6, 188.7; HRMS (EI) calcd for (M⁺), C₇H₆BrNO⁺ 198.9627, found 198.9633.

3-Hydroxy-2, 3-dimethylcyclohex-1-ene-1-carbonitrile (**24**)

A THF solution of MeMgCl (3 M, 3 mmol) was added to a –78 °C THF solution of **21** (0.1 M). After 1 h at –78 °C and 2.5 h at room temperature, MeI was added neat and the resulting solution was stirred overnight. Saturated aqueous NH₄Cl was added and the resultant mixture was extracted with EtOAc. The extracts were combined, washed with brine, and dried (NaSO₄). Concentration and purification of the crude product by radial chromatography (stepped gradient 1:4, 3:7 EtOAc/hexanes) furnished 46 mg (51%) of **24** as a waxy solid. IR (film) 3438, 2213, 1646 cm⁻¹; ¹H NMR (300 MHz, CDCl₃): δ 1.31 (s, 3H), 1.60–1.75 (m, 4H), 2.06 (s, 3H), 2.20–2.22 (m, 2H); ¹³C NMR (300 MHz, CDCl₃): δ 16.7, 19.0, 26.9, 27.8, 38.2, 70.0, 108.7, 118.7, 156.0.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

Financial support of this research from the National Institutes of Health (2R15AI051352), and in part, from the National Science Foundation (CHE 0515715, CRIF 024872 for x-ray facilities, and CHE 0421252 for HRMS instrumentation) are gratefully acknowledged as is assistance from Dr. Derek Beauchamp with the crystallographic analyses.


References

1. Fleming FF, Wang Q. Chem. Rev 2003;103:2035. [PubMed: 12744700]
2. (a) Totleben MJ, Curran DP, Wipf P. J. Org. Chem 1992;57:1740. (b) Yoneda R, Harusawa S, Kurihara T. J. Chem. Soc., Perkin Trans. 1 1988:3163. (c) House HO. Acc. Chem. Res 1976;9:59. (d) House HO, Umen MJ. J. Org. Chem 1973;38:3893.
3. (a) Gololobov YG, Gruber W. Russ. Chem. Rev 1997;66:953. (b) Fleming FF, Pu Y, Tercek F. J. Org. Chem 1997;62:4883. (c) Kung L-R, Tu C-H, Shia K-S, Liu H-J. Chem. Commun 2003:2490. (d) Pan L-R, Tokoroyama T. Chem. Lett 1990;1999 (e) Wallenfels K, Friedrich K, Reiser J, Ertel W, Thieme HK. Angew. Chem. Int. Ed 1976;15:261.


4. (a) Fleming FF, Gudipati S. *Org. Lett* 2006;8:1557. [PubMed: 16597109] (b) Blanchard P, Da Silva AD, El Kortbi MS, Fourrey J-L, Machado AS, Robert-Gero M. *J. Org. Chem* 1993;58:6517. (c) Blanchard P, Da Silva AD, Fourrey J-L, Machado AS, Robert-Gero M. *Tetrahedron Lett* 1992;33:8069. (d) Blanchard P, El kortbi MS, Fourrey J-L, Robert-Gero M. *Tetrahedron Lett* 1992;33:3319. (e) Sarandeses LA, Mourino A, Luche J-L. *J. Chem. Soc. Chem. Commun* 1992:798. (f) Dupuy C, Petrier C, Sarandeses LA, Luche JL. *Synth. Commun* 1991;21:643. (g) Shono T, Nishiguchi I, Sasaki M. *J. Am. Chem. Soc* 1978;100:4314.
5. (a) Jamison TF, Shambayati S, Crowe WE, Schreiber SL. *J. Am. Chem. Soc* 1997;119:4353. (b) Fleming FF, Hussain Z, Weaver D, Norman RE. *J. Org. Chem* 1997;62:1305. (c) Brattesani DN, Heathcock CH. *J. Org. Chem* 1975;40:2165.
6. (a) Fleming FF, Zhang Z, Wang Q, Steward OW. *Angew. Chem. Int. Ed* 2004;43:1126. (b) Fleming FF, Zhang Z, Wang Q, Steward OW. *J. Org. Chem* 2003;68:7646. [PubMed: 14510537] (c) Fleming FF, Gudipati V, Steward OW. *Tetrahedron* 2003;59:5585.
7. Fleming FF, Iyer PS. *Synthesis* 2006:893.
8. Fleming FF, Zhang Z, Wei G. *Synthesis* 2005:3179.
9. Fleming FF, Zhang Z, Wei G, Steward OW. *J. Org. Chem* 2006;71:1430. [PubMed: 16468790]
10. For an excellent overview of terms, steric constraints, and orbital overlap see: Gawley RE. *Tetrahedron Lett* 1999;40:4297.
11. Bare TH, Hershey ND, House HO, Swain CG. *J. Org. Chem* 1972;37:997.
12. Fleming FF, Gudipati S, Zhang Z, Liu W, Steward OW. *J. Org. Chem* 2005;70:3845. [PubMed: 15876069]
13. Fleming FF, Zhang Z. *Tetrahedron* 2005;61:747.
14. Prepared by oxidation of the corresponding cycloalkenecarbonitrile with CrO₃-3,5-dimethylpyrazole.⁸
15. Repetitive alkylations consistently proceed with the same yield, accompanied by the hydroxyalkenenitrile **12** (OMgMe = OH) resulting from carbonyl addition.
16. Comparable alkylations of **11** imply that magnesiated nitriles prefer coordination on carbon rather than on nitrogen (Scheme 2).¹²
17. Nitriles have an extremely small A-value of only 0.2 kcal mol⁻¹: Eliel Ernest L. *Wilensky's Stereochemistry of Organic Compounds* 1994 Wiley, NY 696697
18. The fragmentation of β-alkoxynitriles is often facile and in this instance the formation of a *dioxide* likely facilitates the fragmentation to an even greater extent: (a) Carlier PR, Lo CW-S, Lo MM-C, Wan NC, Williams ID. *Org. Lett* 2000;2:2443. [PubMed: 10956517] (b) Carlier PR, Lam WW-F, Wan NC, Williams ID. *Angew. Chem., Int. Ed* 1998;37:2252.
19. X-ray crystallography of **14b**, and derivatives of **14a**, **14c**, **14d**, confirmed the stereochemical assignments. The authors have deposited the crystallographic data with the Cambridge Crystallographic Data Center. The data can be obtained, on request, from the Director, Cambridge Crystallographic Data Center, 12 Union Road, Cambridge, CB2 1EZ, UK.
20. Bailey WF, Cioffi A. *Magn. Reson. Chem* 1987;25:181.
21. Considerable amounts of starting oxonitrile **1c**, the hydroxyalkenenitrile **i**, and the hydroxyalkenenitrile **ii** are generated, which result from incomplete conjugate addition and alkylation, respectively.




22. Intercepting the magnesiated nitrile **4** with cyclopropanecarboxaldehyde triggers an analogous cyclization when the reaction is allowed to warm to room temperature.⁹
23. Fleming FF, Wei G, Zhang Z, Steward OW. *Org. Lett* 2006;8:4903. [PubMed: 17020332]
24. Beak P, Zajdel WJ. *J. Am. Chem. Soc* 1984;106:1010.
25. ¹H NMR analysis of the crude reaction mixture failed to identify any enamide.
26. For a preliminary account see: Fleming FF, Zhang Z, Wei G, Steward OW. *Org. Lett* 2005;7:447. [PubMed: 15673261]
27. Inoue A, Kitagawa K, Shinkubo H, Oshima K. *J. Org. Chem* 2001;66:4333. [PubMed: 11397172]
28. Simanek E, Huang NL. *Phys. Rev. Lett* 1966;17:699.
29. Gawley RE, Low E, Zhang Q, Harris R. *J. Am. Chem. Soc* 2000;122:3344. Butenyl alkylation is conceptually possible by prior, catalytic radical atom transfer rearrangement to 3-butenyl iodide or through S_N2' alkylation: Alnajjar MS, Smith GF, Kuivila HG. *J. Org. Chem* 1984;49:1271.
30. Presumably the ejection of bromide is faster than reorganization to a C-magnesiated nitrile followed by loss of bromide.
31. Fleming FF, Wang Q, Zhang Z, Steward OW. *J. Org. Chem* 2002;67:5953. [PubMed: 12182627]




Scheme 1.
Electrophile-Dependent Alkylations of the C-Magnesiated Nitrile **4**




Scheme 2.
Stereoselectivity Differences between *N*- and *C*-Metalated Nitriles




Scheme 3.
Grignard Addition-Alkylation with the 5-Membered Oxonitrile **1a**.

**Scheme 4.**Grignard Addition-Acylations with the 7-Membered Oxonitrile **1c**.






















Scheme 5.
Stereodivergent Alkylations of *N*-Metalated Nitriles



Scheme 6.
1,2-1,4-Grignard Addition to Bromooxonitrile **21**

Table 1
Grignard Addition-Alkylations to Cyclic Oxonitriles

				
entry	oxonitrile	electrophile	cyclic nitrile	yield ^a (ratio) ^b
1	 1a	Mel	 14a	49% ^a (3.4:1)
2	 1a		 14b	55% ^a
3	 1a		 14c	51% ^a
4	 1a	 <i>t</i> -Bu	 <i>t</i> -Bu 14d	47% ^a
5	 1a		 14e	49% ^c
6	 1a		 14f	61% ^d (1.8:1)



entry	oxonitrile	electrophile	cyclic nitrile	yield ^a (ratio) ^b
7		Mel		86% ^e
8				61% ^e
9				71% ^e (1.7:1)
10		Mel		50% ^f
11				24%
12				28%

^a Stereochemistry assigned by x-ray crystallography.¹⁹

^b Diastereomeric ratio at the nitrile-bearing carbon with the major isomer shown.


















^c The stereochemical assignment is made by analogy to the acylation with *t*-BuCOCl, entry 4.

^d The stereochemistry of the major isomer is unknown and is made by analogy.

^eIncluded for comparison from reference 9.

^fStereochemical assignment is based on the characteristic ^{13}C shift of the nitrile carbon.²⁰

Table 2
Stereodivergent Oxonitrile Addition-Alkylations

				
entry	R ¹ M	electrophile	cyclic nitrile ^a	yield (ratio) ^b
1	BuLi	MeI	 5d	65% (12:1)
2	BuLi	Br-CH ₂ -CH=CH ₂	 5e	60% (3:1)
3	BuLi	I-CH ₂ -CH ₂ -CH ₂ -CH ₃	 5f	69% (1.4:1)
4	LiF	I-CH ₂ -CH ₂ -CH ₂ -CH ₃	 5f	61% (0:1)
5	BuOLi	I-CH ₂ -CH ₂ -CH ₂ -CH ₃	 5f	52% (1:1)
6	LiO-CH ₂ -CH ₂ -OLi	I-CH ₂ -CH ₂ -CH ₂ -CH ₃	 5f	50% (1.6:1)
7		I-CH ₂ -CH ₂ -CH ₂ -CH ₃	 5f	73% (6:1)
8		Br-CH ₂ -CH=CH ₂	 5e	79% (7.9:1)
9		MeI	 5d	72% (3.3:1)
10		I-CH ₂ -CH ₂ -CH=CH ₂	 5g	69% (5.2:1)
11		MeI	 5h	62% ^c (4:1)

entry	R^1M	electrophile	cyclic nitrile ^a	yield (ratio) ^b
12			 + 5i + 5g	58% (5i : 5g , 1.8:1) ^d

^a Stereochemical assignments are based on x-ray crystallography.

^b Diastereomeric ratio at the nitrile-bearing carbon with the major isomer shown.

^c MeMgCl was employed for the carbonyl addition (1.1 equiv) and CH₂CHMgBr for the conjugate addition.

^d The major diastereomer **5i** predominates by 4.9:1 for the configuration shown and **5g** predominates over the diastereomer by 2.6:1).