



MIT Open Access Articles

Water Overcomes Methyl Group Directing Effects in Epoxide-Opening Cascades

The MIT Faculty has made this article openly available. **Please share** how this access benefits you. Your story matters.

Citation	Morten, Christopher J., and Timothy F. Jamison. "Water Overcomes Methyl Group Directing Effects in Epoxide-Opening Cascades." <i>Journal of the American Chemical Society</i> 131, no. 19 (May 20, 2009): 6678-6679.
As Published	http://dx.doi.org/10.1021/ja9025243
Publisher	American Chemical Society (ACS)
Version	Author's final manuscript
Accessed	Sun Jan 24 00:05:21 EST 2016
Citable Link	http://hdl.handle.net/1721.1/82116
Terms of Use	Article is made available in accordance with the publisher's policy and may be subject to US copyright law. Please refer to the publisher's site for terms of use.
Detailed Terms	

Published in final edited form as:


J Am Chem Soc. 2009 May 20; 131(19): 6678–6679. doi:10.1021/ja9025243.

Water Overcomes Methyl Group Directing Effects in Epoxide-Opening Cascades

Christopher J. Morten and Timothy F. Jamison*

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

Abstract



R ¹	R ²	acid	base	H ₂ O
H	H	~ 1:1	exo	endo
Me	H	exo	endo	endo
H	Me	endo	exo	endo

Water is an effective promoter of the *endo*-selective opening of trisubstituted epoxides, enabling related cascades leading to a variety of substituted ladder polyether structures. When used in conjunction with a tetrahydropyran-templated nucleophile, water can overcome the powerful electronic directing effect of a methyl substituent at either site of the epoxide, making water a uniquely versatile medium and promoter for epoxide opening.

In 1985 Nakanishi advanced a concise and appealing proposal for the biosynthesis of the ladder polyether family of natural products, a synthesis that culminates in a cascade of regio- and stereoselective epoxide openings (Scheme 1).¹ Our group recently reported an emulation of the Nakanishi hypothesis, wherein water serves as the superior promoter of *endo*-selective epoxide-opening cascades.² This earlier account was limited to cascades of *trans*-disubstituted epoxides, and we herein report that water (as solvent) is also a simple and general solution for cascades involving trisubstituted epoxides. Overcome by this method is the well documented, strong directing effect that methyl (Me) groups have on epoxide ring-opening reactions,³ and thus enabled is the rapid assembly of multiple patterns of substituted ladder polyether subunits.

tfj@mit.edu.

Supporting Information Available: Experimental procedures and data for all new compounds (PDF). This material is available free of charge via the Internet at <http://pubs.acs.org>.

An angular Me group is the only substituent other than hydrogen (H) observed at ladder polyether ring junctions, and every structure in this large family of natural products possesses at least one Me group. Nature has conceived two variations of this substitution, requiring two chemically quite different kinds of (*E*)-trisubstituted epoxides in the corresponding polyepoxide precursors. For example, in the hypothesized precursor (**1**) to brevetoxin B (Scheme 1), Me groups are observed both *distal* and *proximal*⁴ to the internal nucleophile; the putative cascade must tolerate both possibilities. In fact, nearly all ladders bearing more than one Me group, including the brevetoxins, maitotoxin, gambierol, and gymnocin B, are proposed to arise from similar polyepoxides bearing an “out-of-register” mixture of both distally and proximally substituted epoxides.

The crux of the problem is that the Me group is generally a strong director of epoxide-opening regioselectivity, particularly under acid catalysis. Valuable methods for *endo*-selective opening, epitomized by those developed by the McDonald group,^{3a–c} take advantage of this directing effect,³ but these necessarily accommodate only *distal* Me substitution. Moreover, distal Me substitution at every epoxide is generally vital for high regioselectivity and yield.⁵ The *endo*-selective opening of epoxides with a Me or other simple alkyl group⁶ *proximal* to the pendent nucleophile has not been documented except under enzyme catalysis⁷ or when a stronger directing group at the distal site of the epoxide was used.⁸ We conjectured that tetrahydropyran (THP)-templated, water-promoted cascades might prove relatively insensitive to the electronic effects and afford a general solution to the problem of Me substitution, particularly for the challenging case of proximal Me substitution noted above.

We accordingly began our investigation with proximally Me-substituted monoepoxide **2b** (Table 1). Unsurprisingly, both Brønsted (CSA) and Lewis (BF₃) acids were highly *exo* selective, affording the undesired **4b**. Conversely, Brønsted base activation by Cs₂CO₃ provided desired bis-THP product **3b**, with moderate *endo* selectivity apparently arising from the alkoxide's preference for the less sterically hindered site of attack. Most striking was water, which effected cyclization with nearly 5:1 selectivity for **3b** over **4b**. Furthermore, the selectivity of cyclizations of **2b** improved to almost 6:1 *endo:exo* in potassium phosphate buffer within a pH range of 8 to 10 (Chart 1); intriguingly, this selectivity drops again as pH increases past 10.

Epoxides with distal Me substituents have been shown to open with high *endo* regioselectivity with a variety of acidic promoters.^{3a–e} Indeed, exposure of epoxy alcohol **2c** to both BF₃•OEt₂ and CSA induced selective cyclization to bis-THP **3c**, but considerable amounts (up to 20%) of isomerization of **2c** to an isopropyl ketone side product were also observed under these conditions. A cleaner reaction was achieved in deionized water, which smoothly transformed **2c** to **3c** with >20:1 *endo:exo* selectivity and no trace of the ketone byproduct. A pH screen (Chart 1) revealed that very high selectivity holds under acidic, neutral, and mildly basic conditions; only at pH >9 does selectivity drop below 10:1.

Optimistic that cascades of substituted epoxides should be possible in aqueous media, we prepared diepoxy alcohol **7** (Table 2) bearing a proximal Me group. Stirring **7** in warm water produced the desired tris-THP triad **8** in 32% yield.⁹ To the best of our knowledge, this transformation represents the first *endo*-selective epoxide-opening cascade to accommodate a proximal Me substituent. Cascades with base or acid in organic solvent supplied no trace of **8**.

A distal Me group was incorporated into diepoxide **9**, and water again proved amenable, affording triad **10** in 67% yield. A somewhat lower 54% yield of triad **12** was obtained in the aqueous reaction of diepoxide **11**, in which both epoxides now bear distal Me substituents. In

promoting cascades of **9** and **11**, CSA and BF₃ were competitive with water, with water slightly better than BF₃ in reactions of **9** and the reverse observed in reactions of **11**.

Reactions of **5**,² the parent system containing two *trans*-disubstituted epoxides, revealed that only water provides a significant quantity (74%) of the desired triad **6**. Thus, water-promoted cyclizations provide a uniquely versatile strategy for the construction of *all three* epoxide substitution patterns found in the Nakanishi hypothesis and clearly proceed by a mechanism fundamentally different from those operating under simple acid or simple base catalysis.

Supplementary Material


Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

C.J.M. thanks the George Büchi Graduate Fellowship for support. We thank Ivan Vilotijevic for cyclization data on **2a**, Dr. Jeffery A. Byers for preparation of **2a**, Dr. Peter Müller (MIT) for X-ray crystallography, and Li Li (MIT) for mass spectrometry.

References

1. Nakanishi K. *Toxicon* 1985;23:473. [PubMed: 3895583]
2. Vilotijevic I, Jamison TF. *Science* 2007;317:1189. [PubMed: 17761875]
3. (a) Bravo F, McDonald FE, Neiwert WA, Do B, Hardcastle KI. *Org Lett* 2003;5:2123. [PubMed: 12790544] (b) Bravo F, McDonald FE, Neiwert WA, Hardcastle KI. *Org Lett* 2004;6:4487. [PubMed: 15548057] (c) Valentine JC, McDonald FE, Neiwert WA, Hardcastle KI. *J. Am. Chem. Soc* 2005;127:4586. [PubMed: 15796519] (d) Wan S, Gunaydin H, Houk KN, Floreancig PE. *J. Am. Chem. Soc* 2007;129:7915. [PubMed: 17547399] (e) Morimoto Y, Nishikawa Y, Ueba C, Tanaka T. *Angew. Chem. Int. Ed* 2006;45:810. (f) Xiong Z, Corey EJ. *J. Am. Chem. Soc* 2000;122:9328.
4. **Distal** and **proximal** Me groups are shown in **red** and **blue**, respectively.
5. In cascades to form polyoxepanes, *trans*-disubstituted epoxides can be accommodated to some extent, generally with lower yields; see references ^{4c} and ^{4d}.
6. *Endo*-selective cyclization with methoxymethyl substitution on the epoxide proximal to the pendent nucleophile was reported by Murai and coworkers; see: (a) Fujiwara K, Tokiwano T, Murai A. *Tetrahedron Lett* 1995;36:8063. (b) Fujiwara K, Mishima H, Amano A, Tokiwano T, Murai A. *Tetrahedron Lett* 1998;39:393.
7. Smith L, Hong H, Spencer JB, Leadlay PF. *ChemBioChem* 2008;9:2967. [PubMed: 19025863]
8. Matsukura H, Morimoto M, Koshino H, Nakata T. *Tetrahedron Lett* 1997;38:5545.
9. A 6,5-fused side product arising from *exo* opening was also collected, in 39% yield; see Supporting Information.



Scheme 1.
Proposed Biosynthetic Cascade to Brevetoxin B

regioselectivity (3 : 4)





Chart 1.
Dependence of Regioselectivity on pH

Table 1

Dependence of Regioselectivity under Various Epoxide-Opening Conditions



epoxide	R^1	R^2	conditions and regioselectivity (3:4) ^a			
			Cs_2CO_3 , MeOH ^b	CSA, CH_2Cl_2 ^c	$BF_3 \cdot OEt_2$, CH_2Cl_2 ^d	H_2O ^e
2a	H	H	1 : 2.7	1 : 1.2	1.4 : 1	10 : 1
2b	Me	H	3.0 : 1	1 : 5.2	1 : 11	4.9 : 1
2c	H	Me	1 : 17	5.8 : 1 ^f	>20 : 1 ^f	>20 : 1

^a ratios determined by 1H NMR spectroscopy.



^b 30 equiv Cs_2CO_3 , rt, 0.02 M.




^c 1 equiv (+/-)-CSA, rt, 0.02 M.

^d 0.25 equiv $BF_3 \cdot OEt_2$, -78° to rt, 0.02 M.

^e deionized water, rt, 0.02 M.

^f Isopropyl ketone side product also isolated, see Supporting Information.

desired product	conditions and isolated yield ^a of desired product:				
	Cs ₂ CO ₃ in MeOH ^b	CSA in CH ₂ Cl ₂ ^c	BF ₃ •OEt ₂ CH ₂ Cl ₂ ^d	H ₂ O ^e	
	0%	trace ^f	trace ^f		74%
	0%	0%	0%		32%

conditions and isolated yield ^a of desired product:					
Cs ₂ CO ₃ in MeOH ^b	CSA in CH ₂ Cl ₂ ^c	BF ₃ •OEt ₂ CH ₂ Cl ₂ ^d	H ₂ O ^e	desired product	
0%	46%	63%	67%	 <p>10</p>	
0%	43%	61%	54%	 <p>11</p>	
				 <p>12</p>	

^a Corrected for diastereomeric purity of starting material (between 7.5:1 and 20:1 for all cases, see Supporting Information); average of at least two experiments.

^b 30 equiv Cs₂CO₃, rt, 0.02 M.

^c 1 equiv (+/-)-CSA, rt, 0.02 M.

^d 0.25 equiv BF₃•OEt₂, -78° to rt, 0.02 M.

^e 60°, 0.02 M.

^f <5% (¹H NMR).