



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Aromatic Heterocycles as Activating Groups for Asymmetric Conjugate Addition Reactions. Enantioselective Copper-Catalyzed Reduction of 2-Alkenylheteroarenes

Citation for published version:

Rupnicki, L, Saxena, A & Lam, HW 2009, 'Aromatic Heterocycles as Activating Groups for Asymmetric Conjugate Addition Reactions. Enantioselective Copper-Catalyzed Reduction of 2-Alkenylheteroarenes' Journal of the American Chemical Society, vol 131, no. 30, pp. 10386–10387., 10.1021/ja904365h

Digital Object Identifier (DOI):

[10.1021/ja904365h](https://doi.org/10.1021/ja904365h)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Journal of the American Chemical Society

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Aromatic Heterocycles as Activating Groups for Asymmetric Conjugate Addition Reactions. Enantioselective Copper-Catalyzed Reduction of 2-Alkenylheteroarenes


Leszek Rupnicki, Aakarsh Saxena, and Hon Wai Lam*

School of Chemistry, University of Edinburgh, Joseph Black Building, The King's Buildings, West Mains Road, Edinburgh EH9 3JJ, United Kingdom.


*h.lam@ed.ac.uk

The 1,4-addition of a nucleophile to an alkene conjugated to an electron-withdrawing group is a fundamental reaction in organic chemistry, and numerous catalytic asymmetric variants (eq 1) of this process are now routinely employed in the synthesis of molecules of interest.^{1,2} The most common functional groups used to activate alkenes toward asymmetric conjugate additions include carbonyls, nitriles, sulfones, phosphonates, and nitro groups. We recently questioned whether other rarely considered yet common functional groups might also be employed in this capacity, and our attention focused on nitrogen-containing aromatic heterocycles. Given that heteroarenes such as oxazoles, thiazoles, pyridines, and others are ubiquitous in biologically active natural products, pharmaceuticals, and agrochemicals, the ability to functionalize these privileged structures through a diverse set of asymmetric conjugate additions of 2-alkenyl derivatives (eq 2) would open up broad-ranging applications.

Catalytic asymmetric conjugate additions:



Aromatic heterocycles as substrates?




Although conjugate additions to 2-vinylheteroarenes ($R^1, R^2 = H$ in eq 2) are relatively common,³ the corresponding reactions of substrates containing a β -substituent are much rarer, presumably for steric reasons.^{4,5} Furthermore, the only report of a catalytic enantioselective variant is limited to poorly selective ($\leq 15\%$ ee) Grignard additions to 4-alkenylpyridines.⁵ Therefore, we recently initiated a program targeted at addressing these deficiencies, and in this Communication, our preliminary findings involving highly enantioselective copper-catalyzed reductions⁶ of β, β' -disubstituted 2-alkenylheteroarenes are presented.

The asymmetric copper-catalyzed conjugate reduction of activated alkenes is a well-established method for the synthesis of various useful chiral building blocks.^{6,7,8,9,10} Whether a nitrogen-containing heteroarene would provide sufficient activation to an adjacent alkene in an analogous reaction was however, uncertain. In addition, it seemed likely that coordination of the Lewis basic nitrogen of the heteroarene to the copper catalyst would occur in such a process, and whether this interaction would be beneficial, inconsequential, or detrimental, was not easy to predict.

Our investigations began with a survey of chiral bisphosphines **L1**–**L6** using 2-alkenylbenzoxazole **1a** as a test substrate (Table 1). Using 10 mol % of Cu(OAc)₂·H₂O, 10 mol % of ligand and 4 equivalents each of PhSiH₃ and *t*-BuOH in toluene at room temperature, biaryl-based ligands **L1**–**L4** proved competent in

Table 1. Ligand Optimization for the Asymmetric Reduction of **1a**^a














^a Reactions were conducted using 0.20 mmol of **1a** in toluene (1 mL). Conversions were determined by GC analysis. Enantioselectivities were determined by chiral HPLC analysis. ^b Reactions complete after 2 h.

promoting conjugate reduction.¹¹ With (*R*)-BINAP (**L1**), both conversion and enantioselectivity were only moderate. However, improved results were observed using (*R*)-MeO-BIPHEP (**L2**) and the SEGPHOS ligands **L3** and **L4**, with 91% ee obtained using (*S*)-SEGPHOS (**L3**). The Josiphos ligands **L5** and **L6** were also effective, providing **2a** in 89% and 87% ee respectively. Of all the ligands, the highest reaction rates were observed with **L4** and **L5** (reactions were complete in 2 h). However, the superior selectivity provided by **L5** prompted us to select this ligand for further optimization and investigation of reaction scope.

Using 5 mol % each of Cu(OAc)₂·H₂O and **L5**, PhSiH₃ (1.5 equiv), and *t*-BuOH (2.0 equiv) at an initial temperature of 0 °C, a range of β, β' -disubstituted 2-alkenylheteroarenes underwent conjugate reduction with generally excellent levels of enantioselection (Table 2).¹¹ In addition to benzoxazole (entries 1–4), other effective nitrogen-containing heteroarenes in this process included 5-phenyloxazole (entry 5), benzothiazole (entry 6), pyridine (entries 7–9), quinoline (entry 10), and pyrazine (entry 11). Tolerated functionality at the β -positions of the alkene included simple aliphatic groups, a phenyl group (entry 4), a benzyl group (entry 9), various oxygenated alkyl groups (entries 2, 3, 7–9, and 11), and a cyclopropane (entry 10). The process is tolerant of lower loadings of copper and ligand. For example, reduction of **1g** on a 1.0 mmol scale using 2 mol % each of Cu(OAc)₂·H₂O and **L5** provided **2g** in 92% yield and 96% ee (entry 7, values in parentheses).


Experiments to explore the origins of reactivity were then

Table 2. Scope of Cu-Catalyzed Asymmetric Conjugate Reduction^a

$ \begin{array}{c} \text{L5 (5 mol \%)} \\ \text{Cu(OAc)}_2 \cdot \text{H}_2\text{O (5 mol \%)} \\ \text{PhSiH}_3 \text{ (1.5 equiv)} \\ \text{t-BuOH (2.0 equiv)} \\ \text{toluene, 0 } ^\circ\text{C to rt, 17 h} \end{array} $				
entry	product	yield (%) ^b	ee (%) ^c	
1		2a	90	93
2		2b	67	94
3		2c	88	95
4 ^d		2d	95	87
5		2e	93	90
6		2f	86	95
7		2g	90 (92) ^e	>99 (96) ^e
8 ^f		2h	81	98
9		2i	90	97
10		2j	90	96
11		2k	89	96

^a Reactions were conducted using 0.20 mmol of **1a–1k** unless otherwise stated. ^b Isolated yield. ^c Determined by chiral HPLC analysis. ^d Using 0.155 mmol of **1d**. ^e Values in parentheses refers to reaction conducted using 1.0 mmol of **1g**, 2 mol % Cu, and 2 mol % **L5**. ^f Using 0.10 mmol of **1h** and 2.0 equiv of PhSiH₃.

conducted. Reduction of 4-alkenylpyridine **3** provided **4** in 60% yield and 94% ee, albeit in a slower reaction that was incomplete even after 4 days (eq 3). This result suggests that alkene reduction by copper hydride can occur without assistance of a directing effect from the nitrogen atom. In contrast, 3-alkenylpyridine **5** was unreactive (eq 4), demonstrating the importance of conjugation of the alkene to a C=N moiety for reactivity.



In summary, copper-catalyzed asymmetric conjugate reductions of β,β' -disubstituted 2-alkenylheteroarenes are reported. In addition to serving as a further demonstration of the power of chiral copper hydride catalysis,⁶ this work has shown nitrogen-containing aromatic heterocycles can provide effective activation of an adjacent alkene for highly enantioselective catalytic conjugate addition reactions. Extension of the general concept to other classes of asymmetric reactions should provide a range of useful tools for chemists working with heteroarenes.

Future studies from our laboratory will be directed towards this goal.


Acknowledgment. This work was supported by the European Commission (Project No. MEST-CT-2005-020744). Drs. Izuru Nagasaki (Takasago), Rudolf Schmid (Hoffmann-La Roche), and Matthias Lotz (Solvias AG) are gratefully acknowledged for supplying the SEGPHOS, MeO-BIPHEP, and Josiphos ligands, respectively, used in this study. We thank the EPSRC National Mass Spectrometry Service Centre at the University of Wales, Swansea, for providing high resolution mass spectra. Dr. Fraser J. White is thanked for assistance with X-ray crystallography.

Supporting Information Available: Experimental procedures, full spectroscopic data for new compounds, and crystallographic data in cif format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) For recent reviews focused on metal-catalyzed enantioselective conjugate addition reactions, see: (a) Thaler, T.; Knochel, P. *Angew. Chem., Int. Ed.* **2009**, *48*, 645–648. (b) Christoffers, J.; Korpelly, G.; Rosiak, A.; Rössle, M. *Synthesis* **2007**, 1279–1300. (c) Lopez, F.; Minnaard, A. J.; Feringa, B. L. *Acc. Chem. Res.* **2007**, *40*, 179–188. (d) Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, *103*, 2829–2844. (e) Alexakis, A.; Benhaim, C. *Eur. J. Org. Chem.* **2002**, 3221–3236. (f) Krause, N.; Hoffmann-Röder, A. *Synthesis* **2001**, 171–196.
- (2) For reviews describing organocatalytic conjugate addition reactions, see: (a) Vicario, J. L.; Badia, D.; Carrillo, L. *Synthesis* **2007**, 2065–2092. (b) Dondoni, A.; Massi, A. *Angew. Chem., Int. Ed.* **2008**, *47*, 4638–4660.
- (3) For representative examples, see: (a) Magnus, G.; Levine, R. *J. Am. Chem. Soc.* **1956**, *78*, 4127–4130. (b) Nelson, J. H.; Howells, P. N.; DeLullo, G. C.; Landen, G. L. *J. Org. Chem.* **1980**, *45*, 1246–1249. (c) Yoneda, A.; Newkome, G. R. *J. Organomet. Chem.* **1991**, *401*, 217–226. (d) Ciufolini, M. A.; Roschinger, F. *J. Am. Chem. Soc.* **1996**, *118*, 12082–12089. (e) Liu, F.; Dalhus, B.; Gundersen, L.-L.; Rise, F. *Acta Chem. Scand.* **1999**, *53*, 269–279. (f) Lautens, M.; Roy, A.; Fukuoka, K.; Fagnou, K.; Martin-Matute, B. *J. Am. Chem. Soc.* **2001**, *123*, 5358–5359. (g) Boy, K. M.; Guernon, J. M. *Tetrahedron Lett.* **2005**, *46*, 2251–2252. (h) Schaaf, G. M.; Mukherjee, S.; Waterson, A. G. *Tetrahedron Lett.* **2009**, *50*, 1928–1933.
- (4) (a) Hoffman, A.; Farlow, M. W.; Fuson, R. C. *J. Am. Chem. Soc.* **1933**, *55*, 2000–2004. (b) Gilman, H.; Gainer, G. C. *J. Am. Chem. Soc.* **1949**, *71*, 2327–2328. (c) Mathis, C.; Hogen-Esch, T. E. *J. Am. Chem. Soc.* **1982**, *104*, 634–635. (d) Dryanska, V.; Ivanov, C. *Synthesis* **1983**, 143–145. (e) Dondoni, A.; Fantin, G.; Fogagnolo, M.; Medici, A.; Pedrini, P. *Tetrahedron* **1988**, *44*, 2021–2031. (f) Epifani, E.; Florio, S.; Troisi, L. *Tetrahedron* **1990**, *46*, 4031–4038. (g) Cliffe, I. A.; Ifill, A. D.; Mansell, H. L.; Todd, R. S.; White, A. C. *Tetrahedron Lett.* **1991**, *32*, 6789–6792.
- (5) For Ni-catalyzed addition of organometallics to 4-alkenylpyridines, see: Houpius, I. N.; Lee, J.; Dorziotis, I.; Molina, A.; Reamer, B.; Volante, R. P.; Reider, P. J. *Tetrahedron* **1998**, *54*, 1185–1195.
- (6) For recent reviews of CuH-catalyzed reactions, see: (a) Deutsch, C.; Krause, N.; Lipshutz, B. H. *Chem. Rev.* **2008**, *108*, 2916–2926. (b) Rendler, S.; Oestreich, M. *Angew. Chem., Int. Ed.* **2007**, *46*, 498–504.
- (7) Selected examples of asymmetric CuH-catalyzed reduction of α,β -unsaturated carbonyls: (a) Appella, D. H.; Moritani, Y.; Shintani, R.; Ferreira, E. M.; Buchwald, S. L. *J. Am. Chem. Soc.* **1999**, *121*, 9473–9474. (b) Moritani, Y.; Appella, D. H.; Jurkauskas, V.; Buchwald, S. L. *J. Am. Chem. Soc.* **2000**, *122*, 6797–6798. (c) Jurkauskas, V.; Buchwald, S. L. *J. Am. Chem. Soc.* **2002**, *124*, 2892–2893. (d) Hughes, G.; Kimura, M.; Buchwald, S. L. *J. Am. Chem. Soc.* **2003**, *125*, 11253–11258. (e) Lipshutz, B. H.; Servesko, J. M. *Angew. Chem., Int. Ed.* **2003**, *42*, 4789–4792. (f) Lipshutz, B. H.; Servesko, J. M.; Taft, B. R. *J. Am. Chem. Soc.* **2004**, *126*, 8352–8353. (g) Rainka, M. P.; Aye, Y.; Buchwald, S. L. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 5821–5823. (h) Lipshutz, B. H.; Servesko, J. M.; Peterson, T. B.; Papa, P. P.; Lover, A. A. *Org. Lett.* **2004**, *6*, 1273–1275. (i) Rainka, M. P.; Milne, J. E.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2005**, *44*, 6177–6180. (j) Movassaghi, M.; E. Ondrus, A. *Org. Lett.* **2005**, *7*, 4423–4426. (k) Lipshutz, B. H.; Tanaka, N.; Taft, B. R.; Lee, C.-T. *Org. Lett.* **2006**, *8*, 1963–1966. (l) Deng, J.; Hu, X.-P.; Huang, J.-D.; Yu, S.-B.; Wang, D.-Y.; Duan, Z.-C.; Zheng, Z. *J. Org. Chem.* **2008**, *73*, 6022–6024.
- (8) Selected examples of asymmetric CuH-catalyzed reduction of conjugated nitroalkenes: (a) Czekelius, C.; Carreira, E. M. *Angew. Chem., Int. Ed.* **2003**, *42*, 4793–4795. (b) Czekelius, C.; Carreira, E. M. *Org. Lett.* **2004**, *6*, 4575–4577.
- (9) Asymmetric CuH-catalyzed reduction of α,β -unsaturated nitriles: (a) Ren, Y.; Xu, X.; Sun, K.; Xu, J. *Tetrahedron: Asymmetry* **2005**, *16*, 4010–4014. (b) Lee, D.; Kim, D.; Yun, J. *Angew. Chem., Int. Ed.* **2006**, *45*, 2785–2787. (c) Lee, D.; Yang, Y.; Yun, J. *Org. Lett.* **2007**, *9*, 2749–2751. (d) Yoo, K.; Kim, H.; Yun, J. *J. Org. Chem.* **2009**, *74*, 4232–4235.
- (10) Asymmetric CuH-catalyzed reduction of α,β -unsaturated sulfones: (a) Llamas, T.; Gómez Arrayás, R. Carretero, J. C. *Angew. Chem., Int. Ed.* **2007**, *46*, 3329–3332. (b) Desrosiers, J.; Charette, A. B. *Angew. Chem., Int. Ed.* **2007**, *46*, 5955–5957.

- (11) The stereochemistries of the products obtained herein were assigned tentatively by analogy with that of **2c**, which was secured through X-ray crystallography of a derivative. See Supporting Information for details.



The versatility of chiral copper hydride catalysis has been demonstrated through development of highly enantioselective 1,4-reductions of 2-alkenylheteroarenes, substrates that have been rarely considered for asymmetric conjugate addition reactions. Both azoles and azines serve as efficient activating groups for this process.
