

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/11606879>

Efficient and versatile synthesis of novel 2alpha-substituted 1alpha,25-dihydroxyvitamin D(3) analogues and their docking to vitamin D receptors.

ARTICLE in THE JOURNAL OF ORGANIC CHEMISTRY · JANUARY 2002

Impact Factor: 4.72 · Source: PubMed

CITATIONS

28

READS

20

9 AUTHORS, INCLUDING:



Yoshitomo Suhara

Shibaura Institute of Technology

81 PUBLICATIONS 2,206 CITATIONS

SEE PROFILE



Atsushi Kittaka

Teikyo University

164 PUBLICATIONS 1,937 CITATIONS

SEE PROFILE



Katsuhiro Konno

University of Toyama

181 PUBLICATIONS 3,320 CITATIONS

SEE PROFILE



Naoki Miyata

Nagoya City University

268 PUBLICATIONS 5,416 CITATIONS

SEE PROFILE

Efficient and Versatile Synthesis of Novel 2 α -Substituted 1 α ,25-Dihydroxyvitamin D₃ Analogues and Their Docking to Vitamin D Receptors

Yoshitomo Suhara,[†] Ken-ichi Nihei,[†] Masaaki Kurihara,[‡] Atsushi Kittaka,[†] Kentaro Yamaguchi,[§] Toshie Fujishima,[†] Katsuhiro Konno,[†] Naoki Miyata,[‡] and Hiroaki Takayama^{*,†}

Faculty of Pharmaceutical Sciences, Teikyo University, Sagamiko, Kanagawa 199-0195, Japan, National Institute of Health Sciences, Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan, and Chemical Analytical Center, Chiba University, Inage-ku, Chiba 263-8522, Japan

hi-takay@pharm.teikyo-u.ac.jp

Received April 11, 2001

Novel 2 α -substituted 1 α ,25-dihydroxyvitamin D₃ analogues with 2 α -alkyl and 2 α -hydroxyalkyl groups were systematically synthesized from D-xylose. Their conformation on binding to the ligand binding domain (LBD) of the vitamin D receptor was analyzed. It has been found that the 2 α -hydroxypropyl group best fits the cavity of the LBD, and the binding activity is three times higher than that for the natural hormone.

Introduction

1 α ,25-Dihydroxyvitamin D₃ (1 α ,25(OH)₂D₃) is the active form of vitamin D in the hormonal system regulating calcium homeostasis as well as cell differentiation, cell proliferation, and immunology and acts by binding to a specific receptor in the target organs, bone, intestine, and kidney.^{1,2} Given the biological responses to this class of compounds, 1 α ,25(OH)₂D₃ might be a potential drug for the treatment of tumors, especially leukemias, breast and prostate cancers, or immunological disorders.^{1,2} However, 1 α ,25(OH)₂D₃ can induce hypercalcaemia.¹ Therefore, the search for a noncalcemic therapeutic agent, and for convenient methods of synthesizing finely modified compounds, has been greatly stimulated by medical needs. Most approaches to date have involved modification of the side chain of the CD-ring moiety, while nonsteroidal vitamin D mimetics have been synthesized quite recently.³

To produce biologically active analogues, we have applied the A-ring modification of **1** using efficient synthetic procedures and investigated conformation–activity relationships. Recently, we reported the synthesis of A-ring diastereomers of 2-methyl-1,25-dihydroxyvitamin D₃ and found that the 2 α -methyl isomer (**2**) was more potent than the native hormone in terms of vitamin D receptor (VDR) binding affinity, elevation of rat serum Ca concentration, and induction of HL-60 cell differentiation.⁴ In addition, the combination of this 2 α -methyl substitution with 20-epimerization, i.e., a double modification, produced a much more potent analogue.⁵ Con-




Figure 1. Structures of 1 α ,25-dihydroxyvitamin D₃ (1 α ,25(OH)₂D₃, **1**) and its 2 α -substituted analogues **2–9**.

sequently, we decided to elucidate the A-ring conformation–activity relationship and the significance of the 2 α -substitution. This time, we focused on the 2 α -substitution of 1 α ,25(OH)₂D₃ and synthesized several analogues into which was introduced a 2 α -alkyl or a 2 α -hydroxyalkyl group into the A-ring. The resulting analogues showed significant biological activities: in particular, introduction of the 2 α -(3-hydroxypropyl) group caused a 3-fold increase in binding activity to VDR and a ca. 500-fold increase in the potency of calcium-mobilizing activity.⁶ Furthermore, we recently reported the synthesis and VDR binding affinity of 2 α -(ω -hydroxyalkoxy)-1 α ,25-(OH)₂D₃,⁷ in which the A-ring epimer of ED-71 was included.⁸ Most of the biological actions of 1 α ,25(OH)₂D₃ are considered to be mediated by the VDR, which belongs to the nuclear receptor superfamily acting as a ligand-

[†] Teikyo University.

[‡] National Institute of Health Sciences.

[§] Chiba University.

(1) (a) *Vitamin D*; Feldman, D., Glorieux, F. H., Pike, J. W., Eds.; Academic Press: New York, 1997. (b) Ettinger, R. A.; DeLuca, H. F. *Adv. Drug Res.* **1996**, *28*, 269.

(2) Bouillon, R.; Okamura, W. H.; Norman, A. W. *Endocr. Rev.* **1995**, *16*, 200.


(3) Boehm, M. F.; Fitzgerald, P.; Zou, A.; Elgort, M. G.; Bischoff, E. D.; Mere, L.; Mais, D. E.; Bissonnette, R. P.; Heyman, R. A.; Nadzan, A. M.; Reichman, M.; Allegretto, E. A. *Chem. Biol.* **1999**, *6*, 265.

(4) (a) Konno, K.; Maki, S.; Fujishima, T.; Liu, Z.-P.; Miura, D.; Chokki, M.; Takayama, H. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 151. (b) Konno, K.; Fujishima, T.; Maki, S.; Liu, Z.-P.; Miura, D.; Chokki, M.; Ishizuka, S.; Yamaguchi, K.; Kan, Y.; Kurihara, M.; Miyata, N.; Smith, C.; DeLuca, F. H.; Takayama, H. *J. Med. Chem.* **2000**, *43*, 4247.

(5) Fujishima, T.; Liu, Z.-P.; Miura, D.; Chokki, M.; Ishizuka, S.; Konno, K.; Takayama, H. *Bioorg. Med. Chem. Lett.* **1998**, *8*, 2145.

(6) Suhara, Y.; Nihei, K.; Tanigawa, H.; Fujishima, T.; Konno, K.; Nakagawa, K.; Okano, T.; Takayama, H. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 1129.

(7) Kittaka, A.; Suhara, Y.; Takayanagi, H.; Fujishima, T.; Kurihara, M.; Takayama, H. *Org. Lett.* **2000**, *2*, 2619.

Scheme 5^a

^a Key: (a) TmCl, DMAP, CH₂Cl₂. (b) NaCN, DMSO. (c) DIBAL-H, CH₂Cl₂. (d) NaBH₄, MeOH. (e) TBSCl, imidazole, DMF. (f) LAH, Et₂O.

ring was opened by a Wittig reaction to make the enyne structure. Elongation of the 2 α -substituent of the A-ring synthon was carried out in a conventional manner by single-carbon elongation. On the other hand, the natural CD-ring portion (**11**) was prepared from vitamin D₃ by a reported method.¹¹ Finally, the 2 α -substituted 1 α ,25-(OH)₂D₃ analogues were synthesized using palladium-catalyzed coupling of the A-ring enyne synthon with the CD-ring portion.¹¹

The pathway to synthesize the A-ring enyne is described in Schemes 2–5. The synthesis of the A-ring portion was started with a commercially available D-xylose derivative (**12**) as shown in Scheme 2. D-Xylose is readily available and useful for stereoselective synthesis corresponding to the A-ring stereochemistry. In the case of 2-methyl analogues, all possible A-ring diastereomers were synthesized; however, we decided to synthesize the 1 α ,2 α ,3 β -isomer stereoselectively. In this way, D-xylose was the most convenient for the synthesis of the desired 2 α -substituted analogues. First, the C-5 primary alcohol of **12** was selectively protected with a pivaloyl group in 86% yield. Then, the C-3 secondary hydroxyl group of **13** was converted to ketone **14** with PCC oxidation in 98% yield. The conventional Wittig reaction afforded an exomethylene compound **13** in a high yield. In this step, we tried several bases, for example, NaH-DMSO, NaOt-Bu, KOt-Bu, and others, because the chemical yield of **15** depends on the base. We found NaHMDS or KHMDS to be the best choice for this reaction. To remove the 1,2-isopropylidene group while simultaneously forming benzyl glycoside, compound **15** was treated with 4.0 M HCl in dioxane/BnOH to afford alcohols **16a** and **16b**. The desired α -isomer **16a** was separated by silica gel column chromatography in 58% yield along with the β -isomer **16b**. For the synthesis of the same configuration corresponding to the 1 α -hydroxyl group of the native hormone 1 α ,25-(OH)₂D₃, the C-2 hydroxyl group of **16a** was inverted via the Mitsunobu reaction to give the 2 β -secondary alcohol **17** in 84% yield. Under these reaction conditions, however, the β -isomer **16b** was not converted to the desired 2 β -hydroxyl compound at all. Presumably, *p*-nitrobenzoate from **16b** was so strained that elimination of the substituent occurred under basic conditions





Figure 2. X-ray crystallographic analysis of compound **19c**.

for the hydrolysis. After protection of the hydroxyl group of **17** with the methoxymethyl (MOM) group, the exomethylene moiety was stereoselectively converted to the 3 β -hydroxymethyl compound **19a** in a high yield by hydroboration with 9-BBN. Our first trial using the hydroboran-THF complex gave a poor stereoselectivity (3 β -isomer **19a**:3 α -isomer **19b** = 1:1) in this step; therefore, a bulky organoboron such as 9-BBN was essential. To determine the 3 β -stereochemistry of **19a**, the ¹H NMR data of the isomer was not enough. Because the coupling constants *J*_{2,3} and *J*_{3,4} were not so different between the 3 α -isomer (**19a**) and the 3 β -isomer (**19b**), we tried to crystallize a derivative from compound **19a** for X-ray crystal analysis. After several examinations, we found that methoxyphenyl ether **19c**, which was derived from **19a** with replacement of the protective groups, could be clearly crystallized from a hexane solution (Scheme 3). Then, we confirmed unambiguously the stereochemistry of **19a** as having the 3 β -hydroxymethyl group on the basis of X-ray crystal analysis of **19c** as shown in Figure 2.¹² Protection of the primary hydroxyl group of **19a** with a *tert*-butyldimethylsilyl (TBS) group afforded the TBS ether **10** in 99% yield. Thus, modifications of the C-2 and

(12) Crystal data of **19c** are as follows: space group *P*2₁ (monoclinic), *Z* = 2, *a* = 8.7115(9) Å, *b* = 22.447(1) Å, *c* = 5.508(2) Å, β = 90.03(1)°, *V* = 1077.1(3) Å³, *D*_c = 1.235 g/cm³.

Scheme 6^a

^a Key: (a) (dba)₃Pd₂·CHCl₃, TPP, TEA/toluene. (b) CSA, MeOH.

C-3 hydroxyl groups of D-xylose, corresponding to the C-1 hydroxyl group and the C-2 alkyl or the C-2 hydroxyalkyl group of vitamin D analogues, respectively, were accomplished.

As shown in Scheme 4, the desired key A-ring synthon, enyne **29**, was obtained from the D-xylose derivative **10**. Hydrogenolysis of benzyl glycoside **10** with a palladium catalyst, followed by ring opening by means of the Wittig reaction, afforded **20** in good yield. In this reaction, LiHMDS gave a better yield than the other bases NaHMDS, KHMDS, KO^tBu, and *n*-BuLi. The olefin **21** was converted to epoxide **22** by sequential treatments with DIBAL-H, 2,4,6-trimethylbenzenesulfonyl chloride (TmCl), and LiHMDS in a high yield. The acetylene unit was introduced by the reaction of **22** with (trimethylsilyl)-acetylene/*n*-BuLi·BF₃·OEt₂ in THF to give the alcohol **23** in 80% yield. Although some groups recently reported the introduction of an acetylene unit into epoxides using only lithium acetylide in THF, our desired compound could not be obtained without a Lewis acid.¹³ In the next step, we changed the MOM protecting group of the hydroxyl group to TBS. The target vitamin D analogues cannot be exposed to a strong acid for deprotection on account of the triene system. The primary hydroxyl group should be distinguished from two secondary hydroxyls for the subsequent modification to synthesize a 2 α -elongated alkyl or hydroxyalkyl group. Removal of both silyl protecting groups (TMS and TBS) by tetrabutylammonium fluoride (TBAF) to afford **24** in 99% yield and subsequent manipulation of the protecting groups by selective pivaloylation, removal of the MOM group, and silylation afforded the fully protected **27** in a good yield. Finally, further manipulation through **28** gave the desired A-ring synthon **29** in a high yield.

Elongation of the hydroxymethyl group of **28** was carried out in a conventional manner as shown in Scheme 5. Sulfonylation and cyanide substitution gave **30** in 82% yield, and reduction of **30** with DIBAL-H followed by further reduction of aldehyde **31** with NaBH₄ furnished the single-carbon-elongated alcohol **32** in a good overall yield. The hydroxyl group was protected to give the second A-ring synthon **33**. The alcohol **32** was further converted to the ethyl derivative **34** through the corresponding tosylate. In the same manner, the double- and triple-carbon-elongated synthons **38**, **39**, **43**, and **44** were prepared from **32** and **37**, respectively. This stepwise

Table 1. Relative Binding Affinity of the 2 α -Alkyl and 2 α -Hydroxyalkyl Series for Bovine Thymus 1 α ,25(OH)₂D₃ Receptor (VDR)

compounds	binding affinity to VDR ^a
1 α ,25(OH) ₂ D ₃	100
3 : hydroxymethyl	20
4 : hydroxyethyl	70
5 : hydroxypropyl	300
6 : hydroxybutyl	120
7 : ethyl	40
8 : propyl	20
9 : butyl	8

^a Potency of 1 α ,25(OH)₂D₃ is normalized to 100.

synthesis has some advantages; for example, deuterium or tritium can be introduced into the molecule for investigating the mechanism of intermolecular interaction between vitamin D₃ analogues and VDR. As a result, we demonstrated the importance of the ω -hydroxyls of the 2 α -substituents for the affinity to VDR.

Finally, palladium-catalyzed coupling of the A-ring synthons **29**, **33**, **38**, **43**, **34**, **39**, and **44** with the CD-ring portion **11**, followed by deprotection with camphorsulfonic acid (CSA) in MeOH gave the target 2 α -alkyl and 2 α -hydroxyalkyl analogues **3–9** as shown in Scheme 6.⁶ Since the deprotection of the 1 α -hydroxyl group was relatively difficult in the case of 2 α -alkyl derivatives, the total yield was not satisfactory. Thus, we have synthesized five novel 2 α -substituted 1 α ,25-dihydroxyvitamin D₃ analogues.

The results of biological evaluations of **3–9** have already been reported in comparison with those of 1 α -25-dihydroxyvitamin D₃ (**1**) and 2 α -methyl-1 α ,25-dihydroxyvitamin D₃ (**2**).⁶ In addition, we newly investigated the VDR binding affinity of 2 α -hydroxybutyl-1 α ,25-dihydroxyvitamin D₃ (**6**) and 2 α -butyl-1 α ,25-dihydroxyvitamin D₃ (**9**) as shown in Table 1.¹⁴

In this study, we found that 2 α -hydroxypropyl analogue (**5**) exhibited the most potent VDR binding affinity in this series. Because of this result, we investigated the three-dimensional structure of 2 α -hydroxypropyl analogue **5** docking at the VDR ligand binding domain. Recently, Moras et al. reported the crystal structure of the natural ligand–VDR complex.¹⁵ Then, we evaluated

(13) Moriety, R. M.; Kim, J.; Brumer, H., III. *Tetrahedron Lett.* **1995**, 36, 51–54.

(14) Imae, Y.; Manaka, A.; Yoshida, N.; Ishimi, Y.; Shinki, T.; Abe, E.; Suda, T.; Konno, K.; Takayama, H.; Yamada, S. *Biochim. Biophys. Acta* **1994**, 1213, 302–308.

(15) Rochel, N.; Wultz, J. M.; Mitschler, A.; Klaholz, B.; Moras, D. *Mol. Cell* **2000**, 5, 173.




Figure 3. (a) Crystal structure of VDR bound to $1\alpha,25(\text{OH})_2\text{D}_3$ **1** by D. Moras et al.¹⁵ (b) Computer modeling of **5** in the VDR ligand binding domain.

the binding of **5** to VDR using Moras' X-ray results, and the preliminary modeling is shown below (Figure 3).¹⁶ The space around the A-ring moiety seems to be mostly filled by the newly introduced 2α -hydroxypropyl group in the binding cavity. The 1α -hydroxyl group retains hydrogen bonds to Ser-237 and Arg-274 as originally formed in the VDR complex with ligand **1**.¹⁵ It is important to note that an additional hydrogen bond to stabilize the complex was observed between the C-2 α terminal hydroxyl group of **5** and Arg-274. In addition, the alkyl chain of the hydroxypropyl group of **5** would contribute to the hydrophobic interaction with the cavity surrounded by Phe-150, Tyr-143, Tyr-147, and Tyr-236. These findings would be the main reasons for the high binding affinity to VDR.

Conclusion

In summary, we have developed an efficient and systematic route for synthesizing new biologically active 2α -substituted analogues of $1\alpha,25(\text{OH})_2\text{D}_3$ (**3–9**) from D-xylose. This method has the advantage that it should be applicable to a variety of 2α -substituted vitamin D analogues. The activity profiles of the synthesized analogues are highly structure sensitive, with even a single-carbon chain difference greatly altering the bioactivities. Consequently, we believe that these analogues will be important for studies on the action mechanism of vitamin D₃ and also as lead compounds for developing therapeutic agents. Further studies are needed to elucidate fully the activity profiles and modes of action of these analogues.

(16) Calculated by molecular dynamics and molecular mechanics energy minimization using MacroModel version 6.5 (Schrodinger, Inc.) on a SGI O2 Workstation.

Experimental Section

General. ¹H NMR spectra were recorded at 400 MHz and ¹³C NMR spectra at 100 MHz using CDCl₃ as a solvent unless otherwise specified. Chemical shifts are given in parts per million (δ) using tetramethylsilane (TMS) as the internal standard. High-resolution mass values were obtained with a high-resolution mass spectrometer at the Faculty of Pharmaceutical Sciences, Teikyo University. FT-IR spectra were recorded using NaCl plates. Ultraviolet spectra were recorded using ethanol as a solvent. Optical rotations were measured at 25 ± 2 °C. Column chromatography was carried out on silica gel 60 (70–230 mesh), and preparative TLC was run on silica gel 60F₂₅₄. Unless otherwise noted, all reagents were purchased from commercial suppliers and used as received.

1,2-O-Isopropylidene-5-O-pivaloyl- α -D-xylofuranoside (13). To a cold (0 °C) and stirred solution of 1,2-O-isopropylidene- α -D-xylofuranose (**12**, 15.0 g, 78.9 mmol) in pyridine (70 mL) was added dropwise over a period of 2 h trimethylacetyl chloride (9.9 g, 82.2 mmol). The resultant solution was stirred at 0 °C for 10 h. The reaction was quenched with MeOH (5 mL), and the mixture was concentrated in vacuo. The crude product was dissolved in Et₂O (500 mL) and washed successively with water (100 mL), saturated aqueous CuSO₄ solution (100 mL), water (100 mL), saturated aqueous NaHCO₃ solution (100 mL), and brine (100 mL). The aqueous layers were extracted with Et₂O (3 \times 100 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (20% EtOAc in hexanes) gave 18.7 g (86%) of pivaloyl ester **13** as a colorless oil: $[\alpha]_D^{25} +16.61$ (c 1.62, CHCl₃); IR (neat) 3480, 2980, 1732, 1375, 1285, 1217, 1165, 1113, 1075, 1017 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.22 (s, 9 H), 1.32 (s, 3 H), 1.51 (s, 3 H), 3.74 (bs, 1 H), 4.10 (d, 1 H, $J = 2.8$ Hz), 4.17 (dd, 1 H, $J = 5.6, 11.2$ Hz), 4.25 (ddd, 1 H, $J = 2.8, 5.6, 7.2$ Hz), 4.50 (dd, 1 H, $J = 7.2, 11.2$ Hz), 4.56 (d, 1 H, $J = 3.6$ Hz), 5.93 (d, 1 H, $J = 3.6$ Hz); ¹³C NMR (100 MHz, CDCl₃) δ 26.2, 26.8, 27.1, 38.9, 60.5, 74.2, 78.4, 84.9, 104.6, 111.8, 179.8; HREIMS calcd for C₁₂H₁₉O₆ (M⁺ - CH₃) 259.1182, found 259.1182.

1,2-O-Isopropylidene-3-keto-5-O-pivaloyl- α -D-xylofuranoside (14). To a stirred solution of pivaloyl ester **13** (10.5 g, 38.3 mmol) in CH₂Cl₂ (800 mL) at room temperature were added freshly activated powdered molecular sieves (4 Å, 75 g) and pyridinium chlorochromate (36.5 g, 169.4 mmol). The resultant brown suspension was stirred for 3 h after which time TLC indicated the disappearance of the starting material. The reaction mixture was diluted with hexanes (800 mL) and filtrated through a pad of silica gel. Filtration and concentration followed by flash chromatography on silica gel (40% EtOAc in hexanes) gave 10.2 g (98%) of ketone **14** as a colorless oil: $[\alpha]_D^{25} +42.60$ (c 1.31, CHCl₃); IR (neat) 2979, 1777, 1734, 1377, 1285, 1221, 1159, 1092, 1005 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.18 (s, 9 H), 1.44 (s, 3 H), 1.48 (s, 3 H), 4.23 (dd, 1 H, $J = 3.2, 11.6$ Hz), 4.37 (dd, 1 H, $J = 1.2, 4.4$ Hz), 4.39 (dd, 1 H, $J = 3.2, 11.6$ Hz), 4.57 (dt, 1 H, $J = 1.2, 3.2$ Hz), 6.10 (d, 1 H, $J = 4.4$ Hz); ¹³C NMR (100 MHz, CDCl₃) δ 27.2, 27.6, 38.7, 63.2, 76.2, 77.2, 103.0, 114.3, 177.4, 207.6; HREIMS calcd for C₁₃H₂₀O₆ (M⁺) 276.1260, found 276.1262.

3-Deoxy-1,2-O-isopropylidene-3-C-methylene-5-O-pivaloyl- α -D-xylo-pentofuranoside (15). Sodium bis(trimethylsilyl)amide (18.0 mL, 1.0 M solution in THF, 18.0 mmol) was slowly added at room temperature to a suspension of methyltriphenylphosphonium bromide (7.7 g, 21.6 mmol) in THF (100 mL). The resultant bright yellow solution was stirred at room temperature for 1.5 h. The solution was then cooled to -78 °C, and a THF (20 mL) solution of ketone **14** (4.3 g, 15.8 mmol) was slowly added. After being stirred for 30 min at -78 °C, the reaction mixture was allowed to warm to room temperature and stirred for another 1 h. The reaction was quenched at 0 °C with MeOH (10 mL), and the mixture was poured into Et₂O (200 mL) and washed with saturated aqueous NH₄Cl solution (3 \times 100 mL) and brine (3 \times 100 mL). The aqueous layers were extracted with Et₂O (3 \times 100 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica

gel (10% EtOAc in hexanes) gave 3.6 g (85%) of olefin **15** as a colorless oil: $[\alpha]_D^{25} +115.79$ (*c* 1.52, CHCl₃); IR (neat) 2997, 1732, 1374, 1282, 1235, 1159, 1071, 1015, 920 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.20 (s, 9 H), 1.39 (s, 3 H), 1.52 (s, 3 H), 4.19 (dd, 1 H, *J* = 4.8, 12.0 Hz), 4.23 (dd, 1H, *J* = 3.6, 12.0 Hz), 4.91 (dd, 1 H, *J* = 1.2, 4.0 Hz), 4.94 (ddd, 1 H, *J* = 2.4, 3.6, 4.8 Hz), 5.22 (t, 1 H, *J* = 1.2 Hz), 5.48 (dd, 1 H, *J* = 1.2, 2.4 Hz), 5.87 (d, 1 H, *J* = 4.0 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 30.3, 30.6, 41.9, 68.3, 80.8, 84.8, 107.8, 115.6, 115.8, 149.1, 181.3; HREIMS calcd for C₁₃H₁₉O₅ (M⁺ - CH₃) 255.1235, found 255.1232.

Benzyl 3-Deoxy-3-C-methylene-5-O-pivaloyl- α -D-xylo-pentofuranoside (16a) and Benzyl 3-Deoxy-3-C-methylene-5-O-pivaloyl- β -D-xylo-pentofuranoside (16b). A cold (0 °C) and stirred mixture of olefin **15** (3.2 g, 11.9 mmol) and benzyl alcohol (8.0 g, 74.1 mmol) in toluene (22 mL) was treated with 4.0 M solution of HCl in dioxane (10 mL, 40.0 mmol). The reaction mixture was warmed to room temperature, and stirring was continued for 16 h. The mixture was poured into Et₂O (200 mL), and the reaction was quenched at 0 °C with saturated aqueous NaHCO₃ solution (100 mL). The mixture was then washed with water (3 \times 50 mL) and brine (3 \times 50 mL). The aqueous layers were extracted with Et₂O (3 \times 50 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (20% EtOAc in hexanes) gave 2.20 g (58%) of alcohol **16a** and 1.22 g (32%) of alcohol **16b** each as a colorless oil. **16a**: $[\alpha]_D^{25} +169.58$ (*c* 1.42, CHCl₃); IR (neat) 3486, 2975, 1732, 1285, 1159, 1090, 1017, 907, 739, 698 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.20 (s, 9 H), 2.29 (d, 1H, *J* = 11.6 Hz), 4.18 (dd, 1 H, *J* = 4.8, 12.0 Hz), 4.22 (dd, 1H, *J* = 3.6, 12.0 Hz), 4.53 (ddt, 1 H, *J* = 2.4, 4.4, 11.6 Hz), 4.60 (d, 1 H, *J* = 12.0 Hz), 4.73 (ddt, 1 H, *J* = 2.4, 3.6, 4.8 Hz), 4.81 (d, 1 H, *J* = 12.0 Hz), 5.15 (d, 1 H, *J* = 4.4 Hz), 5.18 (t, 1 H, *J* = 2.4 Hz), 5.39 (t, 1 H, *J* = 2.4 Hz), 7.34 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 27.2, 38.8, 65.8, 69.2, 73.5, 76.7, 100.0, 108.5, 127.9, 128.0, 128.5, 137.4, 147.6, 178.2; HREIMS calcd for C₁₈H₂₄O₅ (M⁺) 320.1624, found 320.1620. **16b**: $[\alpha]_D^{25} -34.80$ (*c* 1.23, CHCl₃); IR (neat) 3447, 2975, 1730, 1285, 1159, 1105, 1057, 992, 916, 739, 700 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.27 (s, 9 H), 2.61 (bs, 1H), 4.14 (dd, 1 H, *J* = 6.8, 11.2 Hz), 4.19 (dd, 1H, *J* = 4.8, 11.2 Hz), 4.42 (t, 1 H, *J* = 1.2 Hz), 4.49 (d, 1 H, *J* = 11.6 Hz), 4.76 (d, 1 H, *J* = 11.6 Hz), 4.89 (ddt, 1 H, *J* = 1.2, 4.8, 6.2 Hz), 5.03 (s, 1 H), 5.26 (t, 1 H, *J* = 1.2 Hz), 5.49 (t, 1 H, *J* = 1.2 Hz), 7.33 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 27.2, 38.8, 67.7, 69.1, 77.6, 78.1, 106.9, 112.4, 127.0, 127.9, 128.2, 128.3, 128.5, 137.1, 148.7, 178.3; HREIMS calcd for C₁₈H₂₄O₅ (M⁺) 320.1624, found 320.1621.

Benzyl 3-Deoxy-3-C-methylene-5-O-pivaloyl- α -D-lyxo-pentofuranoside (17). A mixture of alcohol **16a** (2.0 g, 6.25 mmol), *p*-nitrobenzoic acid (2.1 g, 12.6 mmol), and triphenylphosphine (3.3 g, 12.6 mmol) in THF (40 mL) was cooled to 0 °C and treated with a 40% solution of azodicarboxylic acid diethyl ester in toluene (5.8 g, 13.9 mmol). After being stirred for 15 min at 0 °C, the resultant solution was allowed to warm to room temperature and stirred for another 3 h. The reaction was quenched with MeOH (10 mL) at 0 °C, and the mixture was poured into Et₂O (100 mL) and washed with saturated aqueous NaHCO₃ solution (3 \times 50 mL) and brine (3 \times 50 mL). The aqueous layer was extracted with Et₂O (3 \times 50 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (10% EtOAc in hexanes) gave crude *p*-nitrobenzoate as a white solid, which was used in the next step without further purification.

To a cold (0 °C) and stirred solution of crude *p*-nitrobenzoate in MeOH (100 mL) was added dropwise 1.0 M aqueous NaOH solution (1.0 mL, 1.00 mmol), and the resultant solution was stirred for 1 h. After 1.0 M aqueous HCl solution (1.5 mL, 1.5 mmol) was added, the reaction mixture was concentrated in vacuo. The crude product was dissolved in Et₂O (200 mL) and washed with saturated aqueous NaHCO₃ solution (3 \times 50 mL) and brine (3 \times 50 mL). The aqueous layer was extracted with Et₂O (3 \times 50 mL), and the combined organic layer was dried over MgSO₄. Filtration and concentration followed by flash

chromatography on silica gel (20% EtOAc in hexanes) gave 1.67 g (84%) of alcohol **17** as a colorless oil: $[\alpha]_D^{25} +97.30$ (*c* 1.15, CHCl₃); IR (neat) 3467, 2975, 1730, 1285, 1161, 1102, 1053, 995, 914, 739, 700 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.22 (s, 9 H), 2.09 (d, 1H, *J* = 7.6 Hz), 4.25 (dd, 1 H, *J* = 4.8, 12.4 Hz), 4.33 (dd, 1H, *J* = 3.2, 12.4 Hz), 4.37 (d, 1H, *J* = 7.6 Hz), 4.55 (d, 1 H, *J* = 11.6 Hz), 4.74 (d, 1 H, *J* = 11.6 Hz), 4.75 (ddd, 1 H, *J* = 1.6, 3.2, 4.8 Hz), 5.10 (s, 1 H), 5.26 (s, 1 H), 5.54 (t, 1 H, *J* = 1.6 Hz), 7.33 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 27.1, 38.8, 65.7, 68.9, 76.8, 76.9, 106.3, 112.6, 127.7, 127.8, 128.0, 128.4, 137.4, 148.0, 178.2; HREIMS calcd for C₁₈H₂₄O₅ (M⁺) 320.1624, found 320.1628.

Benzyl 3-Deoxy-3-C-methylene-2-O-methoxymethyl-5-O-pivaloyl- α -D-lyxo-pentofuranoside (18). To a cold (0 °C) and stirred solution of alcohol **17** (1.0 g, 3.13 mmol) in CH₂Cl₂ (25 mL) were added dropwise diisopropylethylamine (1.24 g, 9.61 mmol) and chloromethyl methyl ether (1.29 g, 16.0 mmol). Solid tetrabutylammonium iodide (360 mg, 975 μ mol) was added to the reaction mixture, and the solution was allowed to warm to room temperature and stirred in the dark for 14 h. The mixture was poured into Et₂O (150 mL) and washed with saturated aqueous NH₄Cl solution (3 \times 25 mL) and brine (3 \times 25 mL). The aqueous layers were extracted with Et₂O (3 \times 50 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (10% EtOAc in hexanes) gave 1.01 g (88%) of MOM ether **18** as a colorless oil: $[\alpha]_D^{25} +90.17$ (*c* 1.15, CHCl₃); IR (neat) 2957, 1732, 1285, 1152, 1103, 1033, 920, 754, 700 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.23 (s, 9 H), 3.35 (s, 3 H), 4.18 (dd, 1 H, *J* = 6.4, 11.2 Hz), 4.22 (dd, 1H, *J* = 5.2, 11.2 Hz), 4.37 (s, 1H), 4.55 (d, 1 H, *J* = 12.0 Hz), 4.60 (d, 1 H, *J* = 6.8 Hz), 4.69 (ddd, 1 H, *J* = 1.6, 5.2, 6.4 Hz), 4.75 (d, 1 H, *J* = 12.0 Hz), 4.76 (d, 1 H, *J* = 6.8 Hz), 5.21 (s, 1 H), 5.36 (s, 1 H), 5.46 (d, 1 H, *J* = 1.6 Hz), 7.33 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 27.2, 38.8, 55.5, 66.5, 68.8, 78.0, 80.2, 94.1, 105.2, 114.0, 127.7, 127.9, 128.4, 137.5, 145.1, 178.3; HREIMS calcd for C₁₃H₂₁O₆ (M⁺ - Bn) 273.1338, found 273.1336.

Benzyl 5-O-Pivaloyl-2-O-methoxymethyl-3-deoxy-3-C-hydroxymethyl- α -D-lyxo-pentofuranose (19a). To a cold (0 °C) and stirred solution of MOM ether **18** (2.0 g, 5.49 mmol) in THF (20 mL) was slowly added a 0.5 M solution of 9-borabicyclo[3.3.1]nonane in THF (20 mL, 10.0 mmol). The resultant solution was warmed to 50 °C, and stirring was continued for 3 h. The reaction mixture was cooled to 0 °C and treated sequentially with a 3.0 M aqueous NaOH solution (6.4 mL, 19.2 mmol) and a 30% aqueous H₂O₂ solution (12.8 mL). The resultant mixture was vigorously stirred at room temperature for 2 h, poured into EtOAc (200 mL), and washed with water (3 \times 50 mL), 5% aqueous Na₂SO₃ solution (3 \times 50 mL), and brine (3 \times 50 mL). The aqueous layers were extracted with EtOAc (3 \times 50 mL), and the combined organic layers were dried over Na₂SO₄. Filtration and concentration followed by flash chromatography on silica gel (30% EtOAc in hexanes) gave 1.78 g (85%) of alcohol **19a** as a colorless oil: $[\alpha]_D^{25} +61.33$ (*c* 1.95, CHCl₃); IR (neat) 3503, 2959, 1730, 1285, 1154, 1113, 1051, 739, 700 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.22 (s, 9H), 2.34 (t, 1H, *J* = 5.6 Hz), 2.89 (tt, 1H, *J* = 5.6, 8.4 Hz), 3.37 (s, 3H), 3.86 (dt, 1H, *J* = 5.6, 11.2 Hz), 3.88 (ddd, 1H, *J* = 5.6, 8.4, 11.2 Hz), 4.19 (dd, 1H, *J* = 7.2, 11.6 Hz), 4.20 (d, 1H, *J* = 5.6 Hz), 4.23 (dd, 1H, *J* = 5.6, 11.6 Hz), 4.38 (ddd, 1H, *J* = 5.6, 7.2, 8.4 Hz), 4.50 (d, 1H, *J* = 12.0 Hz), 4.66 (s, 2H), 4.73 (d, 1H, *J* = 12.0 Hz), 5.15 (s, 1H), and 7.34 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ 27.2, 38.7, 45.0, 55.8, 57.6, 64.9, 68.9, 77.1, 82.4, 97.0, 105.1, 127.6, 127.8, 128.4, 137.7, 178.2; HREIMS calcd for C₁₉H₂₇O₆ (M⁺ - OCH₃) 351.1807, found 351.1810.

Benzyl 5-O-Pivaloyl-2-O-methoxymethyl-3-deoxy-3-C-(tert-butylidimethylsilyloxymethyl)- α -D-lyxo-pentofuranose (10). A solution of alcohol **19a** (2.97 g, 7.77 mmol) in DMF (30 mL) was treated with imidazole (1.06 g, 15.6 mmol) and *tert*-butyldimethylsilyl chloride (1.76 g, 11.7 mmol) at room temperature. The resultant solution was stirred for 3 h, poured into Et₂O (200 mL), and washed with water (3 \times 25 mL) and brine (3 \times 25 mL). The aqueous layers were extracted with Et₂O (3 \times 25 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash

chromatography on silica gel (10% EtOAc in hexanes) gave 3.82 g (99%) of silyl ether **10** as a colorless oil: $[\alpha]_D^{25} +66.61$ (c 1.15, CHCl₃); IR (neat) 2955, 1732, 1283, 1256, 1154, 1096, 1034, 837, 777, 737, 698 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.02 (s, 3H), 0.03 (s, 3H), 0.83 (s, 9H), 1.18 (s, 9H), 2.80 (ddt, 1H, $J = 4.8, 6.8, 8.8$ Hz), 3.28 (s, 3H), 3.70 (dd, 1H, $J = 6.8, 10.0$ Hz), 3.76 (dd, 1H, $J = 8.8, 10.0$ Hz), 4.07 (d, 1H, $J = 4.8$ Hz), 4.09 (dd, 1H, $J = 7.4, 12.0$ Hz), 4.13 (dd, 1H, $J = 4.8, 12.0$ Hz), 4.42 (ddd, 1H, $J = 4.8, 7.4, 8.8$ Hz), 4.46 (d, 1H, $J = 12.0$ Hz), 4.58 (d, 1H, $J = 6.8$ Hz), 4.61 (d, 1H, $J = 6.8$ Hz), 4.69 (d, 1H, $J = 12.0$ Hz), 5.10 (s, 1H), 7.28 (m, 5H); ¹³C NMR (100 MHz, CDCl₃) δ -5.5, 18.1, 25.8, 27.2, 38.7, 45.4, 55.6, 57.4, 65.3, 69.0, 77.1, 81.1, 96.8, 105.3, 127.6, 127.8, 127.8, 137.9, 178.3; HREIMS calcd for C₂₅H₄₁O₆Si (M⁺ - OCH₃) 465.2673, found 465.2666.

(2R,3R,4R)-1-Pivalic Acid 3-(tert-Butyldimethylsilyloxymethyl)-2-hydroxy-4-methoxymethoxyhex-5-enyl Ester (20). A solution of silyl ether **10** (3.98 g, 8.02 mmol) in EtOH (40 mL) was hydrogenated over 20% Pd(OH)₂ on carbon (400 mg) for 12 h. Filtration through Celite and concentration in vacuo gave the desired crude hemiacetal as a colorless oil, which was used in the next step without further purification.

Lithium bis(trimethylsilyl)amide (29 mL, 1.0 M solution in THF, 29.0 mmol) was added dropwise at 0 °C to a suspension of methyltriphenylphosphonium bromide (10.9 g, 30.5 mmol) in THF (50 mL). The resultant bright yellow solution was stirred at room temperature for 40 min and then cooled to 0 °C. After a solution of crude hemiacetal in THF (20 mL) was added dropwise, the reaction mixture was stirred for 40 min at 0 °C. The reaction was quenched at 0 °C with MeOH (5 mL), poured into Et₂O (300 mL), and washed with saturated aqueous NH₄Cl solution (3 × 50 mL) and brine (3 × 50 mL). The aqueous layers were extracted with Et₂O (3 × 50 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (15% EtOAc in hexanes) gave 2.57 g (79%) of olefin **20** as a colorless oil: $[\alpha]_D^{30} -48.96$ (c 1.15, CHCl₃); IR (neat) 3505, 2957, 1732, 1285, 1256, 1157, 1090, 1032, 924, 839, 777 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.05 (s, 3H), 0.07 (s, 3H), 0.89 (s, 9H), 1.21 (s, 9H), 2.74 (ddt, 1H, $J = 3.6, 4.4, 7.2$ Hz), 3.40 (s, 3H), 3.54 (d, 1H, $J = 5.6$ Hz), 3.82 (dd, 1H, $J = 4.4, 10.4$ Hz), 3.90 (dd, 1H, $J = 4.4, 10.4$ Hz), 4.14 (dd, 1H, $J = 4.8, 10.8$ Hz), 4.26 (dd, 1H, $J = 5.6, 10.8$ Hz), 4.28 (ddt, 1H, $J = 3.6, 4.8, 5.6$ Hz), 4.40 (t, 1H, $J = 8.0$ Hz), 4.58 (d, 1H, $J = 6.4$ Hz), 4.72 (d, 1H, $J = 6.4$ Hz), 5.30 (dd, 1H, $J = 1.2, 16.8$ Hz), 5.33 (dd, 1H, $J = 1.2, 10.4$ Hz), 5.70 (ddd, 1H, $J = 8.0, 10.4, 16.8$ Hz); ¹³C NMR (100 MHz, CDCl₃) δ -5.7, -5.6, 18.0, 25.8, 27.2, 38.8, 46.2, 55.9, 60.5, 66.7, 68.9, 76.1, 94.2, 119.1, 136.4, 178.4; HREIMS calcd for C₁₄H₂₉O₄Si (M⁺ - OCH₃) 373.2411, found 373.2421.

(2S,3R,4R)-3-(tert-Butyldimethylsilyloxymethyl)-4-methoxymethoxyhex-5-ene-1,2-diol (21). To a cold (-78 °C) and stirred solution of olefin **20** (3.20 g, 7.92 mmol) in CH₂Cl₂ (50 mL) was added dropwise over a period of 30 min diisobutylaluminum hydride (19.8 mL, 1.0 M solution in toluene, 19.8 mmol). After 10 min, the reaction mixture was treated sequentially with MeOH (1 mL) and saturated aqueous NH₄Cl solution (1 mL), and the resultant suspension was diluted with Et₂O (250 mL) and filtered through a pad of Celite. The filtrate was washed with saturated aqueous NH₄Cl solution (3 × 25 mL) and brine (3 × 25 mL). The aqueous layers were extracted with Et₂O (3 × 25 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (40% EtOAc in hexanes) gave 2.12 g (84%) of diol **21** as a colorless oil: $[\alpha]_D^{25} -60.26$ (c 1.15, CHCl₃); IR (neat) 3443, 2932, 1256, 1156, 1084, 1034, 926, 839, 777 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.05 (s, 3H), 0.06 (s, 3H), 0.89 (s, 9H), 1.77 (ddt, 1H, $J = 3.6, 4.8, 8.0$ Hz), 2.67 (dd, 1H, $J = 4.4, 8.0$ Hz), 3.40 (s, 3H), 3.52 (d, 1H, $J = 6.4$ Hz), 3.67 (ddd, 1H, $J = 4.8, 8.0, 11.2$ Hz), 3.73 (ddd, 1H, $J = 4.4, 6.4, 11.2$ Hz), 3.83 (d, 1H, $J = 4.8$ Hz), 4.14 (ddt, 1H, $J = 3.6, 4.8, 6.4$ Hz), 4.30 (t, 1H, $J = 8.0$ Hz), 4.57 (d, 1H, $J = 6.8$ Hz), 4.71 (d, 1H, $J = 6.8$ Hz), 5.29 (d, 1H, $J = 16.8$ Hz), 5.32 (d, 1H, $J = 10.0$ Hz), 5.69 (ddd, 1H, $J = 8.0, 10.0, 16.8$ Hz); ¹³C NMR (100 MHz, CDCl₃) δ -5.6, 18.1, 25.8,

47.1, 55.9, 61.0, 65.5, 71.1, 76.5, 94.1, 119.3, 136.2; HREIMS calcd for C₁₄H₂₉O₄Si (M⁺ - OCH₃) 289.1835, found 289.1838.

(3R,4S,5R)-4-(tert-Butyldimethylsilyloxymethyl)-3-(methoxymethoxy)-5,6-epoxyhex-1-ene (22). To a solution of diol **21** (3.20 g, 11.6 mmol) in CH₂Cl₂ (40 mL) was added 4-(dimethylamino)pyridine (2.84 g, 23.4 mmol). To the cold (0 °C) and vigorously stirred solution was slowly added 2-mesitylenesulfonyl chloride (3.80 g, 17.4 mmol). After being stirred for 4 h at 0 °C, the reaction mixture was poured into Et₂O (200 mL) and washed with water (3 × 25 mL) and brine (3 × 25 mL). The aqueous layers were extracted with Et₂O (3 × 25 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration gave the desired crude sulfonate as a colorless oil, which was used in the next step without further purification.

The crude sulfonate was dissolved in THF (50 mL), and the solution was cooled to -78 °C and treated with a 1.0 M solution of lithium bis(trimethylsilyl)amide in THF (2.5 mL, 2.50 mmol). After being stirred for 20 min at -78 °C, the reaction mixture was warmed to 0 °C and stirred for another 20 min. The mixture was poured into Et₂O (200 mL) and washed with saturated aqueous NH₄Cl solution (3 × 25 mL) and brine (3 × 25 mL). The aqueous layers were extracted with Et₂O (3 × 25 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (10% EtOAc in hexanes) gave 3.01 g (86%) of epoxide **22** as a colorless oil: $[\alpha]_D^{20} -68.78$ (c 1.15, CHCl₃); IR (neat) 2930, 1259, 1154, 1103, 1034, 918, 837, 777 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.02 (s, 3H), 0.04 (s, 3H), 0.87 (s, 9H), 1.26 (dddd, 1H, $J = 3.2, 4.0, 5.2, 7.2$ Hz), 2.62 (dd, 1H, $J = 2.8, 5.2$ Hz), 2.87 (t, 1H, $J = 5.2$ Hz), 3.04 (ddd, 1H, $J = 2.8, 3.2, 5.2$ Hz), 3.35 (s, 3H), 3.70 (dd, 1H, $J = 4.0, 10.0$ Hz), 3.82 (dd, 1H, $J = 5.2, 10.0$ Hz), 4.31 (t, 1H, $J = 7.2$ Hz), 4.51 (d, 1H, $J = 6.4$ Hz), 4.68 (d, 1H, $J = 6.4$ Hz), 5.24 (d, 1H, $J = 10.0$ Hz), 5.25 (d, 1H, $J = 17.2$ Hz), 5.70 (ddd, 1H, $J = 7.2, 10.0, 17.2$ Hz); ¹³C NMR (100 MHz, CDCl₃) δ -5.6, -5.5, 18.2, 25.9, 48.1, 49.8, 51.0, 55.6, 60.6, 75.8, 94.1, 118.4, 136.6; HREIMS calcd for C₁₄H₂₇O₃Si (M⁺ - OCH₃) 271.1729, found 271.1732.

(4R,5S,6R)-5-(tert-Butyldimethylsilyloxymethyl)-6-(methoxymethoxy)-1-(trimethylsilyl)-oct-7-en-1-yn-4-ol (23). To a cold (0 °C) and stirred solution of ethynyltrimethylsilane (3.88 g, 39.6 mmol) in THF (100 mL) was slowly added a 1.54 M solution of *n*-butyllithium in hexanes (22.7 mL, 36.0 mmol). After being stirred for 15 min at 0 °C, the reaction mixture was cooled to -78 °C and treated with a solution of epoxide **22** (3.01 g, 9.97 mmol) in THF (20 mL). Boron trifluoride diethyl ether complex (1.70 g 12.0 mmol) was added to the reaction mixture, and the solution was allowed to warm to room temperature and stirred for 40 min. The mixture was poured into Et₂O (300 mL) and washed with saturated aqueous NH₄Cl solution (3 × 50 mL) and brine (3 × 50 mL). The aqueous layers were extracted with Et₂O (3 × 50 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (10% EtOAc in hexanes) gave 3.20 g (80%) of enyne **23** as a colorless oil: $[\alpha]_D^{20} -62.26$ (c 1.15, CHCl₃); IR (neat) 3513, 2957, 1251, 1156, 1098, 1030, 926, 841, 777, 762 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.04 (s, 3H), 0.05 (s, 3H), 0.11 (s, 9H), 0.87 (s, 9H), 1.93 (ddt, 1H, $J = 4.0, 5.6, 8.0$ Hz), 2.46 (dd, 1H, $J = 8.0, 16.8$ Hz), 2.63 (dd, 1H, $J = 6.0, 16.8$ Hz), 3.38 (s, 3H), 3.60 (d, 1H, $J = 6.0$ Hz), 3.81 (dd, 1H, $J = 4.0, 10.8$ Hz), 3.92 (dd, 1H, $J = 5.6, 10.8$ Hz), 4.21 (ddt, 1H, $J = 4.0, 6.0, 8.0$ Hz), 4.44 (t, 1H, $J = 8.0$ Hz), 4.58 (d, 1H, $J = 6.4$ Hz), 4.70 (d, 1H, $J = 6.4$ Hz), 5.28 (d, 1H, $J = 17.2$ Hz), 5.29 (d, 1H, $J = 10.8$ Hz), 5.71 (ddd, 1H, $J = 8.0, 10.8, 17.2$ Hz); ¹³C NMR (100 MHz, CDCl₃) δ -5.7, -5.6, 15.8, 16.9, 25.7, 26.6, 46.6, 55.8, 60.1, 69.4, 76.1, 77.1, 86.5, 94.2, 118.6, 136.6; HREIMS calcd for C₁₉H₃₇O₃Si₂ (M⁺ - OCH₃) 369.2282, found 369.2276.

(2S,3S)-2-[(R)-1-(Methoxymethoxy)allyl]hex-5-yn-1,3-diol (24). A cold (0 °C) and stirred solution of enyne **23** (3.20 g, 8.0 mmol) in THF (45 mL) was treated with a 1.0 M solution of tetrabutylammonium fluoride in THF (17.6 mL, 17.6 mmol). The resultant solution was then warmed to room temperature,

and stirring was continued for 1 h. The reaction mixture was poured into EtOAc (200 mL) and washed with saturated aqueous NH₄Cl solution (3 \times 25 mL) and brine (3 \times 25 mL). The aqueous layers were extracted with EtOAc (3 \times 25 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (40% EtOAc in hexanes) gave 1.69 g (99%) of diol **24** as a colorless oil: $[\alpha]_D^{25}$ -96.70 (*c* 1.15, CHCl₃); IR (neat) 3420, 3299, 2857, 1154, 1096, 1030, 924, 637 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.86 (ddt, 1H, *J* = 3.2, 7.2, 4.8 Hz), 2.05 (t, 1H, *J* = 2.8 Hz), 2.50 (ddd, 1H, *J* = 2.8, 6.8, 16.8 Hz), 2.60 (1H, bs), 2.62 (ddd, 1H, *J* = 2.8, 6.8, 16.8 Hz), 3.32 (d, 1H, *J* = 4.4 Hz), 3.42 (s, 3H), 3.83 (dd, 1H, *J* = 4.8, 11.2 Hz), 4.00 (dd, 1H, *J* = 4.8, 11.2 Hz), 4.30 (ddt, 1H, *J* = 3.2, 4.4, 6.8 Hz), 4.44 (tt, 1H, *J* = 1.2, 7.2 Hz), 4.61 (d, 1H, *J* = 6.8 Hz), 4.70 (d, 1H, *J* = 6.8 Hz), 5.33 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.35 (dt, 1H, *J* = 1.2, 17.2 Hz), 5.78 (ddd, 1H, *J* = 7.2, 10.4, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 25.3, 47.2, 56.0, 59.9, 69.5, 70.6, 77.0, 80.8, 94.6, 118.9, 136.1; HREIMS calcd for C₉H₁₃O₃ (M⁺ - CH₂OCH₃) 169.0865, found 169.0875.

1-Pivalic Acid (2S,3S)-3-Hydroxy-2-[(R)-1-(methoxy-methoxy)allyl]hex-5-ynyl Ester (25). The diol **24** (1.62 g, 7.57 mmol) was dissolved in pyridine (1.7 mL) and CH₂Cl₂ (6.8 mL). To the cold (0 °C) and stirred solution was added dropwise over a period of 30 min trimethylacetyl chloride (1.08 g, 8.96 mmol). After being stirred for 1 h at 0 °C, the reaction mixture was allowed to warm to 20 °C and stirred for another 4 h. The mixture was poured into Et₂O (100 mL) and washed with saturated aqueous NH₄Cl solution (3 \times 20 mL) and brine (3 \times 20 mL). The aqueous layers were extracted with Et₂O (3 \times 20 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (20% EtOAc in hexanes) gave 1.92 g (85%) of alcohol **25** as a colorless oil: $[\alpha]_D^{20}$ -87.39 (*c* 1.15, CHCl₃); IR (neat) 3521, 3291, 2975, 1728, 1287, 1156, 1098, 1030, 924, 635 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.22 (s, 9H), 2.03 (t, 1H, *J* = 2.4 Hz), 2.14 (ddt, 1H, *J* = 2.4, 4.8, 5.6 Hz), 2.46 (ddd, 1H, *J* = 2.4, 7.6, 16.8 Hz), 2.58 (ddd, 1H, *J* = 2.4, 6.4, 16.8 Hz), 3.10 (d, 1H, *J* = 2.4 Hz), 3.39 (s, 3H), 4.26 (ddt, 1H, *J* = 2.4, 6.4, 7.6 Hz), 4.29 (dd, 1H, *J* = 5.6, 11.6 Hz), 4.33 (dd, 1H, *J* = 5.6, 11.6 Hz), 4.40 (ddt, 1H, *J* = 1.2, 4.8, 6.8 Hz), 4.57 (d, 1H, *J* = 7.2 Hz), 4.70 (d, 1H, *J* = 7.2 Hz), 5.31 (dt, 1H, *J* = 1.2, 17.6 Hz), 5.34 (d, 1H, *J* = 1.2, 10.4 Hz), 5.71 (ddd, 1H, *J* = 6.8, 10.4, 17.6 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 24.7, 27.2, 38.7, 45.2, 56.1, 61.2, 69.5, 70.4, 77.4, 80.9, 94.3, 118.8, 135.6, 178.4; HREIMS calcd for C₁₅H₂₃O₄ (M⁺ - OCH₃) 267.1596, found 267.1599.

1-Pivalic Acid (2S,3S)-3-Hydroxy-2-[(R)-1-(hydroxy)-allyl]hex-5-ynyl Ester (26). The alcohol **25** (2.10 g, 7.05 mmol) was dissolved in *tert*-BuOH (60 mL). Pyridinium *p*-toluenesulfonate (17.6 g, 70.0 mmol) was added, and the resultant solution was heated at reflux for 12 h. The reaction mixture was cooled to room temperature and concentrated in vacuo. The crude product was dissolved in Et₂O (200 mL) and washed with saturated aqueous NaHCO₃ solution (3 \times 20 mL) and brine (3 \times 20 mL). The aqueous layers were extracted with Et₂O (3 \times 20 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (30% EtOAc in hexanes) gave 1.33 g (74%) of diol **26** as a colorless oil: $[\alpha]_D^{25}$ +15.13 (*c* 1.15, CHCl₃); IR (neat) 3438, 3306, 2977, 1730, 1287, 1163, 1040, 928, 639 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.22 (s, 9H), 2.05 (t, 1H, *J* = 2.4 Hz), 2.06 (dddd, 1H, *J* = 2.4, 4.4, 5.6, 6.8 Hz), 2.43 (ddd, 1H, *J* = 2.4, 7.2, 16.8 Hz), 2.57 (ddd, 1H, *J* = 2.4, 7.2, 16.8 Hz), 2.73 (d, 1H, *J* = 4.4 Hz), 3.07 (d, 1H, *J* = 2.4 Hz), 4.24 (tt, 1H, *J* = 2.4, 7.2 Hz), 4.31 (dd, 1H, *J* = 5.6, 11.6 Hz), 4.43 (dd, 1H, *J* = 6.8, 11.6 Hz), 4.45 (ddt, 1H, *J* = 1.2, 4.4, 5.6 Hz), 5.28 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.40 (dt, 1H, *J* = 1.2, 16.8 Hz), 5.92 (ddd, 1H, *J* = 5.6, 10.4, 16.8 Hz); ¹³C NMR (100 MHz, CDCl₃) δ 25.0, 27.2, 38.8, 45.2, 61.4, 68.9, 70.8, 72.8, 80.7, 116.1, 138.6, 178.9; HREIMS calcd for C₁₀H₁₃O₄ (M⁺ - *t*Bu) 197.0814, found 197.0806.

1-Pivalic Acid (2S,3S)-3-(*tert*-Butyldimethylsilanyloxy)-2-[(R)-1-(*tert*-butyl-dimethylsilanyloxy)allyl]hex-5-ynyl Ester (27). A cold (0 °C) and stirred solution of diol **26** (1.31

g, 5.16 mmol) in CH₂Cl₂ (25 mL) was treated with 2,6-lutidine (2.21 g, 20.6 mmol) and *tert*-butyldimethylsilyl triflate (4.09 g, 15.5 mmol). The resultant solution was stirred at 0 °C for 2 h, poured into Et₂O (150 mL), and washed with saturated aqueous NH₄Cl solution (3 \times 20 mL) and brine (3 \times 20 mL). The aqueous layers were extracted with Et₂O (3 \times 20 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (2.5% EtOAc in hexanes) gave 2.47 g (99%) of silyl ether **27** as a colorless oil: $[\alpha]_D^{25}$ -3.22 (*c* 1.15, CHCl₃); IR (neat) 3316, 2957, 1732, 1285, 1256, 1159, 1089, 924, 837, 776, 639 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.04 (s, 3H), 0.06 (s, 3H), 0.07 (s, 3H), 0.10 (s, 3H), 0.88 (s, 9H), 0.89 (s, 9H), 1.20 (s, 9H), 1.98 (t, 1H, *J* = 2.4 Hz), 2.19 (dq, 1H, *J* = 4.0, 6.0 Hz), 2.45 (ddd, 1H, *J* = 2.4, 4.8, 16.4 Hz), 2.54 (ddd, 1H, *J* = 2.4, 7.2, 16.4 Hz), 3.99 (dd, 1H, *J* = 6.0, 12.0 Hz), 4.16 (ddd, 1H, *J* = 4.0, 4.8, 7.2 Hz), 4.17 (dd, 1H, *J* = 6.0, 7.6 Hz), 4.25 (dd, 1H, *J* = 6.0, 12.0 Hz), 5.09 (d, 1H, *J* = 10.0 Hz), 5.16 (d, 1H, *J* = 17.2 Hz), 5.84 (ddd, 1H, *J* = 7.6, 10.0, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.8, -4.6, -4.1, -3.6, 18.0, 18.2, 25.8, 25.9, 26.5, 27.2, 38.7, 48.5, 61.8, 69.1, 70.5, 74.1, 81.2, 115.7, 139.9, 178.3; HREIMS calcd for C₂₂H₄₁O₄Si₂ (M⁺ - *t*Bu) 425.2543, found 425.2543.

(2S,3S)-3-(*tert*-Butyldimethylsilanyloxy)-2-[(R)-1-(*tert*-butyldimethylsilanyloxy)allyl]hex-5-yn-1-ol (28). To a cold (-78 °C) and stirred solution of silyl ether **27** (2.47 g, 5.12 mmol) in CH₂Cl₂ (20 mL) was added dropwise over a period of 30 min diisobutylaluminum hydride (7.7 mL, 1.0 M solution in toluene, 7.70 mmol). After being stirred for 10 min at -78 °C, the reaction mixture was treated sequentially with MeOH (1 mL) and saturated aqueous NH₄Cl solution (1 mL), and the resultant suspension was diluted with Et₂O (200 mL) and filtered through a pad of Celite. The filtrate was washed with saturated aqueous NH₄Cl solution (3 \times 20 mL) and brine (3 \times 20 mL). The aqueous layers were extracted with Et₂O (3 \times 20 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (10% EtOAc in hexanes) gave 1.96 g (96%) of alcohol **28** as a colorless oil: $[\alpha]_D^{20}$ +0.96 (*c* 1.15, CHCl₃); IR (neat) 3472, 3314, 2932, 1256, 1074, 926, 837, 777, 635 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.06 (s, 3H), 0.10 (s, 3H), 0.11 (s, 3H), 0.13 (s, 3H), 0.90 (s, 18H), 2.01 (t, 1H, *J* = 2.8 Hz), 2.09 (dq, 1H, *J* = 4.8, 7.2 Hz), 2.46 (ddd, 1H, *J* = 2.8, 4.8, 16.8 Hz), 2.51 (ddd, 1H, *J* = 2.8, 6.8, 16.8 Hz), 3.07 (dd, 1H, *J* = 4.8, 7.2 Hz), 3.72 (ddd, 1H, *J* = 4.8, 7.2, 12.4 Hz), 3.82 (dt, 1H, *J* = 7.2, 12.4 Hz), 4.08 (dt, 1H, *J* = 4.8, 6.8 Hz), 4.33 (tt, 1H, *J* = 2.0, 7.2 Hz), 5.20 (dt, 1H, *J* = 2.0, 10.4 Hz), 5.28 (dt, 1H, *J* = 2.0, 17.2 Hz), 5.84 (ddd, 1H, *J* = 7.2, 10.4, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.8, -4.7, -4.4, -3.8, 17.9, 18.1, 25.8, 25.9, 26.8, 50.2, 61.0, 70.7, 70.9, 74.3, 80.5, 116.7, 138.7; HREIMS calcd for C₁₇H₃₃O₃Si₂ (M⁺ - *t*Bu) 341.1968, found 341.1964.

(3R,4S,5R)-3,5-Bis(*tert*-butyldimethylsilanyloxy)-4-(*tert*-butyldimethylsilanyloxymethyl)-oct-1-en-7-yne (29). A solution of alcohol **28** (40.0 mg, 101 μ mol) in DMF (2.0 mL) was treated with imidazole (13.6 mg, 200 μ mol) and *tert*-butyldimethylsilyl chloride (22.6 mg, 150 μ mol) at room temperature. The resultant solution was stirred for 3 h, poured into Et₂O (20 mL), and washed with water (3 \times 5 mL) and brine (3 \times 5 mL). The aqueous layers were extracted with Et₂O (3 \times 5 mL), and the combined organic layers were dried over MgSO₄. Filtration and concentration followed by flash chromatography on silica gel (2.0% toluene in hexanes) gave 46.0 mg (88%) of silyl ether **29** as a colorless oil: $[\alpha]_D^{25}$ +5.13 (*c* 1.15, CHCl₃); IR (neat) 3316, 2932, 1256, 1090, 922, 837, 776, 635 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.03 (s, 6H), 0.04 (s, 3H), 0.06 (s, 6H), 0.10 (s, 3H), 0.89 (s, 9H), 0.90 (s, 18H), 1.93 (t, 1H, *J* = 2.8 Hz), 2.00 (ddt, 1H, *J* = 5.2, 5.6, 6.4 Hz), 2.44 (ddd, 1H, *J* = 2.8, 5.6, 16.8 Hz), 2.57 (ddd, 1H, *J* = 2.8, 5.6, 16.8 Hz), 3.56 (dd, 1H, *J* = 6.4, 10.0 Hz), 3.81 (dd, 1H, *J* = 6.4, 10.0 Hz), 4.09 (dt, 1H, *J* = 5.2, 5.6 Hz), 4.30 (ddt, 1H, *J* = 1.2, 5.6, 6.8 Hz), 5.03 (dt, 1H, *J* = 1.2, 9.6 Hz), 5.11 (dt, 1H, *J* = 1.2, 17.2 Hz), 5.92 (ddd, 1H, *J* = 6.8, 9.6, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -5.4, -4.8, -4.6, -4.2, -4.0, 18.1,

18.2, 25.9, 26.0, 26.3, 51.5, 60.1, 69.2, 69.7, 73.1, 82.4, 114.4, 141.1; HREIMS calcd for $C_{27}H_{56}O_3Si_3$ (M^+) 512.3537, found 512.3533.

(3*S*,4*R*)-4-(*tert*-Butyldimethylsilyloxy)-3-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]hept-6-ynitrile (30**).** To a solution of alcohol **28** (314 mg, 789 μ mol) in CH_2Cl_2 (5.0 mL) was added 4-(dimethylamino)pyridine (372 mg, 3.04 mmol). To the cold (0 °C) and vigorously stirred solution was slowly added 2-mesitylenesulfonyl chloride (582 mg, 2.66 mmol). After being stirred for 12 h at 0 °C, the reaction mixture was poured into Et_2O (100 mL) and washed with water (3 \times 15 mL) and brine (3 \times 15 mL). The aqueous layers were extracted with Et_2O (3 \times 15 mL), and the combined organic layers were dried over $MgSO_4$. Filtration and concentration gave the desired crude sulfonate ester as a colorless oil, which was used in the next step without further purification.

The crude sulfonate ester was dissolved in DMSO (5.0 mL). Sodium cyanide (78.0 mg, 1.59 mmol) was added to the solution, and the resultant solution was heated at 70 °C for 2 h. The reaction mixture was cooled to room temperature, poured into Et_2O (100 mL) and washed with water (3 \times 15 mL) and brine (3 \times 15 mL). The aqueous layers were extracted with Et_2O (3 \times 15 mL), and the combined organic layers were dried over $MgSO_4$. Filtration and concentration followed by flash chromatography on silica gel (5% $EtOAc$ in hexanes) gave 263 mg (82%) of nitrile **30** as a colorless oil: $[\alpha]^{20}_D$ -3.83 (*c* 1.15, $CHCl_3$); IR (neat) 3314, 2932, 2247, 1256, 1090, 936, 836, 777, 639 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 0.06 (s, 3H), 0.09 (s, 3H), 0.12 (s, 3H), 0.13 (s, 3H), 0.90 (s, 9H), 0.91 (s, 9H), 2.03 (t, 1H, *J* = 2.8 Hz), 2.29 (tt, 1H, *J* = 6.0, 6.8 Hz), 2.36 (dd, 1H, *J* = 6.0, 16.4 Hz), 2.46 (ddd, 1H, *J* = 2.8, 4.4, 16.8 Hz), 2.48 (dd, 1H, *J* = 6.0, 16.4 Hz), 2.52 (ddd, 1H, *J* = 2.8, 6.8, 16.8 Hz), 4.04 (dt, 1H, *J* = 4.4, 6.8 Hz), 4.20 (ddt, 1H, *J* = 1.2, 6.4, 6.8 Hz), 5.24 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.31 (dt, 1H, *J* = 1.2, 16.8 Hz), 5.82 (ddd, 1H, *J* = 6.4, 10.4, 16.8 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ -4.8, -4.6, -4.2, -3.9, 13.5, 18.1, 18.2, 25.8, 25.9, 26.5, 46.1, 69.0, 71.3, 74.0, 79.9, 117.8, 120.1, 137.6; HREIMS calcd for $C_{22}H_{41}O_2NSi_2$ (M^+) 407.2676, found 407.2659.

(3*S*,4*R*)-4-(*tert*-Butyldimethylsilyloxy)-3-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]hept-6-ynal (31**).** To a cold (-78 °C) and stirred solution of nitrile **30** (184 mg, 452 μ mol) in CH_2Cl_2 (2.0 mL) was added dropwise over a period of 10 min diisobutylaluminum hydride (530 μ L, 1.0 M solution in toluene, 530 μ mol). After being stirred for 1 h at -78 °C, the reaction mixture was treated sequentially with MeOH (1 mL) and saturated aqueous NH_4Cl solution (1 mL), and the resultant suspension was diluted with Et_2O (100 mL) and filtered through a pad of Celite. The filtrate was washed with saturated aqueous NH_4Cl solution (3 \times 10 mL) and brine (3 \times 10 mL). The aqueous layers were extracted with Et_2O (3 \times 10 mL), and the combined organic layers were dried over $MgSO_4$. Filtration and concentration followed by flash chromatography on silica gel (5% $EtOAc$ in hexanes) gave 157 mg (85%) of aldehyde **31** as a colorless oil: $[\alpha]^{20}_D$ +6.70 (*c* 1.15, $CHCl_3$); IR (neat) 3314, 2932, 2716, 1728, 1256, 1080, 930, 837, 777, 633 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 0.04 (s, 3H), 0.06 (s, 3H), 0.08 (s, 3H), 0.09 (s, 3H), 0.88 (s, 9H), 0.89 (s, 9H), 2.04 (t, 1H, *J* = 2.8 Hz), 2.37 (ddd, 1H, *J* = 2.8, 6.0, 16.8 Hz), 2.40 (ddd, 1H, *J* = 2.8, 6.4, 17.2 Hz), 2.42 (ddd, 1H, *J* = 1.6, 7.2, 17.2 Hz), 2.44 (ddd, 1H, *J* = 2.8, 6.0, 16.8 Hz), 2.68 (ddt, 1H, *J* = 5.2, 6.4, 7.2 Hz), 3.84 (dt, 1H, *J* = 5.2, 6.0 Hz), 4.23 (ddt, 1H, *J* = 1.6, 6.4, 6.8 Hz), 5.18 (dt, 1H, *J* = 1.6, 10.0 Hz), 5.21 (dt, 1H, *J* = 1.6, 16.0 Hz), 5.73 (ddd, 1H, *J* = 6.8, 10.0, 16.8 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ -4.7, -4.6, -4.3, -4.0, 18.0, 18.2, 25.8, 25.9, 26.4, 40.5, 45.4, 70.0, 71.1, 74.3, 80.3, 117.2, 138.1, 202.6; HREIMS calcd for $C_{22}H_{42}O_3Si_2$ (M^+) 410.2673, found 410.2667.

(3*S*,4*R*)-4-(*tert*-Butyldimethylsilyloxy)-3-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]hept-6-yn-1-ol (32**).** A cold (0 °C) and stirred solution of aldehyde **31** (157 mg, 383 μ mol) in MeOH (2.0 mL) was treated with sodium tetrahydroborate (28.0 mg, 741 μ mol). After being stirred for 30 min at 0 °C, the reaction mixture was diluted with Et_2O (100 mL) and

washed with saturated aqueous NH_4Cl (3 \times 10 mL) and brine (3 \times 10 mL). The aqueous layers were extracted with Et_2O (3 \times 10 mL), and the combined organic layers were dried over $MgSO_4$. Filtration and concentration followed by flash chromatography on silica gel (10% $EtOAc$ in hexanes) gave 129 mg (82%) of alcohol **32** as a colorless oil: $[\alpha]^{21}_D$ +12.17 (*c* 1.15, $CHCl_3$); IR (neat) 3420, 3314, 2932, 1256, 1073, 926, 837, 776, 635 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 0.08 (s, 6H), 0.10 (s, 6H), 0.90 (s, 9H), 0.91 (s, 9H), 1.67 (m, 2H), 2.01 (t, 1H, *J* = 2.4 Hz), 2.04 (ddt, 1H, *J* = 5.6, 7.2, 4.8 Hz), 2.41 (ddd, 1H, *J* = 2.4, 5.6, 17.2 Hz), 2.45 (ddd, 1H, *J* = 2.4, 5.6, 17.2 Hz), 2.93 (bs, 1H), 3.65 (m, 2H), 3.85 (q, 1H, *J* = 5.6 Hz), 4.27 (ddt, 1H, *J* = 1.2, 4.8, 6.8 Hz), 5.18 (dt, 1H, *J* = 1.2, 10.8 Hz), 5.21 (dt, 1H, *J* = 1.2, 17.6 Hz), 5.86 (ddd, 1H, *J* = 6.8, 10.8, 17.6 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ -4.8, -4.6, -4.3, -4.1, 18.0, 18.2, 25.8, 26.1, 28.8, 48.2, 62.5, 70.6, 71.6, 75.4, 81.0, 116.6, 138.4; HREIMS calcd for $C_{22}H_{44}O_3Si_2$ (M^+) 412.2829, found 412.2834.

(3*R*,4*S*,5*R*)-3,5-Bis-(*tert*-butyldimethylsilyloxy)-4-[2-(*tert*-butyldimethylsilyloxy)ethyl]oct-1-en-7-yne (33**).** Alcohol **32** (60.0 mg, 146 μ mol) was converted to silyl ether **33** (59.0 mg, 78%) according to the same procedure described above for obtaining compound **29** from **28**. Data for **33**: colorless oil; $[\alpha]^{25}_D$ +7.83 (*c* 1.15, $CHCl_3$); IR (neat) 3316, 2930, 1256, 1082, 924, 835, 776, 627 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 0.03 (s, 3H), 0.04 (s, 6H), 0.05 (s, 3H), 0.07 (s, 3H), 0.09 (s, 3H), 0.89 (s, 27H), 1.55 (m, 2H), 1.84 (ddt, 1H, *J* = 3.6, 5.2, 6.4 Hz), 1.95 (t, 1H, *J* = 2.8 Hz), 2.36 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 2.41 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 3.57 (dt, 1H, *J* = 6.4, 10.0 Hz), 3.69 (dd, 1H, *J* = 6.4, 10.0 Hz), 4.01 (dt, 1H, *J* = 3.6, 6.4 Hz), 4.13 (t, 1H, *J* = 6.4 Hz), 5.11 (d, 1H, *J* = 10.4 Hz), 5.17 (d, 1H, *J* = 17.2 Hz), 5.82 (ddd, 1H, *J* = 6.4, 10.4, 17.2 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ -5.2, -4.6, -4.5, -4.1, -3.7, 18.1, 18.2, 18.4, 25.9, 26.0, 29.1, 46.1, 62.9, 70.0, 71.5, 75.7, 82.0, 121.5, 139.9; HREIMS calcd for $C_{24}H_{49}O_3Si_3$ (M^+ - *t*Bu) 469.2990, found 469.3021.

(3*R*,4*S*,5*R*)-3,5-Bis-(*tert*-butyldimethylsilyloxy)-4-ethyloct-1-en-7-yne (34**).** To a solution of alcohol **32** (100 mg, 243 μ mol) in CH_2Cl_2 (2.0 mL) was added 4-(dimethylamino)pyridine (74.0 mg, 606 μ mol). To the cold (0 °C) and vigorously stirred solution was slowly added 2-mesitylenesulfonyl chloride (106 mg, 485 μ mol). After being stirred for 12 h at 0 °C, the reaction mixture was poured into Et_2O (100 mL) and washed with water (3 \times 10 mL) and brine (3 \times 10 mL). The aqueous layers were extracted with Et_2O (3 \times 10 mL), and the combined organic layers were dried over $MgSO_4$. Filtration and concentration gave the desired crude sulfonate as a colorless oil, which was used in the next step without further purification.

To a cold (0 °C) and stirred solution of crude sulfonate in Et_2O (2.0 mL) was slowly added lithium aluminum hydride (46.0 mg, 1.20 μ mol). After being stirred for 1 h at 0 °C, the reaction mixture was allowed to warm to room temperature and stirred for another 3 h. The mixture was treated with $EtOAc$ (1 mL) at 0 °C and saturated aqueous NH_4Cl solution (1 mL), and the resultant suspension was diluted with Et_2O (100 mL) and filtered through a pad of Celite. The filtrate was washed with saturated aqueous NH_4Cl solution (3 \times 10 mL) and brine (3 \times 10 mL). The aqueous layers were extracted with Et_2O (3 \times 10 mL), and the combined organic layers were dried over $MgSO_4$. Filtration and concentration followed by flash chromatography on silica gel (1% $EtOAc$ in hexanes) gave 77.0 mg (80%) of enyne **34** as a colorless oil: $[\alpha]^{21}_D$ +16.87 (*c* 1.15, $CHCl_3$); IR (neat) 3316, 2957, 1254, 1073, 924, 837, 776, 629 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 0.03 (s, 3H), 0.05 (s, 3H), 0.07 (s, 3H), 0.09 (s, 3H), 0.89 (s, 18H), 0.94 (t, 3H, *J* = 7.6 Hz), 1.36 (m, 2H), 1.68 (ddt, 1H, *J* = 3.6, 5.2, 6.0 Hz), 1.95 (t, 1H, *J* = 2.8 Hz), 2.39 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 2.42 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 4.01 (dt, 1H, *J* = 3.6, 6.4 Hz), 4.14 (ddt, 1H, *J* = 1.2, 5.2, 7.2 Hz), 5.08 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.14 (dt, 1H, *J* = 1.2, 16.8 Hz), 5.86 (ddd, 1H, *J* = 7.2, 10.4, 16.8 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ -4.7, -4.4, -4.2, -3.7, 14.0, 18.1, 19.0, 25.9, 26.1, 51.3, 69.8, 71.5, 75.7, 82.3, 115.2, 140.6; HREIMS calcd for $C_{18}H_{35}O_2Si_2$ (M^+ - *t*Bu) 339.2175, found 339.2194.

(4*S*,5*R*)-5-(*tert*-Butyldimethylsilyloxy)-4-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]oct-7-ynitrile (35). Alcohol **32** (129 mg, 313 μ mol) was converted to nitrile **35** (80.0 mg, 61%) according to the procedure described above for converting **30** from **28**. Data for **35**: colorless oil; $[\alpha]^{20}_D +1.39$ (*c* 1.15, CHCl₃); IR (neat) 3314, 2932, 2247, 1256, 1078, 928, 837, 777, 635 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.05 (s, 3H), 0.08 (s, 6H), 0.10 (s, 3H), 0.90 (s, 18H), 1.73 (m, 2H), 1.96 (q, 1H, *J* = 6.4 Hz), 2.03 (t, 1H, *J* = 2.4 Hz), 2.38 (ddd, 1H, *J* = 2.4, 6.0, 16.8 Hz), 2.43 (ddd, 1H, *J* = 2.4, 4.8, 16.8 Hz), 2.50 (dt, 2H, *J* = 4.0, 8.4 Hz), 3.88 (dt, 1H, *J* = 4.8, 6.0 Hz), 4.18 (dt, 1H, *J* = 1.2, 6.4 Hz), 5.19 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.25 (dt, 1H, *J* = 1.2, 17.2 Hz), 5.80 (ddd, 1H, *J* = 6.4, 10.4, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.7, -4.6, -4.3, -4.2, 17.0, 18.0, 18.2, 22.4, 25.9, 26.0, 26.7, 48.3, 71.0, 71.1, 75.2, 80.5, 116.6, 120.4, 138.2; HREIMS calcd for C₂₃H₄₃O₂NSi₂ (M⁺) 421.2814, found 421.2794.

(4*S*,5*R*)-5-(*tert*-Butyldimethylsilyloxy)-4-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]oct-7-ynal (36). Nitrile **35** (80.0 mg, 190 μ mol) was converted to aldehyde **36** (71.0 mg, 89%) according to the procedure described above for obtaining **31** from **30**. Data for **36**: colorless oil; $[\alpha]^{20}_D +3.22$ (*c* 1.15, CHCl₃); IR (neat) 3314, 2932, 2712, 1728, 1256, 1078, 926, 837, 777, 635 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.04 (s, 3H), 0.06 (s, 3H), 0.07 (s, 3H), 0.10 (s, 3H), 0.89 (s, 18H), 1.69 (m, 2H), 1.87 (ddt, 1H, *J* = 5.2, 5.6, 6.0 Hz), 1.99 (t, 1H, *J* = 2.8 Hz), 2.40 (ddd, 1H, *J* = 2.8, 6.0, 17.2 Hz), 2.43 (ddd, 1H, *J* = 2.8, 6.0, 17.2 Hz), 2.57 (ddd, 1H, *J* = 2.0, 2.8, 6.0 Hz), 2.59 (ddd, 1H, *J* = 2.0, 3.2, 6.0 Hz), 3.94 (dt, 1H, *J* = 4.4, 6.0 Hz), 4.17 (ddt, 1H, *J* = 1.6, 5.6, 6.8 Hz), 5.14 (dt, 1H, *J* = 1.6, 10.4 Hz), 5.21 (dt, 1H, *J* = 1.6, 17.2 Hz), 5.83 (ddd, 1H, *J* = 6.8, 10.4, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.7, -4.5, -4.2, -3.9, 18.2, 18.3, 25.9, 26.4, 30.9, 43.8, 48.4, 70.5, 71.4, 75.6, 81.2, 116.1, 139.2, 203.0; HREIMS calcd for C₂₃H₄₄O₃Si₂ (M⁺) 424.2829, found 424.2829.

(4*S*,5*R*)-5-(*tert*-Butyldimethylsilyloxy)-4-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]oct-7-yn-1-ol (37). Aldehyde **36** (71.0 mg, 167 μ mol) was converted to alcohol **37** (70.0 mg, 98%) according to the procedure described above for obtaining **32** from **31**. Data for **37**: colorless oil; $[\alpha]^{21}_D +10.26$ (*c* 1.15, CHCl₃); IR (neat) 3340, 3314, 2932, 1254, 1067, 924, 837, 776, 635 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.04 (s, 3H), 0.06 (s, 3H), 0.07 (s, 3H), 0.10 (s, 3H), 0.89 (s, 18H), 1.39 (m, 2H), 1.57 (bs, 1H), 1.66 (q, 2H, *J* = 6.4 Hz), 1.85 (ddt, 1H, *J* = 4.8, 5.2, 5.6 Hz), 1.98 (t, 1H, *J* = 2.8 Hz), 2.39 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 2.43 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 3.61 (t, 2H, *J* = 6.4 Hz), 3.97 (dt, 1H, *J* = 4.8, 6.4 Hz), 4.16 (ddt, 1H, *J* = 1.2, 5.6, 7.2 Hz), 5.12 (dt, 1H, *J* = 1.2, 9.6 Hz), 5.17 (dt, 1H, *J* = 1.2, 16.8 Hz), 5.84 (ddd, 1H, *J* = 7.2, 9.6, 16.8 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.8, -4.6, -4.3, -4.1, 18.0, 18.2, 21.5, 25.8, 25.9, 26.1, 32.3, 48.6, 63.0, 70.2, 71.5, 75.7, 81.2, 115.7, 139.0; HREIMS calcd for C₂₃H₄₆O₃Si₂ (M⁺) 426.2985, found 426.2977.

(3*R*,4*S*,5*R*)-3,5-Bis-(*tert*-butyldimethylsilyloxy)-4-[3-(*tert*-butyldimethylsilyloxy)propyl]oct-1-en-7-yne (38). Alcohol **37** (60.0 mg, 141 μ mol) was converted to silyl ether **38** (63.0 mg, 83%) according to the procedure described above for obtaining **33** from **32**. Data for **38**: colorless oil; $[\alpha]^{25}_D +8.70$ (*c* 1.15, CHCl₃); IR (neat) 3316, 2932, 1256, 1102, 924, 837, 776, 633 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.03 (s, 3H), 0.04 (s, 6H), 0.05 (s, 3H), 0.07 (s, 3H), 0.09 (s, 3H), 0.89 (s, 27H), 1.32 (m, 2H), 1.56 (m, 2H), 1.75 (ddt, 1H, *J* = 4.0, 6.4, 6.8 Hz), 1.95 (t, 1H, *J* = 2.8 Hz), 2.38 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 2.42 (ddd, 1H, *J* = 2.8, 6.4, 16.8 Hz), 3.56 (t, 2H, *J* = 6.8 Hz), 4.03 (dt, 1H, *J* = 4.0, 6.0 Hz), 4.12 (dd, 1H, *J* = 6.4, 7.6 Hz), 5.08 (d, 1H, *J* = 10.0 Hz), 5.14 (d, 1H, *J* = 17.2 Hz), 5.84 (ddd, 1H, *J* = 7.6, 10.0, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -5.3, -4.6, -4.4, -4.1, -3.6, 18.1, 18.2, 22.1, 25.9, 26.0, 26.1, 32.6, 49.2, 63.6, 69.9, 71.5, 75.9, 82.1, 115.5, 140.4; HREIMS calcd for C₂₅H₅₁O₃Si₃ (M⁺) 483.3146, found 483.3141.

(3*R*,4*S*,5*R*)-3,5-Bis-(*tert*-butyldimethylsilyloxy)-4-propyl-oct-1-en-7-yne (39). Alcohol **37** (70.0 mg, 164 μ mol) was converted to enyne **39** (54.0 mg, 80%) according to the procedure described above for obtaining **34** from **32**. Data for **39**: colorless oil; $[\alpha]^{22}_D +19.39$ (*c* 1.15, CHCl₃); IR (neat) 3316,

2957, 1254, 1075, 926, 837, 776, 625 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.02 (s, 3H), 0.05 (s, 3H), 0.06 (s, 3H), 0.09 (s, 3H), 0.87 (t, 3H, *J* = 7.2 Hz), 0.89 (s, 18H), 1.27 (m, 2H), 1.36 (m, 2H), 1.76 (ddt, 1H, *J* = 4.0, 5.2, 6.0 Hz), 1.95 (t, 1H, *J* = 2.8 Hz), 2.39 (ddd, 1H, *J* = 2.8, 6.4, 17.6 Hz), 2.41 (ddd, 1H, *J* = 2.8, 6.4, 17.6 Hz), 3.99 (dt, 1H, *J* = 4.0, 6.4 Hz), 4.12 (ddt, 1H, *J* = 6.0, 8.0, 1.2 Hz), 5.07 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.13 (dt, 1H, *J* = 1.2, 17.2 Hz), 5.85 (ddd, 1H, *J* = 8.0, 10.4, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.7, -4.5, -4.2, -3.7, 14.4, 18.0, 18.1, 22.3, 25.8, 25.9, 26.0, 28.2, 49.2, 69.7, 71.6, 75.7, 82.3, 115.2, 140.5; HREIMS calcd for C₁₉H₃₇O₂Si₂ (M⁺) 353.2330, found 353.2332.

(5*S*,6*R*)-6-(*tert*-Butyldimethylsilyloxy)-5-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]non-8-ynitrile (40). Alcohol **37** (470 mg, 1.10 mmol) was converted to nitrile **40** (400 mg, 83%) according to the procedure described above for obtaining **30** from **28**. Data for **40**: colorless oil; $[\alpha]^{20}_D +2.21$ (*c* 1.27, CHCl₃); IR (neat) 3314, 2930, 2361, 1255, 1078, 928, 837, 777, 632 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.05 (s, 3H), 0.06 (s, 3H), 0.72 (s, 3H), 0.09 (s, 3H), 0.89 (s, 18H), 1.43 (m, 2H), 1.73 (m, 3H), 1.99 (t, 1H, *J* = 2.6 Hz), 2.28 (t, 2H, *J* = 7.3 Hz), 2.39 (ddd, 2H, *J* = 2.8, 6.4, 6.4 Hz), 3.88 (dt, 1H, *J* = 2.8, 6.7 Hz), 4.18 (dd, 1H, *J* = 5.8, 7.7 Hz), 5.14 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.19 (dt, 1H, *J* = 1.2, 17.1 Hz), 5.80 (ddd, 1H, *J* = 7.0, 10.4, 17.1 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.7, -4.5, -4.2, -3.8, 17.5, 18.1, 18.2, 25.1, 25.3, 25.86, 25.9, 26.4, 48.5, 70.6, 71.1, 75.5, 81.1, 116.2, 119.8, 139.2; HREIMS calcd for C₂₀H₃₆O₂NSi₂ (M⁺ - *t*Bu) 378.2285, found 378.2298.

(5*S*,6*R*)-6-(*tert*-Butyldimethylsilyloxy)-5-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]non-9-ynal (41). Nitrile **40** (400 mg, 920 μ mol) was converted to aldehyde **41** (320 mg, 82%) according to the procedure described above for obtaining **31** from **30**. Data for **41**: colorless oil; $[\alpha]^{20}_D +6.30$ (*c* 1.78, CHCl₃); IR (neat) 3314, 2930, 2712, 2361, 1730, 1253, 1072, 924, 837, 775, 632 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.04 (s, 3H), 0.06 (s, 3H), 0.07 (s, 3H), 0.10 (s, 3H), 0.830 (s, 9H), 0.833 (s, 9H), 1.3 (m, 2H), 1.69 (m, 2H), 1.82 (m, 1H), 1.96 (t, 1H, *J* = 2.4 Hz), 2.40 (ddd, 1H, *J* = 2.8, 6.0, 17.2 Hz), 2.43 (ddd, 1H, *J* = 2.8, 6.0, 17.2 Hz), 2.37 (m, 3H), 3.96 (m, 1H), 4.12 (dd, 1H, *J* = 5.8, 7.0 Hz), 5.10 (dt, 1H, *J* = 1.6, 10.4 Hz), 5.17 (dt, 1H, *J* = 1.6, 16.5 Hz), 5.83 (ddd, 1H, *J* = 7.3, 10.4, 16.5 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.6, -4.4, -4.2, -3.7, 18.09, 18.17, 25.4, 25.9, 26.3, 44.4, 49.0, 70.3, 71.2, 75.6, 81.5, 115.8, 139.8, 202.8; HREIMS calcd for C₂₀H₃₇O₃Si₂ (M⁺ - *t*Bu) 381.2281, found 381.2278.

(5*S*,6*R*)-6-(*tert*-Butyldimethylsilyloxy)-5-[(*R*)-1-(*tert*-butyldimethylsilyloxy)allyl]non-8-yn-1-ol (42). Aldehyde **41** (320 mg, 731 μ mol) was reduced to alcohol **42** (300 mg, 93%) according to the procedure described above for obtaining **32** from **31**. Data for **42**: colorless oil; $[\alpha]^{21}_D +8.56$ (*c* 0.63, CHCl₃); IR (neat) 3314, 2932, 2361, 1253, 1070, 924, 837, 775, 625 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.04 (s, 3H), 0.06 (s, 3H), 0.07 (s, 3H), 0.10 (s, 3H), 0.89 (s, 18H), 1.37 (m, 2H), 1.54 (m, 2H), 1.78 (m, 1H), 1.97 (t, 1H, *J* = 2.6 Hz), 2.39 (m, 2H), 3.61 (t, 2H, *J* = 6.4 Hz), 3.97 (dt, 1H, *J* = 6.2, 10.0 Hz), 4.13 (t, 1H, *J* = 6.8 Hz), 5.09 (dt, 1H, *J* = 1.2, 10.4 Hz), 5.16 (dt, 1H, *J* = 1.2, 17.1 Hz), 5.84 (ddd, 1H, *J* = 7.3, 10.4, 17.1 Hz); ¹³C NMR (100 MHz, CDCl₃) δ -4.6, -4.4, -4.2, -3.7, 18.1, 18.2, 25.4, 25.7, 25.9, 26.0, 26.2, 33.3, 49.3, 63.0, 70.0, 71.5, 75.7, 82.0, 115.5, 140.3; HREIMS calcd for C₂₄H₄₈O₃Si₂ (M⁺) 440.3142, found 440.3148.

(3*R*,4*S*,5*R*)-3,5-Bis-(*tert*-butyldimethylsilyloxy)-4-[4-(*tert*-butyldimethylsilyloxy)butyl]oct-1-en-7-yne (43). Alcohol **42** (90.0 mg, 205 μ mol) was converted to silyl ether **43** (105 mg, 93%) according to the procedure described above for obtaining **33** from **32**. Data for **43**: colorless oil; $[\alpha]^{25}_D +7.58$ (*c* 1.29, CHCl₃); IR (neat) 3316, 2930, 2359, 1253, 1100, 924, 837, 775, 630 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 0.02 (s, 3H), 0.04 (s, 6H), 0.05 (s, 3H), 0.06 (s, 3H), 0.09 (s, 3H), 0.844 (s, 9H), 0.848 (s, 9H), 0.850 (s, 9H), 1.32 (m, 2H), 1.48 (m, 2H), 1.75 (m, 1H), 1.95 (t, 1H, *J* = 2.5 Hz), 2.39 (dd, 2H, *J* = 2.5, 6.4 Hz), 2.42 (t, 1H, *J* = 6.4 Hz), 3.58 (t, 2H, *J* = 6.4 Hz), 4.03 (dt, 1H, *J* = 6.4, 10.0 Hz), 4.13 (dd, 1H, *J* = 5.5, 7.3 Hz), 5.08 (dt, 1H, *J* = 1.2, 10.0 Hz), 5.14 (dt, 1H, *J* = 1.2, 17.2 Hz), 5.80 (ddd, 1H, *J* = 7.6, 10.0, 17.2 Hz); ¹³C NMR (100 MHz, CDCl₃)

δ -5.2, -4.6, -4.4, -4.2, -3.7, 18.1, 18.2, 18.4, 25.5, 25.89, 25.95, 26.00, 26.10, 33.5, 49.5, 63.3, 69.9, 71.5, 75.7, 82.2, 115.3, 140.5; HREIMS calcd for $C_{30}H_{62}O_3Si_3$ (M^+) 554.4007, found 554.4021.

(3R,4S,5R)-3,5-Bis-(*tert*-butyldimethylsilyloxy)-4-butylnon-1-en-7-yne (44). Alcohol **43** (90.0 mg, 205 μ mol) was reduced to enyne **44** (72.0 mg, 83%) according to the procedure described above for obtaining **34** from **32**. Data for **44**: colorless oil; $[\alpha]_D^{25} +6.69$ (c 0.78, $CHCl_3$); IR (neat) 3314, 2930, 2361, 1255, 1074, 924, 837, 777, 629 cm^{-1} ; 1H NMR (400 MHz, $CDCl_3$) δ 0.02 (s, 3H), 0.05 (s, 3H), 0.06 (s, 3H), 0.09 (s, 3H), 0.87 (t, 3H, J = 7.2 Hz), 0.88 (s, 9H), 0.89 (s, 9H), 1.28 (m, 6H), 1.76 (m, 1H), 1.95 (t, 1H, J = 2.7 Hz), 2.39 (dd, 2H, J = 2.7, 6.1 Hz), 3.99 (dt, 1H, J = 4.0, 6.1 Hz), 4.15 (m, 1H), 5.08 (dt, 1H, J = 1.2, 10.4 Hz), 5.15 (dt, 1H, J = 1.2, 17.2 Hz), 5.83 (ddd, 1H, J = 8.0, 10.4, 17.2 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ -4.7, -4.4, -4.2, -3.7, 14.0, 18.1, 18.2, 23.2, 25.89, 25.95, 26.04, 31.5, 49.5, 69.8, 71.7, 75.7, 82.4, 115.3, 140.5; HREIMS calcd for $C_{24}H_{48}O_2Si_2$ (M^+) 424.3013, found 424.3201.

(5Z,7E)-(1S,2S,3R,20R)-9,10-Seco-5,7,10(19)-cholesta-2,3-dihydroxymethyl-1,3,25-triol (3). The silyl ether **29** (20.0 mg, 39.1 μ mol) and vinyl bromide **11** (20.0 mg, 58.0 μ mol) were dissolved in TEA/toluene (3:1, 2.0 mL), and tris(dibenzylideneacetone)-dipalladium(0)-chloroform adduct (4.0 mg, 3.86 μ mol) and triphenylphosphine (10.0 mg, 38.1 μ mol) were added. After being stirred for 15 min at room temperature, the resultant yellow solution was heated at reflux for 2 h. The reaction mixture was filtered through a pad of silica gel. Concentration followed by preparative thin-layer chromatography on silica gel (20% EtOAc in hexanes) gave the crude protected vitamin as a white solid, which was used in the next step without further purification.

To a cold (0 $^{\circ}C$) and stirred solution of the crude protected vitamin in MeOH (2.0 mL) was added (+)-10-camphorsulfonic acid (10.0 mg, 43.0 μ mol). After being stirred for 1 h at 0 $^{\circ}C$, the reaction mixture was allowed to warm to room temperature and stirred for another 12 h. The resultant solution was diluted with EtOAc (10 mL) and washed with saturated aqueous $NaHCO_3$ solution (3 \times 1 mL) and brine (3 \times 1 mL). The aqueous layer was extracted with EtOAc (3 \times 2 mL), and the combined organic layer was dried over Na_2SO_4 . Filtration and concentration followed by preparative thin-layer chromatography on silica gel (20% MeOH in CH_2Cl_2) gave 7.0 mg (40%) of 2 α -(hydroxymethyl)-1 α ,25-dihydroxyvitamin D₃ **3** as a white solid; $[\alpha]_D^{20} +12.95$ (c 0.0085, $CHCl_3$); UV (EtOH) λ_{max} 269 nm, λ_{min} 226 nm; 1H NMR (400 MHz, $CDCl_3$) δ 0.53 (s, 3H), 0.94 (d, 3H, J = 6.8 Hz), 2.31 (m, 2H), 2.67 (dd, 1H, J = 4.4, 12.8 Hz), 2.73 (bs, 1H), 2.84 (m, 1H), 4.01 (m, 2H), 4.24 (m, 1H), 4.47 (bs, 1H), 5.02 (d, 1H, J = 2.0 Hz), 5.30 (d, 1H, J = 2.0 Hz), 5.98 (d, 1H, J = 10.8 Hz), 6.45 (d, 1H, J = 10.8 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ 12.6, 18.8, 20.8, 22.2, 23.5, 27.7, 29.1, 29.2, 29.4, 36.1, 36.4, 40.5, 44.4, 45.0, 45.9, 50.4, 56.3, 56.5, 63.9, 68.0, 71.1, 115.0, 116.8, 125.4, 132.0, 143.8, 145.9; HREIMS calcd for $C_{28}H_{44}O_3$ ($M^+ - H_2O$) 428.3290, found 428.3287.

(5Z,7E)-(1S,2S,3R,20R)-9,10-Seco-5,7,10(19)-cholesta-2,3-dihydroxyethyl-1,3,25-triol (4). Silyl ether **33** (25.0 mg, 47.5 μ mol) was converted to 2 α -(hydroxyethyl)-1 α ,25-dihydroxyvitamin D₃ **4** (8.5 mg, 39%) according to the procedure described above for **3**. Data for **4**: white solid; $[\alpha]_D^{20} +13.85$ (c 0.00722, $CHCl_3$); UV (EtOH) λ_{max} 268 nm, λ_{min} 228 nm; 1H NMR (400 MHz, $CDCl_3$) δ 0.53 (s, 3H), 0.94 (d, 3H, J = 6.4 Hz), 2.26 (dd, 1H, J = 8.0, 12.8 Hz), 2.53 (bs, 1H), 2.66 (dd, 1H, J = 4.0, 13.2 Hz), 2.83 (m, 1H), 3.79 (m, 2H), 3.94 (m, 1H), 4.37 (d, 1H, J = 2.0 Hz), 5.02 (d, 1H, J = 1.2 Hz), 5.30 (bs, 1H), 6.01 (d, 1H, J = 10.8 Hz), 6.40 (d, 1H, J = 10.8 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ 12.0, 18.8, 20.8, 22.2, 23.5, 27.6, 29.07, 29.14, 29.3, 31.8, 36.1, 36.4, 40.5, 43.9, 44.4, 45.9, 48.6, 56.3, 56.5, 61.6, 70.5, 71.0, 74.9, 113.7, 116.9, 124.8, 132.5, 143.4, 146.4; HREIMS calcd for $C_{29}H_{48}O_4$ (M^+) 460.3553, found 460.3557.

(5Z,7E)-(1S,2S,3R,20R)-9,10-Seco-5,7,10(19)-cholesta-2,3-dihydroxypropyl-1,3,25-triol (5). Silyl ether **38** (25.0 mg, 46.3 μ mol) was converted to 2 α -(hydroxypropyl)-1 α ,25-dihydroxyvitamin D₃ **5** (7.2 mg, 33%) according to the

procedure described above for **3**. Data for **5**: white solid; $[\alpha]_D^{20} +161.29$ (c 0.00186, $CHCl_3$); UV (EtOH) λ_{max} 268 nm, λ_{min} 227 nm; 1H NMR (400 MHz, $CDCl_3$) δ 0.53 (s, 3H), 0.94 (d, 3H, J = 6.4 Hz), 2.25 (dd, 1H, J = 8.4, 13.2 Hz), 2.66 (dd, 1H, J = 4.0, 13.2 Hz), 2.83 (m, 1H), 3.70 (t, 2H, J = 5.6 Hz), 3.90 (dt, 1H, J = 4.0, 8.0 Hz), 4.38 (d, 1H, J = 3.6 Hz), 5.00 (d, 1H, J = 1.6 Hz), 5.28 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 11.2 Hz), 6.40 (d, 1H, J = 11.2 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ 12.2, 18.9, 21.0, 22.4, 23.0, 23.7, 27.8, 29.2, 29.3, 29.5, 30.3, 36.2, 36.5, 40.6, 44.4, 44.5, 46.0, 49.1, 56.4, 56.6, 62.9, 70.4, 71.1, 73.6, 113.5, 116.8, 124.7, 132.5, 143.2, 146.4; HREIMS calcd for $C_{30}H_{50}O_4$ (M^+) 474.3709, found 474.3709.

(5Z,7E)-(1S,2S,3R,20R)-9,10-Seco-5,7,10(19)-cholesta-2,3-dihydroxybutyl-1,3,25-triol (6). Silyl ether **43** (25.0 mg, 45.1 μ mol) was converted to 2 α -(hydroxybutyl)-1 α ,25-dihydroxyvitamin D₃ **6** (8.2 mg, 37%) according to the procedure described above for **3**. Data for **6**: white solid; $[\alpha]_D^{20} +86.15$ (c 0.0650, $CHCl_3$); UV (EtOH) λ_{max} 268 nm, λ_{min} 227 nm; 1H NMR (400 MHz, $CDCl_3$) δ 0.53 (s, 3H), 0.93 (d, 3H, J = 6.6 Hz), 2.24 (dd, 1H, J = 8.4, 13.2 Hz), 2.66 (dd, 1H, J = 4.4, 13.2 Hz), 2.83 (m, 1H), 3.71 (m, 4H), 3.89 (ddd, 1H, J = 4.4, 7.8, 13.2 Hz), 4.38 (t, 1H, J = 3.1 Hz), 4.99 (d, 1H, J = 1.6 Hz), 5.26 (d, 1H, J = 1.6 Hz), 6.00 (d, 1H, J = 11.5 Hz), 6.40 (d, 1H, J = 11.5 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ 12.0, 18.8, 20.8, 22.2, 23.4, 23.5, 26.2, 27.7, 29.1, 29.2, 29.4, 32.8, 36.1, 36.4, 40.5, 44.4, 45.9, 49.5, 56.3, 56.5, 62.8, 70.4, 71.1, 73.5, 113.6, 116.9, 124.8, 132.6, 143.4, 146.6; HREIMS calcd for $C_{31}H_{52}O_4$ (M^+) 488.3866, found 488.3867.

(5Z,7E)-(1S,2S,3R,20R)-9,10-Seco-5,7,10(19)-cholesta-2,3-dihydroxyethyl-1,3,25-triol (7). Enyne **34** (30.0 mg, 75.8 μ mol) was converted to 2 α -ethyl-1 α ,25-dihydroxyvitamin D₃ **7** (6.8 mg, 20%) according to the procedure described above for **3**. Data for **7**: white solid; $[\alpha]_D^{20} +80.79$ (c 0.108, $CHCl_3$); UV (EtOH) λ_{max} 269 nm, λ_{min} 227 nm; 1H NMR (400 MHz, $CDCl_3$) δ 0.53 (s, 3H), 0.94 (d, 3H, J = 6.0 Hz), 0.95 (t, 3H, J = 7.2 Hz), 2.24 (dd, 1H, J = 8.8, 12.8 Hz), 2.66 (dd, 1H, J = 4.0, 13.2 Hz), 2.83 (m, 1H), 3.89 (m, 1H), 4.37 (bs, 1H), 4.99 (d, 1H, J = 1.6 Hz), 5.27 (bs, 1H), 6.00 (d, 1H, J = 11.2 Hz), 6.40 (d, 1H, J = 11.2 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ 11.7, 12.0, 18.8, 19.3, 20.8, 22.2, 23.5, 27.6, 29.07, 29.14, 29.3, 36.1, 36.4, 40.5, 44.4, 45.9, 51.1, 56.3, 56.5, 70.1, 71.0, 73.0, 113.4, 116.9, 124.7, 132.8, 143.3, 146.7; HREIMS calcd for $C_{29}H_{48}O_3$ (M^+) 444.3603, found 460.3604.

(5Z,7E)-(1S,2S,3R,20R)-9,10-Seco-5,7,10(19)-cholesta-2,3-dihydroxypropyl-1,3,25-triol (8). Enyne **39** (30.0 mg, 73.2 μ mol) was converted to 2 α -propyl-1 α ,25-dihydroxyvitamin D₃ **8** (6.2 mg, 18%) according to the procedure described above for **3**. Data for **8**: white solid; $[\alpha]_D^{20} -202.91$ (c 0.06653, $CHCl_3$); UV (EtOH) λ_{max} 269 nm, λ_{min} 227 nm; 1H NMR (400 MHz, $CDCl_3$) δ 0.53 (s, 3H), 0.94 (d, 3H, J = 6.4 Hz), 1.01 (t, 3H, J = 6.8 Hz), 2.24 (dd, 1H, J = 8.8, 12.4 Hz), 2.66 (dd, 1H, J = 4.0, 13.2 Hz), 2.83 (m, 1H), 3.89 (m, 1H), 4.39 (bs, 1H), 4.99 (d, 1H, J = 1.6 Hz), 5.27 (bs, 1H), 6.00 (d, 1H, J = 11.2 Hz), 6.40 (d, 1H, J = 11.2 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ 12.1, 14.4, 18.8, 20.4, 20.8, 22.3, 23.6, 27.7, 28.7, 29.1, 29.2, 29.4, 36.1, 36.4, 40.5, 44.3, 44.4, 45.9, 49.3, 56.4, 56.6, 70.4, 71.1, 73.4, 113.3, 116.9, 124.8, 132.8, 143.3, 146.8; HREIMS calcd for $C_{30}H_{50}O_3$ (M^+) 458.3760, found 458.3755.

(5Z,7E)-(1S,2S,3R,20R)-9,10-Seco-5,7,10(19)-cholesta-2,3-dihydroxybutyl-1,3,25-triol (9). Enyne **44** (25.0 mg, 59.0 μ mol) was converted to 2 α -butyl-1 α ,25-dihydroxyvitamin D₃ **9** (8.40 mg, 30%) according to the procedure described above for **3**. Data for **9**: white solid; $[\alpha]_D^{20} +40.70$ (c 0.36118, $CHCl_3$); UV (EtOH) λ_{max} 269 nm, λ_{min} 227 nm; 1H NMR (400 MHz, $CDCl_3$) δ 0.53 (s, 3H), 2.24 (dd, 1H, J = 8.8, 13.2 Hz), 2.66 (dd, 1H, J = 4.4, 13.8 Hz), 2.82 (m, 1H), 3.72 (m, 2H), 3.89 (ddd, 1H, J = 4.4, 8.2, 13.2 Hz), 4.38 (t, 1H, J = 3.8 Hz), 4.99 (d, 1H, J = 2.2 Hz), 5.26 (d, 1H, J = 1.1 Hz), 6.00 (d, 1H, J = 11.0 Hz), 6.40 (d, 1H, J = 11.0 Hz); ^{13}C NMR (100 MHz, $CDCl_3$) δ 12.1, 14.4, 18.8, 20.4, 20.8, 22.2, 23.1, 23.5, 26.1, 27.7, 29.1, 29.2, 29.4, 29.5, 36.1, 36.4, 40.5, 44.4, 45.9, 49.5, 56.4, 56.6, 70.4, 71.1, 73.4, 113.3, 116.9, 124.8, 132.8, 143.3, 146.8; HREIMS calcd for $C_{31}H_{52}O_3$ (M^+) 472.3916, found 472.3902.

Acknowledgment. We are grateful to Ms. Junko Shimode, Ms. Jinko Nonobe, and Ms. Maroka Kit-sukawa (Teikyo University) for the spectroscopic mea-surements. This study was supported in part by Grants-in-Aid from the Ministry of Education, Science, Sports and Culture of Japan.

Supporting Information Available: ¹H NMR spectra for all new compounds, synthetic route of compound **19c** from **19a**, and elemental analysis data for **19c**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

JO010375I