

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/49740534>

Total Asymmetric Syntheses of β -Hydroxy- δ -lactones via Umpolung with Sulfur Dioxide †

ARTICLE *in* THE JOURNAL OF ORGANIC CHEMISTRY · FEBRUARY 2011

Impact Factor: 4.72 · DOI: 10.1021/jo102035d · Source: PubMed

CITATIONS

6

READS

26

5 AUTHORS, INCLUDING:



Sylvain Laclef

Université de Rouen

14 PUBLICATIONS 99 CITATIONS

SEE PROFILE



Māris Turks

Riga Technical University

87 PUBLICATIONS 404 CITATIONS

SEE PROFILE



Pierre Vogel

École Polytechnique Fédérale de Lausanne

760 PUBLICATIONS 9,926 CITATIONS

SEE PROFILE

Total Asymmetric Syntheses of β -Hydroxy- δ -lactones via Umpolung with Sulfur Dioxide[†]


Claudia J. Exner,^{‡,⊥} Sylvain Laclef,^{‡,⊥} Florent Poli,[‡] Maris Turks,[§] and Pierre Vogel^{*,‡}

[‡]Laboratory of Glycochemistry and Asymmetric Synthesis (LGSA), Swiss Federal Institute of Technology Lausanne (EPFL), Batochime, CH-1015 Lausanne, Switzerland, and [§]Faculty of Material Science and Applied Chemistry, Riga Technical University, LV-1048 Riga, Latvia.

[⊥]These authors contributed equally to this work

pierre.vogel@epfl.ch.

Received October 18, 2010



Cyclic stereotriads and stereotetrads of the β -hydroxy- δ -lactone type, e.g. prelactones B and E, common in polyketides and polypropionates, are prepared via SO_2 -induced oxyallylations of enoxysilanes with (1*E*,3*Z*)-1-(1-phenylethoxy)penta-1,3-dien-3-yl carboxylates. Using (*Z*)- or (*E*)-enoxysilanes both 4,5-*cis*- or 4,5-*trans*- δ -lactones are obtained. Depending on the reduction method applied to the obtained aldol intermediates 5,6-*trans* or 5,6-*cis*-derivatives are formed. The δ -lactones can be prepared in both their enantiomeric forms depending on the (1*R*)- or (1*S*)-configuration of the starting 1-(1-phenylethoxy)penta-1,3-dienes.

Introduction

Polyketides and polypropionates represent an important class of natural compounds¹ with broad potential for pharmacological applications.² Their stereochemical complexity has stimulated intensive research toward the development of chemical^{3,4} and biochemical methods⁵ for their total synthesis. β -Hydroxy- δ -lactones like compounds 1–7 (Chart 1) constitute, as cyclic stereotriads and stereotetrads, a common

structural motif in a large number of natural polyketides and polypropionates and are intermediates in the step-by-step biosynthesis of these compounds.⁶ As such, they have been isolated from different polyketide-producing organisms.

[†] Dedicated to Professor Janine Cossy, ESPCI, Paris, on the occasion of being awarded the Grand Prix Achille LeBel.

(1) (a) O'Hagan, D. *The Polyketide Metabolites*; Ellis Horwood: Chichester, 1991. (b) O'Hagan, D. *Nat. Prod. Rep.* **1995**, *12*, 1–32. (c) Davies-Coleman, M. T.; Garson, M. J. *Nat. Prod. Rep.* **1998**, *15*, 477–493. (d) Paterson, I.; Florence, G. J. *Eur. J. Org. Chem.* **2003**, 2193–2208.



(2) (a) Kim, J.; Park, E. J. *Curr. Med. Chem. Anti-Cancer Agents* **2002**, *2*, 485–537. (b) Watkins, E. B.; Chittiboyina, A. G.; Jung, J.-C.; Avery, M. A. *Curr. Pharm. Des.* **2005**, *11*, 1615–1653. (c) Tan, T. T. *Phytochemistry* **2007**, *68*, 954–979. (d) Mayer, A. M. S.; Gustafson, K. R. *Eur. J. Cancer* (Oxford, England: 1990) **2008**, *44*, 2357–2387; (e) Frenz, J. L.; Kohl, A. C.; Kerr, R. G. *Expert Opin. Therap. Pat.* **2004**, *14*, 17–35. (f) Faulkner, D. J. *Nat. Prod. Rep.* **2000**, *17*, 1–6. (g) Proksch, P.; Edrada, A.; Ebel, R. *Appl. Microbiol. Biotechnol.* **2002**, *59*, 125–134. (h) Andrack, R. B. *Org. Prep. Proced. Int.* **2009**, *41*, 359–383. (i) Bonazzi, S.; Eidam, O.; Guttinger, S.; Wach, J.-Y.; Zemp, I.; Kutay, U.; Gademann, K. *J. Am. Chem. Soc.* **2010**, *132*, 1432–1442.

(3) For reviews, see: (a) Schetter, B.; Mahrwald, R. *Angew. Chem., Int. Ed.* **2006**, *45*, 7506–7525. (b) Li, J.; Menche, D. *Synthesis* **2009**, 2293–2315.

(4) For examples, see: (a) Florence, G. J.; Gardner, N. M.; Paterson, I. *Nat. Prod. Rep.* **2008**, *25*, 342–375. (b) Paterson, I.; Findlay, A. D. *Aust. J. Chem.* **2009**, *62*, 624–638.

(5) (a) Staunton, J. *Angew. Chem., Int. Ed.* **1991**, *30*, 1302–1306. (b) Katz, L. *Chem. Rev.* **1997**, *97*, 2557–2576. (c) Pieper, R.; Ebert-Khosla, S.; Cane, D. E.; Khosla, C. *Biochemistry* **1996**, *35*, 2054–2060. (d) Cane, D. E.; Walsh, C.; Khosla, C. *Science* **1998**, *282*, 63–68. (e) Weissmann, K. J.; Bycroft, M.; Cutter, A. L.; Hanefeld, U.; Frost, E. J.; Timoney, M. C.; Harnis, R.; Handa, S.; Roddis, M.; Staunton, J.; Leadlay, P. F. *Chem. Biol.* **1998**, *5*, 743–754. (f) Ranganathan, A.; Timoney, M.; Bycroft, M.; Cortes, J.; Thomas, I. P.; Wilkinson, B.; Kellenberger, L.; Hanefeld, U.; Galloway, I. S.; Staunton, J.; Leadlay, P. F. *Chem. Biol.* **1999**, *6*, 731–741. (g) Pfeifer, B. A.; Admiraal, S. J.; Gramajo, H.; Cane, D. E.; Khosla, C. *Science* **2001**, *291*, 1790–1792. (h) Nickolson, T. P.; Rudd, B. A. M.; Dawson, M.; Lazarus, C. M.; Simpson, T. J.; Cox, R. J. *Chem. Biol.* **2001**, *8*, 157–178. (i) Beck, B. J.; Aldrich, C. C.; Fecik, R. A.; Reynolds, K. A.; Sherman, D. H. *J. Am. Chem. Soc.* **2003**, *125*, 4682–4683. (j) Regentin, R.; Kennedy, J.; Wu, N.; Carney, J. R.; Licari, P.; Galazzo, J.; Desai, R. *Biotechnol. Prog.* **2004**, *20*, 122–127. (k) Menzella, H. G.; Reid, R.; Carney, J. R.; Chandran, S. S.; Reisinger, S. J.; Patel, K. G.; Hopwood, D. A.; Santi, D. U. *Nat. Biotechnol.* **2005**, *23*, 1171–1176. (l) Chandran, S. S.; Menzella, H. G.; Carney, J. R.; Santi, D. V. *Chem. Biol.* **2006**, *13*, 469–474. (m) Meier, J. L.; Burkart, M. D. *Chem. Soc. Rev.* **2009**, *38*, 2012–2045. (n) Li, Y.; Xu, W.; Tang, T. J. *Biol. Chem.* **2010**, *285*, 22764–22773.

(6) See, e.g.: (a) Kao, C. M.; McPherson, M.; McDaniel, R. N.; Fu, J.; Cane, D. E.; Khosla, C. *J. Am. Chem. Soc.* **1998**, *120*, 2478–2479. (b) Castonguay, R.; He, W. M.; Chen, A. Y.; Khosla, C.; Cane, D. E. *J. Am. Chem. Soc.* **2007**, *129*, 13758–13769. (c) Castonguay, R.; Cane, D. E. *J. Am. Chem. Soc.* **2008**, *130*, 11598–11599. (d) Valenzano, C. R.; Lawson, R. J.; Chen, A. Y.; Khosla, C.; Cane, D. E. *J. Am. Chem. Soc.* **2009**, *131*, 18501–18511.

CHART 1. Different δ -Lactones 1–7, Synthesized Using SO_2 -ChemistrySCHEME 1. One-Pot Synthesis of α,β,γ -*syn,anti*- or *anti,anti*-Stereotriads through SO_2 -Induced Oxyallylation of (*Z*)- or (*E*)-Enoxysilanes^a

^aThe shown structures are obtained for $\text{R}^* = (S)\text{-1-phenethyl}$.

For instance, β -hydroxy- δ -lactone prelactone B (**1**) has been found in the fermentation broth of *Streptomyces* producing concanamycins and bafilomycins (Chart 1),⁷ which stimulated several syntheses of (+)-(**1**) and its stereoisomers using various approaches.⁸ Prelactone E ((-)-**2**), a product of chemical degradation of concanolide derivatives,⁹ has been synthesized recently by two groups applying Evans' aldol chemistry.^{10,11} Using an L-proline-catalyzed aldol reaction Barbas and co-workers obtained lactone (+)-**3** in a two-step process with 11% ee, that could be improved by carrying out the reaction in an ionic liquid.^{12,13} Lactone (-)-**4**, the 5-epimer of (+)-**3**, has been prepared by Cordova and co-workers from propanal in a three-step process with an ee > 99% involving two successive L-proline- and D-proline-catalyzed aldol reactions followed by MnO_2 -oxidation.¹⁴ Compound (-)-**5** was obtained by Hoffmann and co-workers via enantioselective crotylboration of methacrolein followed by diastereoselective hydroboration.^{17b} Chênevert and co-workers made (-)-**5** in a few steps with 58% overall yield via enzymatic desymmetrization

of *meso*-(*anti,anti*)-2,4-dimethyl-1,3,5-pentanetriol.^{15,16} δ -Lactone (+)-**6**, the 5-epimer of (-)-**5**, has been obtained only through biological synthesis applying polyketide synthase.^{6a} Both lactones, (-)-**5** and (+)-**6**, contain an α,β,γ -*anti,anti*-stereotriad subunit, the most elusive to obtain.¹⁷

In this report we propose alternative syntheses of lactones (+)-**1**–(+)-**6**. Our method is general and has been applied also to the synthesis of lactone (-)-**7**, a yet unknown compound.¹⁸



With the use of our SO_2 -reaction cascade that combines electron-rich dienes **8** and (*Z*)- or (*E*)-enoxysilanes **10** via SO_2 -Umpolung, we have developed a one-pot synthesis of α,β,γ -*syn,anti*- and *anti,anti*-stereotriads of types **12** and **13**, respectively (Scheme 1).^{19,20} The starting dienes **8** are readily obtained from pentan-3-one, ethyl formate and inexpensive, enantiomerically enriched (*S*)- or (*R*)-1-phenylethanol, the source of chirality.²¹


A hetero-Diels–Alder addition between diene **8** and SO_2 followed by Lewis acid-assisted ionization gives a zwitterionic

- (7) Bindseil, K. U.; Zeeck, A. *Helv. Chim. Acta* **1993**, *76*, 150–157.
 (8) (a) Hanefeld, U.; Hooper, A. M.; Staunton, J. *Synthesis* **1999**, 401–403. (b) Fournier, L.; Gaudel-Siri, A.; Kocienski, P. J.; Pons, J.-M. *Synlett* **2003**, 107–111. (c) Chakraborty, T. K.; Tapadar, S. *Tetrahedron Lett.* **2003**, *44*, 2541–2543. (d) Pihko, P. M.; Erkkilä, A. *Tetrahedron Lett.* **2003**, *44*, 7607–7609. (e) Csaky, A. G.; Mba, M.; Plumet, J. *Synlett* **2003**, 2092–2094. (f) Enders, D.; Haas, M. *Synlett* **2003**, 2182–2184. (g) Dias, L. C.; Steil, L. J.; Vasconcelos, F. de A. *Tetrahedron: Asymmetry* **2004**, *15*, 147–150. (h) Yadav, J. S.; Reddy, K. B.; Sabitha, G. *Tetrahedron Lett.* **2004**, *45*, 6475–6476. (i) Aggarwal, V. K.; Bae, I.; Lee, H.-Y. *Tetrahedron* **2004**, *60*, 9725–9733. (j) Yadav, J. S.; Reddy, M. S.; Prasad, A. R. *Tetrahedron Lett.* **2005**, *46*, 2133–2136. (k) Salaskar, A. A.; Mayekar, N. V.; Sharma, A.; Nayak, S. K.; Chattopadhyaya, A.; Chattopadhyay *Synthesis* **2005**, 2777–2781. (l) Sellars, J. D.; Steel, P. G. *Org. Bioorg. Chem.* **2006**, *4*, 3223–3224. (m) Srihari, P.; Ravindar, K.; Somaiah, R.; Yadav, J. S. *Synth. Commun.* **2008**, *38*, 1389–1397. (n) Sellars, J. D.; Steel, P. D. *Tetrahedron* **2009**, *65*, 5588–5595.
 (9) Boddien, C.; Gerber-Nolte, J.; Zeeck, A. *Liebigs Ann.* **1996**, 1381–1384.
 (10) Hinterding, K.; Singhanat, S.; Oberer, L. *Tetrahedron Lett.* **2001**, *42*, 8463–8465.
 (11) Sabitha, G.; Padmaja, P.; Reddy, K. B.; Yadav, J. S. *Tetrahedron Lett.* **2008**, *49*, 919–922.
 (12) Chowdari, N. S.; Ramachary, D. B.; Cordova, A.; Barbas, C. F., III. *Tetrahedron Lett.* **2002**, *43*, 9591–9595.
 (13) Cordova, A. *Tetrahedron Lett.* **2004**, *45*, 3949–3952.
 (14) Casas, J.; Enguist, M.; Ibrahim, I.; Kaynak, B.; Cordova, A. *Angew. Chem., Int. Ed.* **2005**, *44*, 1343–1345.

- (15) Chênevert, R.; Courchesne, G.; Caron, D. *Tetrahedron: Asymmetry* **2003**, *14*, 2567–2571.
 (16) (-)-**5** is the precursor of the acid moiety of dolabiferol: (a) Ciavatta, M. L.; Gavagnin, M.; Politi, R.; Cimino, G.; Martinez, E.; Ortea, J.; Mattia, C. A. *Tetrahedron* **1996**, *52*, 12831–12838. For attempted syntheses see also: (b) Dias, L. C.; de Sousa, M. A. *Tetrahedron Lett.* **2003**, *44*, 5625–5628. (c) Pelchat, N.; Caron, D.; Chênevert, R. *J. Org. Chem.* **2007**, *72*, 8484–8438. (d) Lister, T.; Perkins, M. V. *Org. Lett.* **2006**, *8*, 1827–1830.
 (17) (a) Hoffmann, R. W. *Angew. Chem., Int. Ed.* **1987**, *26*, 489–503. (b) Hoffmann, R. W.; Dahmann, G.; Andersen, M. W. *Synthesis* **1994**, 629–638.
 (18) For chemical syntheses of related lactones, see e.g.: (a) Zhen, W.; Chu, K. H.; Rosenblum, M. J. *Org. Chem.* **1997**, *62*, 3344–3454. (b) Tholander, J.; Carreira, E. M. *Helv. Chim. Acta* **2001**, *84*, 613–622. (c) Pichlmair, S.; Marques, M. M. B.; Green, M. P.; Martin, H. J.; Mulzer, J. *Org. Lett.* **2003**, *5*, 4657–4659. (d) Berkenbusch, T.; Brückner, R. *Chem.—Eur. J.* **2004**, *10*, 1545–1557. (e) Yu, Z.; Liu, X.; Dong, Z.; Xie, M.; Feng, X. *Angew. Chem., Int. Ed.* **2008**, *47*, 1308–1311. (f) Magaver, T.; Martin, H. J.; Mulzer, J. *Angew. Chem., Int. Ed.* **2009**, *48*, 6032–6036. (g) Brazeau, J.-F.; Mochirian, P.; Prévost, M.; Guindon, Y. *J. Org. Chem.* **2009**, *74*, 64–74. (h) Magaver, T.; Martin, H. J.; Mulzer, J. *Chem.—Eur. J.* **2010**, *16*, 507–519.
 (19) (a) Vogel, P.; Turks, M.; Bouchez, L. C.; Markovic, D.; Varela-Alvarez, A.; Sordo, J. A. *Acc. Chem. Res.* **2007**, *40*, 931–942. (b) Turks, M.; Exner, C. J.; Hamel, C.; Vogel, P. *Synthesis* **2009**, 1065–1074.
 (20) Laclef, S.; Turks, M.; Vogel, P. *Angew. Chem., Int. Ed.* **2010**, *49*, 825–827.
 (21) Laclef, S.; Exner, C. J.; Turks, M.; Videtta, V.; Vogel, P. *J. Org. Chem.* **2009**, *74*, 8882–8885.

SCHEME 2. Syntheses of Prelactones B ((+)-1) and E ((-)-2)

SCHEME 3. Syntheses of δ -Lactones (+)-3, (-)-4, (-)-5, and (+)-6

SCHEME 4. Synthesis of δ -Lactone (–)-7

species **9**, which undergoes an oxyallylation reaction with alkenes **10** to silylsulfonates **11**. The latter are converted *in situ* into stereotriads **12** and **13**, respectively, in the presence of catalytic $\text{Pd}(\text{OAc})_2/\text{PPh}_3$, involving a highly stereoselective chirality transfer from the ϵ -center in **11** to the γ -center in **12** respective **13** (Scheme 1).^{22,23}

Results and Discussion

Our syntheses of the natural prelactones B ((+)-**1**) and E ((–)-**2**) combine diene **14**²⁴ with (*Z*)-enoxysilanes **15a** and **15b**, respectively. The SO_2 -induced oxyallylation and concomitant desulfinylative desilylation afforded 3:1-mixtures of α,β -*syn*- and α,β -*anti*-stereodiad **16a/17a** (72% yield) and **16b/17b** (55% yield), respectively (Scheme 2). These mixtures were treated with TiCl_4 in CH_2Cl_2 to cleave the phenethyl ether moieties and provided 3:1 mixtures of the corresponding alcohols **18a/19a** (84%) and **18b/19b** (95%). Reduction of these mixtures with $\text{Me}_4\text{NBH}(\text{OAc})_3/\text{AcOH}$ ²⁶ gave 3:1 mixtures of stereotriads **20a/21a** (94%) and **20b/21b** (60%). Ozonolysis of the latter, treatment with Me_2S , and chromatographic purification furnished (+)-**1** (83% based on **20a**) and (–)-**2** (91% based on **20b**).

This synthesis of (+)-**1** gives the final product in four steps and 35% yield based on diene (*S*)-**14** or in eight steps and 13% yield based on propionyl chloride, the starting material of diene (*S*)-**14**.²⁴ Prelactone E ((–)-**2**) was obtained in four steps and 21% overall yield based on diene (*S*)-**14**.

The β -hydroxy- δ -lactones (+)-**3**, (–)-**4**, (–)-**5**, and (+)-**6** (Scheme 3) were obtained by applying our oxyallylation cascade to diene (+)-**22**,^{21,27} and using (*Z*)-**15b** and (*E*)-enoxysilane **23**. This generated the stereotriads (+)-**24**²⁷ and (+)-**25**²⁰ with

diastereoselectivities of 6:1 and 3:1, respectively.²⁸ Treatment of **24** with TiCl_4 in CH_2Cl_2 at -78°C afforded titanium alkoxide **26** which was reacted directly with $\text{BH}_3\cdot\text{Me}_2\text{S}$ ²⁹ to give stereotetrad (+)-**27** in 90% yield. Ozonolysis of the enol ester of (+)-**27** followed by workup with Me_2S provided lactone (+)-**3** (78%). Aqueous workup of **26** furnished alcohol (–)-**28**. Its reduction with $\text{Me}_4\text{NBH}(\text{OAc})_3$ ²⁶ gave (+)-**29** (67%), the ozonolysis of which provided lactone (–)-**4** (78%). The same reaction sequence applied to (+)-**25** furnished stereotetrads (+)-**32** (72%) and (–)-**33** (67%), which were ozonolyzed to produce lactones (–)-**5** (62%) and (+)-**6** (73%), respectively. Structures of lactones (+)-**3**, (–)-**4**, (–)-**5**, and (+)-**6** were proven by their spectral data. Their relative configuration was established by the vicinal $^3J_{\text{H,H}}$ coupling constants in the ^1H NMR spectra. Structures of (–)-**5** and (+)-**6** were also confirmed by single-crystal X-ray diffraction studies.³⁰ The diversity of our methodology is demonstrated by the synthesis of four different stereotetrads, i. e. structures (+)-**27**, (+)-**29**, (+)-**32**, and (–)-**33** (Scheme 3), all using the same diene (*R*)-**22** as starting material.

Lactone (+)-**3** was synthesized in three steps and 48% overall yield from diene (+)-**22**, and (–)-**4** was obtained in four synthetic steps and 29% overall yield based on (+)-**22**. Diastereoisomers (–)-**5** and (+)-**6** were synthesized in four and three steps with 14% and 26% overall yields, respectively, starting from diene (+)-**22**.

Lactone (–)-**7** was derived in a similar way from diene (–)-**34**²⁷ and (*E*)-enoxysilane **35** (Scheme 4). Their SO_2 -mediated condensation produced stereotriad (–)-**36**²⁰ (67%, dr > 10:1). Reduction of ketone (–)-**36** with $\text{Me}_2\text{AlCl}/\text{Bu}_3\text{SnH}$ ³¹ in CH_2Cl_2 (workup with KF) gave stereotetrad (–)-**37**²⁰ in 90% yield (dr > 10:1). FeCl_3 -induced $\text{S}_{\text{N}}1$ -debenzylation of (–)-**37** provided diol (+)-**38** (93%, dr > 10:1). Its relative configuration was confirmed by the ^1H - and ^{13}C NMR

(22) Huang, X.; Craita, C.; Vogel, P. *J. Org. Chem.* **2004**, *69*, 4272–4275.

(23) Vogel, P.; Turks, M.; Bouchez, L.; Craita, C.; Huang, X.; Murcia, M. C.; Fonquerne, F.; Didier, C.; Flowers, C. *Pure Appl. Chem.* **2008**, *80*, 791–805.

(24) Turks, M.; Fonquerne, F.; Vogel, P. *Org. Lett.* **2004**, *6*, 1053–1056.

(25) Turks, M.; Murcia, M. C.; Scopelliti, R.; Vogel, P. *Org. Lett.* **2004**, *6*, 3031–3034.

(26) (a) Evans, D. A.; Chapman, K. T.; Carreira, E. M. *J. Am. Chem. Soc.* **1988**, *110*, 3560–3578. (b) Evans, D. A.; Chapman, K. T. *Tetrahedron Lett.* **1986**, *27*, 5939–5942.

(27) Turks, M.; Huang, X.; Vogel, P. *Chem.—Eur. J.* **2005**, *11*, 465–476.

(28) Compound **25** may be obtained with a better diastereomeric ratio of 5:1 using the *tert*-butyric ester of diene (*S*)-**14** ((1*E*,3*Z*)-2-methyl-1-((*S*)-1-phenylethoxy)penta-1,3-dien-3-yl pivalate). See also ref 20.

(29) Bartoli, G.; Bosco, M.; Marcantoni, E.; Massaccesi, M.; Rinaldi, S.; Sambri, L. *Eur. J. Org. Chem.* **2001**, 4679–4684.

(30) See Supporting Information.

(31) Evans, D. A.; Allison, B. D.; Yang, M. G.; Masse, C. E. *J. Am. Chem. Soc.* **2001**, *123*, 10840–10852.

spectra of the corresponding acetone (+)-**39** obtained by treatment of (+)-**38** with (MeO)₂CMe₂ and catalytic TsOH·H₂O (96%). It was furthermore verified by single-crystal X-ray diffraction studies of crystalline ketone (–)-**40** obtained by treatment of (+)-**39** with MeLi·LiBr in DME/Et₂O (86%).³⁰ Ozonolysis of enol ester (+)-**38** and subsequent treatment with Me₂S provided crystalline lactone (–)-**7**, the structure of which was also proven by single-crystal X-ray diffraction studies.³⁰

Conclusion

Fully substituted 4-hydroxy- δ -lactones containing up to four continuous stereocenters can be prepared applying the oxyallylation of enoxysilanes through SO₂–Umpolung of enantiomerically enriched (1*E*,3*Z*)-1-(1-phenylethoxy)penta-1,3-dien-3-yl carboxylates. These stereotriads and stereotetrad are common motifs in a large number of natural polyketides and polypropionates. The number of synthetic steps, yields, and availability of starting materials are comparable with other well-accepted methods. The present methodology offers an alternative approach and extends the toolbox of chemists chasing cyclic polypropionate structures. We were able to obtain lactone (+)-**6** for the first time using chemical synthesis^{6a} as well as the yet unknown (–)-(3*R*,4*R*,5*S*,6*R*)-4-hydroxy-6-isopropyl-3,5-dimethyltetrahydro-2*H*-pyran-2-one ((–)-**7**).

Experimental Section

(+)-(3*R*,4*R*,5*S*,6*R*)-6-Ethyl-4-hydroxy-3,5-dimethyltetrahydro-2*H*-pyran-2-one ((+)-**6**). O₃ was bubbled into a soln of (–)-**33** (12 mg, 0.037 mmol) in CH₂Cl₂ (2 mL) at –78 °C until persistence of the blue color. After the disappearance of (–)-**33** by TLC, Me₂S (0.1 mL) was added and the mixture stirred at –78 °C for 20 min. The mixture was allowed to warm to 25 °C. Solvent evaporation and flash chromatography on silica gel (PE/EtOAc) gave (+)-**6** (5 mg, 73%) as colorless crystals (X-ray, Supporting Information). *R*_f = 0.23 (PE/EtOAc, 3:2). Mp 79–81 °C. $\alpha^{25}_{\text{D}} = +21$ (CHCl₃, *c* = 0.16). IR (film): ν (cm^{–1}) = 3433, 2971, 2938, 2882, 1707 (s), 1461, 1380, 1212, 1172, 1119, 992, 974. ¹H NMR (CDCl₃, 400 MHz): (ppm) = 1.02 (t, 3H, ³*J* = 7.5 Hz), 1.07 (d, 3H, ³*J* = 6.5 Hz), 1.33 (d, 3H, ³*J* = 7.0 Hz), 1.55 (ddq, 1H, ³*J* = 7.0, 7.5 Hz), 1.82 (ddq, 1H, ³*J* = 3.0, 7.0, 7.5 Hz), 1.87–1.93 (m, 1H), 2.53 (dq, 1H, ³*J* = 3.0, 7.0 Hz), 3.84 (br s, 1H), 4.36 (dd, 1H, ³*J* = 3.0, 7.0, 10.5 Hz). ¹³C NMR (CDCl₃, 100.6 MHz): δ (ppm) = 8.7, 12.9, 14.3, 26.1, 37.7, 42.6, 73.3, 82.1, 173.8. ESI-HRMS: *m/z* calcd for C₉H₁₇O₃⁺ 173.1178, found 173.1175 [M + H⁺].

(–)-(3*S*,4*S*,5*R*,6*S*)-4-Hydroxy-6-isopropyl-3,5-dimethyltetrahydro-2*H*-pyran-2-one ((–)-**7**). O₃ was bubbled through a soln of (–)-**38** (200 mg, 0.62 mmol) in CH₂Cl₂ (5 mL) at –78 °C until persistence of the blue color, then O₂ was bubbled. After the disappearance of (–)-**38** by TLC, Me₂S (0.25 mL) was added and the mixture was allowed to warm to 25 °C overnight. Water (10 mL) was added and the aq phase was extracted with CH₂Cl₂ (3 × 10 mL). The combined organic extracts were washed with brine (2 × 10 mL) and dried (Na₂SO₄), and the solvent was evaporated. Flash chromatography on silica gel (CH₂Cl₂/EtOAc) gave pure (–)-**7** (110 mg, 87%), that was recrystallized from hexane (X-ray, Supporting Information). *R*_f = 0.45 (PE/EtOAc, 7:3). Mp 92–95 °C. $\alpha^{25}_{\text{D}} = -28$ (CHCl₃, *c* = 0.40). IR (film): ν (cm^{–1}) = 3370, 3260, 2970, 2920, 1725, 1695, 1465, 1370, 1345, 1220, 1195, 1170, 1120, 990. ¹H NMR (CDCl₃, 400 MHz): (ppm) = 0.89 (d, 3H, ³*J* = 7.1 Hz), 1.06, 1.11 (2d, 6H, ³*J* = 7.0 Hz), 1.32 (d, 3H, ³*J* = 7.1 Hz), 1.87 (sept, 1H, ³*J* = 7.0 Hz), 1.95 (dq, 1H, ³*J* = 6.6, 10.7 Hz), 2.51 (q, 1H, ³*J* = 7.1 Hz), 3.85 (s, 1H), 4.29 (d, 1H, ³*J* = 10.7 Hz). ¹³C NMR (CDCl₃, 100.6 MHz): δ (ppm) = 12.7,

14.0, 10.0, 28.9, 36.0, 42.3, 73.2, 84.8, 173.9. ESI-HRMS: *m/z* calcd for C₁₀H₁₉O₃⁺ 187.1334, found 187.1336 [M + H⁺].

(2*Z*,4*R*,5*R*,6*R*,7*R*)-5,7-Dihydroxy-4,6-dimethylnon-2-en-3-yl 2-methylpropanoate ((–)-**33**). One molar TiCl₄ in CH₂Cl₂ (3 mL, 3.0 mmol) was added quickly to a stirred soln of (+)-(1*Z*,2*R*,3*S*,4*S*)-1-ethylidene-2,4-dimethyl-5-oxo-3-[(1*R*)-1-phenylethoxy]heptyl-2-methylpropanoate ((+)-**25**) (550 mg, 1.47 mmol). After stirring at –78 °C for 1 h, 1 M BH₃·Me₂S in CH₂Cl₂ (6.7 mL, 6.7 mmol) was added, and the mixture was stirred at –78 °C for two more hours. The reaction mixture was quenched with a sat. aq soln of NaHCO₃ (15 mL). The mixture was extracted with EtOAc (3 × 10 mL). The combined organic extracts were washed with brine (15 mL) and dried (MgSO₄). Solvent evaporation and flash chromatography on silica gel (PE/EtOAc) gave (–)-**33** (269 mg, 67%) as colorless oil. *R*_f = 0.29 (PE/EtOAc, 4:1). $\alpha^{25}_{\text{D}} = -26$ (CHCl₃, *c* = 0.58). IR (film): ν (cm^{–1}) = 3438, 2970, 2935, 2876, 1746 (s), 1691, 1459, 1408, 1337, 1240, 1138, 968. ¹H NMR (CDCl₃, 400 MHz): (ppm) = 0.93 (d, 3H, ³*J* = 7.0 Hz), 0.97 (t, 3H, ³*J* = 7.5 Hz), 1.08 (d, 3H, ³*J* = 6.5 Hz), 1.26 (d, 6H, ³*J* = 7.0 Hz), 1.34–1.49 (m, 1H), 1.46 (d, 3H, ³*J* = 6.5 Hz), 1.58–1.78 (m, 2H), 2.64–2.76 (m, 2H), 3.19 (dd, 1H, ³*J* = 5.0 Hz, ³*J* = 7.5 Hz), 3.51–3.61 (m, 1H), 5.25 (q, 1H, ³*J* = 7.0 Hz). ¹³C NMR (CDCl₃, 100.6 MHz): (ppm) = 10.1, 10.9, 15.3, 16.6, 19.2, 19.3, 27.9, 34.3, 40.1, 44.4, 76.5, 78.5, 114.6, 149.0, 176.2. ESI-HRMS: calcd for C₁₅H₂₉O₄⁺ 273.2066, found 273.2059 [M + H⁺].

(+)-(2*Z*,4*S*,5*S*,6*S*,7*S*)-5,7-Dihydroxy-4,6,8-trimethylnon-2-en-3-yl benzoate ((+)-**38**). To a soln of (2*Z*,4*S*,5*S*,6*S*,7*S*)-7-hydroxy-4,6,8-trimethyl-5-[(1*S*)-1-phenylethoxy]non-2-en-3-yl benzoate ((–)-**37**) (277 mg, 0.65 mmol) in CH₂Cl₂ (50 mL) was added a solution of FeCl₃ (0.2 g, 1.3 mmol) in 20 mL of CH₂Cl₂. The resulting mixture was stirred vigorously for 10 min at 25 °C, and H₂O was added. The aq phase was extracted with CH₂Cl₂ (3 × 30 mL). The combined organic layers were washed with brine (50 mL), dried (Na₂SO₄), filtered, and evaporated. Flash chromatography on silica gel (PE/EtOAc, 9:1) gave (+)-**38** (195 mg, 93%) as colorless oil. *R*_f = 0.48 (PE/AcOEt, 8:2). $\alpha^{25}_{\text{D}} = +16$ (CHCl₃, *c* = 0.22). IR (film): ν (cm^{–1}) = 3441, 3063, 2963, 2930, 2875, 1722, 1715, 1694, 1601, 1462, 1453, 1261, 1176, 1165, 1142, 1105, 1069, 1026. ¹H NMR (CDCl₃, 400 MHz): (ppm) = 0.88, 0.88 (2d, 6H, ³*J* = 6.8 Hz), 1.00, 1.19 (2d, 6H, ³*J* = 6.8 Hz), 1.53 (d, 3H, ³*J* = 6.8 Hz), 1.79–1.90 (m, 2H), 2.91 (quint, 1H, ³*J* = 7.4 Hz), 3.32 (dd, 1H, ³*J* = 7.4, 4.3 Hz), 3.50 (dd, 1H, ³*J* = 8.6, 3.1 Hz), 5.38 (q, 1H, ³*J* = 7.4 Hz), 7.51 (t, 2H, ³*J* = 7.4 Hz), 7.64 (t, 1H, ³*J* = 7.4 Hz), 8.13 (d, 2H, ³*J* = 8.0 Hz). ¹³C NMR (CDCl₃, 100.6 MHz): (ppm) = 11.1, 14.7, 15.4, 16.9, 20.4, 30.2, 37.6, 44.2, 78.7, 79.7, 114.9, 128.7, 129.1, 130.2, 133.8, 149.3, 165.5. ESI-HRMS: calcd for C₁₉H₂₈O₄Na⁺ 343.1885; found 343.1895 [M + Na⁺].

(+)-(2*Z*,4*S*)-4-[(4*S*,5*S*,6*S*)-2,2,5-Trimethyl-6-(propan-2-yl)-1,3-dioxan-4-yl]pent-2-en-3-yl benzoate ((+)-**39**). To a soln of diol (+)-**38** (190 mg, 0.59 mmol) in dimethoxypropane (2 mL) was added *p*-TsOH·H₂O (5.6 mg, 0.03 mmol). The mixture was stirred for 1 h at 25 °C, then neutralized by adding solid NaHCO₃, filtered, and evaporated. Flash chromatography on silica gel (PE/EtOAc, 9:1) gave (+)-**39** (200 mg, 96%) as a colorless oil. $\alpha^{25}_{\text{D}} = +6$ (CHCl₃, *c* = 0.30). IR (film): ν (cm^{–1}) = 2964, 2930, 2875, 2849, 1738, 1687, 1602, 1492, 1453, 1390, 1378, 1261, 1201, 1174, 1154, 1133, 1105, 1026. ¹H NMR (CDCl₃, 400 MHz): (ppm) = 0.72 (d, 3H, ³*J* = 6.5 Hz), 0.88, 0.93 (2d, 6H, ³*J* = 6.4 Hz), 1.18 (s, 3H), 1.21 (d, 3H, ³*J* = 7.1 Hz), 1.30 (s, 3H), 1.54 (d, 3H, ³*J* = 6.8 Hz), 1.80–1.90 (m, 2H), 2.68 (qd, 1H, ³*J* = 7.4, 1.8 Hz), 3.32 (dd, 1H, ³*J* = 10.4, 1.8 Hz), 3.40 (dd, 1H, ³*J* = 9.8, 2.4 Hz), 5.33 (q, 1H, ³*J* = 6.8 Hz), 7.48 (t, 2H, ³*J* = 7.4 Hz), 7.58 (t, 1H, ³*J* = 7.4 Hz), 8.13 (d, 2H, ³*J* = 8.0 Hz). ¹³C NMR (CDCl₃, 100.6 MHz): (ppm) = 10.8, 11.4, 14.1, 15.7, 18.8, 19.8, 27.8, 29.5, 29.7, 32.6, 39.1, 76.4, 77.0, 97.3, 112.7, 128.0, 129.7, 132.6, 148.8, 163.5. ESI-HRMS: calcd for C₂₂H₃₂O₄K⁺ 399.1938; found 399.1937 [M + K⁺]. Anal. calcd for C₂₂H₃₂O₄ (360.49): C, 73.30%; H, 8.95%; O, 17.75%. Found C, 73.22%; H, 8.85%; O, 17.74%.

(-)-(4*S*,5*S*,6*S*,7*S*)-5,7-Isopropylidendioxy-4,6,8-trimethylnona-3-one (-)-**40**. A soln of (+)-**39** (1.00 g, 2.78 mmol) in DME (15 mL) was added to a soln of MeLi·LiBr (2.1 M in Et₂O, 6.6 mL, 13.9 mmol) in Et₂O (10 mL) at -78 °C. The mixture was stirred at -78 °C for 5 h, poured into an ice-cold sat. aq soln of NH₄Cl (30 mL). The aq phase was extracted with Et₂O (4 × 20 mL). The organic layers were washed with brine (20 mL), dried (Na₂SO₄), and evaporated. The residue was purified by recrystallization from hexane, giving (-)-**40** (615 mg, 86%) as colorless crystals. *R*_f = 0.52 (PE/AcOEt, 9:1). Mp = 89–92 °C. α_D²⁵ = -20 (CHCl₃, *c* = 0.15). IR (*film*): ν (cm⁻¹) = 2975, 2959, 2938, 2875, 2841, 1693, 1458, 1412, 1378, 1358, 1346, 1249, 1198, 1164, 1152, 1130, 1101, 1048. ¹H NMR (CDCl₃, 400 MHz): (ppm) = 0.75, 0.80 (2d, 6H, ³*J* = 6.7 Hz), 0.91 (d, 3H, ³*J* = 6.9 Hz), 1.01 (t, 3H, ³*J* = 7.1 Hz), 1.19 (d, 3H, ³*J* = 7.3 Hz), 1.31, 1.36 (2s, 6H), 1.44 (m, 1H), 1.85 (sept, 1H, ³*J* = 6.6 Hz), 2.45, 2.59 (2qd, 2H, ²*J* = 18.1 Hz, ³*J* = 7.2 Hz), 2.71 (m, 1H), 3.27 (d, 1H, ³*J* = 10.0 Hz), 3.56 (d, 1H, ³*J* = 10.3 Hz). ¹³C NMR (CDCl₃, 100.6 MHz): (ppm) = 7.4, 7.6, 11.7, 13.7, 14.2, 19.1, 20.1, 27.9, 29.9, 34.1, 49.7, 76.9, 77.6, 97.8, 214.2.

ESI-HRMS: *m/z* calcd for C₁₅H₂₈O₃Na⁺ 279.1936, found 279.1939 [M + Na⁺].

Acknowledgment. This work was supported by the Swiss National Science Foundation (Grants 200020-116212/1 and 200020_124724) and by the Roche Research Foundation. We are grateful also to Dr. C. Mazet, University of Geneva, Dr. L. Menin, F. Sepulveda, and M. Rey for technical assistance, and Dr. R. Scopelliti for the X-ray diffraction studies.

Supporting Information Available: Further complete experimental procedures and compound characterization data for (+)-**1**, (-)-**2**, (+)-**3**, (-)-**4**, (-)-**5**, **16a**, **16b**, **18a**, **18b**, **20a**, **20b**, (+)-**27**, **28**, (+)-**29**, (-)-**31**, (+)-**32**, (-)-**33** as well as copies of ¹H and ¹³C NMR spectra for all newly reported compounds. X-ray data for (-)-**5**, (+)-**6**, (-)-**7**, and (-)-**40** have been deposited at the Cambridge crystallographic database. This material is available free of charge via the Internet at <http://pubs.acs.org>.