

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/11025661>

Synthesis of Di-Branched Heptasaccharide by One-Pot Glycosylation Using Seven Independent Building Blocks

ARTICLE *in* ORGANIC LETTERS · DECEMBER 2002

Impact Factor: 6.36 · DOI: 10.1021/ol020150+ · Source: PubMed

CITATIONS

50

READS

14

6 AUTHORS, INCLUDING:



Hiroshi Tanaka

Tokyo Institute of Technology

96 PUBLICATIONS 1,327 CITATIONS

SEE PROFILE



Hirokazu Tsukamoto

Tohoku University

63 PUBLICATIONS 813 CITATIONS

SEE PROFILE



Pergamon

N-Acyl 6,7-Dimethoxy-1,2,3,4-tetrahydroisoquinoline: The First Orexin-2 Receptor Selective Non-peptidic Antagonist

Masaaki Hirose*, Shin-ichiro Egashira, Yasuhiro Goto, Takashi Hashihayata, Norikazu Ohtake, Hisashi Iwaasa, Mikiko Hata, Takehiro Fukami, Akio Kanatani and Koji Yamada

Banyu Tsukuba Research Institute, Banyu Pharmaceutical Co., Ltd, Okubo 3, Tsukuba 300-2611, Ibaraki, Japan

Received 16 June 2003; accepted 15 August 2003


Abstract—The identification of potent and selective orexin-2 receptor (OX₂R) antagonists is described based on the modification of *N*-acyl 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline analogue **1**, recently discovered during high throughput screening (HTS). Substitution of an acyl group in **1** with *tert*-Leucine (*tert*-Leu), and introduction of a 4-pyridylmethyl substituent onto the amino function of *tert*-Leu improved compound potency, selectivity, and water solubility. Thus, compound **29** is a promising tool to investigate the role of orexin-2 receptors.

© 2003 Elsevier Ltd. All rights reserved.

Orexin-A and -B are the novel neuropeptides that have been identified as ligands of orphan G protein coupled receptors (GPCR) from rat brain extracts.¹ Evidence continues to indicate that they are involved in the regulation of many neuronal functions, including feeding,¹ sleep/wake cycles,^{2,3} neuroendocrine activity,^{4,5} stress reactions,⁶ and activation of the sympathetic nerve system.⁷ The innervation of orexin neurons in the brain supports diversity in function of the orexins.⁸ The physiological functions of orexins are evoked by two receptor subtypes, orexin-1R (OX₁R) and orexin-2R (OX₂R) which belong to a family of GPCR. Orexin-A is a 33-residue peptide with two intra-molecular disulfide bridges, which possessed potent agonistic activity toward both subtypes of orexin receptors. In contrast, orexin-B, which consists of 28-amino acids, possessed greater affinity for OX₂R than for OX₁R.¹ Moreover, the expression patterns of the two orexin receptors in the brain are different.⁹ The orexin receptor subtypes are likely to be involved in different pharmacological functions. Recently, a small molecular OX₁R selective antagonist was developed as a potential tool to elucidate the functions of OX₁R.¹⁰ Selective agonist of OX₂R, which can address the pharmacological functions of OX₂R, were recently identified;¹¹ however, OX₂R selective antagonists remain critical to understanding the

physiological and pathophysiological roles of OX₂R. Therefore, the development of small molecular OX₂R selective antagonists for use as pharmacological tools is important for clarification of the functions of OX₂R. This communication describes the development of *N*-acyl 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline analogues as the first OX₂R selective non-peptidic antagonist and reports progress in optimizing this structural class, leading to compound **29**, which possesses potency, selectivity, and water solubility. As a part of our program to develop selective OX₂R antagonists, we discovered the non-selective orexin antagonist **1** (human OX₁R (hOX₁R) IC₅₀: 7 μM; human OX₂R (hOX₂R) IC₅₀: 2 μM) through HTS screening.¹²

Substitution and modification of the 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline skeleton in **1** with other acyclic and cyclic amine moieties resulted in loss of potency against both hOX₁R and hOX₂R (data not shown). Thus, the 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline skeleton seems to be an essential core for retaining biological activity. Based on these results, our attention focused on modification of the acyl portion of **1**.



*Corresponding author. Tel.: +81-298-77-2000; fax: +81-298-77-2029; e-mail: hirosems@banyu.co.jp


SAR of 3-Phenylpropanoyl Amides

To understand the structure–activity relationship (SAR) of **1**, initially we prepared *N*-(3-phenylpropanoyl) 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline analogues. Removal of the bromine from the benzene ring of **1** resulted in a 30-fold increase in potency toward hOX₂R (**2**). Removal of the *tert*-butyl group from **2** afforded **3**, which did not possess antagonistic activity against either orexin receptor. The results of functional group substitutions for the *tert*-butyl group of **2** are listed in Table 1. A dimethylamino or benzylamino group at this position was not allowed (**5** and **6**), suggesting that neutral bulky substituents at this position are indispensable for antagonistic activity toward hOX₂R. Although replacement of the *tert*-butyl moiety with a benzoylamino group (**7,8**) led to significant reduction of activity toward hOX₂R compared to **2**, the (*S*)-isomer (**8**) showed relatively high potency than did its racemate (**7**). Introduction of neutral bulky substituent(s) on the benzoyl group in **8** improved potency toward hOX₂R; thus, the (*S*)-3,5-dichlorobenzoylamino analogue (**10**) brought about a 6-fold improvement in hOX₂R potency relative to **8**.

SAR of *N*-Substituted *tert*-Leu Amides

To improve water solubility and develop useful pharmacological tools, we modified the benzyl group in **1**. Introduction of hydrophilic substituents such as a dimethyl amino group on the benzene ring of **1** resulted in loss of potency toward both orexin receptors (data not shown). In some cases, substitution of the benzyl group with a substituted amino group afforded *tert*-Leu amide analogues and resulted in greater water solubility without loss of potency or selectivity (Table 2), especially in the case of the *N*-benzyl substituent (**15**). Reduction or elongation of the carbon chain between the nitrogen atom and phenyl group in **15** resulted in

Table 1. Antagonistic activity of 1-(3-phenylpropanoyl)-tetrahydroisoquinoline analogues against hOX₁R and hOX₂R^a




No.	R ¹	R ²	hOX ₁ R	hOX ₂ R
			(IC ₅₀ , nM) ^b	
1	Br	<i>tert</i> -Butyl	7000	2000
2	H	<i>tert</i> -Butyl	5500	56
3	H	H	> 10,000	> 10,000
4	H	Phenyl	> 10,000	395
5	H	Dimethylamino	> 10,000	> 10,000
6	H	Benzylamino ^c	> 10,000	> 10,000
7	H	Benzoylamino	> 10,000	900
8	H	Benzoylamino ^c	> 10,000	195
9	H	3,5-Dichlorobenzoylamino	5300	36
10	H	3,5-Dichlorobenzoylamino ^c	2300	30
11	H	3-Bromo-4-fluorobenzoylamino	5900	70

^aAll compounds were synthesized as racemic mixture otherwise noted.

^bValues are the mean of more than two independent experiments performed in duplicate.

^c(*S*)-Isomer.

Table 2. Antagonistic activity of 1-(*tert*-leucyl)-tetrahydroisoquinoline analogues against hOX₁R and hOX₂R^a



No.	R	hOX ₁ R	hOX ₂ R
		(IC ₅₀ , nM) ^b	
12	H	> 10,000	> 10,000
13	Phenyl	> 10,000	2100
14	Benzoyl	> 10,000	> 10,000
15	Benzyl	> 10,000	910
16	Benzyl ^c	3850	130
17	Benzyl ^d	> 10,000	2900
18	2-Phenylethyl	> 10,000	> 10,000
19	3-Phenylpropyl	> 10,000	> 10,000
20	Cyclohexylmethyl	> 10,000	> 10,000

^aAll compounds were synthesized as racemic mixture otherwise noted.

^bValues are the mean of more than two independent experiments performed in duplicate.

^c(*S*)-Isomer.


^d(*R*)-Isomer.

loss of potency (**12**, **13**, **18** and **19**). Substitution of a benzyl group in **15** for a cyclohexylmethyl group also resulted in complete loss of potency (**20**). Thus, aromatic functionality in the *N*-benzyl group seemed to be essential. Regarding to the stereochemistry of the *tert*-Leu portion, (*S*)-configuration is more preferable than (*R*)-configuration (**16** vs **17**).

None of the other *N*-benzyl amino acid analogues, such as (*N*-benzyl)phenylalanine **6**, *N*-benzyl-valine **21** or (*N*-benzyl)phenylglycine **22**, possessed greater potency toward hOX₂R than the *tert*-Leu analogue **15** (Table 3). Only the *N*-benzylvaline **21** analogue showed weak activity against hOX₂R (IC₅₀: 3300 nM).

Substitution of the benzyl group in **16** with other arylmethyl groups improved potency (Table 4). Replacement of the benzene ring in **16** with five-membered heteroaromatic rings generally improved hOX₂R potency. In particular, the 2-thienyl analogue **25** led to a

Table 3. Antagonistic activity of *N*-benzyl α-aminoacyl tetrahydroisoquinoline analogues against hOX₁R and hOX₂R^a




No.	R	hOX ₁ R	hOX ₂ R
		(IC ₅₀ , nM) ^b	
6	Benzyl ^c	> 10,000	> 10,000
15	<i>tert</i> -Butyl	> 10,000	910
16	<i>tert</i> -Butyl ^c	3850	130
21	<i>i</i> -Propyl	> 10,000	3300
22	Phenyl ^c	> 10,000	> 10,000

^aAll compounds were synthesized as racemic mixture otherwise noted.

^bValues are the mean of more than two independent experiments performed in duplicate.

^c(*S*)-Isomer.

Table 4. Antagonistic activity of *N*-arylmethyl (*S*)-*tert*-leucyl tetrahydroisoquinoline analogues against hOX₁R and hOX₂R



No.	Ar	hOX ₁ R	hOX ₂ R
		(IC ₅₀ , nM) ^a	
16	Phenyl	3850	130
23	2-(<i>N</i> -Methyl)pyrrolyl	3600	28
24	2-Thiazolyl	> 10,000	59
25	2-Thienyl	1130	25
26	3-Thienyl	3670	35
27	2-Pyridyl	> 10,000	1400
28	3-Pyridyl	> 10,000	240
29	4-Pyridyl	> 10,000	40

^aValues are the mean of more than two independent experiments performed in duplicate.

5-fold improvement in the hOX₂R potency; however, its selectivity against hOX₂R over the hOX₁R was decreased compared to **16**. Nitrogen containing hetero-aromatic analogues were generally showed high OX₁R/OX₂R selectivity. For the pyridine analogues, hOX₂R potency depended heavily on the position of the nitrogen. 4-Pyridyl methyl analogue **29** possessed the most potent activity for hOX₂R compared to other regio isomers (**27** and **28**). Furthermore, **29** showed over 250-fold selectivity for hOX₂R compared with hOX₁R (Table 4), as well as over 50 receptors, ion channels, and transporters (<30% inhibition at 10 μM), which includes G-protein coupled receptors associated with food intake including galanin and neuropeptide Y (data not shown). The high water solubility of **29** was another benefit (0.81 mg/mL at pH 7) to use pharmacological experiment. Full details of additional pharmacological testing of **29** will be described elsewhere.

Synthesis of Compound 29

Compound **29** was prepared according to Scheme 1. Commercially available 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline hydrogen chloride was condensed with *N*-Boc *tert*-leucine using PyBrop¹³ in pyridine in the presence of DMAP followed by treatment of HCl/AcOEt to yield the amine hydrogen chloride **30**. Conversion of **30** to the 4-pyridylmethyl analogue **29** was accomplished by the standard reductive amination procedure using isonicotinaldehyde and sodium triacetoxy-

**Scheme 1.** Synthesis of compound **29**.

borohydride.¹⁴ Other compounds described here were synthesized by similar methods.

In summary, in this communication, the structure–activity relationship of the tetrahydroisoquinoline **1** toward hOX₂R was outlined. 6,7-Dimethoxy-1,2,3,4-tetrahydroisoquinoline was the essential core skeleton for hOX₁R and hOX₂R potency. *N*-Arylmethyl *tert*-leucyl 6,7-dimethoxy-1,2,3,4-tetrahydroisoquinoline analogues generally showed high potency for the hOX₂R. Their selectivity could be enhanced by modifying the arylmethyl motif of the amino functionality. Nitrogen-containing hetero-aromatics were superior for improving both potency and selectivity toward hOX₂R. Finally, introduction of a 4-pyridylmethyl group on the amino function of the *tert*-Leu moiety yielded a potent hOX₂R antagonist **29** with high selectivity and high water solubility.

References and Notes

- Sakurai, T.; Amemiya, A.; Ishii, M.; Matsuzaki, I.; Chemelli, R. M.; Tanaka, H.; Williams, S. C.; Richardson, J. A.; Kozlowski, G. P.; Wilson, S.; Arch, J. R. S.; Buckingham, R. E.; Haynes, A. C.; Carr, S. A.; Annan, R. S.; McNulty, D. E.; Liu, W.-S.; Terrett, J. A.; Elshourbagy, N. A.; Bergsma, D. J.; Yanagisawa, M. *Cell* **1998**, 92, 573.
- Lin, L.; Faraco, J.; Li, R.; Kadotani, H.; Rogers, W.; Lin, X.; Qiu, X.; de Jong, P. J.; Nishino, S.; Mignot, E. *Cell* **1999**, 98, 365.
- Chemelli, R. M.; Willie, J. T.; Sinton, C. M.; Elmquist, J. K.; Scammell, T.; Lee, C.; Richardson, J. A.; Williams, S. C.; Xiong, Y.; Kisanuki, Y.; Fitch, T. E.; Nakazato, M.; Hammer, R. E.; Saper, C. B.; Yanagisawa, M. *Cell* **1999**, 98, 437.
- Pu, S.; Jain, M. R.; Kalra, P. S.; Kalra, S. *Regul. Pept.* **1998**, 78, 133.
- Jaszberenyi, M.; Bujdoso, E.; Pataki, I.; Telegdy, G. *J. Neuroendocrinol.* **2000**, 12, 1174.
- Ida, T.; Nakahara, K.; Katayama, T.; Murakami, N.; Nakazato, M. *Brain Res.* **1999**, 821, 526.
- Antunes, V. R.; Brailoiu, G. C.; Kwok, E. H.; Scruggs, P.; Dun, N. J. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **2001**, 281, R1801.
- Nambu, T.; Sakurai, T.; Mizukami, K.; Hosoya, Y.; Yanagisawa, M.; Goto, K. *Brain Res.* **1999**, 827, 243.
- Trivedi, P.; Yu, H.; MacNeil, D. J.; Van der Ploeg, L. H. T.; Guan, X.-M. *FEBS Lett.* **1998**, 438, 71.
- Porter, R. A.; Chan, W. N.; Coulton, S.; Johns, A.; Hadley, M. S.; Widdowson, K.; Jerman, J. C.; Brough, S. J.; Coldwell, M.; Smart, D.; Jewitt, F.; Jeffrey, P.; Austin, N. *Bioorg. Med. Chem. Lett.* **2001**, 11, 1907.
- Asahi, S.; Egashira, S.; Matsuda, M.; Iwaasa, H.; Kanatani, A.; Ohkubo, M.; Ihara, M.; Morishima, H. *Bioorg. Med. Chem. Lett.* **2003**, 13, 111.
- Brief screening protocol: CHO-K1 cells stably expressing hOX₁R or hOX₂R were seeded into 96-well plates and incubated with a cytoplasmic calcium indicator, Fluo-3 AM. After the cells were washed four times, the intracellular Ca²⁺ mobilization evoked by 0.3 nM of orexins-A was monitored as a change in cell fluorescence intensity by FLIPR (Molecular Devices). Varying concentration of orexin antagonists were added to the plate 5 min prior to the addition of orexin-A. The antagonistic activities were calculated as IC₅₀ values.
- Coste, J.; Le-Nguyen, D.; Castro, B. *Tetrahedron Lett.* **1990**, 31, 205.
- Characterization of **29**: ¹H NMR (300 MHz, CDCl₃) δ 0.97 and 1.02 (s each, 9H), 2.61–2.90 (m, 2H), 3.20–3.51 (m, 3H), 3.62–4.13 (m, 8H), 4.31–4.99 (m, 2H), 6.40, 6.60, 6.60 and 6.65 (s each, 2H), 7.09–7.32 (m, 2H), 8.40–8.56 (m, 2H).