

An Asymmetric Synthesis of Chiral Nifedipine Analogues


A. I. Meyers* and Thomas Oppenlaender

Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523, U.S.A.

The addition of aryl-lithium reagents to chiral pyridyl-dihydro-oxazoles gave (S)-4-aryl-1,4-dihydropyridines in 80–90% enantiomeric excess.

The recent interest in 4-aryl-1,4-dihydropyridines as potent calcium channel blockers has led to the therapeutic use of nifedipine (**1**) for the treatment of angina pectoris and hypertension.¹ Extensive structure–activity relationships have shown that the (–)-enantiomer exhibits enhanced pharmacological properties. Furthermore, a recent report² describes

opposite action (Ca^{2+} blocker or Ca^{2+} enhancer) for each enantiomer as a function of the absolute stereochemistry at the 4-position of the dihydropyridine. We now describe an asymmetric addition to the 4-position of the pyridine nucleus³ (**5**) which carries a chiral dihydro-oxazole at the 3-position leading to 4-aryl-1,4-dihydropyridines (**2**) in 80–90% enan-



Scheme 1. Reagents: (a) MeCOCl-EtOH, 0 °C; (b) (S,S)-PhCH(OH)CH(NH₂)CH₂OMe, ClCH₂CH₂Cl, Et₃N, reflux; (c) FSO₃Me, CH₂Cl₂, 20 °C, NaBH₄, 0 °C; 50% aq. oxalic acid-THF, 25 °C; (d) NaBH₄, THF-EtOH (2:1); (e) 20% aq. KOH in THF; (f) phase-transfer catalysis: KOH (solid), BuⁿBr, THF, Buⁿ₄NBr, 20 °C.

tiomeric excess (e.e.) and opens a route to chiral, non-racemic nefidipine analogues (2) (Scheme 1).

The synthetic route starts with 3-cyano-5-methoxycarbonylpyridine (3) which was transformed initially to the imidate (4) and then to the chiral 3-dihydro-oxazolyl-5-methoxycarbonyl pyridine (5) (oil 76%, [α]_D²⁴ -49.5°, *c* 0.92, CHCl₃). Addition of aryl-lithium reagents (1.2 equiv.) to a tetrahydro-

furan (THF) solution of (5) (0.01–0.05 M) during 2 h at -78 °C and continued stirring (3–4 h) gave a green fluorescing solution which was then treated with methyl chloroformate (5 equiv.) at -78 °C. Aqueous work-up and chloroform extraction produced (6a)[†] or (6b)[†] in diastereoisomeric ratios of 89:11 and 95:5, respectively. The ratios were determined *via* reverse-phase h.p.l.c. (20% MeOH-H₂O). The chiral auxiliary was removed from (6a) *via* the previously reported method of quaternization–reduction–hydrolysis⁴ to give the aldehyde (S)-(7)[†] (oil, 76%, [α]_D²⁴ -4.7°, *c* 1.66, CHCl₃). Reduction to the 3-hydroxymethylpyridine (8) was accomplished with sodium borohydride and removal of the *N*-methoxycarbonyl group using 20% aqueous KOH afforded (S)-(9)[†] (62%, m.p. 152–153 °C, [α]_D²⁴ -116.3°, *c* 0.44, MeCN). Alternatively (S)-(7) was transformed, without isolation of (8), in an overall yield of 62%. *N*-Alkylation of (9) was carried out with *n*-butyl iodide under phase-transfer conditions⁵ furnishing (S)-(10)[†] in 75% yield (m.p. 88–88.5 °C, [α]_D²⁴ -125.7°, *c* 0.37, CHCl₃). Treatment of (S)-(10) with (S)-(+)-α-methoxy-α-trifluoromethylphenylacetyl chloride (pyridine, 0 °C) gave the Mosher ester,⁶ the ¹⁹F n.m.r. spectrum of which showed peaks at δ -72.02 and -72.17 p.p.m. in a ratio of 9:91. Thus, very little racemization at C-4 occurred during the manipulation of the substituents at the 3-position. In summary, we have described a route to nefidipine analogues (6)–(10) with high enantiomeric excess which are reached through simple transformations and could prove significant in pharmacological studies.

The authors thank the National Institutes of Health for funding and the DAAD (West Germany) for a NATO Fellowship (to T. O.). A. I. M. thanks the Alexander von Humboldt Foundation for a Senior Scientist Award (1984–1986) and the Faculty at the University of Wurzburg for their hospitality.

Received, 17th January 1986; Com. 082

References

- F. Bossert, H. Meyer, and E. Wehninger, *Angew. Chem., Int. Ed. Engl.*, 1981, **20**, 762.
- R. P. Hof, U. T. Ruegg, A. Hof, and A. Vogel, *J. Cardiovascular Pharmacol.*, 1985, **7**, 689.
- For related additions to pyridine-containing dihydro-oxazoles see: A. I. Meyers, N. R. Natale, D. G. Wettlaufer, S. Rafii, and J. Clardy, *Tetrahedron Lett.*, 1981, **22**, 5123 (X-ray structure of the MeLi adduct to chiral dihydro-oxazolyl-pyridines); A. I. Meyers and N. R. Natale, *Heterocycles*, 1982, **18**, 13; A. E. Hauck and G. S. Giam, *J. Chem. Soc., Perkin Trans. 1*, 1980, 2070.
- B. A. Barner and A. I. Meyers, *J. Am. Chem. Soc.*, 1984, **106**, 1865.
- J. Palecek and J. Kuthan, *Synthesis*, 1976, 550.
- J. A. Dale, D. L. Dull, and H. S. Mosher, *J. Org. Chem.*, 1969, **34**, 2543.

[†] Satisfactory elemental analyses were obtained.

‡ (6a): [α]_D²⁴ +73.5° (*c* 0.34, CHCl₃); i.r. (CHCl₃) 3052, 3035, 2985, 1751, 1712, 1682, 1638, 1620, 1445, and 1223 cm⁻¹; ¹H (CDCl₃) n.m.r. δ 8.06 (br.s, 1 H), 7.85 (br.s, 1 H), 7.45–7.00 (m, 10 H), 5.26 (d, *J* 6.4 Hz, 1 H), 5.06 (s, 1 H), 4.20–4.00 (m, 3 H), 3.95 (s, 3 H), 3.63–3.42 (m, 2 H), 3.38 (s, 3 H), and 1.22 (t, 3 H). Diastereotopic protons at δ 5.03 and 5.06 showed a ratio of 89:11. (6b): [α]_D²⁴ +119.7° (*c* 0.3, CHCl₃); i.r. (CCl₄) 3055, 3020, 2970, 2945, 2918, 1750, 1713, 1676, 1634, 1612, 1438, and 1218 cm⁻¹; ¹H n.m.r. (CDCl₃) δ 8.03 (br.s, 1 H), 7.82 (br.s, 1 H), 7.35–7.15 (m, 5 H), 7.04 (m, 2 H), 6.79 (m, 2 H), 5.26 (d, *J* 6.3 Hz, 1 H), 5.00 (s, 1 H), 4.30–4.00 (m, 3 H), 3.95 (s, 3 H), 3.77 (s, 3 H), 3.70–3.40 (AB m, 2 H), 3.39 (s, 3 H), and 1.23 (t, 3 H).