

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/221695443>

Competitive [2,3]- and [1,2]-Oxonium Ylide Rearrangements. Concerted or Stepwise?

ARTICLE *in* ORGANIC LETTERS · MARCH 2012

Impact Factor: 6.36 · DOI: 10.1021/ol300213u · Source: PubMed

CITATIONS

10

READS

32

5 AUTHORS, INCLUDING:



Deana M Jaber

Marymount University

5 PUBLICATIONS 122 CITATIONS

SEE PROFILE



Peter Y. Zavalij

University of Maryland, College Park

434 PUBLICATIONS 8,152 CITATIONS

SEE PROFILE



Michael Doyle

University of Texas at San Antonio

472 PUBLICATIONS 14,214 CITATIONS

SEE PROFILE

Published in final edited form as:

Org Lett. 2012 April 6; 14(7): 1676–1679. doi:10.1021/ol300213u.

Competitive [2,3]- and [1,2]-Oxonium Ylide Rearrangements. Concerted or Stepwise?

Deana M. Jaber[‡], Ryan N. Burgin[†], Matthew Helper[†], Peter Y. Zavalij[†], and Michael P. Doyle^{*,†}

Department of Chemistry and Biochemistry, University of Maryland, College Park, Maryland 20742, United States, and Department of Biology and Physical Sciences, Marymount University, Arlington, Virginia 22207, United States

Abstract

The axial-equatorial conformational isomer distribution of the reactant diazoacetoacetate or its metal carbene intermediate is reflected in Rh(II) catalyzed oxonium ylide forming reactions of 3- (*trans*-2-arylvinyl)tetrahydropyranone- 5-diazoacetoacetates that afford diastereoisomeric products for both the symmetry-allowed [2,3]- and the formally symmetry-forbidden [1,2]-oxonium ylide rearrangements.

We have reported that dirhodium (II) catalyzed reactions of racemic *trans*-aryl-substituted tetrahydropyranone diazoacetoacetates **1** produce two diastereoisomeric products (*syn*-**4** and *anti*-**4**) that are proposed to arise from two non-interconvertible oxonium ylide intermediates (**3A** and **3B**) via what appeared to be a concerted [1,2]-rearrangement (Scheme 1).¹ Product ratios were independent of ligand size or electron-withdrawing influence; and, although para-substituents on the aromatic ring affected the yields of the ylide rearrangement products, they had limited effect on their ratios (stereoselectivity). However, increasing the steric bulk of the aryl group (ortho substituents) led exclusively to the formation of one diastereoisomer and suggested that conformational isomers in the reactant and the intermediate metal carbene are responsible for the formation of diastereoisomeric products.

The mechanism of the [1,2]-rearrangement of oxonium ylides has been interpreted as occurring via either a cleavage/recombination (either homolytic or heterolytic) or a concerted pathway.² Because this reaction is formally considered to be symmetry forbidden,³ the rearrangement is regarded to occur by cleavage/recombination,⁴ although other mechanistic possibilities exist.^{5–7} However, convincing evidence regarding the cleavage/recombination pathway is scarce. The most relevant evidence for a homolytic pathway in oxonium ylide reactions comes from detection of a “homo-dimer” in the rhodium acetate catalyzed reactions of 4-benzyloxy-1-diazo-2-butanones,⁸ but similar ylide-producing reactions of the next higher homologue, 5-benzyloxy-1-diazo-2-pentanone, were not reported.⁹ More elaborate evidence for radical pair and/or zwitterionic intermediates comes from ammonium ylide [1,2]-rearrangements^{2,10} where, recently, substituent effects on stereoselectivity (from 4:1 to 20:1 d.r.) were obtained from aryl substituents of aryl diazoacetates in the rhodium acetate catalyzed reactions of configurationally stable

mdoyle3@umd.edu.


[†]University of Maryland

[‡]Marymount University

Supporting Information Available. General experimental procedures, X-ray structures of *syn*-**6a** and *syn*-**7a**, and spectroscopic data for all new compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.


Tröger bases.¹¹ Our proposed mechanism for oxonium ylide formation from **1** and subsequent [1,2]-Stevens rearrangement (Scheme 1) that suggests a concerted reaction is in conflict with a homolytic/heterolytic cleavage/recombination mechanism (i.e., absence of substituent effects, and steric influence on stereo-selectivity), although a very fast cleavage/recombination process that is barely distinguishable from a concerted reaction is also possible. We now report analogous investigations with compounds that are structurally similar to **1** whose oxonium ylides could undergo symmetry-allowed [2,3]-sigmatropic rearrangements that are not subject to orbital symmetry restrictions.

3-(*trans*-Styryl)tetrahydropyranone-5-diazoacetate and its para-substituted derivatives **5a-c** were selected to investigate the competition between the [2,3]-sigmatropic and [1,2]-Stevens rearrangement pathways. We anticipated that treatment of **5** with dinitrogen extrusion catalysts would form oxonium ylide(s), with the resultant reaction(s) leading to either or both products from the [2,3]-sigmatropic or the [1,2]-Stevens rearrangement, expecting the former to dominate.¹² Compounds **5** were prepared solely as the *trans*-diastereoisomers by BF₃·Et₂O-mediated hetero-Diels-Alder reactions between the cinnamaldehyde and Danishefsky's diene¹³ followed by zinc triflate catalyzed Mukaiyama-Michael reaction with methyl 3-(*tert*-butyldimethylsiloxy)-2-diazo-3-butenate (eq 1).¹⁴



(1)

Rhodium(II) catalyzed decomposition of **5a**, using 1.0 mol% of dirhodium octanoate [Rh₂(oct)₄] in refluxing dichloromethane, afforded a mixture of [2,3]-sigmatropic **6a** and [1,2]-Stevens **7a** rearrangement products (eq 2) in 47% and 41% yield, respectively (54:46 ratio). After careful chromatographic separations and spectroscopic analyses of the reaction mixture, two diastereoisomers for both rearrangement products were isolated and identified. The crystal structure of the chromatographically separated major isomer of the [1,2]-Stevens rearrangement product was obtained (Figure 1b), and like compound **4** (Scheme 1) the major isomer was the *syn*-isomer. The minor isomer of the [1,2]-Stevens rearrangement product and the major isomer of the [2,3]-sigmatropic rearrangement product were isolated together and characterized spectroscopically, while the minor isomer of the [2,3]-rearrangement sigmatropic product (*syn*-isomer) was isolated in a separate chromatographic fraction, and its crystal structure was also obtained (Figure 1a). Table 1 reports the negligible variance in the ratio of products from [2,3]- and [1,2]-oxonium ylide rearrangements and



(2)

in the ratios of diastereoisomers formed by each process from rhodium(II) octanoate catalyzed reactions with parasubstituted aryl derivatives **5a-b**. The lower yield and the difference in the ratio of products from the para-methoxy derivative **5c** is due to the formation of elimination products that, although evident in the reaction mixture, were not isolated and fully characterized.

We also investigated ligand effects from dirhodium catalysts on both the [2,3]-/[1,2]-reaction pathways (**6a:7a**) and product diastereoisomer ratios from the [2,3]- and [1,2]-rearrangements to determine if the oxonium ylide intermediate was metal-associated or a free ylide. Results were obtained from reactions of **5a** with a wide spectrum of catalysts (Table 2), and they show minimal dependence on catalyst ligands. No reaction occurred with rhodium caprolactamate under the same conditions. The diastereoisomer product ratio from the [2,3]-sigmatropic rearrangement (**anti-6a:syn-6a**), as well as that from the [1,2]-Stevens rearrangement (**syn-7a:anti-7a**), was invariant with common ligands on dirhodium (piv = pivalate, pfb = perfluorobutyrate, tfa = trifluoroacetate, tpa = triphenylacetate) that cover a broad range of electronic influences, although results with $\text{Rh}_2(\text{tpa})_4$ for (**anti-6a:syn-6a**) suggest a minor steric effect on selectivity. These composite data are consistent with an oxonium ylide intermediate free from association with the ligated metal catalyst during product formation.


Observation of two diastereoisomeric products from the [2,3]-sigmatropic rearrangement with nearly identical diastereoisomer ratios (**anti-6:syn-6**) as those for the [1,2]-Stevens rearrangement (**syn-7:anti-7**) is inconsistent with a cleavage/rearrangement/recombination pathway in products from the [2,3]-sigmatropic rearrangement but not from the Stevens rearrangement. That the diastereomer ratios are nearly identical suggest that the two processes originate from the same intermediate(s). The absence of a substituent effect on the ratio of products from [2,3]- and [1,2]-rearrangements (**6:7**) is unexpected if both sets of products arise from a cleavage/rearrangement/recombination pathway. Instead, the selectivity match between the two pathways is more likely due to the conformational equilibrium in the reactants (Scheme 1) that results in two non-interconvertible oxonium ylide intermediates and a subsequent competition between the symmetry-allowed [2,3]-sigmatropic rearrangement and the formally symmetry- forbidden [1,2]-Stevens rearrangement (Scheme 2).

Competition between [2,3]- and [1,2]-oxonium ylide rearrangements is relatively common, especially with diazoketones.^{15,16} This competition is generally thought to be due to steric influences in structurally rigid systems. Diastereoisomeric products from the [1,2]-Stevens rearrangement have been reported, and their formation has been attributed to a cleavage/rearrangement/recombination pathway. However, the formation of two diastereoisomers for

the [2,3]-sigmatropic rearrangement pathway is virtually unprecedented. The exception is a report from Clark and coworkers in a single example of structurally well-defined diastereoisomeric products for both the [1,2]-Stevens and the [2,3]-sigmatropic rearrangement pathways,¹⁶ but the origin of these diastereoisomeric products was not explained; the explanation that we have advanced adequately accounts for their “fascinating” results.

Based on our previous demonstration that increasing the steric bulk of the aryl group of *trans*-aryl-substituted tetrahydropyranone diazoacetates **1** results in the production of only one diastereoisomer in the [1,2]-Stevens rearrangement, we constructed the di- and triphenyl analogues of **5a** in order to evaluate diastereoisomer formation from the [2,3]-sigmatropic and [1,2]-Stevens rearrangement pathways, as well the influence of size on the [2,3]/[1,2]-rearrangement ratio of products. The syntheses of diazoacetates **9** and **10** were easily accessible by employing the previously described methods.

Rhodium(II) catalyzed reactions of **9** and **10** afforded [1,2]-Stevens rearrangement products **11** and **12**, respectively, in 73% and 80% yield (eq 3). The [2,3]-sigmatropic rearrangement products were not detected by either ¹H NMR spectra of the reaction mixtures or by HPLC analyses. However, unlike the previously established results with di-ortho-substituted aryl derivatives of **1**, two diastereoisomers for each of the [1,2]-Stevens rearrangement products were formed in amounts that corresponded to those found with their mono-phenyl counterpart **5a**. The (*syn:anti*) ratios of **11** and **12** are consistent with those ratios of **7a-c**.



(3)

In addition to the [1,2]-Stevens rearrangement products, 2-methyl-6-vinylpyran-4-ones **13** and **14** were observed as by-products. Although these byproducts were not formed in high yields, they are ubiquitous in catalytic reactions of tetrahydropyranone- and tetrahydrofuranone-diazoacetates.¹⁷

In conclusion, we have discovered that both the [2,3]- and [1,2]-oxonium ylide rearrangement reactions of 3- (*trans*-styryl)tetrahydropyranone-5-diazoacetate and its para-substituted derivatives **5a-c** form diastereoisomeric products in the same ratios and that their diastereoisomer ratios are independent of electronic influences from para substituents or of catalyst ligands. These results are consistent with the involvement of catalyst-free oxonium ylides whose composition reflects the axial-equatorial conformational isomer distribution of the reactant diazoacetate or its metal carbene intermediate. Furthermore, increasing the size of the styryl group of the 3- (*trans*-styryl)tetrahydropyranone-5-diazoacetate with phenyl substituents **9** and **10** removes competition with the [2,3]-oxonium ylide rearrangement, allowing the [1,2]-oxonium ylide rearrangement to occur with production of two diastereoisomers in ratios that are identical with that from **5a-c**.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Support for this research from the National Institutes of Health (GM 46503) and the National Science Foundation (CHE-0748121) is gratefully acknowledged.

References

1. Jaber DM, Burgin RN, Hepler M, Zavalij P, Doyle MP. Chem Commun. 2011; 47:7623.
2. (a) Wang, J. Comprehensive Organometallic Chemistry. Crabtree, RH.; Mingos, DM., editors. Vol. 11. Elsevier; Oxford: 2007. p. 151-178.(b) West, FG. Modern Rhodium-Catalyzed Organic Reactions. Evans, PA., editor. Vol. Chapter 18. Wiley-VCH; New York: 2005. (c) Hodgson DM, Pierard FYTM, Stupp PA. Chem Soc Rev. 2001; 30:50.(d) Doyle, MP.; Mckervery, MA.; Ye, T. Modern Catalytic Methods for Organic Synthesis with Diazo Compounds. Wiley; New York: 1998. (e) Doyle MP, Forbes DC. Chem Rev. 1998; 98:911. [PubMed: 11848918]
3. Woodward RB, Hoffmann R. Angew Chem, Int Ed Engl. 1969; 8:781.
4. (a) Stewart C, McDonald R, West FG. Org Lett. 2011; 13:720. [PubMed: 21241059] (b) Murphy GK, Marmsaeter FP, West FG. Can J Chem. 2006; 84:1470.(c) Murphy GK, West FG. Org Lett. 2005; 7:1801. [PubMed: 15844910] (d) Marmsaeter FP, West FG. J Am Chem Soc. 2001; 123:5144. [PubMed: 11457364] (e) Feldman KS, Wroblewski ML. J Org Chem. 2000; 65:8659. [PubMed: 11112587] (f) Doyle MP, Ene DG, Forbes DC, Tedrow JS. Tetrahedron Lett. 1997; 38:4367.(g) Doyle MP, Griffin JH, Chinn MS, Van Leusen D. J Org Chem. 1984; 49:1917.
5. A concerted mechanism, including those with metal association: (a) Lu CD, Liu H, Chen ZY, Hu WH, Mi AQ. Org Lett. 2005; 7:83. [PubMed: 15624983] (b) Qu Z, Shi W, Wang J. J Org Chem. 2004; 69:217. [PubMed: 14703404] (c) Kitagaki S, Yanamoto Y, Tsutsui H, Anada M, Nakajima M, Hashimoto S. Tetrahedron Lett. 2001; 42:6361.(d) Roskamp EJ, Johnson CR. J Am Chem Soc. 1986; 108:6062. [PubMed: 22175385]
6. Through an oxonium metal enolate: (a) Ji J, Zhang X, Zhu Y, Qian Y, Zhou J, Yang L, Hu W. J Org Chem. 2011; 76:5821. [PubMed: 21667947] (b) Sawada Y, Mori T, Oku A. J Org Chem. 2003; 68:10040. [PubMed: 14682698]
7. Via alkyl/aryl transfer from the metal associated oxonium ylide to the metal, then reductive elimination: (a) West FG, Naidu BN, Tester RW. J Org Chem. 1994; 59:6892.(b) Karche NP, Jachak SM, Dhavale DD. J Org Chem. 2001; 66:6323. [PubMed: 11559182]
8. Eberlein TH, West FG, Tester RW. J Org Chem. 1992; 57:3479.
9. West FG, Naidu BN, Tester RW. J Org Chem. 1994; 59:6892.
10. (a) Sweeney JB. Chem Soc Rev. 2009; 38:1027. [PubMed: 19421580] (b) Tuzina P, Somfai P. Org Lett. 2009; 11:919. [PubMed: 19146452]
11. Sharma A, Guenee L, Naubron J-V, Lacour J. Angew Chem, Int Ed. 2011; 50:3677.
12. Kitagaki S, Yanamoto Y, Tsutsui H, Anada M, Nakajima M, Hashimoto S. Tetrahedron Lett. 2001; 42:6361.
13. Danishefsky S, Larson E, Askin D, Kato N. J Am Chem Soc. 1985; 107:1246.
14. Liu Y, Zhang Y, Jee N, Doyle MP. Org Lett. 2008; 10:1605. [PubMed: 18351770]
15. (a) Marmsater Fredrik P, Vanecko John A, West FG. Org Lett. 2004; 6:1657. [PubMed: 15128260] (b) Clark JS, Bate AL, Grinter T. Chem Commun. 2001:459.(c) Brogan JB, Zercher CK. Tetrahedron Lett. 1998; 39:1691.(d) Brogan JB, Zercher CK, Bauer CB, Rogers RD. J Org Chem. 1997; 62:3902.
16. Clark JS, Walls SB, Wilson C, East SP, Drysdale MJ. Eur J Org Chem. 2006:323.
17. Doyle MP, Dyatkin AB, Autry CL. J Chem Soc, Perkin Trans 1. 1995:619.






Figure 1.
(a) Crystal structure (left) of ***syn-6a*** (minor isomer). (b) Crystal structure (right) of ***syn-7a*** (major isomer).

**Scheme 1.**

Conformation-Dependent Pathway for the Formation of Two Diastereoisomers via [1,2]-Oxonium Ylide Rearrangements.

**Scheme 2.**

Conformation-Dependent Pathway for the Formation of Two Diastereoisomers via [2,3]- and [1,2]- Oxonium Ylide Rearrangements.

Table 1

Effect of ring substitution on product formation in the rhodium(II) octanoate catalyzed reactions of 3-(*trans*-styryl) tetrahydropyranone-5-diazoacetates **5a-c**.^a

	Ar	6:7 ^b	<i>anti</i> -6: <i>syn</i> -6	<i>syn</i> -7: <i>anti</i> -7	yield, % (6+7) ^c
5a	C ₆ H ₅	54:46	78:22	78:22	88
5b	<i>p</i> -NO ₂ C ₆ H ₄	52:48	76:24	76:24	85
5c	<i>p</i> -MeOC ₆ H ₄	69:31	79:21	70:30	60

^aReactions were performed in refluxing CH₂Cl₂ for 2 h using 1.0 mol % of Rh₂(oct)₄. Results reported are averages of two or more reactions $\pm 2\%$.

^bProduct ratio determined by ¹H NMR spectral analysis with a variance of $\pm 2\%$.

^c¹H NMR yield using benzaldehyde as an internal standard.

Table 2

Effect of dirhodium(II) ligands on yields and product ratios from oxonium ylide reactions of 3-(*trans*-styryl)tetrahydropyranone- 5-diazoacetate **5a**.^a

catalyst	(6a : 7a) ^b	<i>anti</i> - 6a : <i>syn</i> - 6a	<i>syn</i> - 7a : <i>anti</i> - 7a	yield, % (6a + 7a) ^c
Rh ₂ (oct) ₄	54:46	78:22	78:22	88
Rh ₂ (OAc) ₄	51:49	80:20	77:23	94
Rh ₂ (pfb) ₄	55:45	80:20	77:23	46
Rh ₂ (piv) ₄	55:45	78:22	76:24	94
Rh ₂ (tpa) ₄	53:47	73:27	75:25	93
Rh ₂ (tfa) ₄	51:49	79:21	75:25	41

^a Reactions were performed in refluxing dichloromethane for 2 h using 1.0 mol % of catalyst. Results reported are averages of two or more reactions.

^b Product ratio determined by ¹H NMR spectral analysis with a variance of ±2%.

^c NMR yield using benzaldehyde as an internal standard.