

# Relevance of tumour necrosis factor- $\alpha$ for the inflammatory and nociceptive responses evoked by carrageenan in the mouse paw

<sup>1</sup>Ana C.C. Rocha, <sup>1</sup>Elizabeth S. Fernandes, <sup>1</sup>Nara L.M. Quintão, <sup>1,2</sup>Maria M. Campos & <sup>\*</sup><sup>1</sup>João B. Calixto

<sup>1</sup>Department of Pharmacology, Centre of Biological Sciences, Universidade Federal de Santa Catarina, Campus Universitário, Florianópolis, Santa Catarina 88049-900, Brazil

**1** The present study evaluated the participation of tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) in the inflammatory and nociceptive responses evoked by carrageenan in the mouse paw.

**2** The intraplantar injection of carrageenan (300  $\mu$ g paw<sup>-1</sup>) induced a marked and biphasic paw oedema formation (peaks at 6 and 72 h), which was accompanied by a long-lasting mechanical allodynia (that remained elevated for up to 72 h) and a significant increase of myeloperoxidase (MPO) activity (peak at 6 h) in both Swiss and C57/BL6 mice.

**3** The paw oedema, the elevation of MPO activity and to a lesser extent the mechanical allodynia elicited by carrageenan were found to be significantly reduced in TNF- $\alpha$  p55 receptor knockout mice.

**4** Of interest, the systemic administration of an anti-TNF- $\alpha$  antibody produced a significant inhibition of paw oedema, mechanical allodynia and MPO activity. A noteworthy decrease in inflammatory and nociceptive responses caused by carrageenan was also observed when mice were previously treated with the preferential inhibitor of TNF- $\alpha$  synthesis, thalidomide.

**5** The present results clearly indicate that the proinflammatory cytokine TNF- $\alpha$  plays a critical role in the oedema formation, as well as in the mechanical allodynia and the neutrophil migration, following carrageenan administration into the mouse paw. Intraplantar injection of carrageenan in mice could constitute a useful model for assessment of the *in vivo* effects of potential inhibitors of TNF- $\alpha$ -related pathways.

*British Journal of Pharmacology* (2006) **148**, 688–695. doi:10.1038/sj.bjp.0706775;

published online 15 May 2006

**Keywords:** Carrageenan; mouse paw; TNF- $\alpha$ ; oedema; nociception; cell migration

**Abbreviations:** MPO, myeloperoxidase; NO, nitric oxide; PBS, phosphate-buffered saline; s.c., subcutaneous; TMB, tetramethylbenzidine; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; VFH, von Frey hair

## Introduction

A great number of studies have attempted to evaluate nociceptive and inflammatory alterations in rodents. Winter *et al.* (1962) introduced the paw oedema model arising from the intraplantar injection of carrageenan in rats. It has been demonstrated that carrageenan produces a monophasic oedematogenic response in the rat paw, which is accompanied by inflammatory cell migration and marked nociceptive alterations. Further studies have shown that carrageenan-induced rat paw oedema is largely associated with the production of several inflammatory mediators such as histamine, prostaglandins, kinins, nitric oxide (NO) and cytokines, among others (Di Rosa *et al.*, 1971; Salvemini *et al.*, 1986; Tabo *et al.*, 1998; Pinheiro and Calixto, 2002; Morris, 2003; Esposito *et al.*, 2005). In mice, the intraplantar injection of carrageenan produces a peculiar oedematogenic response, which possesses a biphasic profile, with peaks at 6 and 72 h (Levy, 1969; Sugishita *et al.*, 1981; Henriques *et al.*, 1987; Ianaro *et al.*, 1994; Posadas *et al.*, 2004). Carrageenan-induced mouse paw oedema is also associated with paw nociceptive changes and

the migration of inflammatory cells to the site of injection (Posadas *et al.*, 2004; Cunha *et al.*, 2005). It has been demonstrated recently that inflammatory responses following carrageenan injection into mice paw are associated with the production of prostaglandins (formed by the action of both COX-1 and COX-2) and NO (derived from both eNOS and iNOS) (Posadas *et al.*, 2004; Bucci *et al.*, 2005).

Tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) is a potent proinflammatory cytokine that possesses multiple effects, including the activation of inflammatory cells, the induction of several inflammatory proteins, cytotoxicity, etc. (Feldmann & Saklatvala, 2001; Haddad, 2002; Hopkins, 2003). The present study was designed to investigate the contribution of TNF- $\alpha$  production to the inflammatory responses (oedema and neutrophil migration) and also the nociceptive response (mechanical allodynia) caused by the intraplantar injection of carrageenan into the mouse paw.

## Methods

### Animals

Male Swiss, C57/BL6 and TNF- $\alpha$  p55<sup>-/-</sup> receptor knockout mice (30–35 g) were used throughout this study. Animals were

\*Author for correspondence; E-mail: calixto@farmaco.ufsc.br or calixto3@terra.com.br

<sup>2</sup>Current address: School of Dentistry, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, RS, Brazil.

housed under conditions of optimum light, temperature and humidity (12 h light–dark cycle,  $22 \pm 1^\circ\text{C}$ , 60–80% humidity), with food and water provided *ad libitum*. Swiss mice were obtained from the Department of Pharmacology, Universidade Federal de Santa Catarina (UFSC, Florianópolis, Brazil). C57/BL6 wild-type or TNF- $\alpha$  p55 $^{-/-}$  receptor knockout mice were obtained from the Universidade Federal de Minas Gerais (UFMG, Belo Horizonte, Brazil). The TNF- $\alpha$  p55 $^{-/-}$  receptor-deficient mice used in the present study were C57/BL6 inbred. Experiments were conducted in accordance with the current guidelines for the care of laboratory animals and ethical guidelines for the investigation of experimental pain in conscious animals laid down by Zimmermann (1983). The Ethics Committee of the Federal University of Santa Catarina approved all the experimental procedures (protocol numbers 262/CEUA and 23080.035334/2003-16/UFSC). The numbers of animals and intensities of noxious stimuli used were the minimum necessary to demonstrate the consistent effects.

### Paw oedema

The experiments were conducted according to the procedures originally described by Henriques *et al.* (1987). Briefly, the animals received a 50  $\mu\text{l}$  subcutaneous (s.c.) injection, into one hindpaw (the right), of phosphate-buffered saline (PBS, composition, in  $\text{mmol l}^{-1}$ : NaCl 137, KCl 2.7 and phosphate buffer 10) containing carrageenan (300  $\mu\text{g paw}^{-1}$ ). The contralateral paw (left paw) received 50  $\mu\text{l}$  of PBS and was used as the control. Paw oedema was measured by means of a plethysmometer, modified for mice (Ugo Basile), at several time points (1/2, 1, 2, 4, 6, 24, 48, 72 and 96 h) after the injection of carrageenan. Paw oedema is expressed in microlitres as the difference between the right and left paws.

### Hindpaw withdrawal response induced by von Frey hairs

The mechanical allodynia was carried out according to the procedures described before by Quintão *et al.* (2005) with minor modifications. The animals received a 50  $\mu\text{l}$  s.c. injection, into the right hindpaw, of PBS containing carrageenan (300  $\mu\text{g paw}^{-1}$ ). Following several intervals of time (1, 3, 4, 6, 24, 48 and 72 h), the mice were placed individually in clear Plexiglas boxes ( $9 \times 7 \times 11\text{ cm}$ ) on elevated wire mesh platforms to allow access to the ventral surface of the right hindpaw. The animals were acclimatized for 30 min before behavioural testing. The withdrawal response frequency was measured following 10 applications (duration of 1 s each) of von Frey hairs (VFH, Stoelting, Chicago, IL, U.S.A.). Stimuli were delivered from below to the plantar surface of the right hindpaw. The VFH of 0.6 g produces a mean withdrawal frequency of about 15%, which is considered to be an adequate value for the measurement of mechanical hypernociception (Bortolanza *et al.*, 2002). Therefore, 0.6 g VFH was used throughout this study. In order to determine the basal mechanical thresholds, all the groups were evaluated before carrageenan injection.

### Neutrophil myeloperoxidase assay

The neutrophil recruitment to the mouse paw was measured by means of tissue myeloperoxidase (MPO) activity, which was determined according to the method described by da Cunha

*et al.* (2004), with some adjustments. The animals received a 50  $\mu\text{l}$  s.c. injection, into the right paw, of PBS containing carrageenan (300  $\mu\text{g paw}^{-1}$ ) and were killed 6 h later. PBS-treated paws were used as control. At the time of killing, the s.c. tissue of the paws was removed, homogenized at 5% (weight volume $^{-1}$ ) in ethylenediaminetetraacetic acid (EDTA)/NaCl buffer (pH 4.7) and centrifuged at 10,000 r.p.m. for 15 min at  $4^\circ\text{C}$ . The pellet was re-suspended in hexadecyltrimethyl ammonium bromide (HTAB) 0.5% buffer (pH 5.4) and the samples were frozen and thawed three times in liquid nitrogen. Upon thawing, the samples were re-centrifuged (10,000 r.p.m., 15 min, at  $4^\circ\text{C}$ ) and 25  $\mu\text{l}$  of the supernatant were used for the MPO assay. The enzymatic reaction was assessed with 1.6 mM tetramethylbenzidine (TMB), 80 mM  $\text{NaPO}_4$  and 0.3 mM hydrogen peroxide. The absorbance was measured at 690 nm and the results are expressed in optical density per milligram of tissue.

### Effect of anti-TNF- $\alpha$ strategies

To evaluate the role of TNF- $\alpha$  in the nociceptive and inflammatory responses induced by carrageenan in the mouse paw, different groups of Swiss mice were systemically treated with an anti-TNF- $\alpha$  antibody (100  $\mu\text{g kg}^{-1}$ , i.v.) or with the known inhibitor of TNF- $\alpha$  synthesis, thalidomide (30  $\text{mg kg}^{-1}$ , s.c.), both administered 5 min before carrageenan (300  $\mu\text{g paw}^{-1}$ ) injection. A separate group of mice received the anti-TNF- $\alpha$  antibody (100  $\mu\text{g kg}^{-1}$ , i.v.) 5 min before and 3 h after the application of carrageenan. The control animals received the vehicle. The doses of inhibitors were chosen on the basis of previous studies (Hattori *et al.*, 1998; Karrow *et al.*, 2003; Arruda *et al.*, 2004; Rocha *et al.*, 2005; Quintão *et al.*, 2006).

In other groups of experiments, the relevance of TNF- $\alpha$  was assessed using TNF- $\alpha$  p55 receptor knockout mice. C57/BL6 mice were used as the control group for this series of experiments. The paw oedema, the mechanical allodynia and the MPO activity were measured as described in the preceding sections.

### Drugs and reagents

The following drugs were used: carrageenan, PBS tablets, EDTA, HTAB, TMB (all from Sigma-Aldrich, St Louis, MO, U.S.A.),  $\text{NaPO}_4$ , hydrogen peroxide and Tween 20 (all from Merck, Germany). Anti-murine neutralizing antibody anti-TNF- $\alpha$  (Lot CT101) was obtained from R&D Systems Inc. (Minneapolis, MN, U.S.A.). Thalidomide was from Tocris (Cookson Inc., Ellisville, MO, U.S.A.). The anti-TNF- $\alpha$  antibody was prepared daily in 0.9% ( $\text{w v}^{-1}$ ) NaCl solution. Thalidomide was diluted in a 1% DMSO/saline solution.

### Statistical analysis

The results are presented as the mean  $\pm$  s.e.m. of 6–10 animals. The percentages of inhibition are reported as the mean  $\pm$  s.e.m. of inhibitions obtained in each individual experiment. Statistical comparison of the data was performed by analysis of variance followed by Dunnett's test or Student's unpaired *t*-test. *P*-values less than 0.05 ( $P < 0.05$ ) were considered significant.

## Results

### Carrageenan-induced paw oedema

As described before (Henriques *et al.*, 1987), the intraplantar injection of carrageenan ( $300 \mu\text{g paw}^{-1}$ ) into the mouse paw produced a biphasic and marked paw oedema formation, which peaked at 6 and 72 h, decreasing after 96 h. The oedematogenic response caused by carrageenan in Swiss mice was similar to that observed in C57/BL6 mice. Interestingly, both the early and the late phases of carrageenan-induced oedema were found to be strikingly reduced in TNF- $\alpha$  p55 receptor knockout mice. The percentages of inhibition were  $89 \pm 4$  and  $50 \pm 3\%$  at 6 and 48 h, respectively (Figure 1). The treatment of animals with a single dose of the anti-TNF- $\alpha$  antibody ( $100 \mu\text{g kg}^{-1}$ , i.v., 5 min) significantly reduced the early, but not the late, phase of carrageenan-evoked oedema, with inhibitions of  $76 \pm 3$  and  $64 \pm 3\%$  at 2 and 4 h, respectively (Figure 2a). In addition, the administration of anti-TNF- $\alpha$  antibody ( $100 \mu\text{g kg}^{-1}$ , i.v.) 5 min before and 3 h after the injection of carrageenan was able to markedly reduce the oedematogenic response caused by administration of carrageenan, an effect that was observed for up to 24 h. The inhibitions observed were  $59 \pm 3$  and  $51 \pm 6\%$ , at 6 and 24 h, respectively (Figure 2b). Finally, the systemic treatment of mice with the preferential blocker of TNF- $\alpha$  synthesis thalidomide ( $30 \text{ mg kg}^{-1}$ , s.c., 5 min) significantly prevented both the early and the late phases of carrageenan-induced paw oedema. The percentages of reduction were, respectively,  $70 \pm 7$  and  $41 \pm 10\%$ , at 6 and 48 h (Figure 3).

### Mechanical allodynia caused by carrageenan

The application of carrageenan ( $300 \mu\text{g paw}^{-1}$ ) caused a sustained mechanical allodynia, as indicated by a large increase from baseline values in response to 0.6 g VFH stimulation, which was observed as early as 1 h and persisted for up to 72 h (Figure 4). The results show that the mechanical allodynia following carrageenan injection was not significantly different between Swiss and C57/BL6 mice, but that it was found significantly reduced in mice with genic deletion of the TNF- $\alpha$  p55 receptor (Figure 4a). The percentages of reduction


observed were  $39 \pm 12$ ,  $37 \pm 3$  and  $25 \pm 2\%$  at 6, 24 and 48 h, respectively. The mechanical allodynia caused by carrageenan was also significantly inhibited in mice pretreated with the anti-TNF- $\alpha$  antibody ( $100 \mu\text{g kg}^{-1}$ , i.v., 5 min before) by  $48 \pm 8$ ,  $42 \pm 10$  and  $36 \pm 14\%$ , at 4, 24 and 48 h, respectively (Figure 5a). The repeated treatment with anti-TNF- $\alpha$  antibody ( $100 \mu\text{g kg}^{-1}$ , i.v.), 5 min before and 3 h after the injection of carrageenan, resulted in a marked decrease of the allodynic response, with percentages of inhibition of  $39 \pm 9$ ,  $82 \pm 6$ ,  $56 \pm 10$  and  $41 \pm 6\%$  at 3, 4, 6 and 24 h, respectively (Figure 5b). Likewise, the pretreatment of animals with thalidomide ( $30 \text{ mg kg}^{-1}$ , s.c., 5 min) was capable of significantly reducing carrageenan-induced mechanical allodynia, with percentages of inhibition of  $44 \pm 5$ ,  $45 \pm 19$ ,  $57 \pm 16$  and  $50 \pm 3\%$  at 4, 6, 24 and 48 h, respectively (Figure 4b).

### Increase in MPO activity


Carrageenan injection is associated with a massive migration of neutrophils, as indicated by an increase of about 10-fold in the MPO activity in the s.c. tissue of the mouse paw at 6 h. The increase of MPO activity induced by carrageenan was not significantly different between Swiss and C57/BL6 mice, but was markedly reduced in TNF- $\alpha$  p55 receptor knockout mice ( $51 \pm 9\%$ ) (Figure 6a). A significant inhibition of increased MPO activity was also achieved by treating animals with anti-TNF- $\alpha$  ( $100 \mu\text{g kg}^{-1}$ ) in either single or repeated schedules of treatment (Figure 6c and d). The percentages of inhibition were  $55 \pm 11$  and  $46 \pm 9\%$ , respectively. Finally, thalidomide ( $30 \text{ mg kg}^{-1}$ , s.c.), given 5 min before carrageenan injection (Figure 6b), also significantly reduced carrageenan-induced increase in MPO activity ( $30 \pm 2\%$ ).

## Discussion


TNF- $\alpha$  is a key proinflammatory cytokine that has been widely implicated in both the establishment and the perpetuation of the inflammatory process. Most of the TNF- $\alpha$  effects are mediated by the interaction with two receptors called p55 (type I) and p75 (type II), and they are clearly dependent on



**Figure 1** Mouse paw oedema formation induced by carrageenan ( $300 \mu\text{g paw}^{-1}$ ) in Swiss, C57/BL6 wild-type or TNF- $\alpha$  p55 $^{-/-}$  receptor knockout mice. Each point represents the mean of 10 animals and vertical lines show the s.e.m. Asterisks denote the significance levels when the values obtained for C57/BL6 wild type were compared to those for TNF- $\alpha$  p55 $^{-/-}$  receptor knockout mice: \*\* $P < 0.01$ .




**Figure 2** Effect of systemic treatment with anti-TNF- $\alpha$  antibody ( $100 \mu\text{g kg}^{-1}$ , i.v.), given 5 min before carrageenan (a) or 5 min before and 3 h after carrageenan (b), on the mouse paw oedema formation induced by carrageenan ( $300 \mu\text{g paw}^{-1}$ ) in Swiss mice. Each point on the curve represents the mean of six animals and vertical lines show the s.e.m. Asterisks denote the significance levels in comparison to control values: \* $P < 0.05$ ; \*\* $P < 0.01$ .




**Figure 3** Effect of systemic treatment with thalidomide ( $30 \text{ mg kg}^{-1}$ , s.c., 5 min before carrageenan) on the mouse paw oedema formation induced by carrageenan ( $300 \mu\text{g paw}^{-1}$ ) in Swiss mice. Each point on the curve represents the mean of six animals and vertical lines show the s.e.m. Asterisks denote the significance levels in comparison to control values: \* $P < 0.05$ ; \*\* $P < 0.01$ .

the activation of transcriptional factors and *de novo* synthesis of several proteins involved in the inflammatory response (Feldmann & Saklatvala, 2001; Haddad, 2002; Hopkins, 2003; Sommer & Kress, 2004). In the present study, we have assessed the role played by TNF- $\alpha$  in the inflammatory and nociceptive responses evoked by carrageenan injection into the mouse paw.

The model of mouse oedema induced by the intraplantar injection of carrageenan has been widely employed in recent years in order to assess the effects of new anti-inflammatory drugs. For this purpose, different concentrations of carrageenan and distinct mouse strains have been employed (Levy, 1969; Sugishita *et al.*, 1981; Henriques *et al.*, 1987; Ianaro *et al.*, 1994; Posadas *et al.*, 2004). From these studies, it is possible to




**Figure 4** Mechanical allodynic response induced by carrageenan ( $300 \mu\text{g paw}^{-1}$ ) in Swiss, C57/BL6 wild-type or TNF- $\alpha$  p55<sup>-/-</sup> receptor knockout mice (a). Effect of systemic treatment with thalidomide ( $30 \text{ mg kg}^{-1}$ , s.c.), given 5 min before carrageenan, on the mechanical allodynia induced by carrageenan injection (b). Each point represents the mean of 6–10 animals and vertical lines show the s.e.m. Asterisks denote the significance levels when the values obtained for C57/BL6 wild type were compared to those for TNF- $\alpha$  p55<sup>-/-</sup> receptor knockout mice or in comparison to Swiss mice (thalidomide experimental group): \* $P < 0.05$ ; \*\* $P < 0.01$ .



**Figure 5** Effect of systemic treatment with anti-TNF- $\alpha$  antibody ( $100 \mu\text{g kg}^{-1}$ , i.v.), given 5 min before carrageenan (a) or 5 min before and 3 h after carrageenan (b), on the mechanical allodynia induced by carrageenan injection. Each point represents the mean of 6–10 animals and vertical lines show the s.e.m. Asterisks denote the significance levels in comparison to control values: \* $P < 0.05$ ; \*\* $P < 0.01$ .

observe that in the mouse, carrageenan evokes a very characteristic oedematogenic response with a clear biphasic profile. In spite of the systematic use of this inflammation model, few studies have evaluated the possible mechanisms implicated in the early and late phases of oedema formation induced by carrageenan in the mouse paw. For instance, Siqueira-Junior *et al.* (2003) demonstrated that treatment with the preferential inhibitor of COX-1 valeryl-salicylate produced a marked reduction of the early phase of carrageenan-induced mouse paw oedema, whereas it significantly increased the late phase. Moreover, it has recently been demonstrated that COX-2 is upregulated only in the second phase of the paw oedema caused by carrageenan (Posadas *et al.*, 2004). Conversely, it has been shown that NO (derived from both the endothelial and inducible NO synthase isoforms) is apparently implicated in the two phases of carrageenan-evoked mouse paw oedema (Posadas *et al.*, 2004; Bucci *et al.*, 2005).

The results of the present study indicate, for the first time, that TNF- $\alpha$  is likely to be involved in both the first and second phases of mouse carrageenan oedema. This conclusion is supported by the results showing that the oedema formation induced by carrageenan was strikingly reduced in TNF- $\alpha$  p55 receptor knockout mice, in comparison to C57/BL6 mice, whereas no significant difference was observed between Swiss and C57/BL6 mouse strains. In addition, the pretreatment with thalidomide (given 5 min before carrageenan) was also able to markedly reduce the early and late stages of the oedematogenic response caused by carrageenan. Of note, thalidomide and its analogues have been shown to largely reduce the increase of TNF- $\alpha$  synthesis, and to a lesser extent, of other proinflammatory cytokines (for review see Teo *et al.*, 2005). Finally, our results showed that treatment of animals with a single dose of anti-TNF- $\alpha$  antibody significantly decreased the early phase of carrageenan oedema, without



**Figure 6** Increase in MPO activity induced by carrageenan ( $300 \mu\text{g paw}^{-1}$ ) in Swiss, C57/BL6 wild-type or TNF- $\alpha$  p55<sup>-/-</sup> receptor knockout mice (a). Effect of systemic treatment with thalidomide ( $30 \text{ mg kg}^{-1}$ , s.c.), given 5 min before carrageenan, on the increase in MPO activity induced by carrageenan injection (b). Effect of systemic treatment with anti-TNF- $\alpha$  antibody ( $100 \mu\text{g kg}^{-1}$ , i.v.), given 5 min before carrageenan (c) or 5 min before and 3 h after carrageenan (d), on the MPO activity induced by carrageenan injection. Each point represents the mean of six animals and vertical lines show the s.e.m. Asterisks denote the significance levels when the values obtained for C57/BL6 wild type were compared to those for TNF- $\alpha$  p55<sup>-/-</sup> receptor knockout mice or in comparison to Swiss mice (anti-TNF- $\alpha$  or thalidomide experimental groups): \*\* $P < 0.01$  and significantly different to PBS-treated group (#) values, ## $P < 0.01$ .

affecting the late phase. Furthermore, the repeated administration of anti-TNF- $\alpha$  antibody, 5 min before and 3 h after carrageenan, markedly reduced the oedema caused by carrageenan for up to 24 h. It is worth noting that anti-TNF- $\alpha$  strategies adopted in the present study were always more effective against the first, in comparison to the second phase of carrageenan oedema. Therefore, the present data allow us to suggest that TNF- $\alpha$  plays an essential role in the early, and to a lesser extent in the late, stage of the paw oedema induced by carrageenan in mice. Further studies are necessary to establish whether TNF- $\alpha$  interferes with the production of other mediators of inflammation following the intraplantar injection of carrageenan into the mouse paw.

It has been widely demonstrated that oedema formation caused by carrageenan is accompanied by a reduction in the threshold to nociceptive stimuli, resulting from primary and central mechanisms of sensitization in both rats and mice (Tabo *et al.*, 1998; Parada *et al.*, 2003; Cunha *et al.*, 2005; Quintão *et al.*, 2005). In the present study, we have demonstrated that carrageenan injection induces a long-lasting

mechanical allodynic response, which is evident after as little as 1 h and remains elevated for up to 72 h. In addition, the results obtained herein clearly indicate the involvement of the proinflammatory cytokine TNF- $\alpha$  in the mechanical allodynia evoked by carrageenan in mice. Our conclusion is derived from two sets of experimental evidence: (1) the mechanical allodynia induced by carrageenan was partially reduced in TNF- $\alpha$  p55 receptor knockout mice, in comparison to C57/BL6 mice, and (2) the pretreatment with both anti-TNF- $\alpha$  antibody (single or repeated administration) and thalidomide prevented, in a significant manner, the allodynic responses evoked by carrageenan in Swiss mice. These data extend and confirm the previous work conducted by Cunha *et al.* (2005) showing that carrageenan-induced mechanical hypernociception is significantly reduced in mice with genic deletion of TNF- $\alpha$  p55 (type 1) receptor, according to assessment with an electronic version of the von Frey test. The relevance of TNF- $\alpha$  to the nociceptive responses elicited by carrageenan has also been demonstrated in rats. For example, both thalidomide and anti-TNF- $\alpha$  antibody are highly effective in preventing the mechanical

hyperalgesia caused by carrageenan in the Randall–Sellito test (Cunha *et al.*, 1992; Ribeiro *et al.*, 2000). It has also been shown that the inhibition of TNF- $\alpha$  by treatment with thalidomide, the anti-TNF- $\alpha$  antibody or the antisense to TNF- $\alpha$  p55 receptor markedly prevented the mechanical hyperalgesia and priming responses induced by carrageenan in rats (Parada *et al.*, 2003). Furthermore, TNF- $\alpha$  has also been deeply implicated in the central and peripheral mechanisms involved in neuropathic pain in both mice and rats (Sommer & Kress, 2004; Quintão *et al.*, 2006). Taken together, the literature and present data clearly indicate the importance of TNF- $\alpha$  for the inflammatory nociception induced by carrageenan.

It has been recently reported that intraplantar injection of carrageenan into the mouse paw produces a marked neutrophil influx to the s.c. tissue, as indicated by a 10-fold increase in the MPO activity, which peaks at 6 h (da Cunha *et al.*, 2004; Posadas *et al.*, 2004). In this connection, another aspect investigated in the present work was the possible relevance of TNF- $\alpha$  for the neutrophil migration induced by carrageenan in the mouse paw. Interestingly, as demonstrated for the oedema and the allodynia, TNF- $\alpha$  also seems to play a relevant role in the neutrophil migration following carrageenan injection. Accordingly, carrageenan-induced increase in MPO activity was reduced to a great extent in mice with genic deletion of the TNF- $\alpha$  p55 receptor, in comparison to C57/BL6 mice. Likewise, the rise in MPO activity in the mouse paw was also markedly reduced by

the single and by the repeated treatment with anti-TNF- $\alpha$ . Interestingly, the increased MPO activity was also prevented by thalidomide. These results are in accordance with previous data showing that neutrophil migration observed in the model of pleurisy induced by carrageenan is significantly blocked by anti-TNF- $\alpha$  strategies (Fröde *et al.*, 2001). Collectively, the results allow us to suggest that TNF- $\alpha$  is a key molecule for the process of cell migration induced by carrageenan in mice.

In summary, the present data clearly indicate the proinflammatory cytokine TNF- $\alpha$  as a crucial mediator for oedema formation, neutrophil migration and allodynic responses following carrageenan injection into the mouse paw. These pieces of evidence cast new light on the mechanisms responsible for mediating the inflammatory and nociceptive actions of carrageenan in the mouse paw.

This work was supported by grants from the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES, Brazil), the Programa de Apoio aos Núcleos de Excelência (PRONEX) (Brazil) and the Fundação de Apoio à Pesquisa de Santa Catarina (FAPESC). A.C.C.R. is an undergraduate medical student receiving a grant from PIBIC/CNPq/BIP/UFSC (Brazil). E.S.F. and N.L.M.Q. are post-graduate students in pharmacology receiving grants from CAPES and CNPq, respectively. M.M.C. holds a post-doctoral fellowship from CAPES. We thank Dr Leda Quercia Vieira very much indeed for her kind donation of C57/BL6 and TNF- $\alpha$  p55<sup>-/-</sup> receptor knockout mice.

## References

- ARRUDA, M.S., RICHINI, V.B., OLIVEIRA, S.M. & VILANI-MORENO, F.R. (2004). Experimental murine mycobacteriosis: evaluation of the functional activity of alveolar macrophages in thalidomide-treated mice. *Braz. J. Med. Biol. Res.*, **37**, 485–492.
- BORTOLANZA, L.B., FERREIRA, J., HESS, S.C., DELLE MONACHE, F., YUNES, R.A. & CALIXTO, J.B. (2002). Anti-allodynic action of the tormentic acid, a triterpene isolated from plant, against neuropathic and inflammatory persistent pain in mice. *Eur. J. Pharmacol.*, **453**, 203–208.
- BUCCI, M., ROVIEZZO, F., POSADAS, I., YU, J., PARENTE, L., SESSA, W.C., IGNARRO, L.J. & CIRINO, G. (2005). Endothelial nitric oxide synthase activation is critical for vascular leakage during acute inflammation *in vivo*. *Proc. Natl. Acad. Sci. U.S.A.*, **102**, 904–908.
- CUNHA, F.Q., POOLE, S., LORENZETTI, B.B. & FERREIRA, S.H. (1992). The pivotal role of tumour necrosis factor alpha in the development of inflammatory hyperalgesia. *Br. J. Pharmacol.*, **107**, 660–664.
- CUNHA, T.M., VERRI JR, W.A., SILVA, J.S., POOLE, S., CUNHA, F.Q. & FERREIRA, S.H. (2005). A cascade of cytokines mediates mechanical inflammatory hypernociception in mice. *Proc. Natl. Acad. Sci. U.S.A.*, **102**, 1755–1760.
- DA CUNHA, F.M., DUMA, D., ASSREUY, J., BUZZI, F.C., NIERO, R., CAMPOS, M.M. & CALIXTO, J.B. (2004). Caffeic acid derivatives: *in vitro* and *in vivo* anti-inflammatory properties. *Free Radic. Res.*, **38**, 1241–1253.
- DI ROSA, M., GIROUD, J.P. & WILLOUGHBY, D.A. (1971). Studies on the mediators of the acute inflammatory response induced in rats in different sites by carrageenan and turpentine. *J. Pathol.*, **104**, 15–29.
- ESPOSITO, E., IACONO, A., RASO, G.M., PACILIO, M., COPPOLA, A., DI CARLO, R. & MELI, R. (2005). Raloxifene, a selective estrogen receptor modulator, reduces carrageenan-induced acute inflammation in normal and ovariectomized rats. *Endocrinology*, **146**, 3301–3308.
- FELDMANN, M. & SAKLATVALA, J. (2001). Proinflammatory cytokines. *Cytokine Ref.*, **1**, 291–305.
- FRÖDE, T.S., SOUZA, G.E. & CALIXTO, J.B. (2001). The modulatory role played by TNF-alpha and IL-1 beta in the inflammatory responses induced by carrageenan in the mouse model of pleurisy. *Cytokine*, **13**, 162–168.
- HADDAD, J.J. (2002). Cytokines and related-receptor mediated signalling pathways. *Biochem. Biophys. Res. Commun.*, **297**, 700–713.
- HATTORI, K., HIRANO, T., MIYAJIMA, H., YAMAKAWA, N., TATENO, M., OSHIMI, K., KAYAGAKI, N., YAGITA, H. & OKUMURA, K. (1998). Differential effects of anti-Fas ligand and anti-tumor necrosis factor alpha antibodies on acute graft-versus-host disease pathologies. *Blood*, **91**, 4051–4055.
- HENRIQUES, M.G., SILVA, P.M., MARTINS, M.A., FLORES, C.A., CUNHA, F.Q., ASSREUY-FILHO, J. & CORDEIRO, R.S. (1987). Mouse paw edema. A new model for inflammation? *Braz. J. Med. Biol. Res.*, **20**, 243–249.
- HOPKINS, S.J. (2003). The pathological role of cytokines. *Legal Med.*, **5**, S45–S47.
- IANARO, A., O'DONNELL, C.A., DI ROSA, M. & LIEW, F.Y. (1994). A nitric oxide synthase inhibitor reduces inflammation, down-regulates inflammatory cytokines and enhances interleukin-10 production in carrageenin-induced oedema in mice. *Immunology*, **82**, 370–375.
- KARROW, N.A., GUO, T.L., ZHANG, L.X., MCCAY, J.A., MUSGROVE, D.L., PEACHEE, V.L., GERMOLEC, D.R. & WHITE JR, K.L. (2003). Thalidomide modulation of the immune response in female B6C3F1 mice: a host resistance study. *Int. Immunopharmacol.*, **3**, 1447–1456.
- LEVY, L. (1969). Carrageenan paw edema in the mouse. *Life Sci.*, **8**, 601–606.
- MORRIS, C.J. (2003). Carrageenan-induced paw edema in the rat and mouse. *Methods Mol. Biol.*, **225**, 115–121.
- PARADA, C.A., YEH, J.J., JOSEPH, E.K. & LEVINE, J.D. (2003). Tumor necrosis factor receptor type-1 in sensory neurons contributes to induction of chronic enhancement of inflammatory hyperalgesia in rat. *Eur. J. Neurosci.*, **17**, 1847–1852.

- PINHEIRO, R.M. & CALIXTO, J.B. (2002). Effect of the selective COX-2 inhibitors, celecoxib and rofecoxib in rat acute models of inflammation. *Inflamm. Res.*, **51**, 603–610.
- POSADAS, I., BUCCI, M., ROVIEZZO, F., ROSSI, A., PARENTE, L., SAUTEBIN, L. & CIRINO, G. (2004). Carrageenan-induced mouse paw oedema is biphasic, age-weight dependent and displays differential nitric oxide cyclooxygenase-2 expression. *Br. J. Pharmacol.*, **142**, 331–338.
- QUINTÃO, N.L.M., BALZ, D., SANTOS, A.R.S., CAMPOS, M.M. & CALIXTO, J.B. (2006). Long-lasting neuropathic pain induced by brachial plexus injury in mice: role triggered by the pro-inflammatory cytokine TNF $\alpha$ . *Neuropharmacology*, **50**, 614–620.
- QUINTÃO, N.L.M., MEDEIROS, R., SANTOS, A.R.S., CAMPOS, M.M. & CALIXTO, J.B. (2005). Effects of diacerhein on mechanical allodynia in inflammatory and neuropathic models of nociception in mice. *Anest. Analg.*, **101**, 1763–1769.
- RIBEIRO, R.A., VALE, M.L., FERREIRA, S.H. & CUNHA, F.Q. (2000). Analgesic effect of thalidomide on inflammatory pain. *Eur. J. Pharmacol.*, **391**, 97–103.
- ROCHA, A.C.C., FERNANDES, E.S., PASSOS, G.F., CALIXTO, J.B. & CAMPOS, M.M. (2005). Assessment of TNF $\alpha$  contribution to the functional up-regulation of kinin B<sub>1</sub> receptors in the mouse paw after treatment with LPS. *Int. Immunopharmacol.*, **5**, 1593–1600.
- SALVEMINI, D., WANG, Z.Q., WYATT, P.S., BOURDON, D.M., MARINO, M.H., MANNING, P.T. & CURRIE, M.G. (1986). Nitric oxide: a key mediator in the early and late phase of carrageenan-induced rat paw inflammation. *Br. J. Pharmacol.*, **118**, 829–838.
- SIQUEIRA-JUNIOR, J.M., PETERS, R.R., BRUM-FERNANDES, A.J. & RIBEIRO-DO-VALLE, R.M. (2003). Effects of valeryl salicylate, a COX-1 inhibitor, on models of acute inflammation in mice. *Pharmacol. Res.*, **48**, 437–443.
- SOMMER, C. & KRESS, M. (2004). Recent findings on how proinflammatory cytokines cause pain: peripheral mechanisms in inflammatory and neuropathic hyperalgesia. *Neurosci. Lett.*, **361**, 184–187.
- SUGISHITA, E., AMAGAYA, S. & OGIHARA, Y. (1981). Anti-inflammatory testing methods: comparative evaluation of mice and rats. *J. Pharmacobiodyn.*, **4**, 565–575.
- TABO, E., EISELE JR, J.H. & CARSTENS, E. (1998). Force of limb withdrawals elicited by graded noxious heat compared with other behavioral measures of carrageenan-induced hyperalgesia and allodynia. *J. Neurosci. Methods*, **81**, 139–149.
- TEO, S.K., STIRLING, D.I. & ZELDIS, J.B. (2005). Thalidomide as a novel therapeutic agent: new uses for an old product. *Drug Discov. Today*, **10**, 107–114.
- WINTER, C.A., RISLEY, E.A. & NUSS, G.W. (1962). Carrageenan-induced edema in hind paw of the rat as an assay for anti-inflammatory drugs. *Proc. Soc. Exp. Biol.*, **111**, 544–547.
- ZIMMERMANN, M. (1983). Ethical guidelines for investigations of experimental pain in conscious animals. *Pain*, **16**, 109–110.

(Received December 2, 2005

Revised March 16, 2006

Accepted April 4, 2006

Published online 15 May 2006)