

# Gold(I)-Catalyzed Intramolecular Hydroamination of Alkyne with Trichloroacetimidates


Ji-Eun Kang, Hak-Beom Kim, Jin-Woo Lee, and Seunghoon Shin\*

Department of Chemistry, Hanyang University, 17 Haengdang-dong, Seongdong-gu, Seoul 133-791, Korea

sshin@hanyang.ac.kr

Received May 29, 2006

## ABSTRACT



A study on the gold (I)-catalyzed intramolecular hydroamination of trichloroacetimidates derived from propargyl and homopropargyl alcohols is described. In the presence of 2–5 mol % of cationic Au(I) complex, a variety of trichloroacetimidates undergo efficient hydroamination under an exceptionally mild condition. An orthogonality of the current catalytic protocol with those using a stoichiometric electrophile as well as a preliminary synthetic application as a stable precursor of 2-acylamino-1,3-diene has been demonstrated.

A formal addition of an N–H bond across C–C multiple bonds, collectively known as hydroamination, has gained a great deal of attention as a simple and atom-economical protocol for N-functionalization.<sup>1</sup> While hydroamination catalyzed by lanthanide, alkali metal, and early transition metal complexes typically takes place through activation of an N–H bond,<sup>2</sup> those catalyzed by late transition metal complexes,<sup>3</sup> such as Pd, Ru, Pt, Ir, Rh, Ni, Ag, and Au, can occur through activation of C–C multiple bonds, followed by attack of the N-nucleophile. Recently, gold catalysts are emerging as efficient  $\pi$ -group activators,<sup>4</sup> and a number of

N-nucleophiles have been reported for Au-catalyzed hydroamination of alkenes and alkynes,<sup>5</sup> including free amine,<sup>5a</sup> aniline,<sup>5b</sup> indole,<sup>5c</sup> sulfonamide,<sup>5d</sup> and carbamate.<sup>5e</sup> Considering that many late transition metals require high reaction temperature, we were prompted to explore new types of N-donors that react in mild reaction conditions under Au catalysis. Herein we report that trichloroacetimidates derived from propargyl and homopropargyl alcohols undergo an efficient Au(I)-catalyzed intramolecular hydroamination to alkyne under exceptionally mild conditions.

In continuation of our interest in gold-catalyzed activation of  $\pi$ -bonds as a catalytic alternative to using a stoichiometric electrophile, such as  $\text{IBr}$ ,<sup>6</sup> we set out to explore the feasibility of imidate as a “soft” nucleophile that can N-functionalize an alkyne that is activated by a catalytic amount of Au(I)

(1) (a) Beller, M.; Seayad, J.; Tillack, A.; Jiao, H. *Angew. Chem., Int. Ed.* **2004**, *43*, 3368 and references therein. (b) Muller, T. E.; Beller, M. *Chem. Rev.* **1998**, *98*, 675. (c) Hong, S.; Marks, T. J. *Acc. Chem. Res.* **2004**, *37*, 673.


(2) (a) Aspinall, H. C. *Chem. Rev.* **2002**, *102*, 1807. (b) Tzalis, D.; Koradin, C.; Knochel, P. *Tetrahedron Lett.* **1999**, *40*, 6193. (c) Ackermann, L.; Bergman, R. G.; Loy, R. N. *J. Am. Chem. Soc.* **2003**, *125*, 11956.

(3) (a) Utsunomiya, M.; Hartwig, J. F. *J. Am. Chem. Soc.* **2003**, *125*, 14286. (b) Shimada, T.; Yamamoto, Y. *J. Am. Chem. Soc.* **2003**, *125*, 6646. (c) Karstedt, D.; Bell, A. T.; Tilley, T. D. *J. Am. Chem. Soc.* **2005**, *127*, 12640. (d) Ahmed, M.; Seayad, A. M.; Jackstell, R.; Beller, M. *J. Am. Chem. Soc.* **2003**, *125*, 10311.

(4) For recent reviews, see: (a) Hashmi, A. S. K. *Gold Bull.* **2004**, *37*, 51. (b) Hoffmann-Roder, A.; Krause, N. *Org. Biomol. Chem.* **2005**, *3*, 387.

(5) (a) Morita, N.; Krause, N. *Org. Lett.* **2004**, *6*, 4121. (b) Mizushima, E.; Hayashi, T.; Tanaka, M. *Org. Lett.* **2003**, *5*, 3349. (c) Li, Z.; Shi, Z.; He, C. *J. Organomet. Chem.* **2005**, *690*, 5049. (d) Zhang, J.; Yang, C.-G.; He, C. *J. Am. Chem. Soc.* **2006**, *128*, 1798. (e) Han, X.; Widenhoefer, R. A. *Angew. Chem., Int. Ed.* **2006**, *45*, 1747.

complex. Although cyclization of imidates of allyl and homoallyl alcohols promoted by a stoichiometric electrophile or mercuric salt is well documented,<sup>7</sup> the corresponding catalytic reaction of the triple bond has received little attention. Two most closely related examples are thermal 5-*exo-dig* cyclization of propargylic benzimidate into 4-methylene-4,5-dihydrooxazole and [3,3]-sigmatropic rearrangement of propargylic trichloroacetimidates, leading to 1-amino-1,3-diene (Scheme 1).<sup>8</sup>




In addition to their limited scope, these thermal reactions require a high temperature (~110 °C). On the other hand, current Au(I) catalysis occurs under exceptionally mild conditions (0 °C to room temperature) to give a Markovnikov product. Thus, trichloroacetimidates **1** and **3** cyclize in 5-*exo-dig* and 6-*exo-dig* mode to give **2** and **4**, respectively, having the usual 4-*exo*-methylene unit. Furthermore, we were intrigued by the possibility of **4** acting as a masked 2-acylamino-1,3-diene for subsequent Diels–Alder cycloaddition. The latter type of diene has scarce literature precedents, due to its thermal instability.<sup>9</sup>

We initiated our study by examining cyclization of propargylic trichloroacetimidates. Using Au(PAr<sub>3</sub>)SbF<sub>6</sub> (Ar = C<sub>6</sub>F<sub>5</sub>) as a catalytic precursor prepared in situ,<sup>6b</sup> a variety

of propargylic imidates cyclized efficiently without further necessity for optimization (Table 1). The 5-*exo-dig* process

**Table 1.** Cycloisomerization of **1**




entry	<b>1</b>	R	catalyst loading	conditions <sup>a</sup>	yield (%)
1	<b>1a</b>	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	1%	0 °C, 3 h	87
2	<b>1b</b>	<i>n</i> -C <sub>7</sub> H <sub>15</sub>	2%	0 °C, 5 h	92
3	<b>1c</b>	<i>i</i> -Pr	2%	0 °C, 8 h	74
4	<b>1d</b>	<i>t</i> -Bu	2%	0 °C, 2 h	89
5	<b>1e</b>	PhCH <sub>2</sub>	2%	0 °C, 5 h	97
6	<b>1f</b>	H	2%	0 °C, 9 h	98

<sup>a</sup> Catalyst was prepared by mixing Au[P(C<sub>6</sub>F<sub>5</sub>)<sub>3</sub>]Cl (5 mol %) and AgSbF<sub>6</sub> (5 mol %) in situ.

proceeded with remarkable efficiency and only 1–2 mol % of catalyst was sufficient to give 4-methylene-4,5-dihydrooxazoles **2a–f** in good yields. Internal alkynes or aryl substrates (R = Ar in **1**), however, were not viable substrates (not shown). It is noteworthy that thermodynamically more stable oxazole compounds **5** (i.e., double bond isomerization) were not observed in the course of the reaction and purification.<sup>10</sup> The lack of isomerization to the oxazole is in sharp contrast to the related cycloisomerizations,<sup>11</sup> clearly demonstrating the mildness of the current protocol.

Next, we examined the reaction parameters for the cyclization of homopropargylic trichloroacetimidates using substrate **3a**. Selected optimization data are shown in Table 2. Using Au(PPh<sub>3</sub>)NTf<sub>2</sub> (5%) as catalyst precursor, various

**Table 2.** Optimization of Cyclization of **3a**



catalyst <sup>a</sup>	solvent	temp (°C)	time	yield <sup>b</sup> (%)
Au(PPh <sub>3</sub> )NTf <sub>2</sub>	CH <sub>3</sub> CN	40	24 h	29
Au(PPh <sub>3</sub> )NTf <sub>2</sub>	CH <sub>3</sub> NO <sub>2</sub>	40	12 h	15
Au(PPh <sub>3</sub> )NTf <sub>2</sub>	DCE	40	4 h	49
AuCl <sub>3</sub>	DCE	40	12 h	37
Au(PAr <sub>3</sub> )SbF <sub>6</sub>	DCE	40	24 h	NR
Au(PAr <sub>3</sub> )NTf <sub>2</sub>	DCE	40	12 h	messy
Au(PPh <sub>3</sub> )OTf	DCE	40	30 min	7
Au(PPh <sub>3</sub> )BF <sub>4</sub>	DCE	40	10 min	57 (62)
Au(PPh <sub>3</sub> )BF <sub>4</sub>	DCE	0	30 min	91
Au(PPh <sub>3</sub> )BF <sub>4</sub>	DCE	0	12 h	79

<sup>a</sup> Catalyst was prepared by mixing Au(PAr<sub>3</sub>)Cl (5 mol %) with appropriate AgX salt in situ. <sup>b</sup> Isolated yields (brsm in parentheses).

(6) (a) Kang, J. E.; Lee, E. S.; Park, S. I.; Shin, S. *Tetrahedron Lett.* **2005**, 72, 4334. (b) Kang, J. E.; Shin, S. *Synlett* **2006**, 717. (c) Shin, S. *Bull. Korean Chem. Soc.* **2005**, 26, 1925. (d) Buzas, A.; Gagosz, F. *Org. Lett.* **2006**, 8, 515. For IBP-promoted reaction, see: (e) Marshall, J. A.; Yanik, M. M. *J. Org. Chem.* **1999**, 64, 3798.

(7) (a) Cardillo, G.; Orena, M. *Tetrahedron* **1990**, 46, 3321. (b) Kang, S. H.; Jun, H. S. *Chem. Commun.* **1998**, 1929. (c) Kang, S. H.; Kim, J. S. *Chem. Commun.* **1998**, 1353.





(8) (a) Overman, L. E.; Tsuboi, S.; Angle, S. *J. Org. Chem.* **1979**, 44, 2323. (b) Overman, L. E.; Clizbe, L. A.; Freerks, R. L.; Marlowe, C. K. *J. Am. Chem. Soc.* **1981**, 103, 2807. For application of 1-acylamino-1,3-diene in Diels–Alder reaction, see: (c) Overman, L. E.; Freerks, R.; Petty, C. B.; Clizbe, L.; Ono, R. K.; Taylor, G. F.; Jessup, P. J. *J. Am. Chem. Soc.* **1981**, 103, 2816.

(9) (a) Chou, S.-S. P.; Chow, T. J.; Hsu, C.-H.; Yang, C.; Long, S.-H.; Lin, C.-D. *J. Chem. Soc., Chem. Commun.* **1997**, 1691. (b) McIntosh, J. M.; Pillon, L. Z. *Can. J. Chem.* **1984**, 62, 2089. See also ref 8a.

solvents were tested and 1,2-dichloroethane was found to be the best suited for the current cyclization, giving **4a** in 49% yield. We then screened a variety of catalysts. Use of  $\text{PAr}_3$  ( $\text{Ar} = \text{C}_6\text{F}_5$ ) ligand that was advantageously used in the formation of enol carbonates and the cycloisomerization of **1** was not effective at all. Interestingly, we found there is a significant counteranion effect in this reaction. While  $\text{NTf}_2$ ,  $\text{SbF}_6$ , and  $\text{OTf}$  as counteranions were ineffective, change of counteranion to  $\text{BF}_4$  led to 62% (brsm) yield of **4a** in just 10 min at 40 °C. Cooling the reaction mixture resulted in a cleaner conversion, giving 91% of **4a**. Lowering the catalyst loading to 2% still led to a reasonable yield, albeit in a prolonged reaction time.

The generality of the current method using the above optimized protocol is demonstrated in Table 3. Aliphatic and

**Table 3.** Cyclization of Homopropargylic Imidates

					
entry	sub	R <sup>1</sup>	R <sup>2</sup>	conditions <sup>a</sup>	yield <sup>b</sup> (%)
1	<b>3b</b>	<i>m</i> -CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	H	0 °C, 3 h	84
2	<b>3c</b>	<i>m</i> -MeO-C <sub>6</sub> H <sub>4</sub>	H	0 °C, 2 h	88
3	<b>3d</b>	<i>p</i> -CN-C <sub>6</sub> H <sub>4</sub>	H	0 °C, 1.5 h	99
4	<b>3e</b>	<i>p</i> -Cl-C <sub>6</sub> H <sub>4</sub>	H	0 °C, 3 h	91
5	<b>3f</b>	H	H	0 °C, 10 min	84
6	<b>3g</b>	<i>c</i> -C <sub>6</sub> H <sub>11</sub>	H	0 °C, 2 h	85
7	<b>3h</b>	<i>n</i> -C <sub>3</sub> H <sub>7</sub>	H	0 °C, 30 min	74
8	<b>3i</b>	<i>t</i> -Bu	H	0 °C, 30 min	78
9	<b>3j</b>	Ph	Ph	0 °C, 4 h	95
10	<b>3k</b>	Ph	SiMe <sub>3</sub> <sup>c</sup>	rt, 24 h	84
11	<b>3l</b>			0 °C, 2 h	80
12	<b>3m</b>			0 °C, 2 h	85
13	<b>3n</b>			0 °C, 2 h	73

<sup>a</sup>  $\text{Au}(\text{PPh}_3)\text{BF}_4$  (5 mol %) in DCE (0.2 M). <sup>b</sup> Isolated yield. <sup>c</sup>  $[\text{Au}(\text{n-Bu}_3\text{P})]\text{BF}_4$  (5 mol %) was used instead.

aromatic groups at R<sup>1</sup> having various steric and electronic demand were well accommodated (entries 1–8). An internal alkyne having a phenyl group at R<sup>2</sup> was also a viable substrate providing (*Z*)-**4j** in 95% isolated yield (entry 9).<sup>12</sup> Substrate **3k** having a TMS group at R<sup>2</sup> required change of ligand to  $\text{P}(n\text{-Bu})_3$  and gave a clean conversion to **4k** (entry

(10) Upon keeping **2a** at room temperature for ~3 days, we started to observe formation of **5**. However, the dihydrooxazoles **2** could be kept for ca. 3 weeks at –20 °C without any decomposition.


(11) In a closely related cycloisomerization of propargyl amide, the intermediate dihydrooxazole with *exo*-methylene was observed only as an intermediate by NMR spectroscopy. See: Hashmi, A. S. K.; Weyrauch, J. P.; Frey, W.; Bats, J. W. *Org. Lett.* **2004**, *6*, 4391 and references therein.

(12) Based on NOE experiments. See Supporting Information.

10). Substrate having a propargylic substituent, **3l** or **3m**, also underwent smooth reaction (entries 11 and 12). Excellent functional group tolerance is exemplified in the formation of **4n**, where the epoxide ring remained intact, underscoring high chemoselectivity of the current Au(I) catalysis (entry 13).


Chemoselectivity is a prime issue in organic chemistry, and we prepared substrates **3o** derived from homoallyl homopropargyl alcohol to further test this selectivity.<sup>13</sup> We found that IBr (2.5 equiv at –78 °C) and  $\text{Au}(\text{PPh}_3)\text{BF}_4$  (5 mol % at 0 °C) showed completely orthogonal reactivities in the activation of alkene and alkyne, providing **4o** and **6**, respectively. This implies that the two reaction conditions could be employed complementarily (Scheme 2).

**Scheme 2**



Finally, we demonstrate the utility of heterocyclic product **4a** as a precursor for 2-acylamino-1,3-diene as the Diels–Alder cycloaddition partner. A preliminary result toward this goal is shown in Scheme 3. Treating **4a** with dimethylacety-

**Scheme 3**



lenedicarboxylate in the presence of  $\text{Zn}(\text{OTf})_2$  (0.2 equiv) in toluene at 70 °C after 2 days provided cycloadduct **7** in 57% yield after chromatography.

In summary, we have demonstrated that trichloroacetimidates derived from propargyl and homopropargyl alcohols undergo exceptionally mild cycloisomerization under Au(I) catalysis to provide 4,5-dihydrooxazoles or 5,6-dihydro-1,3-oxazine with an unusual *exo*-methylene unit. A preliminary application of the 5,6-dihydro-1,3-oxazine as a

(13) Maddess, M. L.; Lautens, M. *Org. Lett.* **2005**, *7*, 3557 and references therein.

stable 2-acylamino-1,3-diene precursor is presented. A study directed at applying this N-functionalization protocol in the context of total synthesis of natural product is currently underway in this laboratory.

**Acknowledgment.** Financial support was provided by the Korean Science and Engineering Foundation (KOSEF, R01-2006-000-11283-0). J.E.K. and H.B.K. thank the BK21 program for a fellowship. The authors gratefully thank the

reviewers for helpful comments on the manuscript organization.

**Supporting Information Available:** Representative experimental procedures for the formation **2** and **4** as well as characterization of compounds **2a–f**, **4a–o**, **6**, and **7**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL061307R