

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/6746567>

# S- Adenosylhomocysteine Analogues with the Carbon-5' and Sulfur Atoms Replaced by a Vinyl Unit

ARTICLE *in* ORGANIC LETTERS · NOVEMBER 2006

Impact Factor: 6.36 · DOI: 10.1021/ol062026m · Source: PubMed

---

CITATIONS

13

---

READS

21

## 2 AUTHORS:



Daniela Andrei

Dominican University

9 PUBLICATIONS 140 CITATIONS

SEE PROFILE



Stanislaw F Wnuk

Florida International University

162 PUBLICATIONS 2,125 CITATIONS

SEE PROFILE

Published in final edited form as:

Org Lett. 2006 October 26; 8(22): 5093–5096. doi:10.1021/ol062026m.

## S-Adenosylhomocysteine Analogues with the Carbon-5' and Sulfur Atoms Replaced by a "Vinyl Unit"

Daniela Andrei and Stanislaw F. Wnuk\*

Department of Chemistry and Biochemistry, Florida International University, Miami, Florida 33199.

### Abstract

Cross-metathesis of suitably protected 5'-deoxy-5'-methyleneadenosines with racemic and chiral *N*-Boc protected six-carbon amino acids bearing a terminal double bond in the presence of the Hoveyda-Grubbs catalyst gave adenosylhomocysteine analogues with the C5'–C6' double bond. Bromination with pyridinium tribromide and dehydrobromination with DBU followed by standard deprotections yielded the 5'-(bromo)vinyl analogue.

The enzyme *S*-adenosyl-L-homocysteine (AdoHcy) hydrolase (EC 3.3.1.1) effects hydrolytic cleavage of AdoHcy to adenosine (Ado) and L-homocysteine (Hcy).<sup>1</sup> The cellular levels of AdoHcy and Hcy are critical because AdoHcy is a potent feedback inhibitor of crucial transmethylation enzymes.<sup>1,2</sup> Also elevated plasma levels of Hcy in humans have been shown to be a risk factor in coronary artery disease.<sup>3</sup>

The X-ray crystal analysis of human AdoHcy hydrolase inactivated with 9-(dihydroxycyclopentene)-adenine<sup>4a</sup> and neplanocin A,<sup>4b</sup> as well as AdoHcy hydrolase from rat liver<sup>4c</sup> established the presence of a water molecule in the active site of the enzyme. This observation made it a high priority to prepare analogues of AdoHcy that closely resemble the natural substrate that bind tightly to the enzyme. Such compounds should form "stable" complexes with the enzyme that would help to identify key binding groups at the active site of the enzyme that interact with the Hcy moiety and participate in subsequent elimination and 'hydrolytic' activity steps.

Based on the previous finding that AdoHcy hydrolase is able to add the enzyme-sequestered water molecule across the 5',6'-double bond of 5'-deoxy-5'-(halo or dihalomethylene) adenosines causing covalent binding inhibition,<sup>5,6</sup> we now describe the synthesis of AdoHcy analogues **A** (X = H) with the 5',6'-olefin motif incorporated in place of the carbon-5' and sulfur atoms (Figure 1). The analogues **A** or **B** (X = halogen) should be substrates for the oxidative activity of the enzyme, and the resulting 3'-keto products might be substrates for the 'hydrolytic' activity. Enzyme-mediated addition of water might occur at C5' or C6' of **A** or **B** to generate new species with hydroxyl or keto (after  $\beta$ -elimination of HBr) binding sites within the enzyme. X-Ray structures of such 'oxidation' and/or 'hydrolytic' activity-bound products might provide important information regarding key residues in the protein and their interactions with substrates (Hcy unit) and/or the sequestered water molecule.

Retrosynthetic analysis indicates that AdoHcy analogue **A** (X = H) can be prepared by construction of a new C5'–C6' double bond via Wittig or metathesis reactions. For example, condensation of adenosine 5'-aldehyde with Wittig-type reagent or cross-metathesis between 5'-deoxy-5'-methyleneadenosine and the appropriate amino acid-derived terminal alkenes

should give **A**. Since nucleoside 5'-aldehydes are unstable in the presence of strong bases required for the generation of non-stabilized phosphorane-Wittig reagents,<sup>7</sup> we decided to target an AdoHcy analogue of type **A** via the cross-metathesis reaction. Another possibility is Pd-catalyzed cross-coupling approaches between  $sp^2$ - $sp^3$  hybridized carbons to form a new C6'-C7' single bond as a key step.<sup>8,9</sup>

Alkylation of protected glycine **1** with 4-bromo-1-butene followed by hydrolysis of the resulting Schiff base derivative<sup>10</sup> **2** yielded racemic 2-amino-5-hexenoate **3** (Scheme 1). Attempted cross-metathesis<sup>11</sup> between 5'-deoxy-2',3'-*O*-isopropylidene-5'-methyleneadenosine **9a**<sup>5a,7</sup> with *N*-benzoyl **4** or *N*-Boc **5** protected amino acids bearing terminal double bond in the presence of 1<sup>st</sup> and 2<sup>nd</sup> (2-imidazolidinylidene-Ru) generation Grubbs catalysts<sup>11c,e</sup> failed to give desired products **10a** or **11a** (Scheme 2). Also, metathesis of the 6-*N*-benzoyl adenosine substrate **9b** with **4** or **5** was unsuccessful. It is noteworthy that metathesis between 5'-deoxy-2',3'-*O*-isopropylidene-5'-methylenuridine<sup>12</sup> and **4** ( $\text{CH}_2\text{Cl}_2$ /2<sup>nd</sup> generation Grubbs catalyst) afforded the desired product of type **10** (i.e., B = U; 62%)<sup>13</sup> in addition to two dimers resulting from the self-metathesis of nucleoside<sup>14</sup> and amino acid<sup>15</sup> (e.g. **17**) substrates.

We found however that treatment of **9b** with **4** in the presence of Hoveyda-Grubb's catalyst<sup>16</sup> (*o*-isopropoxy-phenylmethylene-Ru) led to the formation of metathesis product **10b** (51%) in addition to dimer **17** (11%) while self-metathesis of **9b** was not observed. Metathesis of the 6-*N,N*-dibenzoyl **9c** with **4** gave **10c** in 60% yield in addition to dimer **17** (18%). The protection of 6-amino group of the adenine ring seems to be necessary because metathesis between **9a** and **4** or **5** in the presence of Hoveyda-Grubbs catalyst did not yield the corresponding product **10a** or **11a**.

Metathesis of **9b** and **9c** with *N*-Boc protected **5** gave **11b** (61%) and **11c** (76%) in higher isolated yields. Moreover, by-products of the self-metathesis of amino acid or nucleoside substrates were not isolated. The cross-metathesis products **10** and **11** were found to be predominantly the *trans* isomers.<sup>17</sup> Purification on a silica gel column afforded **10** and **11** as 5'*E* isomers of a ~1:1 mixture of 9'*R/S* diastereomers. The *E* stereochemistry for **10** and **11** was established from <sup>1</sup>H NMR spectra based on the magnitude of  $J_{\text{H}5'-\text{H}6'}$ . For example, the 5' proton in **11c** appears at  $\delta$  5.58 (dd,  $J_{\text{H}5'-\text{H}4'} = 7.3$  Hz and  $J_{\text{H}5'-\text{H}6'} = 15.2$  Hz) while the 6' proton resonates at  $\delta$  5.73 (dt,  $J_{\text{H}6'-\text{H}7'/7''} = 6.5$  Hz and  $J_{\text{H}5'-\text{H}6'} = 15.2$  Hz).

Deprotection of **10** or **11** turned out to be more challenging than we expected. Thus, treatment of **11c** (or **11b**) with a 1:1 mixture of saturated (at ~0 °C) methanolic ammonia solution and methanol for 48 h at ~5 °C removed the 6-*N*-benzoyl group(s) and produced a partially separable mixture of methyl **12** and ethyl **13** esters (~3:2, ~92% total yield). Using diluted  $\text{NH}_3/\text{MeOH}$  minimized formation of the amidation byproducts (~5%). Acid-catalyzed deprotection of **12** and **13** with an aqueous solution of trifluoroacetic acid (TFA) effected the removal of both Boc and the isopropylidene protection groups to give **14** and **15** in high yields. It is important to perform debenzoylation of **11c** (or **11b**) as initial deprotection step, because treatment of **11c** (or **11b**) with TFA/ $\text{H}_2\text{O}$  resulted in the substantial cleavage of the glycosylic bond. Saponification of **14** and **15** with NaOH in  $\text{H}_2\text{O}/\text{MeOH}$  solution and purification on RP-HPLC afforded the sodium salt of **16** [67%; *E*, 9'*R/S* (~1:1)].

Since separation of 9'*R/S* diastereomers in products **10–16** was difficult, we attempted the synthesis of analogue **A** with 9'*S* configuration employing a chiral amino acid precursor e.g., (*S*)-homoallylglycine. Given that the methods available for the preparation of enantiomerically pure unnatural amino acids usually require multistep synthesis,<sup>18</sup> we chose the enantioselective hydrolysis of racemic **5** as a way to provide chiral (*S*)-homoallylglycine. Thus, treatment of **5** with  $\alpha$ -chymotrypsin in phosphate buffer (24 h, 37 °C)<sup>19</sup> gave the unreacted

(*R*)-ester **5** (~50%) and (*S*)-acid **6** (~50%, Scheme 1). Enantiomeric purity of the **5-R** was established using the Mosher test.<sup>20a</sup> Thus, treatment of **5-R** with TFA/H<sub>2</sub>O followed by acylation with (*R*)-2-methoxy-2-trifluoromethyl-2-phenylacetyl chloride (MTPA-Cl)<sup>20</sup> gave **8-R/S**. (Note that the absolute configuration at the chiral carbon in Mosher reagent is the same but the *R/S* descriptors change owing to the change in Cahn-Ingold-Prelog priority.) Analysis of the <sup>19</sup>F NMR spectra [ $\delta$  -69.16 (s, 0.02F) and -69.55 (s, 0.98F)] established the stereochemistry for **5** as *R* (ee 96%) in agreement with Mosher's empirical formula.<sup>20a</sup> Since metathesis of the "free" carboxylic acid precursor **6-S** with **9b** or **9c** in the presence of Hoveyda-Grubbs catalyst failed, the **6-S** was converted into the methyl ester **7-S** with diazomethane.

Cross-metathesis of **9c** with **7-S** afforded **18-S** (77%; Scheme 3). Sequential deprotections of **18-S** with NH<sub>3</sub>/MeOH (to give **12-S**, 91%) and TFA/H<sub>2</sub>O gave the enantiomerically pure **14-S** (90%) as the single *E* isomer (Scheme 3). On the other hand metathesis of **9c** with **5-R** gave ethyl ester **11c-R**. Contrary to products **10-16** obtained from racemic homoallylglycine, the <sup>13</sup>C NMR spectra for the products obtained from (*S*)- and (*R*)- homoallylglycine substrates showed a single set of peaks.

Finally, we attempted the synthesis of bromovinyl analogue **B** by the bromination-dehydrobromination strategy. Treatment of **11c** with pyridinium tribromide<sup>21</sup> gave the 5',6'-dibromo diastereomers **19** which were dehydrobrominated with 1,8-diazobicyclo[5.4.0]undec-7-ene (DBU) to yield **20** as a single isomer (one of the 6-*N*-benzoyl protective group was also partially cleaved) in 70% yield (Scheme 4). Standard deprotections with NH<sub>3</sub>/MeOH and TFA/H<sub>2</sub>O followed by saponification with NaOH and HPLC purification gave **23** (*E*, 54% overall).

The regioselectivity of HBr elimination and the position of bromine (at 5') were assigned based on the presence of a triplet signal for the olefinic hydrogen (H6') in <sup>1</sup>H NMR spectra [ $\delta$  6.40 (t,  $J_{6-7'/7''} = 7.6$  Hz) for **23**]. This assignment was also supported by COSY experiment. The product *E* configuration is expected from a specific anti-addition in the pyridinium tribromide bromination of the *E* alkene **11c** followed by an E2 (anti elimination) process. This was also supported by NOESY analysis of **23** in which the cross-peaks between H4' and H7'/7'' were observed.

In summary, we have developed a synthesis of AdoHcy analogues in which the carbon-5' and sulfur atoms is replaced by a "vinyl unit" utilizing cross-metathesis reactions between 5'-deoxy-5'-methyleneadenosine analogues and homoallylglycine in the presence of Hoveyda-Grubbs catalyst. The 5'-(bromo)vinyl AdoHcy analogue has been prepared via the bromination-dehydrobromination strategy. Enzymatic studies with AdoHcy hydrolase and our attempts to synthesize 6'-(halo)vinyl analogues **B** via cross-coupling approaches will be published elsewhere.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgment


We thank NIH/NIGMS program (S06 GM08205) for supporting this research, US ARO (W911NF-04-1-0022) for financial help to purchase 600 MHz NMR spectrometer and Materia Company for a gift of Hoveyda-Grubbs catalyst.

## References


1. Yuan, C.-S.; Liu, S.; Wnuk, S.F.; Robins, M.J.; Borchardt, R.T. *Advances in Antiviral Drug Design*. De Clercq, E., editor. Vol. 2. Greenwich: JAI Press; 1996. p. 41-88. (b) Turner MA, Yang X, Yin D,

- Kuczera K, Borchardt RT, Howell PL. *Cell Biochem. Biophys* 2000;33:101–125. [PubMed: 11325033] (c) Wnuk SF. *Mini-Rev. Med. Chem* 2001;1:307–316. [PubMed: 12369977]
2. (a) Ueland PM. *Pharmacol. Rev* 1982;34:223–253. [PubMed: 6760211] (b) Chiang PK. *Pharmacol. Ther* 1998;77:115–134. [PubMed: 9578320]
  3. (a) Nehler MR, Taylor LM, Porter JM. *Cardiovasc. Surgery* 1997:559–567. (b) Refsum H, Ueland PM, Nygard O, Vollset SE. *Annu. Rev. Med* 1998;49:31–62. [PubMed: 9509248] (c) Schynder G, Roffi M, Pin R, Flammer Y, Lange H, Eberli FR, Meier B, Turi ZG, Hess OM. *N. Eng. J. Med* 2001;345:1593–1600.
  4. (a) Turner MA, Yuan C-S, Borchardt RT, Hershfield MS, Smith GD, Howell PL. *Nature Struct. Biol* 1998;5:369–376. [PubMed: 9586999] (b) Yang X, Hu Y, Yin DH, Turner MA, Wang M, Borchardt RT, Howell PL, Kuczera K, Schowen RL. *Biochemistry* 2003;42:1900–1909. [PubMed: 12590576] (c) Hu Y, Komoto J, Huang Y, Gomi T, Ogawa H, Takata Y, Fujioka M, Takusagawa F. *Biochemistry* 1999;38:8323–8333. [PubMed: 10387078]
  5. (a) Wnuk SF, Yuan C-S, Borchardt RT, Balzarini J, De Clercq E, Robins MJ. *J. Med. Chem* 1994;37:3579–3587. [PubMed: 7932585] (b) Wnuk SF, Mao Y, Yuan C-S, Borchardt RT, Andrei J, Balzarini J, De Clercq E, Robins MJ. *J. Med. Chem* 1998;41:3078–3083. [PubMed: 9685247] (c) Yuan C-S, Wnuk SF, Robins MJ, Borchardt RT. *J. Biol. Chem* 1998;273:18191–18197. [PubMed: 9660780]
  6. For other examples on the hydrolytic activity of AdoHcy hydrolase see (a) Ref 1c (b) Yang X, Yin D, Wnuk SF, Robins MJ, Borchardt RT. *Biochemistry* 2000;39:15234–15241. [PubMed: 11106503] (4'-haloacetyleneadenosine analogues) (c) Guillerme G, Guillerme D, Vandenplas-Witkowski C, Roginaux H, Carte N, Leize E, Van Dorsselaer A, De Clercq E, Lambert C. *J. Med. Chem* 2001;44:2743–2752. [PubMed: 11495586] (5'-S-allenyl-5'-thioadenosine) (d) Jeong LS, Yoo SJ, Lee KM, Koo MJ, Choi WJ, Kim HO, Moon HR, Lee MY, Park JG, Lee SK, Chun MW. *J. Med. Chem* 2003;46:201–203. [PubMed: 12519056] (fluoro-neplanocin A) (e) Wnuk SF, Lewandowska E, Sacasa PR, Crain LN, Zhang J, Borchardt RT, De Clercq E. *J. Med. Chem* 2004;47:5251–5257. [PubMed: 15456269] (4'-enyne adenosine analogues) (f) Guillerme G, Muzard M, Glapski C. *Bioorg. Med. Chem. Lett* 2004;14:5799–5802. [PubMed: 15501043] (haloethyl esters of homoadenosine-6'-carboxylic acid) (g) Guillerme G, Muzard M, Glapski C, Pilard S, De Clercq E. *J. Med. Chem* 2006;49:1223–1226. [PubMed: 16480257] [5'-deoxy-5'-(cyanomethylene)adenosine]
  7. Wnuk SF, Robins MJ. *Can. J. Chem* 1991;69:334–338.
  8. For example couplings between 5'-deoxy-5'-(iodomethylene)-adenosine<sup>5a</sup> and suitable alkylzinc bromides produced analogues of type A: Wnuk SF, Lalama J, Andrei D, Garmendia C, Robert J, S-Adenosylhomocysteine and S-ribosylhomocysteine analogues with sulfur atom replaced by the vinyl unit. Abstracts of Papers, Carbohydrate Division 2005 March 22–29th National Meeting of the American Chemical Society San Diego, CA 1317 CARB-035
  9. (a) Pd-catalyzed alkylation of the 5'-deoxy-5'-(dihalomethylene)-adenosine<sup>5b</sup> precursors employing recently reporting selective monoalkylation of the unactivated 1,1-dichloro-1-alkenes<sup>9b</sup> or 1-fluoro-1-halo-1-alkenes<sup>9c</sup> might give direct access to analogues B. (b) Tan Z, Negishi E-I. *Angew. Chem., Int. Ed* 2006;45:762–765. (c) Andrei D, Wnuk SF. *J. Org. Chem* 2006;71:405–408. [PubMed: 16388671]
  10. (a) O'Donnell MJ, Wojciechowski K. *Synthesis* 1984:313–315. (b) O'Donnell MJ, Polt RL. *J. Org. Chem* 1982;47:2663–2666.
  11. (a) Grubbs RH, Chang S. *Tetrahedron* 1998;54:4413–4450. (b) Fürstner A. *Angew. Chem., Int. Ed* 2000;39:3012–3043. (c) Trnka TM, Grubbs RH. *Acc. Chem. Res* 2001;34:18–29. [PubMed: 11170353] (d) Chatterjee AK, Choi T-L, Sanders DP, Grubbs RH. *J. Am. Chem. Soc* 2003;125:11360–11370. [PubMed: 16220959] (e) Nicolaou KC, Bulger PG, Sarlah D. *Angew. Chem., Int. Ed* 2005;44:4490–4527.
  12. Wnuk SF, Robins MJ. *Can. J. Chem* 1993;71:192–198.
  13. Sacasa, Pablo R. M.Sc. Thesis. Florida International University; 2003.
  14. (a) For recent reviews on application of metathesis towards synthesis of nucleoside analogues see (a) Agrofoglio LA, Nolan SP. *Curr. Top. Med. Chem* 2005;5:1541–1558. [PubMed: 16378491] Amblard F, Nolan SP, Agrofoglio LA. *Tetrahedron* 2005;61:7067–7080. (b) For an example on self metathesis reaction of carbohydrate derived terminal olefins (e.g., 5,6-dideoxy-1,2-O-isopropylidene- $\alpha$ -D-ribo-hex-5-enofuranose) see: Hadwiger P, Stütz AE. *Synlett* 1999:1787–1789

15. (a) Gibson SE, Gibson VC, Keen SP. *Chem. Commun* 1997;1107–1108. (b) Biagini SCG, Gibson SE, Keen SP. *J. Chem. Soc., Perkin Trans. 1* 1998;2485–2499. (c) Vasbinder MM, Miller SJ. *J. Org. Chem* 2002;67:6240–6242. [PubMed: 12182670]
16. (a) Garber SB, Kingsbury JS, Gray BL, Hoveyda AH. *J. Am. Chem. Soc* 2000;122:8168–8179. (b) Gessler S, Randl S, Blechert S. *Tetrahedron Lett* 2000;41:9973–9976.
17.  $^1\text{H}$  NMR analysis of the crude reaction mixtures showed the presence of other isomers in variable quantities of ~2–6%.
18. (a) Dunn MJ, Jackson RFW, Pietruszka J, Turner D. *J. Org. Chem* 1995;60:2210–2215. (b) Waelchli R, Beerli C, Meigel H, Revesz L. *Bioorg. Med. Chem. Lett* 1997;7:2831–2836. (c) Löhr B, Orlich S, Kunz H. *Synlett* 1999:1139–1141. (d) Bachmann S, Knudsen KR, Jorgensen KA. *Org. Biomol. Chem* 2004;2:2044–2049. [PubMed: 15254632]
19. Schricker B, Thirring K, Berner H. *Bioorg. Med. Chem. Lett* 1992;2:387–390.
20. (a) Sullivan GR, Dale JA, Mosher HS. *J. Org. Chem* 1973;38:2143–2147. (b) Oh SS, Butler WM, Koreeda M. *J. Org. Chem* 1989;54:4499–4503.
21. Husstedt U, Schäfer HJ. *Tetrahedron Lett* 1981:623–624.




**Figure 1.**  
*S*-Adenosyl-L-homocysteine and analogues with sulfur atom replaced by the "vinyl unit".




Scheme 1.






Scheme 2.



Scheme 3.



Scheme 4.