

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/41547422>

# Tricyclic Alkaloid Core Structures Assembled by a Cyclotrimerization–Coupled Intramolecular Nucleophilic Substitution Reaction

ARTICLE in ORGANIC LETTERS · FEBRUARY 2010

Impact Factor: 6.36 · DOI: 10.1021/ol100177u · Source: PubMed

---

CITATIONS

36

---

READS

32

## 2 AUTHORS:



[Andrew L Mciver](#)

University of North Carolina at Chapel Hill

10 PUBLICATIONS 126 CITATIONS

SEE PROFILE



[Alexander Deiters](#)

University of Pittsburgh

135 PUBLICATIONS 5,049 CITATIONS

SEE PROFILE

# Tricyclic Alkaloid Core Structures Assembled by a Cyclotrimerization—Coupled Intramolecular Nucleophilic Substitution Reaction


Andrew L. McIver and Alexander Deiters\*

North Carolina State University, Department of Chemistry,  
Raleigh, North Carolina 27695-8204

alex\_deiters@ncsu.edu

Received January 24, 2010

## ABSTRACT



A facile approach to tricyclic alkaloid core structures was developed by sequencing a pyridine-forming [2 + 2 + 2] cyclotrimerization reaction with an intramolecular nucleophilic substitution. This methodology enabled the facile assembly of the spiroindolinone framework of citrinadins A and B, and cyclopiamine B.

Tricyclic alkaloid structures are present in a wide range of natural products, many of which have important biological activity. Figure 1 shows six examples of these types of natural products with the central tricyclic alkaloid core highlighted in red. The citrinadins A (**1**) and B (**2**) are recently isolated marine derived pentacyclic spiroindolinone alkaloids containing a dodecahydrocyclopenta[b]quinolizine core.<sup>1,2</sup> They both exhibit important cytotoxic activities against various cancer cell lines with IC<sub>50</sub>'s in the low micromolar range.<sup>1,2</sup> No total synthesis of **1** or **2** has been reported to date, but one stereoselective approach to the spirooxindole A,B,C-ring system in seven steps has been accomplished recently.<sup>3</sup> Cyclopiamine B (**3**) is another fungal spiroindolinone alkaloid<sup>4,5</sup> containing six rings including a central tricyclic decahydro-1*H*-cyclopenta[*f*]indolizine struc-

ture. The biological function of **3** is unknown, and no total synthesis of **3** has been reported.

Other notable natural products containing similar tricyclic alkaloid structures are veraflorizine (**4**), a steroidal cevanine alkaloid<sup>6</sup> that has also not been synthesized, roserine (**5**), a pyrrolophenanthridinium alkaloid<sup>7</sup> that has been synthesized once,<sup>8</sup> and xylopinine (**6**), a protoberberine alkaloid, which is part of a large group of isoquinoline alkaloids that have attracted considerable synthetic interest due to their diverse biological activities.<sup>9</sup>

Building onto our recent applications of microwave-mediated [2 + 2 + 2] cyclotrimerization reactions in the

(1) Tsuda, M.; Kasai, Y.; Komatsu, K.; Sone, T.; Tanaka, M.; Mikami, Y.; Kobayashi, J. *Org. Lett.* **2004**, *6*, 3087–3089.

(2) Mugishima, T.; Tsuda, M.; Kasai, Y.; Ishiyama, H.; Fukushi, E.; Kawabata, J.; Watanabe, M.; Akao, K.; Kobayashi, J. *J. Org. Chem.* **2005**, *70*, 9430–9435.

(3) Pettersson, M.; Kneuppel, D.; Martin, S. F. *Org. Lett.* **2007**, *9*, 4623–4626.

(4) Bond, R. F.; Boeyens, J. C. A.; Holzapfel, C. W.; Steyn, P. S. *J. Chem. Soc., Perkin Trans. 1* **1979**, *7*, 1751–1761.


(5) Steyn, P. S.; Vleggaar, R.; Rabie, C. J. *Phytochemistry* **1981**, *20*, 538–539.

(6) Kaneko, K.; Kawamura, N.; Kuribayashi, T.; Tanaka, M.; Mitsuhashi, H.; Koyama, H. *Tetrahedron Lett.* **1978**, *48*, 4801–4804.

(7) Bastida, J.; Codina, C.; Viladomat, F.; Rubiralta, M.; Quirion, J. C.; Weniger, B. *J. Nat. Prod.* **1992**, *55*, 134–136.

(8) Bunnell, A. E.; Flippin, L. A.; Liu, Y. Z. *J. Org. Chem.* **1997**, *62*, 9305–9313.

(9) Chrzanowska, M.; Rozwadowska, M. D. *Chem. Rev.* **2004**, *104*, 3341–3370.



**Figure 1.** Selected natural products with tricyclic alkaloid core structures shown in red.


synthesis of alkaloids,<sup>10,11</sup> we are reporting the sequencing of a cyclotrimerization with an intramolecular pyridinium formation via a nucleophilic substitution to rapidly access a variety of tricyclic alkaloid structures; including the ones present in the natural products shown in Figure 1. Moreover, we are reporting the synthesis of the pentacyclic spiroindolinone core of citrinadin A (1), citrinadin B (2), and cyclopamine B (3).

The classical [2 + 2 + 2] cyclotrimerization reaction toward pyridines involves the reaction of two alkynes and a nitrile.<sup>12–14</sup> In order to avoid chemoselectivity issues in the cyclotrimerization step, the two alkynes are often tethered together, leading to the synthesis of fused pyridine rings.<sup>12–16</sup> These cyclotrimerization reactions are typically conducted under cobalt catalysis.<sup>12–17</sup> Recently, it was discovered by others<sup>18–22</sup> and us<sup>23,24</sup> that microwave irradiation<sup>25–27</sup> greatly enhances the rates and yields of [2 + 2 + 2] cyclotrimerization reactions. In the case of Co-catalyzed reactions, reaction times are reduced from days to minutes without the necessity of catalyst activation through additives or light irradiation.

We speculated that a [2 + 2 + 2] cyclotrimerization reaction of the commercially available diynes **7** and **8** with

the nitrile **9** tethered to a leaving group X would deliver the intermediate fused pyridine **10** (Scheme 1). A tandem intramolecular S<sub>N</sub>2 reaction would directly form the tricyclic pyridinium compounds **11**, which could subsequently be reduced, e.g., with NaBH<sub>4</sub>, to give the tricyclic structures **12**. This reaction sequence could provide the alkaloid core structures found in the natural products in Figure 1 in as little as two steps.


**Scheme 1.** [2 + 2 + 2] Cyclotrimerization Reaction Coupled with an Intramolecular S<sub>N</sub>2 Reaction Enables the Rapid Assembly of Tricyclic Pyridinium Ions **11** and a Subsequent Reduction Delivers the Alkaloid Core Structures **12**<sup>a</sup>



<sup>a</sup> X = Br, I, OSO<sub>2</sub>CH<sub>3</sub>; m, n ≥ 1.

The investigation of this approach commenced with commercially available 1,6-heptadiyne (**7**) or 1,7-octadiyne (**8**) by reacting each with either 4-bromobutyronitrile (**13**) or the corresponding cyano mesylate **14**<sup>28</sup> to directly produce the tricyclic pyridinium structures **15–18** (Scheme 2). The tricyclic molecules **15–18** were obtained in 22–45% yield (40 min). Increasing the reaction times to 1 h or increasing the amount of the nitrile **13** or **14** to >10 equiv did not afford improved yields.

**Scheme 2.** Tandem [2 + 2 + 2] Cyclotrimerization–Substitution Reactions Delivering the Pyridinium Compounds **15–18** with Bromide and Mesylate Counterions




m	X	compd (yield)
1	Br	<b>15</b> (33%)
2	Br	<b>16</b> (45%)
1	OMs	<b>17</b> (22%)
2	OMs	<b>18</b> (36%)

We suspected that the modest yields for **15–18** were caused by decomposition of the pyridinium salts due to their strong absorption of microwave irradiation based on their ionic nature<sup>25–27</sup> leading to localized heating in the microwave reactor. Additionally, the intermolecular reaction of the pyridine intermediate **10** with an excess alkyl bromide

- (10) Teske, J. A.; Deiters, A. *J. Org. Chem.* **2008**, *73*, 342–345.  
 (11) McIver, A.; Young, D. D.; Deiters, A. *Chem. Commun.* **2008**, 39, 4750–4752.  
 (12) Schore, N. E.; Trost, B. M.; Fleming, I.; Paquette, L. A. [2 + 2 + 2] Cycloadditions. In *Comprehensive Organic Synthesis*; Pergamon Press: Oxford, 1991.  
 (13) Kotha, S.; Brahmachary, E.; Lahiri, K. *Eur. J. Org. Chem.* **2005**, 22, 4741–4767.  
 (14) Chopade, P. R.; Louie, J. *Adv. Synth. Catal.* **2006**, *348*, 2307–2327.  
 (15) Yamamoto, Y. *Curr. Org. Chem.* **2005**, *9*, 503–519.  
 (16) Schore, N. E. *Chem. Rev.* **1988**, *88*, 1081–1119.  
 (17) Aalbersberg, W. G. L.; Barkovich, A. J.; Funk, R. L.; Hillard, R. L.; Vollhardt, K. P. C. *J. Am. Chem. Soc.* **1975**, *97*, 5600–5602.  
 (18) Shanmugasundaram, M.; Aguirre, A. L.; Leyva, M.; Quan, B.; Martinez, L. E. *Tetrahedron Lett.* **2007**, *48*, 7698–7701.  
 (19) Zhou, Y.; Porco, J. A.; Snyder, J. K. *Org. Lett.* **2007**, *9*, 393–396.  
 (20) Hrdina, R.; Kadlcikova, A.; Valterova, I.; Hodacova, J.; Kotora, M. *Tetrahedron: Asymmetry* **2006**, *17*, 3185–3191.

**13** or mesylate **14** could potentially lead to a competing side reaction. In order to solve this problem, we employed a two-step cyclotrimerization–substitution process. Here, the diynes **7** and **8** underwent a smooth cyclotrimerization reaction using the  $\text{CpCo}(\text{CO})_2$  catalyst in toluene under microwave irradiation (300 W) for 40 min with the known cyano alcohols **19** and **20**.<sup>28</sup> This afforded the fused bicyclic pyridine rings **21–24**, bearing an  $\varepsilon$ -hydroxyalkyl chain (Scheme 3), in

**Scheme 3.** Two-Step [2 + 2 + 2] Cyclotrimerization–Substitution Reaction Followed by Reduction to the Indolizines and Quinolizines **29–32**




89–95% yield. The hydroxy group was then converted into a mesylate in situ using  $\text{MsCl}$  and polymer-bound piperidine as the base, thus obviating purification and affording clean products **25–28** in almost quantitative yield. The overall yield for formation of the tricyclic pyridinium compounds **25–28** from the diynes **7–8** was greater than 80%, making this two-step reaction favorable over the tandem one-step reaction depicted in Scheme 2. The reduction of the pyridinium rings was accomplished using  $\text{NaBH}_4$ <sup>29</sup> (Scheme 3) to afford the amines **29–32** that display the tricyclic motif found in the natural products **1–6** (Figure 1). The remaining double bond represents a valuable handle for potential further functionalization toward the installation of the substituents present in citrinadin A and B (**1** and **2**), and cyclopiamine B (**3**).

In order to demonstrate the generality of the developed approach, another reaction sequence was performed that would lead to the core structure of xylopinine (**6**). The nitrile **33** was synthesized starting from commercially available 2-iodophenylacetic acid, which was converted to the methyl

ester in 94% yield, followed by a reaction with  $\text{CuCN}$ <sup>30</sup> delivering the nitrile **33** in 85% yield. The nitrile **33** was reacted in a Co-catalyzed cyclotrimerization reaction with the two diynes **7** and **8** in toluene under microwave irradiation (300 W) for 40 min to give the pyridines **34** and **35** in 93% and 86% yield, respectively (Scheme 4). This

**Scheme 4.** Three-step Cyclotrimerization–Substitution Reaction to form the Tetracyclic Pyridinium Compounds **38–39** Followed by Reduction to **40–41**



was followed by a  $\text{LiAlH}_4$  reduction of the esters **34** and **35** to the alcohols **36** and **37**. The pyridinium formation was conducted under the previously developed cyclization conditions using  $\text{MsCl}$  and polymer bound piperidine to afford the tricyclic compounds **38** and **39** in near-quantitative yields. Reduction of **38** and **39** with  $\text{NaBH}_4$  delivered the tetracyclic molecules **40** and **41** in quantitative yield (Scheme 4).

The developed route shown in Scheme 3 was applied to the synthesis of the core structure of citrinadins A (**1**) and B (**2**) and cyclopiamine B (**3**). The synthesis commenced with the known ester **42** (Scheme 5A).<sup>31</sup> Reduction of the nitro

(21) Saaby, S.; Baxendale, I. R.; Ley, S. V. *Org. Biomol. Chem.* **2005**, *3*, 3365–3368.

(22) Efskind, J.; Undheim, K. *Tetrahedron Lett.* **2003**, *44*, 2837–2839.

(23) Young, D. D.; Deiters, A. *Angew. Chem., Int. Ed.* **2007**, *46*, 5187–5190.

(24) Young, D. D.; Sripada, L.; Deiters, A. *J. Comb. Chem.* **2007**, *9*, 735–738.

(25) Kappe, C. O.; Dallinger, D. *Nat. Rev. Drug Discovery* **2006**, *5*, 51–63.

(26) Kappe, C. O. *Angew. Chem., Int. Ed.* **2004**, *43*, 6250–6284.


(27) Loupy, A., *Microwaves in organic synthesis*, 2nd ed.; Wiley-VCH: Weinheim, 2006.

(28) Deutsch, H. M.; Ye, X. C.; Shi, Q.; Liu, Z. Z.; Schweri, M. M. *Eur. J. Med. Chem.* **2001**, *36*, 303–311.

(29) Blagg, J.; Coote, S. J.; Davies, S. G.; Mobbs, B. E. *J. Chem. Soc., Perkin Trans. 1* **1986**, *12*, 2257–2261.

(30) Le, T. N.; Cho, W. J. *Bull. Korean Chem. Soc.* **2007**, *28*, 763–766.

**Scheme 5.** (A) Synthesis of Racemic **49** and **50**, the Core Structures of Citrinadin A (**1**) and B (**2**), as Well as Cyclopamine B (**3**).<sup>a</sup> (B) Reaction of the Terminal Diyne **51** Produced the Pyridine **52** in Good Yield



<sup>a</sup> brsm = based on recovered starting material.

group with zinc and ammonium chloride<sup>31,32</sup> followed by a Mitsunobu reaction<sup>33</sup> with methanol produced the known methoxy oxindole **43** in 60% yield over both steps. Treatment of **43** with 3-bromo-1-trimethylsilyl-1-propyne and NaH generated the diyne **44** in 65% yield, which set the stage for the [2 + 2 + 2] cyclotrimerization reaction. The cyclotrimerization reaction was accomplished by reacting the diyne **44** with 4-hydroxypentanenitrile (**20**, n = 2) for the citrinadin A (**1**) and B (**2**) core and with 4-hydroxybutanenitrile (**19**, n = 1) for the cyclopamine B (**3**) core in the presence of

CpCo(CO)<sub>2</sub> under microwave irradiation (300 W, 90 min). This delivered the pyridines **45** and **46** in 42% and 41% yield, respectively. In addition, 57% and 50% of the starting material **44** was recovered, but extending reaction times led to unidentified by-products. Gratifyingly, both compounds were obtained as single regioisomers, since the bulky trimethylsilyl (TMS) group on **44** directs the formation of the desired pyridine regioisomer.<sup>34–36</sup> However, the sterically demanding TMS group also leads to the moderate cyclotrimerization yields, as previously observed.<sup>34,37–39</sup> As expected, the cyclotrimerization reaction with the terminal diyne **51** delivered the pyridine **52** in a much higher yield of 83% (Scheme 5B). The majority of the TMS moiety was removed from the cyclotrimerization product of **44** under the cyclotrimerization microwave conditions, and any remaining silyl groups were cleaved by a subsequent treatment with potassium fluoride under microwave irradiation (300 W, 2 min) to give the desired products **45** and **46**. The substitution and reduction sequence following Scheme 3 was then employed by treating the alcohols **45** and **46** with MsCl in the presence of polymer-bound piperidine to produce the pyridinium compounds **47** and **48** in excellent yields. Reduction with NaBH<sub>4</sub> completed the pentacyclic spiroindolinone framework **49** of citrinadin A (**1**) and B (**2**) in 72% yield and the alkaloid core structure **50** of cyclopamine B (**3**) in 82% yield. The overall yield of the assembly of **49** and **50** from the common diyne **44** was 30% and 28%, respectively.

In summary, we developed an expedient route to tricyclic alkaloid core structures by conducting a microwave-mediated [2 + 2 + 2] cyclotrimerization/intramolecular nucleophilic substitution/reduction sequence. This methodology was demonstrated to deliver several tricyclic frameworks in good to excellent yield. These represent the core structures in a variety of natural alkaloids with important biological activities.

**Acknowledgment.** Financial support from the National Science Foundation (CAREER Award 0846756 to A.D.) is acknowledged.

**Supporting Information Available:** Detailed experimental procedures, analytical data, and copies of NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL100177U

(31) Somei, M.; Sato, H.; Komura, N.; Kaneko, C. *Heterocycles* **1985**, 23, 1101–1106.

(32) Somei, M.; Yamada, F.; Kurauchi, T.; Nagahama, Y.; Hasegawa, M.; Yamada, K.; Teranishi, S.; Sato, H.; Kaneko, C. *Bull. Korean Chem. Soc.* **2001**, 49, 87–96.

(33) Henmi, T.; Sakamoto, T.; Kikugawa, Y. *Heterocycles* **1997**, 44, 157–163.

(34) Vollhardt, K. P. C. *Angew. Chem., Int. Ed.* **1984**, 23, 539–556.

(35) Hillard, R. L.; Parnell, C. A.; Vollhardt, K. P. C. *Tetrahedron* **1983**, 39, 905–911.

(36) Parnell, C. A.; Vollhardt, K. P. C. *Tetrahedron* **1985**, 41, 5791–5796.

(37) Bonnemant, H. *Angew. Chem., Int. Ed.* **1985**, 24, 248–262.

(38) Saa, C.; Crofts, D. D.; Hsu, G.; Vollhardt, K. P. C. *Synlett* **1994**, 7, 487–489.

(39) Varela, J. A.; Castedo, L.; Saa, C. *J. Org. Chem.* **1997**, 62, 4189–4192.