

N-(3,4-Dihydro-4-oxoquinazolin-3-yl)pyridinium Imides: Aziridinating Agents for Alkenes: Measurement of Rotational Barriers around each N–N Bond in an N–N–N System

Robert S. Atkinson,^a Emma Barker,^a Paul J. Edwards^a and Gordon A. Thomson^b

^a Department of Chemistry, Leicester University, Leicester, UK LE1 7RH

^b Zeneca Specialties, PO Box 42, Hexagon House, Blackley, Manchester, UK M9 8ZS

Pyridinium imides **3–5** and **8** are obtained as crystalline solids from the reaction of 3-acetoxyaminoquinazolinones **2** and **10** with the corresponding pyridines; barriers to rotation around each of the N–N bonds in **3** have been measured by NMR spectroscopy, and the imides **3–5** react both with styrene and with diethyl fumarate to give the corresponding aziridines.

Oxidation of 3-aminoquinazolinones, *e.g.* **1**, with lead tetraacetate in dichloromethane gives the 3-acetoxyaminoquinazolinone **2** which is stable in solution at $-20\text{ }^{\circ}\text{C}$.¹ Addition of pyridine (10 mol equiv.) gives the pyridinium imide **3** as a pale yellow crystalline solid, mp $134\text{--}136\text{ }^{\circ}\text{C}$ (decomp.). Similarly, the pyridinium imides **4** (53%) and **5** (55%) were obtained from reaction of **2** with 2-methylpyridine and 3-methylpyridine, respectively.[†] These imides are not stable to silica chromatography and were purified by crystallisation after removal of the bulk of the pyridine.

In the NMR spectra of the imides **3** and **5**, broadened signals are observed at room temperature for the methyl groups in the quinazolinone 2-isopropyl substituent and from the two *ortho*-protons on the pyridinium ring.

Examination of a number of NMR spectra of **3** at 400 MHz between $-95\text{ }^{\circ}\text{C}$ and ambient temperature reveals the involvement of *two* dynamic processes in the broadening of the signals from the two types of proton referred to above. In anticipation of what follows, the preferred stereostructure proposed for this imide (**3'** \rightleftharpoons **3''**) is that shown in Fig. 1.

In the NMR spectrum of **3** at $-40\text{ }^{\circ}\text{C}$, the broadened isopropyl methyl signals have separated and sharpened: from their separation and coalescence temperature, a barrier $\Delta G^{\ddagger} = 60\text{ kJ mol}^{-1}$ was calculated. This barrier is assigned to that for rotation around the $\text{N}_{\alpha}\text{--N}_{\beta}$ bond in Fig. 1. (**3'** \rightleftharpoons **3''**).

The signals from the *ortho*-protons of the pyridinium ring in **3** are separated at $-90\text{ }^{\circ}\text{C}$ (CD_2Cl_2 solution) although not sharpened fully. From their separation and coalescence temperature, a barrier $\Delta G^{\ddagger} = 48\text{ kJ mol}^{-1}$ was calculated and this is assigned to that for rotation around the $\text{N}_{\beta}\text{--N}_{\gamma}$ bond in Fig. 1.


Interconversion of the magnetic environments of the two *ortho*-protons of the pyridinium ring in **3** would result from direct inversion via **6** at the divalent nitrogen N_{β} in **3'** (Fig. 1) in which the three nitrogens are linear. However, the lower energy barrier measured above is *not* that associated with this inversion at N_{β} since such a process (**3'** \rightleftharpoons **3''**) would, at the same time, also interconvert the magnetic environments of the isopropyl methyl groups. This inversion at N_{β} via **6** could account for the

higher energy barrier referred to above; however, this transition state requires a higher energy sp^2 -hybridisation for N_{β} and unfavourable interaction between lone pairs contained in p-orbitals on N_{α} and N_{β} .[‡] Rotation around this $\text{N}_{\alpha}\text{--N}_{\beta}$ bond, on the other hand, can proceed via **7** (Fig. 1) having sp^3 hybridisation for N_{β} and less unfavourable interactions between lone pairs on N_{α} and N_{β} .

Support for this interpretation comes from the NMR spectra at various temperatures of the pyridinium imide **8**, prepared from 4-cyanopyridine and the 3-acetoxyaminoquinazolinone **10** derived from **9**. From the separation and coalescence of the *ortho*-protons on the cyanopyridine ring, the barrier for $\text{N}_{\beta}\text{--N}_{\gamma}$ rotation in **8** is higher ($\Delta G^{\ddagger} = 62\text{ kJ mol}^{-1}$) than that in **3**, corresponding to greater resonance delocalisation with the pyridine ring and hence more double bond character in this $\text{N}_{\beta}\text{--N}_{\gamma}$ bond. The barrier to rotation around the $\text{N}_{\alpha}\text{--N}_{\beta}$ is also raised ($\Delta G^{\ddagger} > 77\text{ kJ mol}^{-1}$)[§] which is consistent with a rotation around this bond which is coupled with the rotation around the $\text{N}_{\beta}\text{--N}_{\gamma}$ bond, *i.e.* the transition state for the $\text{N}_{\alpha}\text{--N}_{\beta}$ rotation has an orientation around the $\text{N}_{\beta}\text{--N}_{\gamma}$ bond as illustrated in **7**.

These imides **3–5** function as aziridinating agents: heating **3** with styrene (3 mol equiv.) at $135\text{ }^{\circ}\text{C}$ gave aziridine **11** (44%) identical with a sample prepared by aziridination of styrene using **2**. Likewise, aziridination of diethyl fumarate (3 mol equiv.) using **5** at $145\text{ }^{\circ}\text{C}$ gave aziridine **12** (33%); the same aziridine **12** was obtained in 56% yield by heating the 2-methylpyridine-derived imide **4** in diethyl fumarate at $150\text{ }^{\circ}\text{C}$.[¶] Aziridine **12** (81%) was also prepared by aziridination of diethyl fumarate using **2**.

Oxidation of **1** in the presence of 3,5-lutidine gave an oily impure product which could not be purified. However, if the oil obtained on work-up was heated directly with diethyl fumarate



Scheme 1 Reagents and conditions: i, $\text{Pb}(\text{OAc})_4$, CH_2Cl_2 , $-20\text{ }^{\circ}\text{C}$




Fig. 1


at 145 °C, aziridine **12** was isolated (30%) after chromatography.

A large number of pyridinium imides are known,² the majority having the additional substituent on the divalent nitrogen as CO₂R, COR, SO₂Ar or Ar. Barriers to rotation around the N–N bond in these compounds have not been reported; the symmetry likely to be present in many of them means that the two *ortho*-protons of the pyridinium ring are magnetically equivalent. Some of these pyridinium imides act

as nitrene precursors and on photolysis bring about the aziridination of alkenes.²

We thank Zeneca Specialities for a CASE award (P. J. E.), the SERC (P. J. E. and E. B.) for support and Dr O. Howarth (University of Warwick) for the 400 MHz spectra.

Received, 29th December 1994; Com. 4/07862D



Scheme 2

Footnotes

† All new compounds have been fully characterised.

‡ Some alleviation of the lone-pair interaction in **6** is possible by inversion *via* a rotamer around the N_α–N_β bond in which these lone pairs are not eclipsed.

§ The increase in this higher barrier is difficult to account for in terms of inversion at N_β *via* a transition state analogous to **6**.

¶ The cyano-substituted imide **8** was unreactive towards either styrene or diethyl fumarate even at higher temperatures.

References

- 1 R. S. Atkinson, M. J. Grimshire and B. J. Kelly, *Tetrahedron*, 1989, **45**, 2875; R. S. Atkinson, P. J. Edwards and G. A. Thomson, *J. Chem. Soc., Chem. Commun.*, 1992, 1256.
- 2 Reviews: A. Katritzky and J. N. Lam, *Heterocycles*, 1992, **33**, 1011; J. Streith, *Chimia*, 1991, **45**, 65; see also C. R. Arnold, J. Melvin, W. M. Nelson and G. B. Schuster, *J. Org. Chem.*, 1992, **57**, 3316; R. Poe, K. Schnapp, M. J. T. Young, J. Grayzar and M. S. Platz, *J. Am. Chem. Soc.*, 1992, **114**, 5054.