

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/225051619>

# Total Synthesis of (–)-Galanthamine and (–)-Lycoramine via Catalytic Asymmetric Hydrogenation and Intramolecular Reductive Heck Cyclization

ARTICLE *in* ORGANIC LETTERS · MAY 2012

Impact Factor: 6.36 · DOI: 10.1021/ol300913g · Source: PubMed

---

CITATIONS

27

---

READS

52

5 AUTHORS, INCLUDING:



Jian-Hua Xie

Nankai University

104 PUBLICATIONS 2,587 CITATIONS

SEE PROFILE

# Total Synthesis of (–)-Galanthamine and (–)-Lycoramine via Catalytic Asymmetric Hydrogenation and Intramolecular Reductive Heck Cyclization


Ji-Qiang Chen, Jian-Hua Xie,\* Deng-Hui Bao, Sheng Liu, and Qi-Lin Zhou\*

State Key Laboratory and Institute of Elemento-organic Chemistry, Nankai University,  
Tianjin 300071, China

jhxie@nankai.edu.cn; qlzhou@nankai.edu.cn

Received April 9, 2012


## ABSTRACT



A synthetic strategy featuring efficient ruthenium-catalyzed asymmetric hydrogenation of racemic  $\alpha$ -aryloxy cyclic ketone via dynamic kinetic resolution and palladium-catalyzed intramolecular reductive Heck cyclization has been developed for the asymmetric total synthesis of (–)-galanthamine (20.1%, 12 steps) and (–)-lycoramine (40.2%, 10 steps)

The galanthamine-type alkaloids including (–)-galanthamine (**1**) and (–)-lycoramine (**2**) (Figure 1), isolated from the bulbs of different species of the Amaryllidaceae family, have attracted much attention of synthetic chemists because of their intriguing structures and potent biological activities.<sup>1</sup> (–)-Galanthamine (**1**) is a selective, reversible, and competitive acetylcholinesterase inhibitor and has been used in the early treatment of Alzheimer's disease.<sup>2</sup> (–)-Lycoramine (**2**) has a similar, albeit less potent, activity as an acetylcholinesterase inhibitor and an allosteric potentiating ligand.<sup>3</sup> Because of the limited supplies of these alkaloids from natural sources, a number of synthetic strategies have been developed for the syntheses of galanthamine and its analogues since Barton and Kirby initiatively reported the total synthesis of galanthamine

in the early 1960s.<sup>4</sup> However, most of the reported synthetic strategies provided the alkaloids in racemic form.<sup>5</sup>



**Figure 1.** Representative galanthamine-type alkaloids and their core structure.

(1) Cordell, G. A., Ed. *The Alkaloids: Chemistry and Biology*; Amsterdam; Elsevier 2010 Vol. 68,.

(2) Sramek, J. J.; Frackiewicz, J. E.; Cutler, N. R. *Expert Opin. Invest. Drugs* **2000**, 9, 2393.

(3) Han, S.; Sweeney, J. E.; Bachman, E. S.; Schweiger, E. J.; Forloni, G.; Coye, J. T.; Davis, B. M.; Joullie, M. M. *Eur. J. Med. Chem.* **1992**, 27, 673.

(4) (a) Barton, D. H. R.; Kirby, G. W. *J. Chem. Soc.* **1962**, 806. (b) Barton, D. H. R.; Kirby, G. W.; Taylor, J. B. *J. Chem. Soc.* **1962**, 4545.

(5) For recent reviews on the synthesis of galanthamine and its analogues, see: (a) Marco-Contelles, J.; do Carmo Carreiras, M.; Rodríguez, C.; Villarroja, M.; García, A. G. *Chem. Rev.* **2006**, 106, 116. (b) Marco-Contelles, J.; Perez-Mayoral, E.; van Nhien, A. N.; Postel, D. *Targets Heterocycl. Syst.* **2007**, 11, 365. (c) Zhong, J. *Nat. Prod. Rep.* **2009**, 26, 363. (d) Fang, L.; Gou, S.; Zhang, Y. *Chin. J. Org. Chem.* **2011**, 31, 286.


Galanthamine-type alkaloids feature a unique tricyclic benzofuran core structure with a chiral arylated-quaternary carbon center, and the enantioselective construction of this sterically congested quaternary carbon center is a major challenge in the synthesis of these alkaloids. Thus, a number of strategies have been developed to construct the chiral arylated-quaternary carbon center for the synthesis of galanthamine-type alkaloids by means of phenolic oxidative coupling,<sup>6</sup> intramolecular Heck reaction,<sup>7</sup> Claisen rearrangement,<sup>8</sup> Brich–Cope sequence,<sup>9</sup> phenolic oxidative coupling and crystallization-induced chiral conversion,<sup>10</sup> and organocatalyzed Michael addition.<sup>11</sup> However, although being one of the most promising methods for the synthesis of chiral compounds, the asymmetric catalysis has been rarely used for the enantioselective synthesis of galanthamine-type alkaloids, and the reported catalytic asymmetric syntheses of (–)-galanthamine (**1**) and (–)-lycoramine (**2**) were not very efficient.<sup>7,11</sup>

During the study on the catalytic asymmetric synthesis of chiral natural products, we found that the intramolecular reductive Heck reaction is a convenient access to the construction of tricyclic dihydrobenzofuran ring with a quaternary stereocenter.<sup>12</sup> This intramolecular reductive Heck reaction combining with the asymmetric hydrogenation of  $\alpha$ -aryloxy cyclic ketone via a dynamic kinetic resolution (DKR), recently developed in our laboratory,<sup>13</sup> provides a highly efficient strategy for the enantioselective synthesis of (–)-galanthamine (**1**, 20.1%, 12 steps) and (–)-lycoramine (**2**, 40.2%, 10 steps).

Our strategy is outlined in Scheme 1. We expected that the target molecules **1** and **2** could be synthesized from ester (S,R)-**3** via formation of the seven-membered aza ring through several steps including amidation and Pictet–Spengler cyclization. The ester (S,R)-**3** could be obtained

from  $\alpha,\beta$ -unsaturated ester (S)-**4** containing an  $\alpha$ -halogenated phenoxyl group via an intramolecular reductive Heck cyclization. The  $\alpha,\beta$ -unsaturated ester (S)-**4** could be easily prepared from  $\alpha$ -aryloxy cyclohexanone (S)-**5** via a Horner–Wadsworth–Emmons reaction, and the optically pure (S)-**5** could be obtained by ruthenium-catalyzed asymmetric hydrogenation of *rac*-**5** via DKR<sup>13</sup> followed by a Swern oxidation.

**Scheme 1.** Synthetic Strategy for (–)-Galanthamine (**1**) and (–)-Lycoramine (**2**)



Since Larock<sup>14</sup> and Trost<sup>7a,c</sup> had reported that the intramolecular Heck reaction of aryl allyl ethers often suffers from a competitive palladium-catalyzed ionization of the aryloxy group, producing a phenol, the prevention of this unwanted side reaction inevitably became one of the

(6) (a) Tomioka, K.; Shimizu, K.; Yamada, S.; Koga, K. *Heterocycles* **1977**, *6*, 1752. (b) Shimizu, K.; Tomioka, K.; Yamada, S.; Koga, K. *Heterocycles* **1977**, *8*, 277. (c) Kodama, S.; Hamashima, Y.; Nishide, K.; Node, M. *Angew. Chem., Int. Ed.* **2004**, *43*, 2659. (d) Node, M.; Kodama, S.; Hamashima, Y.; Katoh, T.; Nishide, K.; Kajimoto, T. *Chem. Pharm. Bull.* **2006**, *54*, 1662.

(7) (a) Trost, B. M.; Toste, F. D. *J. Am. Chem. Soc.* **2000**, *122*, 11262. (b) Trost, B. M.; Tang, W. *Angew. Chem., Int. Ed.* **2002**, *41*, 2759. (c) Trost, B. M.; Tang, W.; Toste, F. D. *J. Am. Chem. Soc.* **2005**, *127*, 14785. Catalytic asymmetric synthesis of (–)-galanthamine (10 steps, 8% overall yield).

(8) (a) Tanimoto, H.; Kato, T.; Chida, N. *Tetrahedron Lett.* **2007**, *48*, 6267. (b) Kato, T.; Tanimoto, H.; Yamada, H.; Chida, N. *Heterocycles* **2010**, *82*, 563.

(9) Malachowski, W. P.; Paul, T.; Phounsavath, S. *J. Org. Chem.* **2007**, *72*, 6792.

(10) Magnus, P.; Sane, N.; Fauber, B. P.; Lynch, V. *J. Am. Chem. Soc.* **2009**, *131*, 16045.

(11) Chen, P.; Bao, X.; Zhang, L.-F.; Ding, M.; Han, X.-J.; Li, J.; Zhang, G.-B.; Tu, Y.-Q.; Fan, C.-A. *Angew. Chem., Int. Ed.* **2011**, *50*, 8186. Catalytic asymmetric synthesis of (–)-galanthamine (16 steps, 2.8% overall yield) and (–)-lycoramine (14 steps, 8.1% overall yield).

(12) Reviews for reductive Heck reaction, see: (a) Oestreich, M., Ed. *The Mizoroki–Heck Reaction*; Wiley: Chichester, 2009. (b) Link, J. T.; Overman, L. E. *Metal-catalyzed Cross Coupling Reaction*; Diederick, F., Stang, P. J., Eds.; Wiley-VCH: Weinheim, 1998, Chapter 6. (c) Tsuji, J. *Palladium Reagent and Catalysts. Innovation in Organic Synthesis*; John Wiley: New York, 1995. Selected papers for intramolecular reductive Heck reaction for the construction of dihydrobenzofuran ring: (d) Trost, B. M.; Thiel, O. R.; Tsui, H.-C. *J. Am. Chem. Soc.* **2003**, *125*, 13155. (e) Trost, B. M.; Thiel, O. R.; Tsui, H.-C. *J. Am. Chem. Soc.* **2002**, *124*, 11616.

(13) Bai, W.-J.; Xie, J.-H.; Li, Y.-L.; Liu, S.; Zhou, Q.-L. *Adv. Synth. Catal.* **2010**, *352*, 81.

**Table 1.** Palladium-Catalyzed Cyclization of *rac*-**4** to *rac*-**3**<sup>a</sup>

entry	X	[Pd]	reductive reagent (equiv)	temp (°C)	time (h)	yield <sup>b</sup> (%)
1	Br	Pd(OAc) <sub>2</sub>	HCO <sub>2</sub> H (2.0)/Et <sub>3</sub> N (2.5)	80	10	20
2	Br	Pd(OAc) <sub>2</sub>	HCO <sub>2</sub> H (2.0)/Et <sub>3</sub> N (2.5)	100	10	40
3	Br	Pd(OAc) <sub>2</sub>	HCO <sub>2</sub> H (2.0)/Et <sub>3</sub> N (2.5)	120	10	32
4	Br	Pd(OAc) <sub>2</sub>	HCO <sub>2</sub> Na (2.0)	100	10	43
5	Br	Pd <sub>2</sub> (dba) <sub>3</sub> · CHCl <sub>3</sub>	HCO <sub>2</sub> Na (2.0)	80	3	49
6 <sup>c</sup>	I	Pd <sub>2</sub> (dba) <sub>3</sub> · CHCl <sub>3</sub>	HCO <sub>2</sub> Na (2.0)	60	3	95

<sup>a</sup> Reaction conditions: 10 mmol of *rac*-**4**, 5 mol % of Pd catalyst, 10 mol % of PPh<sub>3</sub>, DMF as solvent. <sup>b</sup> Isolated yield. <sup>c</sup> Without PPh<sub>3</sub>.

major challenges of our synthetic strategy. We first evaluate the intramolecular reductive Heck cyclization of *rac-4a* (X = Br), which was prepared in good yield (67%, two steps) from 2-bromo-6-methoxyphenol and 7-bromo-1,4-dioxaspiro[4.5]decan-8-one<sup>15</sup> (see the Supporting Information). When the reaction was performed in the presence of 5 mol % of Pd(OAc)<sub>2</sub>, 10 mol % of PPh<sub>3</sub>, 2.0 equiv of HCO<sub>2</sub>H, and 2.5 equiv of Et<sub>3</sub>N at 80 °C in DMF for 10 h, *rac-4a* was cyclized to *rac-3* in 20% yield accompanied by a significant amount of ionization product, 2-bromo-6-methoxyphenol (Table 1, entry 1). Increasing the reaction temperature led to a higher yield of *rac-3*, but still lower than 40% (entries 2 and 3). Use of HCO<sub>2</sub>Na (2 equiv), instead of HCO<sub>2</sub>H/Et<sub>3</sub>N, as a reductive reagent resulted in a slight improvement of yield to 49% (entry 5). Fortunately, when the iodinated  $\alpha,\beta$ -unsaturated ester *rac-4b* (X = I) was subjected to the reaction the ionization of aryloxy group was strongly suppressed and the desired reductive Heck cyclization product *rac-3* was obtained in 95% yield under milder conditions (60 °C, 3 h) (entry 6).<sup>16</sup> This palladium-catalyzed intramolecular reductive Heck

**Scheme 2.** Palladium-Catalyzed Cyclization of *rac-7* and *rac-8*




cyclization is also highly efficient for the cyclization of *rac-7* and *rac-8*, the analogues of *rac-4* derived from  $\alpha$ -(2-iodophenoxy)cycloalkanones with five or seven-membered ring to the corresponding benzofurans *rac-9* (93%) and *rac-10* (84%) in high yields (Scheme 2).

Naturally, after the establishment of a highly efficient intramolecular reductive Heck cyclization to create the tricyclic benzofuran core structure of galanthamine-type alkaloids, we then tried the synthesis of (*S,R*)-**3** in optically pure form. To address this topic, the enantioselective synthesis of the optically pure  $\alpha$ -aryloxy cyclohexanone (*S*)-**5b** is the key. Recently, we developed a highly efficient ruthenium-catalyzed asymmetric hydrogenation of racemic  $\alpha$ -aryloxy cyclohexanones via DKR, producing chiral  $\beta$ -aryloxy cycloalkanols in excellent enantioselectivity.<sup>13</sup> This provides an method for us to try the synthesis of the optically pure  $\alpha$ -aryloxy cyclohexanone (*S*)-**5b** with a bulky ethylene ketal group at the 4-position of the cyclohexane ring. By using chiral ruthenium catalyst

RuCl<sub>2</sub>-(*S*)-SDP/(*R,R*)-DPEN,<sup>17</sup> *rac-5b* (obtained from the reaction of 2-iodo-6-methoxyphenol with 7-bromo-1,4-dioxaspiro[4.5]decan-8-one in 75% yield, see the Supporting Information) was hydrogenated to chiral  $\beta$ -aryloxy cyclohexanol (*S,R*)-**6** in high yield (99%) with excellent enantioselectivity (97% ee) and *cis*-selectivity (*cis/trans* > 99:1) (Scheme 3), indicating that the ethylene ketal group in the substrate has a negligible effect to the reaction. The  $\beta$ -aryloxy cyclohexanol (*S,R*)-**6** was then converted into  $\alpha$ -aryloxy cyclohexanone (*S*)-**5b** in 95% yield without loss of its optical purity by Swern oxidation. Subsequently, a condensation of (*S*)-**5b** with ethyl 2-(dimethoxy-phosphoryl)

**Scheme 3.** Enantioselective Synthesis of (*S,R*)-**3** (DMSO = Dimethyl Sulfoxide)



acetate by a Horner-Wadsworth-Emmons reaction gave  $\alpha,\beta$ -unsaturated esters (*S*)-**4b** in 96% yield with (*E*)-configuration as the major isomer (*Z/E*  $\approx$  1:5). The *Z,E*-mixture of (*S*)-**4b** was then submitted to the palladium-catalyzed intramolecular reductive Heck cyclization to give (*S,R*)-**3** in 95% yield.

With ester (*S,R*)-**3** in hand, we then focused our attention on its conversion into (–)-galanthamine (**1**) and (–)-lycoramine (**2**). As shown in Scheme 4, the ester (*S,R*)-**3** was hydrolyzed with sodium hydroxide in methanol/H<sub>2</sub>O, followed by activation with ethyl chloroformate (ClCO<sub>2</sub>Et) in the presence of triethylamine at –15 °C and amidation with aqueous methylamine (MeNH<sub>2</sub>) to offer amide (*S,R*)-**11** in 83% yield (2 steps). Subsequent Pictet-Spengler cyclization of (*S,R*)-**11** with paraformaldehyde furnished the tetracyclic intermediate (*S,R*)-**12** with a seven-membered azepine ring in 89% yield.<sup>18</sup> The intermediate (*S,R*)-**12** was then subjected to a selective

(14) Larock, R. C.; Stinn, D. E. *Tetrahedron Lett.* **1988**, 29, 4687.

(15) Laukhtien, G.; Volk, F.-J.; Frahm, A.-W. *Tetrahedron: Asymmetry* **1997**, 8, 3457.


(16) Ester *rac-3* can also be synthesized in 65% yield by using radical cyclization (AIBN, *n*-Bu<sub>3</sub>SnH, benzene, reflux, 3 h). However, the use of toxic metal tin compound made the radical cyclization less applicable. Selected papers for radical cyclization: (a) Parker, K. A.; Fokas, D. *J. Org. Chem.* **2006**, 71, 449. (b) Parker, K. A.; Kim, H.-J. *J. Org. Chem.* **1992**, 57, 752.

(17) (a) Xie, J.-H.; Wang, L.-X.; Zhu, S.-F.; Fu, Y.; Fan, B.-M.; Duan, H.-F.; Zhou, Q.-L. *J. Am. Chem. Soc.* **2003**, 125, 4404. (b) Xie, J.-H.; Liu, S.; Huo, X.-H.; Cheng, X.; Duan, H.-F.; Fan, B.-M.; Wang, L.-X.; Zhou, Q.-L. *J. Org. Chem.* **2005**, 70, 2967.

(18) Guillou, C.; Beunard, J.-L.; Gras, E.; Thal, C. *Angew. Chem., Int. Ed.* **2001**, 40, 4745.

reduction of the ketone group to the hydroxyl using K-Selectride and a further reduction of the amide motif to amine with triethoxysilane ((EtO)<sub>3</sub>SiH) in the presence of Zn(OAc)<sub>2</sub> as a catalyst<sup>19</sup> to give (–)-lycoramine (2) in 84.6% yield (2 steps).<sup>20</sup> The NMR spectroscopic data and the optical rotation ([α]<sub>D</sub><sup>20</sup> = –102 (c 0.35, EtOH), lit. [α]<sub>D</sub><sup>20</sup> = –100 (c 0.35, EtOH),<sup>9</sup> [α]<sub>D</sub><sup>20</sup> = –92.7 (c 0.35, EtOH)<sup>11</sup>) of our synthetic (–)-lycoramine are identical to those reported in the previous synthesis. Thus, enantioselective synthesis of (–)-lycoramine was achieved in 40.2% overall yield over 10 steps from the commercially available 2-iodo-6-methoxyphenol via a ruthenium-catalyzed asymmetric hydrogenation and a palladium-catalyzed intra-molecular reductive Heck cyclization as the key steps.


**Scheme 4.** Enantioselective Synthesis of (–)-Lycoramine (2) (K-Selectride = Potassium Tri-*sec*-butyl Borohydride)



The enantioselective synthesis of (–)-galanthamine (1) is outlined in Scheme 5. Using the procedure of Saegusa–Ito oxidation,<sup>21</sup> the tetracyclic intermediate (S,R)-12 was treated with TBSOTf, followed by palladium-mediated oxidation to provide the desired enone (S,S)-14 in 51% yield (two steps).<sup>22</sup> The (S,S)-14 was then reduced sequentially with K-Selectride and (EtO)<sub>3</sub>SiH/Zn(OAc)<sub>2</sub> to yield (–)-galanthamine (1) in 83% yield. The data of NMR spectroscopy and optical rotation of our synthetic (–)-galanthamine (1) are identical to those reported in the literature ([α]<sub>D</sub><sup>20</sup> = –119.5 (c 0.30, EtOH), lit.<sup>6c</sup> [α]<sub>D</sub><sup>20</sup> = –121.7 (c 0.30, EtOH)). The enantioselective

synthesis of (–)-galanthamine (1) was thus accomplished in 12 steps with 20.1% overall yield.

**Scheme 5.** Enantioselective Synthesis of (–)-Galanthamine (1) (TBSOTf = *tert*-Butyldimethylsilyl Trifluoromethanesulfonate)



In conclusion, we have developed highly efficient enantioselective syntheses of (–)-galanthamine (1) and (–)-lycoramine (2) based on a ruthenium-catalyzed asymmetric hydrogenation and a palladium-catalyzed intramolecular reductive Heck cyclization as the key steps. The (–)-galanthamine (1) was synthesized in twelve steps with 20.1% overall yield and the (–)-lycoramine (2) was synthesized in ten steps with 40.2% overall yield from commercially available starting materials. These high yielding syntheses of (–)-galanthamine and (–)-lycoramine have good reproducibility. These results demonstrated that this catalytic enantioselective synthetic strategy is efficient for the asymmetric construction of the polycyclic benzofuran ring systems bearing an all-carbon quaternary center and has high potential for application to the syntheses of other galanthamine-type, morphine-type, and lunarine-type biologically significant alkaloids.

**Acknowledgment.** We thank the National Natural Science Foundation of China, the National Basic Research Program of China (973 Program) (2012CB821600), the “111” project (B06005) of the Ministry of Education of China, and the Program for New Century Excellent Talents in University (NCET-08-0292) for financial support.

**Supporting Information Available.** Experimental procedures, characterization data for the products, and HPLC spectra for (S)-5b and (S,R)-6. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.

(19) Das, S.; Addis, D.; Zhou, S.; Junge, K.; Beller, M. *J. Am. Chem. Soc.* **2010**, *132*, 1770.

(20) The reduction of the amide group of (S,S,R)-9 with LiAlH<sub>4</sub> gave 2 in 81% yield.

(21) Ito, Y.; Hirao, T.; Saegusa, T. *J. Org. Chem.* **1978**, *43*, 1011.

(22) For improving the transformation of (S,R)-12 to (S,S)-14 other methods, such as DDQ oxidation of silyl enol ester, treatment with PhSeCl and oxidation with H<sub>2</sub>O<sub>2</sub>, and bromination followed with elimination of HBr, have also been tried; however, the desired product was obtained in very low yields.