

J Nat Prod. Author manuscript; available in PMC 2013 April 27

Published in final edited form as:

J Nat Prod. 2012 April 27; 75(4): 771–773. doi:10.1021/np200862y.

Antineoplastic Agents. 587. Isolation and Structure of 3-Epipancratistatin from Narcissus cv. Ice Follies1

George R. Pettit*, Rui Tan, Guan-Hu Bao, Noeleen Melody, Dennis L. Doubek, Song Gao, Jean-Charles Chapuis, and Lee Williams

Cancer Research Institute and Department of Chemistry and Biochemistry, Arizona State University, PO Box 871604, Tempe, AZ 85287-1604, United States

> Bioassay-guided (cancer cell line) separation of an extract prepared from Narcissus cv. Ice Follies (from the Netherlands) led to isolation of a new Amaryllidaceae isocarbostiryl, 3epipancratistatin (1b), as well as narciclasine (2). This Narcissus cultivar was found to be a good source of narciclasine. The structure of 1b was established by high resolution mass and high-field 2D NMR spectroscopic analyses. Against a panel of murine and human cancer cell lines, 3-epipancratistatin (1b) led to cell growth inhibition (GI₅₀ 2.2–0.69 μ g/mL) some 100× less than that found for pancratistatin (1a) and narciclasine (2), thereby revealing an important configurational requirement in 1a for strong cancer cell growth inhibition.

The Amaryllidaceae plant family genera and particularly the Narcissus genus have a long and quite notable place in the history of traditional and western medicine.² Research concerning the discrete constituents had a formal beginning in 1877 with a description of lycorine from Narcissus pseudonarcissus. 3a Presently, well over 100 Amaryllidaceae nitrogen containing compounds are known, and some of the most promising members with antineoplastic properties have been nicely reviewed.³ Current progress in preclinical development of the isocarbostyril pancratistatin (1a) has been summarized in other important reviews, 4 and efforts towards efficient total syntheses continue.⁵

Importantly, the biological foundation for advancement of pancratistatin (1a), in the form of its very soluble phosphate prodrug, 6 has continued to increase rapidly. Illustrative are the recent experimental results confirming that 1a very selectively sends a range of cancer cell types, including metastatic prostate cancer, breast cancer and colon carcinoma, as well as ex vivo models of leukemia, into apoptosis without toxicity to noncancerous counterpart cell lines; the mode of action appears to involve selective targeting of cancer cell mitochondria.⁷

While investigating Narcissus varieties from horticultural sources in order to increase the availability of narciclasine (2)⁸ for use in our semisyntheses of 1a,⁹ we found *Narcissus* cv. Ice Follies¹⁰ to be a new and useful source of 2. During a scale-up isolation of 2, we detected the presence of a new isocarbostyril clearly related to pancratistatin that offered the prospect of finding useful anticancer activity and new SAR data. The isolation and structure elucidation proceeded as follows.

The bulbs (4,000 kg) of Narcissus cv. Ice Follies were extracted with CH₃OH. The extract was partitioned between 9:1 CH₃OH-H₂O and hexane, and the hexane-soluble lipid fraction was removed. The aqueous phase was then partitioned between 3:2 CH₃OH–H₂O and

^{*}Corresponding Author To whom correspondence should be addressed. Tel: (480) 965-3351. Fax: (480) 965-2747.

Supporting Information. ¹H and ¹³C NMR spectra of compound 1b and a table of the data (Table 2). This material is available free of charge via the Internet at http://pubs.acs.org.

Pettit et al. Page 2

> CH₂Cl₂, and the CH₂Cl₂ fraction was separated. After removal of CH₃OH from the aqueous layer, the remaining H₂O phase was extracted with EtOAc followed by n-butanol. The CH₂Cl₂, EtOAc, and n-butanol fractions significantly inhibited the murine lymphocytic leukemia cell line (P388: ED₅₀ 0.24, 0.034, and 0.92 µg/mL, respectively). Bioassay guided (P388) separation of these three fractions was conducted using a series of gel permeation and partition chromatographic separations on Sephadex LH-20 columns, followed by final purification by crystallization, to afford 3-epipancratistatin (1b, 50 mg, 1.25×10^{-6} %, P388 ED_{50} 0.69 μ g/mL) and narciclasine (2, 26 g, 6.5 × 10⁻⁴ %, P388 ED_{50} 0.018 μ g/mL).

1a, $3-\beta$ -OH, (+)-pancratistatin

2, narciclasine

1b, $3-\alpha$ -OH

3-Epipancratistatin (1b) crystallized from MeOH as a colorless solid. The molecular formula of 1b was established by high-resolution mass spectroscopy as C₁₄H₁₅NO₈, which is also the molecular formula of pancratistatin (1a). The ¹H NMR, ¹³C NMR and HMQC spectra of **1b** indicated the presence of six aromatic carbons, including one methine carbon ($\delta_{\rm H}$ 6.49, $\delta_{\rm C}$ 97.7) and five quaternary carbons ($\delta_{\rm C}$ 152.1, 145.4, 135.8, 131.7, 107.4), as well as one OCH₂O group ($\delta_{\rm H}$ 6.02, 6.04, $\delta_{\rm C}$ 103.4), one amide unit ($\delta_{\rm NH}$ 7.49, $\delta_{\rm CO}$ 169.4) and six sp^3 methine groups, of which four were oxygenated ($\delta_{\rm C}/\delta_{\rm H}$ 67.3/4.27, 72.6/3.82, 70.4/3.55, 71.9/3.54, 54.0/3.37 and 38.3/2.94). The six exchangeable protons in the ¹H NMR spectrum were removed by addition of D_2O , revealing the presence of a phenolic OH group (δ_H 13.02), an amide NH group ($\delta_{\rm H}$ 7.49), and four additional OH groups ($\delta_{\rm H}$ 5.11, 4.94, 4.62, and 5.03). Thus all the atoms of 1b by NMR spectroscopy matched the molecular formula C₁₄H₁₅NO₈ from its high-resolution mass spectrum.

In an analysis of the H-H COSY spectrum, the six sp^3 protons were found to interact in a sequence that indicated a six-membered ring. The four OH signals at $\delta_{\rm H}$ 5.11, 4.94, 4.62, and 5.03 were assigned to positions 1 to 4 according to the H-H COSY correlations between the OH groups and the adjacent protons. The aromatic ring, six-membered ring and amide segments were further connected by analysis of cross signals in the HMBC spectrum, which led to the conclusion that compound 1b had the same planar structure as pancratistatin (1a).

The relative configuration of 1b was elucidated by interpretation of the ROESY spectrum and coupling constant values. The large coupling constant between H-10b and H-4a $(J_{\text{H-}10\text{h/H-}4a} = 12.6 \text{ Hz})$ suggested a trans-fused B/C ring system, with each proton having an axial orientation as in pancratistatin (1a). In the ROESY spectrum of 1b, cross peaks for axial oriented H-10b with the 2-OH group and H-4 were observed ($\delta_H/\delta_H = 2.94/4.94$ and 2.94/3.54, respectively). An unambiguous relationship was observed between the axialoriented 1-OH group and H-3 ($\delta_{\rm H}/\delta_{\rm H}$ 5.11/3.55). Although the water peak interfered, ROESY cross peaks were noted for H-4a ($\delta_{\rm H}$ 3.37) with 1-OH and H-3. For comparison, a NOESY experiment using an authentic pancratistatin (1a) sample was performed, which showed the NOE effects between H-4a and 1-OH (δ_H/δ_H 3.70/4.82), H-4a and 3-OH (3.70/5.07), and H-4 and 2-OH (3.72/5.36). All these results described 1-3-5 space effects in a chair conformation of the six membered ring and implied an a (equatorial) assignment for the 3 OH group of 1b, the reverse of the β (axial) 3-OH in pancratistatin (1a). Furthermore, a closer comparison of the ¹H NMR spectra of **1b** and **1a** clearly showed that

Pettit et al. Page 3

the difference in configuration of the 3 OH group in **1b** appeared in chemical shift changes for H-2, H-3, H-4, and H-5, as well as the 1-OH and 3-OH groups, relative to those of **1a**. Also, a 1–3 effect resulted in an upfield shift of 4 ppm for the C-4a signal in the 13 C spectrum (from δ_C 50 for **1a** to δ_C 54 for **1b**).

The growth inhibitory properties of *Narcissus* cv. Ice Follies constituent **1b** were evaluated using the murine P388 lymphocytic leukemia cell line and a selection of human cancer cell lines (Table 1) and compared to those of its C-3 epimer, compound **1a**. As we have consistently found, pancratistatin (**1a**) exhibited strong inhibition against growth of the murine P388 lymphocytic leukemia and a minipanel of human tumor cell lines. The new isocarbostyril 3-epipancratistatin (**1b**) also exhibited significant growth inhibitory activity against the same cancer cell lines but was over $100 \times$ less inhibitory than **1a**, thereby revealing a very important configurational requirement for stronger cancer cell growth inhibitory activity.

EXPERIMENTAL SECTION

General Experimental Procedures

All solvents used in the extraction and isolation processes were redistilled prior to use. Both gel and partition column chromatography was performed with Sephadex LH-20 from Pharmacia. The uncorrected melting points were observed using a Fisher-Johns apparatus. The IR spectra were obtained with a Thermo Nicolet Avatar 360 E.S.P. spectrometer. ¹H and ¹³C NMR spectra were recorded using a Varian Inova 500 MHz instrument. High resolution APCI mass spectra were obtained employing a JEOL LC Mate LCMS system.

Plant Material

The bulbs of *Narcissus* cv. Ice Follies were grown commercially (K. Van Bourgondien and Sons, P.O. Box 2000, Virginia Beach, VA 23450) in 1997. A voucher specimen (B849350) was deposited in the herbarium of the Cancer Research Institute at Arizona State University.

Extraction and Isolation

All the separations were guided by bioassay results using the murine P388 lymphocytic leukemia (P388) and a minipanel of human cancer cell lines. The dry Narcissus bulbs (4,000 kg) were placed in barrels and soaked in CH3OH for at least a week, and an extract was prepared^{8b} and processed in 1-kg batches as follows. The bulb extract was partitioned between 9:1 CH₃OH-H₂O (8 L) and hexane (8 L), and the CH₃OH-H₂O layer was further extracted with hexane (4 × 8 L). The hexane layers were combined and dried, and water (4 L) was added to the methanolic phase to give a 3:2 CH₃OH-H₂O layer, which was extracted with CH₂Cl₂ (3 × 8 L). The CH₂Cl₂ fractions were combined and dried. The CH₃OH was removed from the aqueous layer in vacuo, and the remaining H²O phase was made up to 4.5 L and extracted with EtOAc (5 × 4.5 L), followed by n-BuOH (4 × 4.5 L), to give EtOAc and 1-butanol fractions. This process was repeated and the final extracts combined to give CH_2Cl_2 (3.12 kg, P388 ED50 0.24 μ g/mL), EtOAc (935 g, P388 ED₅₀ 0.034 μ g/mL), and n-butanol (3.45 kg, P388 ED₅₀ 0.92 μ g/mL) extracts. These fractions contained narciclasine (2, detected by TLC), and each was dissolved in CH₃OH and subjected to gel permeation column chromatography on Sephadex LH-20 (100 × 15 cm). Continued elution with CH₃OH provided 45 g of a crude narciclasine containing fraction, which was subjected to repeated partition column chromatography on Sephadex LH 20 (5 × 110 cm) employing hexane-CH₂Cl₂-CH₃OH (3:1:1) as eluent to afford 50 mg of 3-epipancratistatin (**1b**) and 26 g of pure narciclasine (2).

Pettit et al. Page 4

3-Epipancratistatin (*1b*): colorless solid that crystallized from CH₃OH: mp 293–295 °C; IR (film) ν_{max} 3421, 3227, 2924, 1670, 1463, 1342, 1088, 1032 cm⁻¹; ¹H NMR (DMSO- d_6) δ 13.02 (1H, s, 7-OH), 7.49 (1H, s, 5-NH), 6.49 (1H, s, H-10), 6.04, 6.02 (1H each, d, J= 1.5 Hz, H-8a), 5.11 (1H, d, J= 4.8 Hz, 1-OH), 5.03 (1H, d, J= 3.6 Hz, 4-OH), 4.94 (1H, d, J= 3.6 Hz, 2-OH), 4.62 (1H, d, J= 4.5 Hz, 3-OH), 4.27 (1H, br s, H-1), 3.82 (1H, br s, H-2), 3.55 (1H, br s, H-3), 3.54 (1H, br s, H-4), 3.37 (1H, dd, J= 13.5, 9.6 Hz, H-4a), 2.94 (1H, d, J= 12.6 Hz, H-10b); ¹³C NMR (DMSO- d_6) δ 169.4 (C-6), 152.1 (C-9), 145.4 (C-7), 135.8 (C-10a), 131.7 (C-8), 107.4 (C-6a), 101.8 (C-8a), 97.7 (C-10), 72.6 (C-2), 71.9 (C-4), 70.4 (C-3), 67.3 (C-1), 54.0 (C-4a), 38.3 (C-10b); HRAPCI m/z 326.0987 [M + H]⁺ (calcd for C₁₄H₁₆NO₈, 326.0876). Compound **1b** recrystallized from DMSO-H₂O: mp 297-299 °C; anal. C 46.91%, H 4.78%, N 3.82%, calcd for C₁₄H₁₅NO₈.2H₂O, C 46.50%, H 5.26%, N 3.87%.

Cancer Cell Line Procedures

Inhibition of human cancer cell line growth was determined using the NCI's sulforhodamine B assay as previously described. 12 In summary, cells in a 5% fetal bovine serum/RPMI1640 medium solution were inoculated in 96-well plates and incubated for 24 h. The compounds were then added as serial dilutions. After 48 h, the plates were fixed with trichloroacetic acid, stained with sulforhodamine B, and then read with an automated microplate reader. Growth inhibition of 50% (GI $_{50}$ or the drug concentration causing a 50% reduction in the net protein increase) was calculated from optical density data with Immunosoft software.

The murine lymphocytic leukemia P388 cells¹³ were incubated for 24 h in a 10% horse serum/Fisher medium solution followed by a 48-h incubation with serial dilutions of each compound. Cell growth inhibition (ED_{50}) was then calculated using a Z1 Beckman/Coulter particle counter.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial assistance was received with appreciation from research grants CA-10612-02 to CA-10612-3 and 5 R01 CA090441-07 from the Division of Cancer Treatment and Diagnosis, NCI, DHHS; the Arizona Biomedical Research Commission (formerly the Arizona Disease Control Research Commission); Dr. Alec D. Keith; the J. W. Kieckhefer Foundation; the Margaret T. Morris Foundation; Polly J. Trautman; Dr. John C. Budzinski; and the Robert B. Dalton Endowment Fund. For other assistance, we are pleased to thank Drs. Fiona Hogan, and John C. Knight, as well as Felicia Craciunescu and Michael J. Dodson.

DEDICATION In memory of Professor Norman Farnsworth (1930–2011), a world class pharmaceutical and natural products scientist who will be deeply missed.

REFERENCES

- (1). For Antineoplastic Agents part 586, refer to Pettit RK, Pettit GR, Xu JP, Weber CA, Richert LA. Planta Med. 2010; 76:500–501. [PubMed: 19941263]
- (2)(a). Pettit GR, Gaddamidi V, Cragg GM, Herald DL, Sagawa Y. J. Chem. Soc. Chem. Commun. 1984:1693–1694.(b) Ingrassia L, Lefranc F, Dewelle J, Pottier L, Mathieu V, Spiegl-Kreinecker S, Sauvage S, El Yazidi M, Dehoux M, Berger W, Van Quaquebeke E, Kiss R. J. Med. Chem. 2009; 52:1100–1114. [PubMed: 19199649]
- (3)(a). Evidente A, Kornienko A. Phytochem. Rev. 2009; 8:449–459.(b) Kornienko A, Evidente A. Chem. Rev. 2008; 108:1982–2014. [PubMed: 18489166]
- (4)(a). Grothaus PG, Cragg GM, Newman DJ. Curr. Org. Chem. 2010; 14:1781–1791.(b) Kinghorn AD, Chin Y-W, Swansom SM. Curr. Opin. Drug Discov. Devel. 2009; 12:189–196.(c) Cragg,

Pettit et al. Page 5

- GM.; Newman, DJ. In Anticancer Agents from Natural Products. 2nd ed. Cragg, GM.; Kingston, DGI.; Newman, DJ., editors. CRC; Boca Raton: 2011.
- (5)(a). Dam JH, Madsen R. Eur. J. Org. Chem. 2009:4666–4673.(b) Manpadi M, Kornienko A. Org. Prep. Proc. Intl. 2008; 40:107–161.(c) Chapleur Y, Chretien F, Ahmed S. Ibn, Khaldi M. Curr. Org. Syn. 2006; 3:341–378.(d) Li M, Wu A, Zhou P. Tetrahedron Lett. 2006; 47:3707–3710.
- (6)(a). Pettit GR, Melody N, Herald DL. J. Nat. Prod. 2004; 67:322–327. [PubMed: 15043403] (b) Pettit GR, Orr B, Ducki S. Anti Cancer Drug Des. 2000; 15:389–395.
- (7)(a). Griffin C, McNulty J, Pandey S. Int. J. Oncol. 2011; 38:1549–1556. [PubMed: 21424119] (b) Griffin C, Karnik A, McNulty J, Pandey S. Mol. Cancer Ther. 2011; 10:57–68. [PubMed: 21220492] (c) Siedlakowski P, McLachlan-Burgess A, Griffin C, Tirumalai SS, McNulty J, Pandey S. Cancer Biol Ther. 2008; 7:376–384. [PubMed: 18075307]
- (8)(a). Pettit GR, Pettit GR III, Groszek G, Backhaus RA, Doubek DL, Barr RJ, Meerow AW. J. Nat. Prod. 1995; 58:756–759. [PubMed: 7623049] (b) Pettit GR, Pettit GR III, Backhaus RA, Boettner FE. J. Nat. Prod. 1995; 58:37–43. [PubMed: 7760076] (c) Pettit GR, Pettit GR III, Backhaus RA. J. Nat. Prod. 1993; 56:1682–1687. [PubMed: 8277308] (d) Ceriotti G. Nature. 1967; 213:595–596. [PubMed: 5340258]
- (9). Pettit GR, Melody N, Herald DL. J. Org. Chem. 2001; 66:2583–2587. [PubMed: 11304174]
- (10). For a report of other alkaloids isolated from this variety, see Moraes-Cerdeira RM, Bastos JK, Burandt CL Jr. Nanayakkara NPD, Mikell J, McChesney JD. Planta Med. 1997; 63:92–93. [PubMed: 17252336]
- (11). Pettit GR, Melody N, Herald DL. J. Nat. Prod. 2004; 67:322–327. [PubMed: 15043403]
- (12). Monks A, Scudiero D, Skehan P, Shoemaker R, Paull K, Vistica D, Hose C, Langley J, Cronise P, Vaigro-Wolff A, Gray-Goodrich M, Campbell H, Mayo J, Boyd M. J. Natl. Cancer Inst. 1991; 83:757–766. [PubMed: 2041050]
- (13). Suffness, M.; Douros, J. Methods in Cancer Research. DeVita, VT., Jr.; Busch, H., editors. Vol. Vol. 16. Academic; New York: 1979. p. 73-126.

Pettit et al.

Table 1

Murine P388 Lymphocytic Leukemia (ED $_{50}$) and Human Cancer Cell Line Values (GI $_{50}$ $\mu g/mL$)

cance	cancer cell line a,b	a,b					
	P388	BXPC-3	MCF-7	SF-268	P388 BXPC-3 MCF-7 SF-268 NCI-H460 KM20L2 DU-145	KM20L2	DU-145
$1a^{11}$	$1a^{11}$ 0.017 0.02	0.02	0.023	0.014	0.032	0.025	0.015
1b	1b 0.69 1.7	1.7	2.2	0.74	0.85	2.3	1.3

^aIn DMSO.

Page 6

bancer type: P388 (murine lymphocytic leukemia); BXPC-3 (pancreas adenocarcinoma); MCF-7 (breast adenocarcinoma; SF 268 (CNS glioblastoma); NCI-H460 (lung large cell); KM 20L2 (colon adenocarcinoma); DU-145 (prostate carcinoma).