

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/24176760>

# Total Synthesis of Phomactin A

ARTICLE *in* ORGANIC LETTERS · MAY 2009

Impact Factor: 6.36 · DOI: 10.1021/ol900237e · Source: PubMed

---

CITATIONS

26

---

READS

15

5 AUTHORS, INCLUDING:



Grant S Buchanan

10 PUBLICATIONS 158 CITATIONS

SEE PROFILE

Published in final edited form as:


Org Lett. 2009 April 2; 11(7): 1591–1594. doi:10.1021/ol900237e.

## Total Synthesis of Phomactin A.†

Yu Tang, Kevin P. Cole, Grant S. Buchanan, Gang Li, and Richard P. Hsung\*

Division of Pharmaceutical Sciences and Department of Chemistry, University of Wisconsin, Madison, WI 53705

### Abstract



A total synthesis of (±)-phomactin A is described to highlight the final completion of a complex natural product target that had commenced with an intramolecular *oxa*-[3 + 3] annulation strategy in the construction of the ABD-tricycle. These efforts reveal structural intricacies of this ABD-tricycle with an illustrative example being the conformational analysis that was ultimately critical for the C5a-homologation.

In the last decade, phomactin A,<sup>1–4</sup> a structurally unique natural product isolated from the culture filtrate of a parasitic fungus *Phoma* sp. [SANK 11486] found on the shell of *Chinoecetes opilio*, has captured an impressive array of synthetic efforts.<sup>5,6</sup> Although possessing only modest inhibition against platelet-activating factor<sup>7</sup> [PAF] induced platelet aggregation [IC<sub>50</sub> = 10 μM], phomactin A embodies a new class of PAF antagonists. The most active member is D and was synthesized by Yamada.<sup>8</sup> Phomactin A is the only known tetracycle [discounting epoxides] in the phomactin family with all other members lacking either the B-ring or C-ring. Thus, phomactin A represents structurally the most complex member. To date, two elegant total syntheses of (±)- and (+)-phomactin A were completed by Pattenden<sup>9</sup> and Halcomb,<sup>10</sup> respectively. Recently, Wulff<sup>11</sup> reported the synthesis of phomactin B2.

We approached<sup>12</sup> (±)-phomactin A with an intent to feature our intramolecular *oxa*-[3 + 3] annulation strategy<sup>13–16</sup> en route to ABD-tricycle **2**, which possesses a unique structural topology [Scheme 1].<sup>17</sup> We recently designed a 12-step asymmetric synthesis of the annulation precursor **3**<sup>18</sup> entailing Suzuki-Miyaura coupling of vinyl bromide **4** with the derivative of A-ring **5**, which was assembled from an asymmetric Diels-Alder cycloaddition<sup>19,20</sup> of **6**. We communicate here our total synthesis of (±)-phomactin A.

An immense amount of effort<sup>17</sup> was exerted to succeed in oxidizing the C3-3a olefin of ABD-tricycle **2** into its proper oxidation states. Ultimately only a singlet-oxygen Diels-Alder cycloaddition could be achieved selectively to give endo-peroxide **8**<sup>21</sup> [Scheme 2] without

†This paper is dedicated to Professor Bill Wulff on the very special occasion of his 60<sup>th</sup> birthday.

E-mail: rhsung@wisc.edu.

**Supporting Information Available:** Experimental procedures as well as NMR spectra, characterizations, and X-ray structural files are available for all new compounds and free of charge via Internet <http://pubs.acs.org>.

significant competition from [2 + 2] cycloaddition or ene reaction with the C3'-4' olefin in D-ring or on the "belt." After failing an array of reductive protocols [i.e., Lindlar's [H], thiourea, or Ph<sub>3</sub>P] to cleave the weak endo-peroxide bond, KOAc and 18-c-6 successfully opened the endo-peroxide bridge via a deprotonation pathway to give ene-dione **9**. Treatment of **9** with *p*-TsOH in MeOH isomerized the lactol motif to methyl ether **10**, proceeding through a vinyl oxocarbenium intermediate that was trapped by MeOH at C3a position.

We recognized that while the singlet-oxygen Diels-Alder cycloaddition sets up the desired stereochemistry for the C3-OH group, ring-opening of the endo-peroxide bond through the deprotonation pathway effectively destroyed this valuable stereochemical information. Consequently, with the knowledge of the "belt" blocking the bottom face, we chose a small hydride source such as NaBH<sub>4</sub> to reduce the C3 ketone in **10**, but only to attain a mixture of isomers with a 4:1 ratio in favor of the wrong alcohol diastereomer **11-α**. Surprisingly, when using L-Selectride™, we isolated only the desired isomer **11-β**. In hindsight, by examining the model of **10** [Figure 1 – left side], it would appear that the pseudo-axial C3a-OMe group likely plays a bigger role than the "belt" in the facial differentiation of the reduction. With a more bulky hydride, the C3a-OMe group was able to better prevent the hydride approaching from the top face.

Reduction at C8a was relatively less eventful. As shown in Scheme 3, capping of C3-OH in **11-β** with TESCl followed by demethylation with BBr<sub>3</sub> led to vinylogous ester **12**. However, no condition that we screened [i.e., L-Selectride™, CuI/LAH, or Na/IPA] was capable of reducing the vinylogous ester motif in a 1,4-manner. Realizing that vinylogous ester **12** may not be sufficiently electron deficient, we oxidized C3a-OH using Dess-Martin periodinane reagent, and an ensuing reduction effectively gave diketone **13**, which is isolable, but with extended reduction time at temperatures slightly greater than rt, hydroxy ketone **14** was obtained in 96% overall yield with completely selective reduction at C3a.

With hydroxy ketone **14**, we completed our efforts in transforming B-ring into its proper oxidation states, and a single-crystal X-ray structure of **14** [Figure 1 – right side] further affirms our success. However, we were concerned about the reactivity of the C5a carbonyl group because the challenge of C5a-homologation lies ahead. It became obvious that no reduction of the C5a carbonyl group in **10** had occurred when using L-Selectride™, and nor did NaBH<sub>4</sub> touch the C5a carbonyl group in **13**. While one could concede that reduction in **10** involved a vinylogous ester, lack of reduction in **13** was quite disconcerting.

Upon examination of the minimized Spartan™ model of **10** and X-ray structure of **14**, we found some unique conformational elements. In **10**, the α-Me group in the A ring [red] is pseudo-equatorial with the β-Me group [blue] being pseudo-axial, thereby blocking any incoming nucleophiles toward the C5a carbonyl group. On the other hand, hydroxy ketone **14** assumes a very different conformation with its AB-ring junction being both sp<sup>3</sup>-hybridized instead of sp<sup>2</sup> as in **10**. In this case, the β-Me group [blue] is now pseudo-equatorial with the α-Me group [red] turning to occupy the pseudo-axial position, thereby hindering the attack of the C5a carbonyl group.

In contrast, the AB-ring junction of endo-peroxide **8** consists of sp<sup>3</sup>-hybridized C8a and sp<sup>2</sup>-hybridized C8b [Figure 2]. This set of hybridizations leads to yet another conformation in which the β-Me group [blue] remains pseudo-equatorial as in **14**, but the α-Me group [red] shifts away versus its respective position in **14**. We hoped that this minor shift would provide just enough opening to allow C5a to be accessible for homologation.

To test this hypothesis, we elected to construct enone **15** and epoxy ketone **16** for which both ring-junctions contain one sp<sup>3</sup>- and one sp<sup>2</sup>-hybridized carbon.<sup>22</sup> As shown in Scheme 4, after failing to eliminate the C3a-OH group via dehydrative protocols, we isolated sulfite **17** during

an attempt to chlorinate at C3a. A retro-Diels-Alder process in refluxing toluene would extrude SO<sub>2</sub> and afford the desired enone **15**. To our relief, we could add various one-carbon nucleophiles such as ROCH<sub>2</sub>Li [R = PBMB or MOM] to afford ene-diols **18a/b**, thereby succeeding what had appeared to be a daunting task in C5a-homologation. However, ene-diols **18a/b** were not useful for the total synthesis. Consequently, we prepared epoxy ketone **16**, but in three steps, because epoxidation of enone **15** would not take place unless the TES group was removed. Homologation of **16** via addition of MeLi followed by elimination gave vinyl epoxide **19**.

Nucleophilic ring-opening of vinyl epoxide **19** at C5 via a S<sub>N</sub>2' pathway would have been truly welcome at this stage [Scheme 5]. Instead, a S<sub>N</sub>1-like process occurred with H<sub>2</sub>O adding at C8b with retention of stereochemistry when using Mg(OTf)<sub>2</sub> in wet CH<sub>3</sub>CN,<sup>23</sup> leading to 1,2-diol **20** after re-silylating C3-OH. Allylic alcohol transposition using Dauben's protocol<sup>24</sup> led to epoxy diol **21**.<sup>25,26</sup> Subsequent treatment of **21** with Ph<sub>3</sub>P-I<sub>2</sub><sup>27</sup> followed by Luche reduction of the enal intermediate **22** gave 1,4-diol **23**, which was confirmed through its X-ray structure.

After failing a number of approaches for constructing the C-ring using either enal **22** or 1,4-diol **23**, mainly due to our inability to consistently oxidize the C3a-OH group, we managed to first acrylate C5-OH. Subsequently, we found that oxidation of the C3a-OH group employing Dess-Martin periodinane reagent at a warmer temperature gave enone **25**. An ensuing de-protection sequence allowed for the formation of the lactol C-ring and the final completion of our total synthesis of (±)-phomactin A in 24 steps from ABD-tricycle **2**.

We have described a total synthesis of (±)-phomactin A that highlights the final completion of a complex natural product target that had commenced with an intramolecular *oxa*-[3 + 3] annulation strategy in the construction of an ABD-tricycle. Our efforts reveal structural intricacies of this ABD-tricycle with an illustrative example being the conformational analysis that was ultimately critical for the C5a-homologation.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


We thank NIH [NS38049] for funding.

## References


1. Sugano M, Sato A, Iijima Y, Oshima T, Furuya K, Kuwano H, Hata T, Hanzawa H. J Am Chem Soc 1991;113:5463.
2. (a)Sugano M, Sato A, Saito K, Takaishi S, Matsushita Y, Iijima Y. J Med Chem 1996;39:5281. [PubMed: 8978857](b)Sugano M, Sato A, Iijima Y, Furuya K, Hata T, Kuwano H. J Antibiot 1995;48:1188. [PubMed: 7490234](c)Sugano M, Sato A, Iijima Y, Furuya K, Haruyama H, Yoda K, Hata T. J Org Chem 1994;59:564.
3. Koyama K, Ishino K, Takatori K, Takashi Sugita T, Kinoshita K, Takahashi K. Tetrahedron Lett 2004;45:6947.
4. Also see: Zhu X, Lambertino AT, Houghton TJ, McGilvra JD, Xu C, Rawal VH, Leff AR. Life Sciences 2003;73:3005. [PubMed: 14519449]
5. For reviews on synthetic efforts toward phomactins, see; (a)Cole KP, Hsung RP. ChemTracts 2003;16:811.(b)Goldring WPD, Pattenden G. Acc Chem Res 2006;39:354. [PubMed: 16700534]
6. For synthetic approaches, see: (a)Foote KM, Hayes CJ, Pattenden G. Tetrahedron Lett 1996;37:275. (b)Chen D, Wang J, Totah NI. J Org Chem 1999;64:1776. [PubMed: 11674261](c)Seth PP, Totah NI. J Org Chem 1999;64:8750.(d)Seth PP, Totah NI. Org Lett 2000;2:2507. [PubMed: 10956533](e)

- Kallan NC, Halcomb RL. *Org Lett* 2000;2:2687. [PubMed: 10990428](f)Chemler SR, Danishefsky SJ. *Org Lett* 2000;2:2695. [PubMed: 10990430](g)Seth PP, Chen D, Wang J, Gao X, Totah NI. *Org Lett* 2000;56:10185.(h)Foote K, John M, Pattenden G. *Synlett* 2001:365.(i)Mi B, Maleczka R. *Org Lett* 2001;3:1491. [PubMed: 11388849](j)Chemler SR, Iserloh U, Danishefsky SJ. *Org Lett* 2001;3:2949. [PubMed: 11554815](k)Houghton T, Choi S, Rawal VH. *Org Lett* 2001;3:3615. [PubMed: 11700095](l)Mohr PJ, Halcomb RL. *Org Lett* 2002;4:2413. [PubMed: 12098260](m)Balnaves AS, McGowan G, Shapland PDP, Thomas EJ. *Tetrahedron Lett* 2003;44:2713.(n)Cheing JWC, Goldring WPD, Pattenden G. *Chem Commun* 2003:2788.(o)Ryu K, Cho YS, Jung SI, Cho CG. *Org Lett* 2006;8:3343. [PubMed: 16836401](p)Huang J, Wang H, Wu C, Wulff WD. *Org Lett* 2007;9:2799. [PubMed: 17580880](q)Seth PP, Chen D, Wang J, Gao X, Totah NI. *Tetrahedron Lett* 2007;48:4605. [PubMed: 18575571](r)Peng W, Lee CS. *Synlett* 2008:142.
7. Platelet-activating factor is a phospholipid mediator released in the body by several cell types that play a role in causing inflammatory diseases. See: (a)Koltai M, Braquet PG. *Clin Rev Allergy* 1994;12:361. [PubMed: 7743462](b)Xhu X, Muñoz NM, Kim KP, Sano H, Cho W, Leff AR. *J Immunol* 1999;163:3423. [PubMed: 10477614]
  8. For total synthesis of (+)-phomactin D, see: Miyaoka H, Saka Y, Miura S, Yamada Y. *Tetrahedron Lett* 1996;37:7107.
  9. For total synthesis of (±)-phomactin A, see: (a)Goldring WPD, Pattenden G. *Chem Commun* 2002:1736.(b)Diaper CM, Goldring WPD, Pattenden G. *Org Biomol Chem* 2003;1:3949. [PubMed: 14664384](c)Foote KM, Hayes CJ, John MP, Pattenden G. *Org Biomol Chem* 2003;1:3917. [PubMed: 14664383]
  10. For total synthesis of (+)-phomactin A, see: Mohr PJ, Halcomb RL. *J Am Chem Soc* 2003;125:1712. [PubMed: 12580592]
  11. For total synthesis of (±)-phomactin B2, see: Huang J, Wu C, Wulff WD. *J Am Chem Soc* 2007;129:13366. [PubMed: 17929921]
  12. Cole KP, Hsung RP. *Org Lett* 2003;5:4843. [PubMed: 14653688]
  13. For reviews, see: (a)Harrity JPA, Provoost O. *Org Biomol Chem* 2005;3:1349. [PubMed: 15827625] (b)Hsung RP, Kurdyumov AV, Sydorenko N. *Eur J Org Chem* 2005;1:23.(b)Hsung RP, Cole KP, Harmata M. *Strategies and Tactics in Organic Synthesis* Elsevier Science, Pergamon PressOxford, UK2004;4:41.
  14. For our work, see: (a)Kurdyumov AV, Lin N, Hsung RP, Gullickson GC, Cole KP, Sydorenko N, Swiderski J. *J Org Lett* 2006;8:191.(b)Shen HC, Wang J, Cole KP, McLaughlin MJ, Morgan CD, Douglas CJ, Hsung RP, Coverdale HA, Gerasyuto AI, Hahn JM, Liu J, Wei L-L, Sklenicka HM, Zehnder LR, Zifcsak CA. *J Org Chem* 2003;68:1729. [PubMed: 12608785](c)Hsung RP, Wei L-L, Sklenicka HM, Douglas CJ, McLaughlin MJ, Mulder JA, Yao L. *Org Lett* 1999;1:509.(d)Hsung RP, Shen HC, Douglas CJ, Morgan CD, Degen SJ, Yao LJ. *J Org Chem* 1999;64:690. [PubMed: 11674130]
  15. For recent related studies, see: (a)Brioche JCR, Goodenough KM, Whatrup DJ, Harrity JPA. *J Org Chem* 2008;73:1946. [PubMed: 18254645](b)Yavari I, Sabbaghan M, Hossaini Z. *Synlett* 2008:1153.(c)Hubert C, Moreau J, Batany J, Duboc A, Hurvois JP, Renaud JL. *Adv Syn Cat* 2008;350:40.(d)Zhu M, Wei Q, Gong L. *Adv Synth Catal* 2008;350:1281.(e)Zhong W, Lin F, Chen R, Su W. *Synthesis* 2008:2561.(f)Brioche JCR, Goodenough KM, Whatrup DJ, Harrity JPA. *Org Lett* 2007;9:3491. [PubMed: 17685530](g)Lee YR, Xia L. *Synthesis* 2007:3240.(h)Epstein OL, Rovis T. *J Am Chem Soc* 2006;128:16480. [PubMed: 17177379]
  16. For applications in natural product syntheses, see: (a)Kurdyumov AV, Hsung RP. *J Am Chem Soc* 2006;128:6272. [PubMed: 16683764](b)Kurdyumov AV, Hsung RP, Ihlen K, Wang J. *Org Lett* 2003;5:3935. [PubMed: 14535747](c)Hsung RP, Cole KP, Zehnder LR, Wang J, Wei LL, Yang XF, Coverdale HA. *Tetrahedron* 2003;59:311.(d)Zehnder LR, Hsung RP, Wang J, Golding GM. *Angew Chem Int Ed* 2000;39:3876.(e)Malerich JP, Trauner D. *J Am Chem Soc* 2003;125:9554. [PubMed: 12903998](f)Olson BS, Trauner D. *Synlett* 2005:700.(g)Sunazuka T, Handa M, Nagai K, Shirahata T, Harigaya Y, Otoguro K, Kuwajima I, Ōmura S. *Tetrahedron* 2004;60:7845.(h)Hu H, Harrison TJ, Wilson PD. *J Org Chem* 2004;69:3782. [PubMed: 15153009](h)Lee YR, Wang X. *Bull Korean Chem Soc* 2005;26:1933.
  17. Cole KP, Hsung RP. *Chem Commun* 2005:5784.

18. You L, Hsung RP, Bedermann AA, Kurdyumov AK, Tang Y, Buchanan GS, Cole KP. *Adv Syn Cat* 2008;350:2885.
19. (a) Kozmin SA, Rawal VH. *J Org Chem* 1997;62:5252. (b) Kozmin SA, Rawal VH. *J Am Chem Soc* 1999;121:9562.
20. (a) Kozmin SA, Green MT, Rawal VH. *J Org Chem* 1999;64:8045. (b) Huang Y, Iwama T, Rawal VH. *J Am Chem Soc* 2000;122:7843. (c) Huang Y, Iwama T, Rawal VH. *Org Lett* 2002;4:1163. [PubMed: 11922808] (d) Kozmin SA, Iwama T, Huang Y, Rawal VH. *J Am Chem Soc* 2002;124:4628. [PubMed: 11971711]
21. See Supporting Information.
22. Spartan<sup>TM</sup> models of enone 15 and epoxy ketone 16 reveal similar conformation as in endo-peroxide 8.
23. For related protocols, see: (a) Boyer FD, Hanna I. *J Org Chem* 2005;70:1077. [PubMed: 15675876] (b) Iranpoor N, Shekarriz M, Shiriny F. *Syn Commun* 1998;28:347.
24. Dauben WG, Michno DM. *J Org Chem* 1977;42:682.
25. For a recent example, see: (a) Chai Y, McIntosh MC. *Tetrahedron Lett* 2004;45:3269.. Also see: (b) Sundararaman P, Herz W. *J Org Chem* 1977;42:813. (c) Chu A, Mander LN. *Tetrahedron Lett* 1988;29:2727.
26. To circumvent this epoxide formation, we also attempted MeReO<sub>3</sub> that was successful in a related 1,3-transposition, see: (a) John M, Hutchison JM, Lindsay HA, Dormi SS, Jones GD, Vicic DA, McIntosh MC. *Org Lett* 2006;8:3663. [PubMed: 16898786]. Also, see: (b) Jacob J, Espenson JH, Jensen JH, Gordon MS. *Organometallics* 1998;17:1835. (c) Morrill C, Grubbs RH. *J Am Chem Soc* 2005;127:2842. [PubMed: 15740106]
27. Wydra H, Paryzek Z. *Tetrahedron Lett* 1984;25:2601.




**Figure 1.**  
Conformation Analysis for **10** and **14**.




**Figure 2.**  
Rationale for Choosing **15** and **16**.







**Scheme 1.**  
A Synthetic Plan Toward Phomactin A.


**Scheme 2.**

Oxidation of C3-3a Olefin: An Endo-Peroxide Route.



**Scheme 3.**  
Reduction of C8a at the AB-Ring Junction.

**Scheme 4.**C5a-Homologation of Enone **15** and Epoxy Ketone **16**.



**Scheme 5.**  
Completing a Total Synthesis of (±)-Phomactin A.