

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/23674837>

Iron-Catalyzed 1,4-Addition of α -Olefins to Dienes

ARTICLE in ORGANIC LETTERS · JANUARY 2009

Impact Factor: 6.36 · DOI: 10.1021/ol802524r · Source: PubMed

CITATIONS

73

READS

32

3 AUTHORS:



Benoit Moreau

Blend Therapeutics

18 PUBLICATIONS 310 CITATIONS

SEE PROFILE



Jessica Wu

University of Liverpool

10 PUBLICATIONS 259 CITATIONS

SEE PROFILE



Tobias Ritter

Harvard University

99 PUBLICATIONS 4,841 CITATIONS

SEE PROFILE

Iron-Catalyzed 1,4-Addition of α -Olefins
to Dienes


Benoît Moreau, Jessica Y. Wu, and Tobias Ritter*

Department of Chemistry and Chemical Biology, Harvard University,
Cambridge, Massachusetts 02138

ritter@chemistry.harvard.edu

Received October 31, 2008

ABSTRACT



A new intermolecular, stereo- and regioselective iron-catalyzed 1,4-addition of α -olefins to 1,3-dienes using as low as 1 mol % of an iminopyridine–ferrous chloride complex was developed. Importantly, both double bonds of the linear 1,4-diene addition products are obtained with absolute stereocontrol.

Iron catalysis has the potential to form valuable carbon–carbon bonds starting from simple starting materials such as hydrocarbons.¹ In this communication we present a functional-group-tolerant, stereo- and regioselective iron-catalyzed 1,4-addition reaction of α -olefins to 1,3-dienes (eq 1). Previously, in the 1960's, Hata investigated the addition of butadiene and isoprene to ethylene and propylene in an autoclave at >500 psi catalyzed by an Fe(acac)₃/Et₃Al mixture.² Using the same Fe(acac)₃/Et₃Al mixture but in the presence of 2,2'-bipyridine, Takacs observed formal [4 + 4] ene reactions with 1,3-dienes.³ The iron-diimine-catalyzed dimerization of 1,3-dienes to afford cyclooctadienes and vinyl cyclohexenes was developed by tom Dieck in the 1980's.⁴ Within the past 5 years, Fürstner has shown that a well-defined anionic iron(0) complex, supported by a cyclopentadienyl ligand, can catalyze intramolecular ene reactions,⁵ intramolecular cycloadditions, and cycloisomerizations.⁶ In 2006, Chirik demonstrated the intramolecular [2 + 2] cycloaddition of 1,6-dienes catalyzed by a redox-active bisiminopyridine–iron complex.⁷

We have investigated iminopyridine-ligated iron complexes that provide access to linear 1,4-diene products that cannot be obtained with tom Dieck's or Takacs' catalysts or by any other hydrovinylation reaction reported to date.⁸ Our 1,4-addition reaction complements the functional-group-tolerant cobalt-catalyzed 1,4-hydrovinylation reaction, pioneered by Hilt, which typically affords branched 1,4-dienes.⁹ Notable features of the carbon–carbon bond-forming reaction presented herein include the regioselective introduction of prenyl and geranyl substituents, the stereospecific formation of (*E*)-trisubstituted double bonds, and the functionalization of unactivated commodity chemicals using a simple low-valent iron catalyst.

The readily available iron(II) complex **1**·FeCl₂ shown in eq 1 can be reduced in situ with activated magnesium metal

(1) For recent reviews of iron catalysis, see: (a) Bolm, C.; Legros, J.; Le Pailh, J.; Zani, L. *Chem. Rev.* **2004**, *104*, 6217. (b) Enthaler, S.; Junge, K.; Beller, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 3317. (c) Correa, A.; Mancheno, O. G.; Bolm, C. *Chem. Soc. Rev.* **2008**, *37*, 1108. (d) Sherry, B. D.; Fürstner, A. *Acc. Chem. Res.* **2008**, *41*, 1500.

(2) (a) Hata, G. *J. Am. Chem. Soc.* **1964**, *86*, 3903. (b) Hata, G.; Aoki, D. *J. Org. Chem.* **1967**, *32*, 3754. (c) Hata, G.; Miyake, A. *Bull. Chem. Soc. Jpn.* **1968**, *41*, 2762. For related publications, see: (d) Iwamoto, M.; Yuguchi, S. *J. Org. Chem.* **1966**, *31*, 4290. (e) Iwamoto, M.; Yuguchi, S. *Bull. Chem. Soc. Jpn.* **1966**, *39*, 2001.

(3) (a) Takacs, J. M.; Anderson, L. G.; Madhavan, G. V. B.; Creswell, M. W.; Seely, F. L.; Devroy, W. F. *Organometallics* **1986**, *5*, 2395. (b) Takacs, J. M.; Anderson, L. G.; Madhavan, G. V. B.; Seely, F. L. *Angew. Chem., Int. Ed.* **1987**, *26*, 1013. (c) Takacs, J. M.; Anderson, L. G.; Newsome, P. W. *J. Am. Chem. Soc.* **1987**, *109*, 2542. (d) Takacs, J. M.; Myoung, Y.-C.; Anderson, L. G. *J. Org. Chem.* **1994**, *59*, 6928. (e) Takacs, J. M.; Newsome, P. W.; Kuehn, C.; Takusagawa, F. *Tetrahedron* **1990**, *46*, 5507.

(4) (a) tom Dieck, H.; Dietrich, J. *Chem. Ber.-Recueil* **1984**, *117*, 694. (b) tom Dieck, H.; Dietrich, J. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 781.

(5) Fürstner, A.; Martin, R.; Majima, K. *J. Am. Chem. Soc.* **2005**, *127*, 12236.

(6) Fürstner, A.; Majima, K.; Martin, R.; Krause, H.; Kattnig, E.; Goddard, R.; Lehmann, C. W. *J. Am. Chem. Soc.* **2008**, *130*, 1992.

(7) Bouwkamp, M. W.; Bowman, A. C.; Lobkovsky, E.; Chirik, P. J. *J. Am. Chem. Soc.* **2006**, *128*, 13340.


Table 1. Iron-Catalyzed 1,4-Addition of Olefins to 1,3-Dienes

$\text{R}^1\text{CH=CH}_2 + \text{CH}_2=\text{C}(\text{Me})\text{CH}(\text{R}^2)\text{CH}(\text{R}^3)\text{CH}_2 \xrightarrow[\text{Et}_2\text{O (1 M)}]{\text{1 or 2 + FeCl}_2 \text{ (2 mol \%)} \text{ Mg (4 mol \%)}} \text{R}^1\text{CH=CH-CH}(\text{R}^2)\text{CH}(\text{R}^3)\text{CH}_2$									
entry	alkene	diene	product	X	ligand	yield	regioselectivity ^a	1-E/Z	4-E/Z
1				H (3)		94 ^b	—	>99:1	—
2				F (4b)		86	—	>99:1	—
3				Cl (4c)		78	—	>99:1	—
4				OMe (4d)		85	—	>99:1	—
5				CF ₃ (4e)		90	—	>99:1	—
6				^t Bu (4f)		92	—	>99:1	—
7				H (4a)		60 ^c	90:10	>99:1	—
8		isoprene		F (4g)		74 ^c	91:9	>99:1	—
9				H (4h)		66 ^d	93:7	>99:1	>99:1
10		myrcene		OMe (4i)		51 ^d	96:4	>99:1	>99:1
11	Ph			4j	1	77	>98:2	>99:1	>99:1
12	Ph			4k	1	79 ^e	—	>99:1	—

^a Regioisomeric ratio as described (eq 2). ^b 1 mol % 1•FeCl₂ used. ^c 20 mol % 2•FeCl₂ used. ^d 10 mol % 2•FeCl₂ used. ^e 5 mol % 1•FeCl₂ used.

to catalyze carbon–carbon bond formation between styrene and 2,3-dimethylbutadiene to afford 1,4-diene **3** in 94% yield after 6 h at 23 °C. Control experiments showed that the presence of iron, together with reducing agent and ligand, is essential for productive catalysis. A 1:1 complex of ferrous chloride and iminopyridine ligand¹⁰ (for X-ray, see Supporting Information) is responsible for the exclusive formation of the intermolecular addition product **3**, with no traces of diene dimerization or subsequent observable double bond isomerization. A selection of 18 other complexes derived from bidentate nitrogenous ligands afforded the desired linear 1,4-diene **3** in less than 50% yield, provided significant amounts of diene dimerization products as observed by tom Dieck,^{4a} or gave double bond isomerization to form a conjugated 1,3-diene subsequent to addition (see Supporting Information). As shown in Table 1, 2,3-dimethylbutadiene

can be added to electron-rich and -poor styrenes bearing various functional groups such as ethers, an ester, and halides. This practical procedure can be scaled to 100 mmol using 2 mol % of catalyst, 100 mL of solvent, and commercially available styrene and isoprene (eq 2).



(8) (a) RajanBabu, T. V. *Chem. Rev.* **2003**, *103*, 2845. (b) Bogdanovi, B.; Pauling, H.; Wilke, G.; Meister, B.; Henc, B. *Angew. Chem., Int. Ed.* **1972**, *11*, 1023. (c) Zhang, A. B.; RajanBabu, T. V. *J. Am. Chem. Soc.* **2006**, *128*, 54. (d) Shirakura, M.; Sugimoto, M. *J. Am. Chem. Soc.* **2008**, *130*, 5410. (e) Saha, B.; Smith, C. R.; RajanBabu, T. V. *J. Am. Chem. Soc.* **2008**, *130*, 9000.

(9) (a) Hilt, G.; du Mesnil, F. X.; Lüers, S. *Angew. Chem., Int. Ed.* **2001**, *40*, 387. (b) Hilt, G.; Lüers, S. *Synthesis* **2002**, 609. (c) Hilt, G.; Lüers, S.; Schmidt, F. *Synthesis* **2004**, 634. The functional-group-tolerant Co-catalyzed 1,4-hydrovinylation typically affords the branched 1,4-dienes; if acrylates are used as enes, linear 1,4-diene products are observed, see ref 9c.

(10) Gibson, V. C.; O'Reilly, R. K.; Wass, D. F.; White, A. J. P.; Williams, D. J. *Dalton Trans.* **2003**, 2824. For a related review of bisiminopyridine–iron complexes, see: (b) Gibson, V. C.; Redshaw, C.; Solan, G. A. *Chem. Rev.* **2007**, *107*, 1745.

Although the reaction shown in eq 2 afforded 1,4-addition product in 79% yield, two regioisomeric products were obtained due to bond formation at either of the two different olefin termini of isoprene. The regioselectivity in favor of prenylation product **4a** induced by ligand **1** was 86:14. The introduction of a prenyl group upon 1,4-addition to isoprene is synthetically useful,¹¹ but control over regioselectivity is challenging. To increase the regioselectivity, we evaluated different iminopyridine ligands and found a strong dependence of the regioselectivity on the ligand. The highest ratio of 10:1 in favor of the desired linear 1,4-diene product was observed with the trimethylsilyl-substituted ligand **2** (Table 1). Methyl substitution at the 6-position of the pyridyl ligand



can invert the regioselectivity to 16:84 to afford **5a** as the major isomer (see Supporting Information). Substitution of the 1,3-diene starting material with substituents larger than methyl, such as that in the commodity chemical myrcene, increased the regioselectivity to up to >20:1 (entry 10, **4i**). Using this procedure, the geranyl substituent can be introduced regioselectively. A diene with branched substitution afforded a single regioisomer as determined by ^1H NMR (entry 11, **4j**). Although the regioselectivity was increased using ligand **2**, higher catalyst loadings were required to achieve full conversion of starting material.

The geometry of both double bonds of the 1,4-diene products was controlled, and all addition products were observed as single double bond isomers. Importantly, the trisubstituted double bond that is generated upon 1,4-addition is formed as its *E*-isomer exclusively (entries 9–11); trisubstituted double bonds with three alkyl substituents are otherwise difficult to form as *E*-isomers with high selectivity.¹² The aliphatic α -olefin allylbenzene afforded a single addition product after double bond transposition (entry 12, **4k**). Addition of olefins to 1-substituted 1,3-dienes afforded addition products, albeit in lower yields. Overall, the iminopyridine iron complexes control stereo- and regioselectivity and afford linear 1,4-diene addition products selectively over potential dimerization, isomerization, or branched products.

Although 1,4-addition reactions proceeded with aliphatic α -olefins, product mixtures were generally observed. In addition to the desired linear 1,4-dienes, branched 1,4-dienes⁹ and dienes resulting from double bond isomerization subsequent to addition were detected. Therefore, the presented addition reaction is currently limited to styrene derivatives and selected aliphatic α -olefins (e.g., **4k**, Table 1).

A mechanistic hypothesis that is consistent with our experimental observations is detailed in Scheme 1. We propose formation of iron(II) alkyl allyl complex **8** after reduction¹³ of the Fe(II) complex **1**•FeCl₂ and oxidative

coupling with styrene. We cannot yet rationalize the influence of the catalyst on the regioselectivity that may be determined during oxidative coupling. A π - σ rearrangement to form the seven-membered η^1 -allyl ferracycle **9**, in which the phenyl substituent is pseudoequatorial, positions only one β -hydride (H_Z) *syn* to iron. Hence, β -hydride elimination can only afford a single double bond isomer and delivers the 1,2-disubstituted *E* double bond as part of the iron(II) alkyl hydride **10**. Assuming that subsequent reductive elimination proceeds without σ - π - σ rearrangement of **10**, the new trisubstituted double bond is formed with *E* geometry stereospecifically. To substantiate the assumption that stereospecific reductive elimination from **10** is faster than isomerization, we designed two deuterium-labeling experiments. The deuterium atoms of β,β -dideuterostyrene and (*E*)- β -deuterostyrene in the 1,4-addition reactions to isoprene were observed exclusively at the positions indicated in **11** and **12**, respectively, consistent with our mechanistic hypothesis (eqs 3 and 4).




In conclusion, we have developed an intermolecular iron-catalyzed 1,4-addition of α -olefins to 1,3-dienes using readily available iminopyridine–ferrous chloride complexes. Importantly, both regioselectivity and double bond geometry of the addition products can be controlled. Furthermore, synthetically useful products such as prenyl- and geranyl-substituted molecules are synthesized from simple commodity chemicals.

Acknowledgment. We thank Merck & Co. for unrestricted support, FQRNT for a postgraduate fellowship for B.M., Novartis for a graduate fellowship for J.Y.W., Materia for a gift of Grubbs catalyst, Dr. Douglas M. Ho (Harvard University) for X-ray crystallographic analysis, and Rebecca Loy and Stephan Zuend (Harvard University) for GC analysis.

Supporting Information Available: Detailed experimental procedures and spectroscopic data for all new compounds, as well X-ray data in CIF format. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL802524R

Scheme 1. Proposed Catalytic Cycle for 1,4-Addition to Dienes



(11) *Isoprenoids Including Carotenoids and Steroids*; Cane, D. E., Ed.; *Comprehensive Natural Products Chemistry*; Barton, D., Nakanishi, K., Eds.; Elsevier: Oxford, 1999; Vol. 2.

(12) Yu, J. S.; Kleckley, T. S.; Wiemer, D. F. *Org. Lett.* **2005**, 7, 4803.

(13) A related (1,2-diimine)iron(diene) complex was obtained similarly: Bart, S. C.; Hawrelak, E. J.; Lobkowsky, E.; Chirik, P. J. *Organometallics* **2005**, 24, 5518.