

## A radical thia-Brook rearrangement†‡

Béatrice Quiclet-Sire\* and Samir Z. Zard\*

Cite this: *Chem. Commun.*, 2014, 50, 5990Received 5th March 2014,  
Accepted 14th April 2014

DOI: 10.1039/c4cc01683a


www.rsc.org/chemcomm

**Geminal mercapto trialkyl- and trialkoxy-silanes undergo an efficient radical chain rearrangement, whereby the silyl group migrates from carbon to sulfur; the starting materials are readily obtained by exploiting the peroxide initiated radical addition of dithiocarbonates (xanthates) to trialkyl- or trialkoxy-vinylsilanes.**

We recently found that various xanthates readily add to vinyl trialkoxysilanes allowing access to numerous functional trialkoxysilane derivatives **3** (Scheme 1).<sup>1</sup> This peroxide initiated radical chain addition is flexible, modular, and experimentally very simple to implement.<sup>2</sup> Trialkoxysilanes are of key importance in numerous areas: material sciences, sol gels and organogelators, surface modification and monolayer formation, especially on metal oxides and silica surfaces, supported catalysts *etc.*<sup>3</sup> In the course of this work we stumbled upon an unexpected and very efficient radical thia-Brook rearrangement we now describe.

While the xanthate group in adduct **3** may be reductively removed if needed, it is in fact a protected form of the corresponding thiol **4**.<sup>4</sup> The presence of the thiol would indeed offer some further interesting possibilities, either by itself as a cross-linking handle (*via* the disulfide) or as a springboard for a host of transformations through ionic (*e.g.* alkylation) or radical (*e.g.* addition to alkenes, the so-called “thiol click reaction”<sup>5</sup>) processes. With these considerations in mind, we attempted to generate the thiol through the Chugaev reaction<sup>6,7</sup> by simply heating adduct **3a** in diphenyl ether at 200 °C. The thermolysis takes place under neutral conditions that should not affect groups sensitive to nucleophilic attack, such as the phthalimido group present in **3a**.

In the event, heating xanthate **3a** in diphenyl ether at 200 °C for 1 h furnished after purification two inseparable products, one of which was indeed the expected thiol **4a** but the other



Scheme 1 A possible route to geminal mercapto trialkoxysilanes.

turned out to be rearranged trialkoxysilane **5a**, as determined by analysis of the NMR spectrum of the mixture. We noticed, furthermore, that, while the combined yield of **4a** and **5a** was generally good, their relative yield varied significantly with the exact experimental conditions. In one experiment, the ratio of **4a**:**5a** by NMR was approximately 2:1 in a combined yield of 82% (reaction time 1 h).

The unexpected formation of rearranged trialkoxysilane **5a** may be rationalised by the mechanism displayed in Scheme 2 and proceeding through a radical thia-Brook rearrangement.<sup>8</sup> In one experiment, we succeeded in isolating a pure sample of thiol **4a** in 53% yield and, indeed, exposing it to di-*t*-butyl peroxide (DTBP) in refluxing chlorobenzene for 1 h afforded a quantitative yield of rearranged silane **5a**. This mechanism also accounts for the variability in the relative yield of **4a** and **5a** initially observed, since the efficiency of the chain reaction leading to the latter depends on the presence of adventitious radical initiators (oxygen, traces of metallic salts, *etc.*).


Whereas the radical Brook rearrangement involving silicon migration from carbon to oxygen is well documented,<sup>9</sup> only one instance of a radical thia-Brook rearrangement has been reported as far as we know.<sup>10</sup> It involves the cleavage of a silicon–silicon bond in going from sulfur radical **10** to silicon radical **11** (Scheme 3). This step was incorporated in a radical sequence allowing the use of thiol **9** in ingenious tin-free reductive dehalogenations and Barton–McCombie deoxygenations.

Rupture of a carbon–silicon bond in a thia-Brook rearrangement, as in the present case, appears to be unprecedented.


Laboratoire de Synthèse Organique, CNRS UMR 7652, Ecole Polytechnique, F-91128 Palaiseau, France. E-mail: beatrice.sire@polytechnique.edu, samir.zard@polytechnique.edu; Fax: +33 169335972; Tel: +33 169335971

† This paper is dedicated with respect to the memory of Professor Adrian G. Brook (University of Toronto).

‡ Electronic supplementary information (ESI) available. See DOI: 10.1039/c4cc01683a



Scheme 2 An unexpected radical thia-Brook rearrangement.




Scheme 3 Cleavage of a silicon-silicon bond by a thia-Brook rearrangement.

The driving force may be the formation of a carbon radical **7a** stabilised by a sulfur atom, with the possible equilibrium between intermediates **6a** and **7a** being finally driven by the irreversible hydrogen abstraction from another thiol molecule **4a**.


A similar sequence could be accomplished starting from the addition product of xanthate **1a** to vinyl trimethylsilane **2a'** (Scheme 2). Thus, thermolysis of **3a'** indeed furnished the corresponding rearranged derivative **5a'**, but this compound was too labile to chromatographic purification and decomposed to the free thiol **8**, which was isolated in 60% yield. A small amount of un-rearranged thiol **4a'** (*ca.* 10%) was also observed by NMR of the crude reaction mixture. In view of the hydrolytic lability of trimethylsilyl derivatives, the remainder of the study was conducted with the tri(*t*-butoxy)silyl derivatives.

Our next task was to examine the scope of the reaction. Addition of chloropyridinyl xanthates **1b** to vinyl tri(*t*-butoxy)silane **2a** afforded adduct **3b** in 68% yield (Scheme 4). Thermolysis for 1 h gave a mixture of thiol **4b** (42%) and thia-Brook product **5b** (28%). Repetition of the experiment and exposure of the crude product from the thermolysis to DTBP in refluxing chlorobenzene for 2 h furnished **5b** in a better yield (78%). In the case of adduct **3c**, derived from xanthate **1c**, no attempt was made to separate the intermediate thiol. The crude mixture was treated directly with DTBP in refluxing chlorobenzene to furnish the rearranged material **5c** in 62% yield.




Scheme 4 Further examples of the radical thia-Brook rearrangement.

An unexpected problem was encountered with substrate **3d**, prepared using xanthate **1d** (Scheme 5). The thermolysis step gave, after chromatographic purification, thiolactone **12** (39%) and a polar mixture of compounds. This mixture was simply



Scheme 5 Unexpected formation of a thialactone.



Scheme 6 Additional examples of the radical thia-Brook rearrangement.

subjected to the action of DTBP in refluxing chlorobenzene. Purification then afforded aminopyridine **13** (41%) and the radical thia-Brook rearrangement product **5d** (55%). Aminopyridine **13** is the leaving group in the formation thiolactone **12**. This latter compound would be difficult to obtain by more conventional routes and is interesting in its own right, for example as a cross-linking agent in material science; however, in the present context, its formation by attack of the thiol sulfur on the activated amide is clearly in competition with the desired radical thia-Brook rearrangement.

To circumvent this complication, we resorted to a more traditional cleavage of the xanthate group by aminolysis with 1,2-ethylenediamine.<sup>4</sup> Thus, treatment with xanthate **3d** with 1,2-ethylenediamine in a 1 : 1 (v/v) mixture of ethanol and ether at room temperature gave the crude thiol **4d**, which was not purified but directly heated in refluxing chlorobenzene with DTBP. This gave the expected rearranged product **5d** in good yield. No thiolactone **12** or the corresponding aminopyridine **13** were observed under these conditions. Because of the lower temperature (heating in refluxing chlorobenzene at 130 °C vs. thermolysis in diphenyl ether at 200 °C) and, especially, the presence of the DTBP initiator, the radical chain process overcomes the intramolecular ionic ring-closure leading to thiolactone **12**.

The *O*-isopropyl group in the xanthate is now not needed any more and can be replaced by the simpler *O*-ethyl analogue.

Thus, adduct **3c'**, obtained in 52% yield by the radical addition of benzotriazole xanthate **1c'**<sup>11</sup> to vinyl tri(*t*-butoxy)silane **2a**, was cleaved by 1,2-ethylenediamine into thiol **4c** (76% yield) and the latter rearranged quantitatively into tri(*t*-butoxy)silyl sulfide **5c** by heating in refluxing chlorobenzene in the presence of DTBP initiator (Scheme 6).

In the same manner, but without purification of the intermediate thiols, xanthates **3e–g**<sup>1</sup> underwent conversion into the corresponding radical thia-Brook rearrangement products **5e–g** (Scheme 6). The possibility of introducing a *geminal* trifluoromethyl acetamido motif is worthy of note.

In summary, we have described a hitherto unknown migration of a silicon group from carbon to sulfur by a radical chain mechanism, a process that may be viewed as a formal radical thia-Brook rearrangement. This provides a route to a plethora of otherwise inaccessible functionalised silyl sulfides **5**. The possibility of capturing intermediate carbon radical **7** (Scheme 2) before hydrogen atom abstraction from the thiol has occurred, for example by cyclisation to a suitably located internal alkene, could also be of some synthetic interest. Studies along these lines are underway.

## Notes and references

- 1 B. Quiclet-Sire, Y. Yanagisawa and S. Z. Zard, *Chem. Commun.*, 2014, **50**, 2324.
- 2 For reviews of the xanthate transfer, see: (a) S. Z. Zard, *Angew. Chem., Int. Ed. Engl.*, 1997, **36**, 672; (b) B. Quiclet-Sire and S. Z. Zard, *Top. Curr. Chem.*, 2006, **264**, 201; (c) B. Quiclet-Sire and S. Z. Zard, *Chem. – Eur. J.*, 2006, **12**, 6002; (d) B. Quiclet-Sire and S. Z. Zard, *Pure Appl. Chem.*, 2011, **83**, 519; for an account of the discovery of the basic process, see: (e) S. Z. Zard, *Aust. J. Chem.*, 2006, **59**, 663.
- 3 (a) S. Fujita and S. Inagaki, *Chem. Mater.*, 2008, **20**, 891; (b) H. Zou, S. Wu and J. Shen, *Chem. Rev.*, 2008, **108**, 3893; (c) M. Llusar and C. Sanchez, *Chem. Mater.*, 2008, **20**, 782; (d) D. B. Cordes, P. D. Lickiss and F. Rataboul, *Chem. Rev.*, 2010, **110**, 2081; (e) S. Onclin, B. J. Ravoo and D. N. Reinhoudt, *Angew. Chem., Int. Ed.*, 2005, **44**, 6282; (f) A. Ulman, *Chem. Rev.*, 1996, **96**, 1533; (g) A. P. Wight and M. E. Davis, *Chem. Rev.*, 2002, **102**, 3589; (h) Z.-L. Lu, E. Lindner and H. A. Mayer, *Chem. Rev.*, 2002, **102**, 3543.
- 4 (a) F. Duus, in *Comprehensive Organic Chemistry*, ed. D. H. R. Barton and W. D. Ollis, Pergamon Press, Oxford, 1979, vol. 3, pp. 373–487; (b) S. Ramachandra Rao, *Xanthates and Related Compounds*, Marcel Dekker Inc., New York, 1971.
- 5 (a) C. E. Hoyle and C. N. Bowman, *Angew. Chem., Int. Ed.*, 2010, **49**, 1540; (b) A. K. Tucker-Schwartz, R. A. Farrell and R. L. Garrell, *J. Am. Chem. Soc.*, 2011, **133**, 11026; (c) E. L. Tyson, M. S. Ament and T. P. Yoon, *J. Org. Chem.*, 2013, **78**, 2046.
- 6 (a) L. Chugaev, *Chem. Ber.*, 1899, **32**, 3332; (b) H. R. Nace, *Org. React.*, 1962, **12**, 57.
- 7 (a) K. K. K. Goh, S. Kim and S. Zard, *J. Org. Chem.*, 2013, **78**, 12274; (b) B. Quiclet-Sire and S. Z. Zard, *Org. Lett.*, 2013, **15**, 5886.
- 8 (a) A. G. Brook, *Acc. Chem. Res.*, 1974, **7**, 77; for general reviews, see: (b) M. A. Brook, *Silicon in Organic, Organometallic, and Polymer Chemistry*, Wiley, New York, 1999; (c) H. Moser, *Tetrahedron*, 2001, **57**, 2065.
- 9 For general reviews on organosilanes in radical chemistry, see: (a) C. Chatgililoglu, *Acc. Chem. Res.*, 1992, **25**, 188; (b) C. Chatgililoglu, *Chem. – Eur. J.*, 2008, **14**, 2310; (c) C. Chatgililoglu, *Organosilanes in Radical Chemistry*, Wiley, Chichester, 2003.
- 10 M. Ballestri, C. Chatgililoglu and G. Seconi, *J. Organomet. Chem.*, 1991, **408**, C1.
- 11 A. R. Katritzky, M. A. C. Button and S. N. Denisenko, *Heterocycles*, 2001, **54**, 301.