

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/5604013>

Palladium(II)-Directed Self-Assembly of Dynamic Donor-Acceptor [2]Catenanes

ARTICLE *in* ORGANIC LETTERS · APRIL 2008

Impact Factor: 6.36 · DOI: 10.1021/ol702830p · Source: PubMed

CITATIONS

53

READS

27

4 AUTHORS, INCLUDING:



Yi Liu

Dalian University of Technology

455 PUBLICATIONS 9,259 CITATIONS

SEE PROFILE



Jiu-Ming He

Chinese Academy of Medical Sciences

63 PUBLICATIONS 985 CITATIONS

SEE PROFILE

Palladium(II)-Directed Self-Assembly of Dynamic Donor–Acceptor [2]Catenanes


Yi Liu,^{*,†} Angela Bruneau,[†] Jiuming He,[‡] and Zeper Abliz[‡]

The Molecular Foundry, Lawrence Berkeley National Laboratory, One Cyclotron Road, Berkeley, California 94720, and Institute of Materia Medica, Chinese Academy of Medical Sciences and Peking Union Medical College, Beijing 100050, China

yliu@lbl.gov

Received November 22, 2007

ABSTRACT




Highly efficient syntheses of donor–acceptor [2]catenanes were developed using a combination of templation and reversible metal–ligand coordination. The desired [2]catenanes were obtained within minutes through a five-component assembly, involving a donor-containing crown ether, an acceptor-containing ligand, two Pd(II) metal centers, and a dipyrindyl ligand. The [2]catenane formation was characterized by ¹H NMR and UV–vis spectroscopies and cold-spray ionization mass spectrometry. In particular, great translational selectivity was observed when a crown ether with two different donor units was employed.

[2]Catenanes¹ have attracted great attention owing to the nontrivial interlocking topology² and nanoscale motions³ associated with their movable macrocyclic components. The advancement on templated synthesis⁴ has led to multistable catenanes that function as reconfigurable molecular switches⁵ and artificial molecular motors.⁶ Indeed, bistable donor–

acceptor [2]catenanes⁷ have become one of the representative species in the molecular based electronic devices.⁸ A majority of donor–acceptor [2]catenanes was assembled through a *kinetically* controlled process to contain two interlocked macrocyclic components, one being a π -electron-donating crown ether and the other being a tetracationic cyclophane. Meanwhile, assembly of donor–acceptor [2]catenanes through *thermodynamic* processes was exercised to a lesser extent. An elegant example was reported⁹ recently where the donor–

[†] Lawrence Berkeley National Laboratory.[‡] Chinese Academy of Medical Sciences and Peking Union Medical College.(1) (a) Schill, G. *Catenanes, Rotaxanes, and Knots*; Academic Press: New York, 1971. (b) Sauvage, J.-P.; Dietrich-Buchecker, C., Eds. *Molecular Catenanes, Rotaxanes, and Knots*; Wiley-VCH: Weinheim 1999.(2) Amabilino, D. B.; Stoddart, J. F. *Chem. Rev.* **1995**, 95, 2725–2828.(3) (a) Balzani, V.; Credi, A.; Raymo, F. M.; Stoddart, J. F. *Angew. Chem., Int. Ed.* **2000**, 112, 3484–3530. (b) Dietrich-Buchecker, C. O.; Jimenez-Molero, M. C.; Sartor, V.; Sauvage, J.-P. *Pure Appl. Chem.* **2003**, 75, 1383–1393. (c) Kay, E. R.; Leigh, D. A.; Zerbetto, F. *Angew. Chem., Int. Ed.* **2007**, 46, 72–191.(4) (a) Dietrich-Buchecker, C. O.; Sauvage, J. P. *Chem. Rev.* **1987**, 87, 795–810. (b) Diederich, F.; Stang, P. J., Eds. *Templated Organic Synthesis*; Wiley-VCH: Weinheim, 1999.(5) (a) Green, J. E.; Choi, J. W.; Boukai, A.; Bunimovich, Y.; Johnston-Halperin, E.; DeIonno, E.; Luo, Y.; Sherif, B. A.; Xu, K.; Shin, Y. S.; Tseng, H.-R.; Stoddart, J. F.; Heath, J. R. *Nature* **2007**, 445, 414–417. (b) Flood, A. H.; Stoddart, J. F.; Heath, J. R. *Science* **2004**, 306, 2055–2056. (c) Feringa, B. L., Ed. *Molecular Switches*; Wiley-VCH: Weinheim, 2001.(6) (a) Serreli, V.; Lee, C.-F.; Kay, E. R.; Leigh, D. A. *Nature* **2007**, 445, 523–527. (b) Balzani, V.; Credi, A.; Ferrer, B.; Silvi, S.; Venturi, M. *Top. Curr. Chem.* **2005**, 262, 1–27. (c) Mandl, C. P.; König, B. *Angew. Chem., Int. Ed.* **2004**, 43, 1622–1624. (d) Hernandez, J. V.; Kay, E. R.; Leigh, D. A. *Science* **2004**, 306, 1532–1537.(7) (a) Liu, Y.; Klivansky, L. M.; Zhang, X. *Org. Lett.* **2007**, 9, 2577–2580. (b) Asakawa, M.; Ashton, P. R.; Balzani, V.; Credi, A.; Hamers, C.; Mattersteig, G.; Montalti, M.; Shipway, A. N.; Spencer, N.; Stoddart, J. F.; Tolley, M. S.; Venturi, M.; White, A. J. P.; Williams, D. J. *Angew. Chem., Int. Ed.* **1998**, 37, 333–337. (c) Balzani, V.; Credi, A.; Mattersteig, G.; Matthews, O. A.; Raymo, F. M.; Stoddart, J. F.; Venturi, M.; White, A. J. P.; Williams, D. J. *J. Org. Chem.* **2000**, 65, 1924–1936.(8) Collier, C. P.; Mattersteig, G.; Wong, E. W.; Luo, Y.; Beverly, K.; Sampaio, J.; Raymo, F. M.; Stoddart, J. F.; Heath, J. R. *Science* **2000**, 289, 1172–1175.(9) Miljanic, O. S.; Stoddart, J. F. *Proc. Nat. Acad. Sci. U.S.A.* **2007**, 104, 12966–12970.

Scheme 1. Assembly of Donor–Acceptor [2]Catenanes from Crown Ethers, Pyridyl Ligands, and Pd(II) Metal Ions



acceptor [2]catenanes were obtained from two isolated ring components through iodide-catalyzed dynamic covalent chemistry (DCC).¹⁰ Besides the DCC approach, metal–ligand coordination also proves¹¹ to be an efficient synthetic means for thermodynamically controlled self-assembly of catenanes.¹² Examples are rare, however, where metal–ligand coordination operates cooperatively with donor–acceptor templation to give donor–acceptor [2]catenanes.^{13,14} Here we report (Scheme 1) the highly efficient syntheses of donor–acceptor [2]catenanes through a reversible self-assembly approach by utilizing the versatile Pd(II)–pyridine coordination.¹⁵ Ideally, donor–acceptor interactions between the crown ether and the bipyridinium **1**²⁺ would lead to the formation of an inclusion complex, while the involvement

of two [Pd(en)(II)]²⁺ (en = ethylenediamine) **2**²⁺ and one neutral bipyridine **3** would close the macrocycle to give a rectangular cavity that is optimal for alternative arrangement of the π -systems.




Figure 1. Partial ¹H NMR (500 MHz, 298 K) spectra of (a) a mixture of **1**·2PF₆, **2**, and **3** (1:2:1) in CD₃CN/D₂O (9:1), (b) the [2]catenane **7**·6PF₆ in CD₃CN, and (c) the DNP-containing crown ether **4** in CD₃CN.

A mixture of **1**·2PF₆, Pd(en)(NO₃)₂ (**2**), **3**, and dioxynaphthalene (DNP) containing crown ether **4** in the ratio of 1:2:1:1 was dissolved in CD₃CN/D₂O (9:1) after sonication. ¹H NMR spectroscopy of the resulting purple solution revealed that (Figure 1b) the [2]catenane **7**⁶⁺ was formed quantitatively as the sole product, which could be isolated in 90% yield after exchanging the anions to PF₆[−]. The pyridyl protons of the coordination macrocyclic component resonated as 12 well-resolved peaks in the aromatic region. The breakdown of symmetry resulted from the local C₂ symmetry imposed by the nearby DNP ring systems. In contrast to the free crown ether **4** (Figure 1c), the two DNP ring systems in **7**⁶⁺ were now resonating at higher field as two sets of doublet–triplet–doublets, indicating a slow site exchange for the “inside” and “outside” aromatic rings on the ¹H NMR time scale. Characteristically, the H_{4/8} of the inside DNP ring system appeared as a doublet at δ = 2.67 ppm as a result of a shielding effect from the nearby pyridine ring. For comparison, the assembly was also tested in the absence of the crown ether **4** under otherwise identical conditions. The ¹H NMR spectrum (Figure 1a) indicated only a complex mixture of oligomeric species. In addition, it appeared that neither CD₃CN nor D₂O was a suitable solvent by itself; rather a combination of both worked effectively. The efficient catenation appeared to be a result of an interplay of donor–

(10) (a) Rowan, S. J.; Cantrill, S. J.; Cousins, G. R. L.; Sanders, J. K. M.; Stoddart, J. F. *Angew. Chem., Int. Ed.* **2001**, *41*, 898–952. (b) Corbett, P. T.; Leclaire, J.; Vial, L.; West, K. R.; Wietor, J.-L.; Sanders, J. K. M.; Otto, S. *Chem. Rev.* **2006**, *106*, 3652–3711.

(11) (a) Fujita, M.; Ibukuro, F.; Hagihara, H.; Ogura, K. *Nature* **1994**, *367*, 720–723. (b) Fujita, M.; Aoyagi, M.; Ibukuro, F.; Ogura, K.; Yamaguchi, K. *J. Am. Chem. Soc.* **1998**, *120*, 611–612. (c) Dietrich-Buchecker, C. O.; Colasson, B.; Fujita, M.; Hori, A.; Geum, N.; Sakamoto, S.; Yamaguchi, K.; Sauvage, J.-P. *J. Am. Chem. Soc.* **2003**, *125*, 5717–5725. (d) Hori, A.; Sawada, T.; Yamashita, K.; Fujita, M. *Angew. Chem., Int. Ed.* **2005**, *44*, 4896–4899. (e) Yamashita, K.; Kawano, M.; Fujita, M. *J. Am. Chem. Soc.* **2007**, *129*, 1850–1851.


(12) (a) Kidd, T. J.; Leigh, D. A.; Wilson, A. J. *J. Am. Chem. Soc.* **1999**, *121*, 1599–1600. (b) Leigh, D. A.; Lusby, P. J.; Teat, S. J.; Wilson, A. J.; Wong, J. K. Y. *Angew. Chem., Int. Ed.* **2001**, *40*, 1538–1543. (c) Hogg, L.; Leigh, D. A.; Lusby, P. J.; Morelli, A.; Parsons, S.; Wong, J. K. Y. *Angew. Chem., Int. Ed.* **2004**, *43*, 1218–1221. (d) Lam, R. T. S.; Belenguer, A.; Roberts, S. L.; Naumann, C.; Jarroson, T.; Otto, S.; Sanders, J. K. M. *Science* **2005**, *308*, 667–669. (e) Blight, B. A.; Wisner, J. A.; Jennings, M. C. *Angew. Chem., Int. Ed.* **2007**, *46*, 2835–2838.

(13) For examples of donor–acceptor metallocatenanes, see: (a) Try, A. C.; Harding, M. M.; Hamilton, D. G.; Sanders, J. K. M. *Chem. Commun.* **1998**, 723–724. (b) Chas, M.; Blanco, V.; Peinador, C.; Quintela, J. M. *Org. Lett.* **2007**, *9*, 675–678.

(14) During the preparation of this manuscript, donor–acceptor [3]catenanes and related metallomacrocycles based on a similar concept were reported; see: (a) Blanco, V.; Chas, M.; Abella, D.; Peinador, C.; Quintela, J. M. *J. Am. Chem. Soc.* **2007**, *129*, 13978–13986. (b) Chas, M.; Abella, D.; Blanco, V.; Pia, E.; Blanco, G.; Fernandez, A.; Platas-Iglesias, C.; Peinador, C.; Quintela, J. M. *Chem. Eur. J.* **2007**, *13*, 8572–8582.

(15) (a) Fujita, M.; Tominaga, M.; Hori, A.; Therrien, B. *Acc. Chem. Res.* **2005**, *38*, 371–380. (b) Leininger, S.; Olenyuk, B.; Stang, P. J. *Chem. Rev.* **2000**, *100*, 853–908.

Scheme 2. Chemical Structures and Labeling of [2]Catenanes 8^{6+} and 9^{6+}



acceptor interactions and the reversible Pd(II)–pyridine coordination, as well as proper solvation.

When the hydroquinone (HQ)-containing crown ether **5** was used for the [2]catenane formation, 3 equiv of **5** were required to reequilibrate the oligomeric Pd(II)–pyridine complexes toward the assembly of [2]catenane 8^{6+} (Scheme 2). The ^1H NMR spectra (Figure 2b) of an equilibrated mixture of **1**·2PF₆, **2**, **3**, and **5** in the ratio of 1:2:1:3 revealed the presence of [2]catenane 8^{6+} as the only product. Pure [2]catenane 8^{6+} could be isolated as a PF₆[−] salt in 90% after anion exchange and removal of excess **5** by rinsing with CH₂Cl₂. Partial dissociation of **8**·6PF₆ was observed when redissolved in CD₃CN (2.4 mM). The ^1H NMR spectrum (Figure 2c) indicated not only the presence of **8**·6PF₆ as the




Figure 2. (a) Illustration of the ring-exchange process. ^1H NMR spectrum of 8^{6+} (b) before anion exchange in CD₃CN/D₂O (9:1), together with excess **5**, (c) after anion exchange (CD₃CN, 2.4 mM), and (d) after addition of crown ether **4** (1.4 equiv). The blue asterisks in (c) indicate the resonances from the free coordination macrocycle, and the purple dots in (d) indicate the DNP protons from excess crown ether **4**.

major specie but also an equal amount of free crown ether **5** and the free coordination metallocyclophane. An association constant was calculated to be $1.8 \times 10^4 \text{ M}^{-1}$ using a single-point method.¹⁶ In contrast, no dissociation was observed for the DNP-based [2]catenane 9^{6+} at 1.0 mM, attesting that the DNP ring system is a better donor than the HQ ring system in holding the π -accepting coordination macrocycle around itself. Such differences in the donor–acceptor interactions prompted us to test a ring-exchange experiment reminiscent of a magician’s magic ring trick (Figure 2a). When DNP-containing crown ether **4** (1.4 equiv) was added to the above solution of 8^{6+} , the color of the solution changed immediately from red to purple. The resulting ^1H NMR spectrum revealed (Figure 2d) the disappearance of 8^{6+} , together with the emergence of 7^{6+} and the released crown ether **5**. The ring-exchange experiment clearly demonstrated the dynamic yet selective nature of the five-component self-assembly process.




Figure 3. Partial ^1H NMR spectrum of 9^{6+} at (a) 298 K and (b) 228 K.

Furthermore, crown ether **6**^{7b} bearing both DNP and tetrathiafulvalene (TTF) π -donating ring systems was tested in the self-assembly process. A green product **9**·6PF₆ (Scheme 2) could be isolated in 92% yield after counterion exchange. Noticeably, translational selectivity^{7b,17} was well maintained where the TTF unit sits inside the cavity of the coordination macrocycle. ^1H NMR spectroscopy (Figure 3a) at 298 K indicated one single specie in the solution. The absence of DNP protons between 2–3 ppm clearly indicated the “alongside” location of the DNP ring system with respect to the coordination macrocycle. When temperature was lowered to 228 K, both TTF and DNP resonances remained sharp (Figure 3b), while splitting and broadening of the resonances of the pyridyl protons in **1**·2PF₆ and the 4,4′-dipyridyl protons were observed as a result of restrained

(16) Fyfe, M. C. T.; Stoddart, J. F. *Adv. Supramol. Chem.* **1999**, 5, 1–53.

(17) Ashton, P. R.; Perez-Garcia, L.; Stoddart, J. F.; White, A. J. P.; Williams, D. J. *Angew. Chem., Int. Ed. Engl.* **1995**, 34, 571–574.

rotation at low temperature. No resonances corresponding to two translational isomers were observed, suggesting the absence of site exchange, and indeed the translational selectivity was maintained throughout. It is worth noting that out of the usual *cis/trans* TTF positional isomers, only one was observed, which was probably the *trans* one.^{7b-c,18}




Figure 4. CSI-TOFMS spectrum of **9·6PF₆**. (Inset) Isotopic distribution of the **[9·6PF₆-2PF₆]²⁺** ion.

The formation of the [2]catenanes was further verified¹⁹ by elemental analysis, cold-spray ionization mass spectrometry (CSI-MS),²⁰ and UV-vis spectra. The significant results agreed well with their empirical structures. For example, the CSI-TOFMS spectrum of **9·6PF₆** indicated (Figure 4) ions at *m/z* 1076 and 669, corresponding to the doubly and triply charged **[M - 2PF₆]²⁺** and **[M - 3PF₆]³⁺** ions, respectively.¹⁹ The inset in Figure 4 revealed the isotopic distribution for the fragment **[M - 2PF₆]²⁺**, which was consistent with the theoretical distribution. The UV-vis spectra of the [2]-catenanes **7–9·6PF₆** indicated (Figure 5) charge-transfer absorption bands centered at 535, 463, and 821 nm, corresponding respectively to the purple, red, and green color of the individual donor–acceptor [2]catenanes.

In summary, highly efficient synthesis of [2]catenanes was developed using a combination of donor–acceptor templation

(18) However, the sample stored in the dry solid form for more than three months was found to partially isomerize to give a mixture of *cis* and *trans* isomers. See Supporting Information for a comparison of the ¹H NMR spectra before and after storage. For a similar observation of TTF *cis/trans* isomerization in the solid state, see: Liu, Y.; Flood, A. H.; Moskowicz, R. M.; Stoddart, J. F. *Chem. Eur. J.* **2005**, *11*, 369–385.

(19) See Supporting Information for details of characterization data.

(20) (a) Yamaguchi, K. *J. Mass Spectrom.* **2003**, *38*, 473–490. (b) Yamanoi, Y.; Sakamoto, Y.; Kusakawa, T.; Fujita, M.; Sakamoto, S.; Yamaguchi, K. *J. Am. Chem. Soc.* **2001**, *123*, 980–981.




Figure 5. UV-vis spectra of **7·6PF₆** (2.0×10^{-4} M), **8·6PF₆** (2.8×10^{-4} M), and **9·6PF₆** (4.4×10^{-5} M) in MeCN, showing the charge-transfer absorption maximum at $\lambda = 535$ nm ($\epsilon = 840$ M⁻¹ dm⁻¹), 463 nm ($\epsilon = 560$ M⁻¹ dm⁻¹), and 821 nm ($\epsilon = 1700$ M⁻¹ dm⁻¹), respectively.

and metal–ligand coordination. The [2]catenanes were easily obtained within minutes through a five-component assembly process from three crown ethers. The [2]catenane formation was fully characterized by ¹H NMR and UV-vis spectroscopies and cold-spray ionization mass spectrometry. The dynamic nature of the [2]catenanes was further demonstrated in a ring-exchange experiment. In particular, translational selectivity was maintained when a crown ether with two different donor units was employed. The current approach is remarkable in the perspective of bistable molecular devices as it could potentially provide a fast, mild, convenient, and high-yielding means to install bistability in molecular species. Such an assembly process could be utilized in high-throughput screenings of matching donor–acceptor pairs for the design of sophisticated multistable molecular switches.

Acknowledgment. This work was supported by the Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under contract No. DE-AC02-05 CH11231. We thank financial support from the New Analysis Technology and Method Research Program of the Chinese Ministry of Science and Technology (Grant No. 2005JG200050).

Supporting Information Available: Experimental details for the preparation of **1·2PF₆**, **7–9·6PF₆** and spectroscopic data thereof. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL702830P