

# Synthesis of Naturally Occurring 5-Allyl-2-aryl-7-methoxybenzofuran and 2-Aryl-5-(3-hydroxypropyl)-7-methoxybenzofurans†

Raghao S. Mali\* and Archana Patience Massey

Garware Research Centre, Department of Chemistry, University of Pune, Pune-7, India


*J. Chem. Research (S)*,  
1998, 230–231  
*J. Chem. Research (M)*,  
1998, 1109–1120

A convenient and general procedure is described for the synthesis of 5-allyl-2-aryl-7-methoxybenzofurans (**8a–e**) from 2-allyloxy-3-methoxybenzaldehyde (**3**). The compounds **8a** and **8b** on hydroboration followed by oxidation provide the naturally occurring benzofurans (**1a** and **1b**).

A few compounds, containing the 2-arylbenzofuran nucleus (**1a**, **1b**, **2** and **8a**), have been isolated from plants.<sup>2–4</sup> Egonol (**1a**) and homoegonol (**1b**) were isolated<sup>2,3</sup> from the seeds of *Styrax japonicum* and *Styrax officinalis* L. respectively, while (±)-machicendiol **2** was isolated<sup>4</sup> from the leaf extracts of *Machilus glaucescens*, which are used for the treatment of asthma, rheumatism and ulcers. These compounds are also reported for their cytostatic activity against human leukemic HL-60 cells.<sup>5</sup>

In view of the natural occurrence and valuable biological activities associated with **1a**, **1b** and **2**, several methods have been developed for their synthesis.<sup>7–9,11</sup> Four approaches are known<sup>7–9,11</sup> for egonol **1a** and one<sup>8,9</sup> each for homoegonol **1b** and neolignan **8a**.

We report herein a convenient, general approach (Scheme 1) for the synthesis of egonol **1a**, homoegonol **1b** and neolignan **8a**, starting from 2-allyloxy-3-methoxybenzaldehyde<sup>13</sup> **3**. When a solution of aldehyde **3** in *N,N*-dimethylaniline was irradiated in a microwave oven for 10 min, 5-allyl-2-hydroxy-3-methoxybenzaldehyde **4** was obtained in 65% yield along with minor amount (15%) of 2-allyl-6-methoxyphenol. The aldehyde **4** on reduction with sodium tetrahydroborate in ethyl acetate solution, gave the benzyl alcohol **5** as a thick liquid in 76% yield. Reaction of **5** with thionyl chloride in methylene chloride, followed by treatment with triphenylphosphine in benzene solution yielded the phosphonium salt **6**, which on reaction with benzoyl chlorides in toluene solution, in the presence of



Scheme 1

\*To receive any correspondence (e-mail: rsmali@chem.unipune.ernet.in).

†Dedicated to Professor Dr Dieter Seebach on the occasion of his 60th birthday.

triethylamine, furnished the 2-arylbenzofurans (**8a–e**) in 60–89% yields via the intermediacy of phosphonium salts **7a–e**.

The present work thus describes the total synthesis of neolignan **8a** and the related compounds **8b–e**. Conversion of 2-arylbenzofuran **8a** into egonol **1a** has already been reported<sup>8</sup> in the literature using the hydroboration approach. The compound **8b** on similar reaction provided homoeegonol **1b**, mp 121 °C (lit.,<sup>3</sup> mp 120–122 °C) in 70% yield, which is another natural product.

We thank Professor N. S. Narasimhan for critical reading of the manuscript and valuable discussions. A.P.M. thanks CSIR, New Delhi for the award of Senior Research Fellowship.

Techniques used: IR, <sup>1</sup>H NMR, elemental analyses, TLC and column chromatography

References: 15

Received, 17th October 1997; Accepted, 14th January 1998  
Paper E/7/07498K

#### References cited in this synopsis

- 2 H. Okada, *J. Pharm. Soc. Jpn.*, 1915, 657.
- 3 R. Segal, I. M. Goldzweig, S. Sokoloff and D. V. Zaitschek, *J. Chem. Soc. C*, 1967, 2402.
- 4 B. Talaparta, T. Ray and S. Talaparta, *Indian J. Chem., Sect. B*, 1976, **14**, 613.
- 5 T. Hirano, M. Goto and K. Oka, *Life Sci.*, 1994, **55**, 1061.
- 7 S. Kawai, T. Nakamura and N. Sugiyama, *Ber. Dtsch. Chem. Ges.*, 1939, **72**, 1146.
- 8 E. Ritchie and W. C. Taylor, *Aust. J. Chem.*, 1969, **22**, 1329.
- 9 F. G. Schreiber and R. Stevenson, *J. Chem. Soc., Perkin Trans. I*, 1976, 1514.
- 11 Y. Aoyagi, T. Mizusaki, A. Hatori, T. Asakura, T. Aihara, S. Inaba, K. Hayatsu and A. Ohta, *Heterocycles*, 1995, **41**, 1077.
- 13 R. S. Mali and A. P. Massey, *Indian J. Chem., Sect. B*, 1995, **34**, 686.