Org Lett. Author manuscript; available in PMC 2011 March 19.

Published in final edited form as:

Org Lett. 2010 March 19; 12(6): 1256–1259. doi:10.1021/ol1001126.

## Diastereoselective Allylstannane Additions to (S)-5,6-Dihydro-2 H-5-phenyloxazin-2-one. A Concise Synthesis of (S)- $\beta$ -Methylisoleucine

Julie A. Pigza<sup>§,‡</sup> and Tadeusz F. Molinski<sup>\*,§,†</sup>

Department of Chemistry and Biochemistry, and Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California San Diego, 9500 Gilman Drive MC0358, La Jolla, CA, 92093

### **Abstract**

The addition of allyl stannanes to (S)-4,5-dihydro-5-phenyl-2 H-oxazinone (3) was achieved under Brønsted acid catalysistogive 2-allylmorpholinones with high diastereoselectivity (up to dr14.2:1). The product of dimethylallyltributylstannane addition to 3 was converted to L- $\beta$ -methylisoleucine, an  $\alpha$ -amino acid residue found in the complex, biologically-active marine-derived peptides polytheonamides A and B and polydiscamides A–C.

Marine peptides often contain highly modified amino acids, including chlorinated amino acids,  $^1$   $\beta$ -amino acids $^2$  and highly-substituted  $\beta$ ,  $\beta$ -dimethyl- $\alpha$ -amino acids.  $^3$  L-tert-Leucine (1a) and L-tert-amylglycine ( $\beta$ -methylisoleucine, 1b) occur in the 48-mers polytheonamides A and B; two highly cytotoxic  $\beta$ -helix, membrane-pore forming peptides from the marine sponge, Theonella swinhoei.  $^4$  The peptides polydiscamide A from Discodermia $^{5a}$  (2, Figure 1), and the polydiscamides B–C from the sponge Ircinia $^{5b}$  which are the first non-endogenous inhibitors of sensory neuron-specific G-protein coupled receptors (SNSRs), also contain 1a and 1b.  $^6$  The amino acid  $\gamma$ -hydroxy-tert-leucine (L-pantonine, 1c) was proposed as an intermediate in the biosynthesis of pantoic acid.  $^7$  With an eye to the synthesis of highly biologically active, cyclic peptides, we sought to fulfill a need for a general synthesis of highly-branched  $\alpha$ -amino acids.

Racemic syntheses of *tert*-alkyl- $\alpha$ -amino acids have been reported. Asymmetric preparation of L-**1a** and L-**1b** was achieved by tandem-enzyme coupled reductive amination of the

Supporting Information Available. Experimental procedures, full spectroscopic data and <sup>1</sup>H, <sup>13</sup>C NMR spectra for all new compounds. This material is available free of charge via the Internet at http://pubs.acs.org.

tmolinski@ucsd.edu.

Department of Chemistry and Biochemistry

<sup>†</sup>Skaggs School of Pharmacy and Pharmaceutical Sciences

<sup>‡</sup>Present address, Queensborough Community College 222-05, 56th Avenue, Bayside, NY 11364

corresponding  $\alpha$ -ketoacids. However, this biotechnology is not amenable to preparation of the D-antipode due to the enantiospecificity of the enzymes. In this report, we demonstrate a concise *asymmetric* synthesis of L-**1b** which exploits an efficient allylstannane addition to highly electrophilic 2*H*-oxazinone, **3**, a chiral glycine equivalent, and is amenable to preparation of D-**1** or other highly branched amino acids.

SeO<sub>2</sub>-promoted oxidative rearrangement of 2-substituted oxazolines ii (Figure 2) to 5,6-dihydro-2H-1,4-oxazin-2-ones (e.g. 3, hereafter, referred to as 'oxazinones'),  $^{10}$  followed by hydrogenation-hydrogenolysis,  $^{11}$  allows convenient access to a wide variety of  $\alpha$ -amino acids, iii.  $^{12}$  Thus, the conversion of carboxylic acid i to iii, constitutes a highly useful transformation; formal preparation of amino acids by oxidative insertion of NH<sub>2</sub> to the  $\alpha$ -carbon of a carboxylic acid  $^{12}$  of either configuration by choice of an appropriate chiral auxillary,  $^{13}$  R- or S-phenylglycinol, obtained readily from the corresponding commercially available phenylglycines.

The highly-electrophilic 2-*unsubstituted* oxazinone  $3^{14}$  is particularly attractive as a chiral glycine equivalent that can add a variety of carbon-centered nucleophiles at the C=N bond to give morpholinone amino acid precursors, however, the diastereoselectivity and yield of these additions can be variable. For example, addition of MeMgBr or *t*-BuMgBr to 3 in the presence of BF<sub>3</sub>•Et<sub>2</sub>O gave one detectable diastereomer in poor yield (34% and 33%, respectively). We now describe synthesis of 3-allylmorpholinones by highly diastereoselective allyl stannane additions to 3 promoted by Brønsted acid to give 4, and subsequent conversion to  $\beta$ -methylisoleucine (1b).

Oxazoline (ii, R = H, Figure 2) was prepared in two steps from S-phenylglycinol<sup>16,17</sup> in 80%. <sup>10,18</sup> The original procedure for SeO<sub>2</sub>-promoted rearrangement of the oxazoline to (S)-oxazinone  ${\bf 3}^{10}$  required refluxing 1,4-dioxane for 2 h. Instead, short-exposure of the substrate (~1 mmole scale) to SeO<sub>2</sub> in a microwave reactor (10 min, 300 W, 110 °C), adapted from Snider's procedure for SeO<sub>2</sub>-promoted allylic oxidations, <sup>19</sup> improved the yield of  ${\bf 3}$  (74%) and reduced byproducts.

As reported earlier, <sup>20</sup> BF<sub>3</sub>•Et<sub>2</sub>O-promoted additions of allyltrimethylsilane, methallyltrimethylsilane and dimethylallyltrimethylsilane to (S)-3 (entries 1–3, Table 1) gave only modest diastereoselectivity and/or low yields of 4. The diastereoselectivity of BF<sub>3</sub>•Et<sub>2</sub>Opromoted allyltrimethylsilane addition to 3 was 8:1 to give 4a (entry 1, 60% yield), but with the more hindered nucleophiles, dimethylallyltrimethylsilane and methallyltrimethylsilane the diastereoselectivity diminished to 5:1 (entry 2, 25% yield) and 2:1 (entry 3, 60%), respectively. <sup>21</sup> Addition of allyltributylstannane in the presence of either BF<sub>3</sub>•Et<sub>2</sub>O or TFA also gave **4a** with low diastereoselectivities (dr 2.6:1 and 3.2:1, respectively, entries 5, 6),<sup>22</sup> as did use of methallyltributylstannane23 to give 4c (37%, dr 1.4:1, entry 11). Gratifyingly, addition of dimethylallyltributylstannane to 3 in the presence of TFA (-78 °C, CH<sub>2</sub>Cl<sub>2</sub>, 1 h, entry 9) gave a dramatic increase in both the yield and diastereoselectivity for 4b (dr 14.8:1, 80% yield). The major diastereomer (3S,5S)-4b was purified from the diastereomeric mixture by selective recrystallization from Et<sub>2</sub>O/pentane (dr >20:1 Scheme 1).<sup>24</sup> A similar outcome was observed when the addition was carried out in acetonitrile at -30 °C (dr 12.5:1, 62% yield, entry 10), but diastereoselectivity eroded when the reaction was conducted with TFA in CH<sub>2</sub>Cl<sub>2</sub> at higher temperatures (-20 °C, dr 6.2:1, 68% yield, entry 8).

The surprising difference in the outcome of additions of the two dimethylallyltrialkyl silane (entry 2) and stannanes (entries 9, 10) to oxazinone 3 deserves some comment. The role of the Brønsted or Lewis acid is activation of the imine to an iminium ion (Figure 3). For electronic reasons, both dimethylallyltrialkyl-silane and the corresponding stannane add at their more substituted sp<sup>2</sup> olefinic carbon. Consequently, the higher diastereoselectivity in formation of

(3*S*,5*S*)-**4b** from the stannane in the presence of TFA may be due to relaxed steric congestion with the ion pair **3**•H<sup>+</sup> TFA<sup>-</sup> compared to the corresponding bulkier **3**•BF<sub>3</sub> complex. Under these conditions, tighter association of the vinyl bond to the C=N bond is allowed, serving to amplify differences in energies between top and bottom facial additions (Figure 3(a) and (b), respectively) in the respective transition states and favoring approach of the nucleophilic allyl equivalent from the side opposite the Ph group.

The synthesis of (–)-β-methylisoleucine (**1b**) was accomplished by conversion of the dimethylallylated morpholinone (3*S*,5*S*)-**4b** as follows (Scheme 1). Hydrogenation of (3*S*, 5*S*)-**4b** under acidic conditions (6 atm, Pd(OH)<sub>2</sub>, MeOH, 1M HCl) followed by exhaustive acid hydrolysis (refluxing HCl) provided the optically pure amino acid salt (L-**1b**•HCl, Scheme 1) which was converted to the free amino acid by ion-exchange chromatography (elution with 2M NH<sub>4</sub>OH) to neutral L-**1b** (96% yield, two steps).

Additions of dimethylallyl anion equivalent to **3** should find wider applicability in synthesis of other  $\beta$ , $\beta$ -dimethyl-substituted amino acids. The versatile vinyl 'handle' in intermediate (3S,5S)-**4b** may find uses for preparation of amino acid derivatives related to **1b**. For example, selective hydrogenation of the terminal vinyl group of (3S,5S)-**4b** (Scheme 3, H<sub>2</sub>, Pd-C, 96%) to give **5** models a potential route to preparation of specifically-labeled [<sup>2</sup>H]- and [<sup>3</sup>H]- (S)-**1b**. Oxidative or reductive modifications of the terminal vinyl group or olefin metathesis should provide access to other highly modified natural and non-natural amino acids including the  $\gamma$ -methysulfinyl-*tert*-leucine residue found in polytheonamide A.<sup>4</sup>

The optical rotation of L-**1b** ( $[\alpha]_D^{22}$  +12.7 (c 0.48, 1M HCl), lit. Error! Bookmark not defined. +9.9 (c 1M, 1M HCl) and  $^1$ H and  $^{13}$ C NMR data  $^{6,25}$  matched the literature values. Thus, L-**1b** was obtained in three steps from the chiral glycine equivalent (S)-**3** in an overall yield of 58%.

In summary, the versatility of chiral glycine synthon **3** for  $\alpha$ -amino acid synthesis has been extended to highly diastereoselective additions dimethylallylstannane and a concise conversion of the product (3*S*,5*S*)-**4b** to the highly branched amino acid (–)- $\beta$ -methylisoleucine (L-**1b**). The configurations of amino acids derived through allylstannane additions to **3** complement those from hydrogenation of oxazinones <sup>10,11</sup> (Figure 2, 3 R = alkyl). Consequently, antipodal amino acids can be obtained in high yield from oxazinones derived from a common phenylglycinol chiral auxilllary.

## **Supplementary Material**

Refer to Web version on PubMed Central for supplementary material.

### Acknowledgments

We thank Y. Su (UCSD) and R. New (UC Riverside) for HRMS measurements. The 500 MHz NMR spectrometers were purchased with a grant from the NSF (CRIF, CHE0741968). This work was supported by the NIH (AI-039987).

### References


- 1. Jiménez JI, Vansach T, Yoshida WY, Sakamoto B, Pörzgen P, Horgen FD. J Nat Prod 2009;72:1573. [PubMed: 19739598]
- (a) Zabriskie TM, Klocke JA, Ireland CM, Marcus AH, Molinski TF, Faulkner DJ, Xu C, Clardy J. J Am Chem Soc 1986;108:3123.
   (b) Crews P, Manes LV, Boehler M. Tetrahedron Lett 1986;27:2797.
- 3. Coleman JE, Desilva ED, Kong FM, Andersen RJ, Allen TM. Tetrahedron 1995;51:10653.
- 4. Hamada T, Matsunaga S, Yano G, Fusetani N. J Am Chem Soc 2005;127:110. [PubMed: 15631460]
- (a) Gulavita NK, Gunasekera SP, Pomponi SA, Robinson EV. J Org Chem 1992;57:1767.
   (b) Feng Y, Carroll AR, Pass DM, Archbold JK, Avery VM, Quinn RJ. J Nat Prod 2008;71:8. [PubMed: 18163586]

 Curiously, the first identification of naturally occuring 1b was from an extraterrestrial source; the Murchison meteorite, a carbonaceous chondrite. Cronin JR, Pizzarello S. Geochim Cosmochim Acta 1986;50:2419. [PubMed: 11542028]

- 7. Ackermann WW, Kirby H. J Biol Chem 1948:483. [PubMed: 18873328]
- 8. (a) Barker J, Cook SL, Lasterra-Sanches E, Thomas SE. J Chem Soc Chem Commun 1992:830. (b) Bretschneider T, Miltz W, Munster P, Steglich W. Tetrahedron 1988;44:5403.
- 9. Krix G, Bommarius AS, Drauz K, Kottenhahn M, Schwarm M, Kula MR. J Biotech 1997;53:29.
- 10. (a) Shafer CM, Molinski TF. J Org Chem 1996;61:2044. (b) Shafer CM, Morse DI, Molinski TF. Tetrahedron 1996;52:14475.
- 11. Cox GG, Harwood LM. Tetrahedron: Asymmetry 1994;5:1669.
- 12. (a) Pigza JA, Quach T, Molinski TF. J Org Chem 2009;74:5510. [PubMed: 19537688] (b) Liu C, Masuno MN, MacMillan JB, Molinski TF. Angew Chem Int Ed 2004;43:5951.
- 13. (a) Harwood LM, Macro J, Watkin D, Williams CE, Wong LF. Tetrahedron: Asymmetry 1992;3:1127. (b) Ager D, Cooper N, Cox GG, Garro-Helion F, Harwood LM. Tetrahedron: Asymmetry 1996;7:2563.
- 14. The 5,6-diphenylmorpholinone **3a** has been used both as a nucleophilic glycine equivalent and precursor for in situ prepartion of the electrophilic **3b**. (a) Williams RM. Aldrichimica Acta 1992;25:11. (b) Williams RM, Hendrix JA. Chem Rev 1992;92:889. and references, therein. (c) Williams RM, Sinclair PJ, Zhai D, Chen D. J Am Chem Soc 1988;110:1547.**3b** and **3c** have been converted to arylglycines by Mannich reaction with arenes. (d) Tohma S, Rikimaru K, Endo A, Shimamoto K, Kan T, Fukuyama T. Synthesis 2004:909. (e) Chen YJ, Lei F, Liu L, Wang D. Tetrahedron 2003;59:7609.**3b-c** were made in 5–6 steps from phenylglycine, and **3a** from benzoin
- Harwood LM, Tyler SNG, Anslow AS, MacGilp ID, Drew MGB. Tetrahedron Asymmetry 1997;8:4007.
- 16. McKennon MJ, Meyers AI, Drauz K, Schraum M. J Org Chem 1993;58:3568.
- 17. Hsiao Y, Hegedus LS. J Org Chem 1997;62:3586.
- Gorunova ON, Keuseman KJ, Goebel BM, Kataeva NA, Churakov AV, Kuz'mina LG, Dunina VV, Smoliakova IP. J Organomet Chem 2004;689:2382.
- 19. Zou Y, Chen CH, Taylor CD, Foxman BM, Snider BB. Org Lett 2007;9:1825. [PubMed: 17407302]
- 20. Shafer CM, Pigza JA, Molinski TF. Lett Org Chem 2009;6:224.
- 21. Diastereomers of **4** were assigned by nOe measurements. Ref. <sup>10, 12</sup> and conversion of **4b** to (S)-**1b**.
- 22. Side-product formation lead to diminished yields of the desired 4a.
- 23. Naruta Y, Nishigaichi Y, Maruyama K. Chem Lett 1986:1857.
- 24. Removal of Sn-containing byproducts was conveniently effected by chromatography over SiO<sub>2</sub> impregnated with KF. Harrowven DC, Guy IL. Chem Commun 2004:1968.
- 25. Barker J, Cook SL. Inorg Chim Acta 1994;220:137.

Figure 1. Structures of L-*tert*-leucine (1a), L- $\beta$ -methylisoleucine (*tert*-amylglycine, 1b), pantonine (1c) and polydiscamide A (2), a marine-derived peptide containing  $\beta$ , $\beta$ -dimethyl-substituted  $\alpha$ -amino acids.

**Figure 2.** Oxazoline-oxazinone oxidative rearrangement.



**Figure 3.** Possible transition states for reaction of 3•H<sup>+</sup> with dimethylallyl-silane or -stannane

H<sub>2</sub> (1 atm),  
Pd-C, toluene, r.t.  
96%

5 Ph

i. H<sub>2</sub> (6 atm)
Pd(OH)<sub>2</sub>,
1M HCl, MeOH, r.t.
ii. HCl (conc), 100 °C
iii. Dowex 50X2
ion exch.
3 (3S,5S)-4b 96%, 3 steps (S)-1b

80%, dr 14.8:1 recryst.
76%, dr >20:1 Et<sub>2</sub>O/pentane (
$$\alpha$$
 0.48, 1M HCl)

Scheme 1. Conversion of morpholinone (3*S*,5*S*)-4b to L- $\beta$ -methylisoleucine 1b.

Page 9 Pigza and Molinski

Table 1


promoted AllyIsilane and AllyIstannane Additions to Oxazinone (S)-3.

	ref.	20	20	20
		8:1	5:1	2:1
	solvent temp, time yield $(\%)^d$ $d_rb$	73c	25c,d,e	<sub>2</sub> 09
	temp, time	$BF_3 \bullet Et_2O  CH_2Cl_2  -78 \stackrel{\circ}{\circ}C, 0.5  73^{\mathcal{C}}$	BF <sub>3</sub> •Et <sub>2</sub> O CH <sub>2</sub> Cl <sub>2</sub> -78 °C, 0.5 25c.d.e	BF₃•Et₂O CH₂Cl₂ −78 °C, 1 h
, , ,	solvent	$\mathrm{CH}_2\mathrm{Cl}_2$	CH <sub>2</sub> Cl <sub>2</sub>	$\mathrm{CH}_2\mathrm{Cl}_2$
O = N	Brønsted or Lewis acid	$\mathrm{BF_3}ullet \mathrm{E}_{t_2}\mathrm{O}$	$\mathrm{BF_3}ullet \mathrm{Et_2O}$	BF₃•Et₂O
or	product	4a	4b	4
OHNH dt				
o HHN HH PAR				
allyl equivalent conditions	allyl equivalent	SiMe <sub>3</sub>	SiMe <sub>3</sub>	SiMe <sub>3</sub>
O = N - H (8)				

Org Lett. Author manuscript; available in PMC 2011 March 19.

Page 10

Pigza and Molinski		
	ref.	<b>∞</b>
	$dr^{b}$	2.6:1
	p(°)	




Org Lett. Author manuscript; available in PMC 2011

<i>&gt;</i>	— ≟ Σ 3	o-\£				
product	Brønsted or Lewis acid	solvent	temp, time	yield (%) <sup><i>a</i></sup>	$dr^{b}$	ref.
48	TFA	CH <sub>2</sub> Cl <sub>2</sub>	–78°C, 1 h	<sub>2</sub> 09	5:1	00
4 <del>p</del>	$\mathrm{BF_3}ullet \mathrm{Et_2O}$	$\mathrm{CH}_2\mathrm{Cl}_2$	–78 °C, 0.5 h	64(75) <sup>f</sup>	7.1:1	00
4b	TFA	$\mathrm{CH}_2\mathrm{Cl}_2$	$CH_2Cl_2$ -20 °C, 1 h 68 $f$	£89	6.2:1	00

# .Sn"Bu


allyl equivalent


\ \ \



Org Lett. Author manust ipt; vailable in PMC 2011 March 19.

o= NH 5
o NH d4
o⇒NH & &
allyl equivalent conditions
o=\~£ §





. 1	1	ı
ref.	∞	
$dr^{b}$	1.4:1	
solvent temp, time yield $(\%)^d$ $dr^b$	37¢.e	
temp, time	CH <sub>2</sub> Cl <sub>2</sub> -78 °C, 1 h 37c,e	
	CH <sub>2</sub> Cl <sub>2</sub>	
Brønsted or Lewis acid	TFA	
Brønsted or Lewis product acid	4	
oso allyl equivalent	Lett. Author manuscript; available in MG2011 M	arc

yield after SiO2 colum<u>n</u>chromatography (yields in parentheses are based on recovered starting material). ்

<sup>1</sup>H NMR integration.

mers not separated.

ized.

on to an unidentified by product (see Ref. 20 and footnote 23).

stereomer separated by crystallization.

e = this work.