

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8124135>

# Synthesis of Zwitterionic 4-Hydroxycoumarin Derivatives through a Unique Reaction of 4-Hydroxycoumarins with p - Benzoquinone and Pyridine

ARTICLE in ORGANIC LETTERS · JANUARY 2005

Impact Factor: 6.36 · DOI: 10.1021/ol048109p · Source: PubMed

---

CITATIONS

21

---

READS

32

## 8 AUTHORS, INCLUDING:



Zhishu Huang

Sun Yat-Sen University

175 PUBLICATIONS 2,350 CITATIONS

SEE PROFILE



Xian-Zhang Bu

Sun Yat-Sen University

82 PUBLICATIONS 983 CITATIONS

SEE PROFILE



Lin Ma

Sun Yat-Sen University

105 PUBLICATIONS 1,364 CITATIONS

SEE PROFILE



Albert S. C. Chan

Hong Kong Baptist University

404 PUBLICATIONS 9,483 CITATIONS

SEE PROFILE

# Synthesis of Zwitterionic 4-Hydroxycoumarin Derivatives through a Unique Reaction of 4-Hydroxycoumarins with *p*-Benzoquinone and Pyridine


Sheng-Ling Zhang,<sup>†,‡,§</sup> Zhi-Shu Huang,<sup>\*,†</sup> Lin-Kun An,<sup>†</sup> Xian-Zhang Bu,<sup>†</sup> Lin Ma,<sup>‡</sup> Yue-Ming Li,<sup>||</sup> Albert S. C. Chan,<sup>||</sup> and Lian-Quan Gu<sup>\*,†,‡</sup>

School of Pharmaceutical Science, Sun Yat-Sen University, Guangzhou, 510275 People's Republic of China, School of Chemistry and Chemical Engineering, Sun Yat-Sen University, Guangzhou, 510275 People's Republic of China, and Department of Applied Biology & Chemical Technology, The Hong Kong Polytechnic University, Kowloon, Hong Kong

huangzshu@hotmail.com; cedc42@zsulink.zsu.edu.cn

Received September 17, 2004

## ABSTRACT



A unique reaction of 4-hydroxycoumarins with *p*-benzoquinone and pyridine was found, and through the reaction, six zwitterionic 4-hydroxycoumarin derivatives were synthesized. The structures of these compounds were determined by IR, MS(ESI), <sup>1</sup>H NMR, <sup>13</sup>C NMR, and single-crystal X-ray diffraction studies.

4-Hydroxycoumarin is an important component of many synthetic and natural products<sup>1</sup> with wide ranging biological activities that include anticoagulant and HIV protease inhibition effects.<sup>2</sup> These special properties have stimulated considerable interest in this class of compounds, and various 2,3- or 3,4-fused polycycles or open-chain derivatives have been synthesized. A number of 4-hydroxycoumarin derivatives can be synthesized through the Michael addition of 4-hydroxycoumarins to  $\alpha,\beta$ -unsaturated carbonyl compounds utilizing the nucleophilicity of C-3.<sup>3</sup>

The reaction of 4-hydroxycoumarins with quinone is illustrative in this regard. It has been reported that 4-hy-

<sup>†</sup> School of Pharmaceutical Science, Sun Yat-Sen University.  
<sup>‡</sup> School of Chemistry and Chemical Engineering, Sun Yat-Sen University.

<sup>§</sup> Shaoguan College.


<sup>||</sup> The Hong Kong Polytechnic University.

(1) (a) Feuer, G. *Prog. Med. Chem.* **1973**, *10*, 85. (b) Obaseki, A. O.; Porter, W. R. *J. Heterocycl. Chem.* **1982**, *19*, 385.

(2) (a) O'Reilly, R. A. *Pharmacology* **1972**, *8*, 181. (b) Bourinbaier, A. S.; Tan, X.; Nagorny, R. *Acta Virol.* **1993**, *37*, 241. (c) Tummino, P. J.; Ferguson, D.; Hupe, D. *Biochem. Biophys. Res. Commun.* **1994**, *201*, 290. (d) Kirkiacharian, S.; Thuy, D. T.; Sicsic, S.; Bakhchinian, R.; Kurkjian, R.; Tonnaire, T. *IF Farmaco* **2002**, *57*, 703. (e) Thaisrivongs, S.; Tomich, P. K.; Watenpugh, K. D.; Chong, K. T.; Howe, W. J.; Yang, C. P.; Strohbach, J. W.; Turner, S. R.; McGrath, J. P.; Bohanon, M. J.; Lynn, J. C.; Mulichak, A. M.; Spinelli, P. A.; Hinshaw, R. R.; Pagano, P. J.; Moon, J. B.; Ruwart, M. J.; Wilkinson, K. F.; Rush, B. D.; Zipp, G. L.; Dalga, R. J.; Schwinde, F. J.; Howard, G. M.; Padbury, G. E.; Toth, L. N.; Zhao, Z.; Koeplinger, K. A.; Kakuk, T. J.; Cole, S. L.; Zaya, R. M.; Piper, R. C.; Jeffrey, P. *J. Med. Chem.* **1994**, *37*, 3200. (f) Thaisrivongs, S.; Watenpugh, K. D.; Howe, W. J.; Tomich, P. K.; Dolak, L. A.; Chong, K. T.; Tomich, C. S. C.; Tomasselli, A. G.; Turner, S. R.; Strohbach, J. W.; Mulichak, A. M.; Janakiraman, M. N.; Moon, J. B.; Lynn, J. C.; Horng, M. M.; Hinshaw, R. R.; Curry, K. A.; Rothrock, D. J. *J. Med. Chem.* **1995**, *38*, 3624.

(3) Trivedi, K. N.; Madhava Rao, S. S.; Mistry, S. V.; Desai, S. M. *J. Indian Chem. Soc.* **2001**, *18*, 579 and references cited therein.


droxycoumarins react with *o*-quinone to give the corresponding coumestans.<sup>4</sup> In contrast, the reaction of 4-hydroxycoumarin with *p*-quinone has been less well studied with only a few products having been reported from such reactions (Figure 1).<sup>5</sup>



**Figure 1.** Structures of the products from the reaction of 4-hydroxycoumarin with *p*-benzoquinone and naphthoquinone, respectively.

In the present work we now report an interesting reaction of 4-hydroxycoumarins with *p*-benzoquinone that is mediated by excess pyridine in aqueous acetone (v:v = 1:1) (Scheme 1). The reaction of 4-hydroxycoumarin with *p*-benzoquinone

**Scheme 1.** Preparation of Compounds **2**



and pyridine gives a pale yellow product **2a**,<sup>6</sup> which we have shown to be a pyridinium-zwitterionic compound by IR, MS- (ESI), <sup>1</sup>H NMR, <sup>13</sup>C NMR, and X-ray crystallographic studies.

Pyridinium zwitterions are often reactive species that are widely used in organic synthesis.<sup>7</sup> The preparation of zwitterions via the addition of pyridine derivatives to reactive double bonds or carbenes has been reported.<sup>8</sup> In most cases, the negative charge in the zwitterionic compound is on the atom adjacent to that attached to the pyridinium nitrogen and is delocalized by the presence of electron-withdrawing

substituents to improve the stability of the compound. Yet, the result is usually unsatisfactory. Recently, charge-separated pyridinium zwitterions have been prepared and proved to be stable.<sup>9</sup>

Under similar reaction conditions for the synthesis of **2a**, five other zwitterions were also prepared. The proposed zwitterionic structures for these compounds are consistent with the <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) spectroscopic data.<sup>10</sup> The unusual downfield signals of the pyridine ring of **2b** showed the presence of the pyridinium structure. The presence of the oxyanion on the lactone ring of **2b** was corroborated by the absence of a broad OH signal at δ 12.37 ppm, which appears in the <sup>1</sup>H NMR of 8-methyl-4-hydroxycoumarin. The quadruple peaks of the AB system for two protons of the phenolic ring at δ 6.95–6.98 ppm indicated that the pyridinium moiety and the 4-hydroxycoumarin moiety were on the same side of the hydroquinone ring. The different shift of the two α-protons located on the pyridine ring of **2b**, at δ 8.95 and 8.59 ppm, respectively, was caused by the uneven shielding of the negative ion attached to the pyrano ring. Similarly, the signals of the two β-protons were also unequal. Thus, these compounds might exist as atropisomers due to the restricted rotation about the C6–N1 bond.

The unambiguously determined structure of **2a** via single-crystal X-ray diffraction also lent support to the proposed structures of the other related compounds.<sup>11</sup> (Figure 2)

The positive charge located on the pyridinium cation and the negative charge was delocalized on the 1,3-dicarbonyl structure of the 4-hydroxycoumarin fragment. The delocalization of the negative charge on the 1,3-dicarbonyl structure

(6) **Preparation of Compounds 2a–f: General Procedure.** A mixture of 4-hydroxycoumarin **1a** (0.81 g, 5 mmol), *p*-benzoquinone (1.08 g, 10 mmol), and pyridine (0.79 g, 10 mmol) in 60 mL aqueous acetone (v:v = 1:1) was magnetically stirred at room temperature for 24 h. The reaction mixture was filtered to afford a brown crude product that was purified by column chromatography (silica gel, methanol/trichloromethane = 1:10) to give the yellow compound **2a** (0.79 g, 2.3 mmol) in 46% yield: <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 9.89 (1H, s), 9.62 (1H, s), 8.97 (1H, d, *J* = 6 Hz), 8.57 (1H, d, *J* = 6 Hz), 8.51 (1H, t, *J* = 8 Hz), 8.06 (1H, t, *J* = 7 Hz), 7.86–7.89 (2H, m), 7.42 (1H, t, *J* = 7 Hz), 7.17 (1H, t, *J* = 8 Hz), 7.07 (1H, d, *J* = 8 Hz), 6.98 ppm (2H, q, δ<sub>Ha</sub> = 6.99 ppm, δ<sub>Hb</sub> = 6.97 ppm, *J*<sub>AB</sub> = 9 Hz); <sup>13</sup>C NMR (125 MHz, DMSO-*d*<sub>6</sub>) δ 173.7, 162.2, 153.2, 150.3, 150.1, 146.7, 145.7, 142.9, 130.8, 130.4, 126.2, 125.9, 124.8, 122.4, 121.9, 121.9, 120.7, 115.4, 115.1, 92.6 ppm; IR (KBr) 3064, 1626, 1599, 1522, 1502 cm<sup>−1</sup>; MS (ESI) 346 (M − 1)<sup>−</sup>.


(7) Jursic, B. S.; Neumann, D. M.; Moore, Z.; Stevens, E. D. *J. Org. Chem.* **2002**, *67*, 2372 and references cited therein.

(8) (a) Visser, P.; Zuhse, R.; Wong, M. W.; Wentrup, C. *J. Am. Chem. Soc.* **1996**, *118*, 12598. (b) Kuhn, A.; Plüg, C.; Wentrup, C. *J. Am. Chem. Soc.* **2000**, *122*, 1945.

(9) (a) Jursic, B. S.; Neumann, D. M.; Martin, K. L.; Stevens, E. D. *Org. Lett.* **2002**, *4*, 811. (b) Jursic, B. S.; Neumann, D. M.; Moore, Z.; Stevens, E. D. *J. Org. Chem.* **2002**, *67*, 2372. (c) Sosnovskikh, V. Y.; Usachev, B. I.; Sizov, A. Y.; Vorontsov, I. I.; Shklyayev, Y. V. *Org. Lett.* **2003**, *5*, 3123.

(10) **Compound 2b:** <sup>1</sup>H NMR (500 MHz, DMSO-*d*<sub>6</sub>) δ 9.87 (1H, s), 9.58 (1H, s), 8.95 (1H, d, *J* = 6 Hz), 8.59 (1H, d, *J* = 6 Hz), 8.51 (1H, t, *J* = 7.5 Hz), 8.06 (1H, t, *J* = 8 Hz), 7.89 (1H, t, *J* = 8 Hz), 7.70 (1H, d, *J* = 8 Hz), 7.28 (1H, d, *J* = 7 Hz), 7.05 (1H, t, *J* = 7.5 Hz), 6.96 (2H, q, AB system, δ<sub>Ha</sub> = 6.97 ppm, δ<sub>Hb</sub> = 6.95 ppm, *J*<sub>AB</sub> = 9 Hz), 2.21 ppm (3H, s); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>) δ 173.9, 162.3, 151.6, 150.3, 149.9, 146.9, 145.6, 142.9, 131.8, 130.4, 126.2, 126.0, 123.9, 122.5, 121.9, 121.848, 121.4, 120.6, 114.9, 92.4, 15.1 ppm; IR (KBr) 3067, 1623, 1596, 1523, 1504 cm<sup>−1</sup>; MS (ESI) 360 (M − 1)<sup>−</sup>.

(11) **Crystal Data for 2a:** C<sub>20</sub>H<sub>15</sub>NO<sub>6</sub>, MW = 365.33, monoclinic P2<sub>1</sub>/c, *a* = 7.142(10), *b* = 16.314(2), *c* = 16.054(2) Å, *a* = 90°, *β* = 111.45(3)°, *γ* = 90°, *V* = 1741.0(4) Å<sup>3</sup>, *Z* = 4, *D*<sub>calcd</sub> = 1.394 g/cm<sup>3</sup>, *R*<sub>1</sub> = 0.047, *wR*<sub>2</sub> = 0.110 [*I* > 2σ(*I*)], *R*<sub>1</sub> = 0.085, *wR*<sub>2</sub> = 0.128 (all data), *S* = 1.03.




**Figure 2.** X-ray structure of compound **2a**.

was consistent with the observation that the C(20)–O(5) bond length [1.251(2) Å] was similar to the C(13)–O(3) bond length [1.255(2) Å] and shorter than the C–O bond length [1.348(5) Å] of the enol form in the 4-hydroxycoumarin fragment.<sup>12</sup> Neither the pyridine ring nor the pyranone ring of the 4-hydroxycoumarin fragment was found to conjugate to the hydroquinone ring. The dihedral angles for C(11)–C(6)–N(1)–C(1) and C(10)–C(11)–C(12)–C(20) were found to be 65.2 and 62.5°, respectively.

A possible mechanism of **2a** formation is presented in Scheme 2. The Michael addition of 4-hydroxycoumarin into *p*-benzoquinone first gives unstable intermediate **3**, which is subsequently autooxidized to generate quinone derivative **4**. **4** is attacked by pyridine at the 3-position to form **2a**, which is in a hydroquinone form due to the electron-withdrawing effect of the pyridinium moiety.

(12) Dolmella, A.; Gatto, S.; Girardi, E.; Bandoli, G. *J. Mol. Struct.* **1999**, *513*, 177.

**Scheme 2.** Possible Mechanism of Adduct Formation of **2a**



In summary, we described here an unexpected and interesting reaction of 4-hydroxycoumarins with *p*-benzoquinone and pyridine in aqueous acetone (v:v = 1:1). Six stable zwitterionic compounds have been synthesized via this reaction.

**Acknowledgment.** We thank the National Science Foundation of China (2027085), the Guangdong Provincial Science Foundation (031594), and the Hong Kong Polytechnic University ASD Fund for financial support of this study.

**Supporting Information Available:** X-ray data for compound **2a**; spectral data of compounds **2a–f**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL048109P