

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/8179180>

Unusually Stable Molecular Capsule Formation of a Tetraphenyleneurea Cavitand

ARTICLE *in* ORGANIC LETTERS · DECEMBER 2004

Impact Factor: 6.36 · DOI: 10.1021/ol048446y · Source: PubMed

CITATIONS

27

READS

7

6 AUTHORS, INCLUDING:



Heung-Jin Choi

Kyungpook National University

93 PUBLICATIONS 1,089 CITATIONS

SEE PROFILE



Chan Sik Cho

Kyungpook National University

216 PUBLICATIONS 3,249 CITATIONS

SEE PROFILE



Hong Sung Kim

Pusan National University

133 PUBLICATIONS 966 CITATIONS

SEE PROFILE

Unusually Stable Molecular Capsule Formation of a Tetraphenyleneurea Cavitand


Heung-Jin Choi,* Yeon Sil Park, Chan Sik Cho, Kwangnak Koh,[†]
Sung-Hong Kim,[‡] and Kyungsoo Paek[§]

Department of Applied Chemistry, Kyungpook National University,
Daegu 702-701, Korea, College of Pharmacy, Pusan National University,
Busan 609-735, Korea, Korea Basic Science Institute, Daegu Branch,
Daegu 702-701, Korea, and CAMDRC and Department of Chemistry,
Soongsil University, Seoul 156-743, Korea

choihj@knu.ac.kr

Received August 5, 2004

ABSTRACT



An unusually stable molecular capsule was formed by heating phenyleneurea-spanned resorcinarene cavitand with 4-methyl-*N*-*p*-tolylbenzamide. The molecular capsule behaved as a discrete molecular entity showing a cylindrical D_{4d} structure and showed no guest exchange in toluene- d_8 even at 100 °C.

Self-assembled molecular capsules held together by hydrogen bonds or metal–ligand interaction have attracted much interest.^{1–3} The successful application of the complementary hydrogen-bonding theme was first introduced by the groups

of Rebek and de Mendoza for self-assembling glycoluril-derived capsules such as “tennis balls” and “softballs”.^{4–10}

Cavitand **1** reported by Rebek and co-workers self-assembled to form a cylinder-shape capsule **1•1** by eight bifurcated hydrogen bonds between an imide hydrogen atom in a cavitand with two neighboring carbonyl oxygen atoms as shown in Figure 1.^{11–23} Capsule **1•1** can accommodate various guests such as 4-methyl-*N*-*p*-tolylbenzamide **4** in a

[†] Pusan National University.

[‡] Korea Basic Science Institute.

[§] Soongsil University.

(1) For reviews, see: (a) Hof, F.; Craig, S. L.; Nuckolls, C.; Rebek, J., Jr. *Angew. Chem., Int. Ed.* **2002**, *41*, 1489. (b) Conn, M. M.; Rebek, J., Jr. *Chem. Rev.* **1997**, *97*, 1647.

(2) (a) Vysotsky, M. O.; Pop, A.; Broda, F.; Thondorf, I.; Böhmer, V. *Chem. Eur. J.* **2001**, *7*, 4403. (b) Chapman, R. G.; Olovsson, G.; Trotter, J.; Sherman, J. C. *J. Am. Chem. Soc.* **1998**, *120*, 6252. (c) MacGillivray, L. R.; Atwood, J. L. *Nature* **1997**, *389*, 469. (d) Zadnád, R.; Junkers, M.; Schrader, T.; Grawe, T.; Kraft, A. *J. Org. Chem.* **2003**, *68*, 6511.

(3) (a) Holliday, B. J.; Mirkin, C. A. *Angew. Chem., Int. Ed.* **2001**, *40*, 2022. (b) Fochi, F.; Jacopozzi, P.; Wegelius, E.; Rissanen, K.; Cozzini, P.; Marastoni, E.; Fiscaro, E.; Manini, P.; Fokkens, R.; Dalcanele, E. *J. Am. Chem. Soc.* **2001**, *123*, 7539. (c) Levi, S. A.; Guatter, P.; van Veggel, F. C. J. M.; Vancso, G. J.; Dalcanele, E.; Reinhoudt, D. N. *Angew. Chem., Int. Ed.* **2001**, *40*, 1892. (d) Ikeda, A.; Udzu, H.; Zhong, Z.; Shinkai, S.; Sakamoto, S.; Yamaguchi, K. *J. Am. Chem. Soc.* **2001**, *123*, 3872.

(4) Wyler, R.; de Mendoza, J.; Rebek, J., Jr. *Angew. Chem., Int. Ed.* **1993**, *32*, 1699.

(5) Branda, N.; Wyler, R.; Rebek, J., Jr. *Science* **1994**, *263*, 1267.

(6) Meissner, R. S.; Rebek, J., Jr.; de Mendoza, J. *Science* **1995**, *270*, 1485.

(7) Kang, J.; Rebek, J., Jr. *Nature* **1996**, *382*, 239.

(8) Conn, M. M.; Rebek, J., Jr. *Chem. Rev.* **1997**, *97*, 1647.

(9) Martín, T.; Obst, U.; Rebek, J., Jr. *Science* **1998**, *281*, 1842.

(10) Rebek, J., Jr. *Acc. Chem. Res.* **1999**, *32*, 278.

(11) Heinz, T.; Rudkevich, D. M.; Rebek, J., Jr. *Nature* **1998**, *394*, 764.

(12) Heinz, T.; Rudkevich, D. M.; Rebek, J., Jr. *Angew. Chem., Int. Ed.* **1999**, *38*, 1136.




Figure 1. Hydrogen bonding modes of self-assembled pyrazin-imide capsule **1·1** and phenyleneurea capsule **2·2**. The bond lengths were calculated from the energy-minimized structures (Spartan 04, V1.0.1, AM1 semiempirical).

nonpolar mesitylene-*d*₁₂. The inclusion complexes **G@1·1** show interesting guest-exchange phenomena and unprecedented isomerism.^{11–23}

Phenyleneurea cavitands **2**, **3**, and other analogues were first reported by de Mendoza et al.²⁴ These cavitands aggregated, forming different self-organized structures such as vesicles or filaments, depending on the nature and length of the four alkyl feet. Cavitand **2** formed large reverse vesicles through side to side extensive stacking. In contrast, cavitand **3** formed dimeric capsules with carboxylic acids.

We report the observation of the unusually stable capsule formation of cavitand **2**. At elevated temperature guest molecules could template cavitand **2** to form a dimeric capsule **2·2** by reorganizing the intermolecular hydrogen bonds of the aggregate of cavitand **2**.

Cavitand **2** was directly prepared as reported by de Mendoza,²⁴ and the intermediates, i.e., the corresponding octanitro cavitand and octaamino cavitand reported by Rebek^{23,25–27} were similarly prepared by the method reported by Cram et al.²⁸

At room temperature, cavitand **2** was barely soluble in CH₂Cl₂, CHCl₃, toluene, or mesitylene but reasonably soluble in DMF or DMSO, as reported by de Mendoza.²⁴ However, at elevated temperature it became soluble in these nonpolar solvents and then stayed in homogeneous solution at room temperature.

The encapsulation studies of cavitand **2** and 4-methyl-*N*-*p*-tolylbenzamide **4** were performed in mesitylene as devised by Rebek.^{12,13} A mixture of cavitand **2** and guest **4** in mesitylene remained as a heterogeneous mixture at room temperature even after 5 days, but at above 100 °C the mixture was slowly homogenized. The encapsulation complex **4@2·2** was prepared by heating the mixture under reflux until it became homogeneous and then removing the solvent by vacuum distillation at 70 °C. The solid residue was dried at 100 °C under vacuum. The solid complex **4@2·2** is then soluble in CDCl₃, DMSO-*d*₆, or toluene-*d*₈ at room temperature.

An equimolar mixture of cavitand **2** and guest **4** (1.67 mM: 1.67 mM) in mesitylene gave a soluble complex in a relatively short period (30 min at 170 °C), but incomplete encapsulation was observed by ¹H NMR spectroscopy, which showed three different chemical shifts of the N–H of urea moieties for cavitand **2** (10.35, 10.36, and 10.40 ppm). The chemical shift of the N–H of cavitand **2** was a good indicator of whether the encapsulation complex **4@2·2** was formed completely or partially: the chemical shift of **2·2** in toluene-*d*₈ was observed at 10.35 ppm.

A complete encapsulated complex **4@2·2** was prepared from a mixture of cavitand **2** and guest **4** (1.67 mM:6 mM, respectively) in mesitylene by heating under reflux for 30 min. The ¹H NMR spectrum of **4@2·2** in toluene-*d*₈ at 100 °C showed the chemical shifts for guest **4** at 5.41 (d, H_c), 5.20 (d, H_e), 3.29 (d, H_b), 3.14 (d, H_f), –2.33 (s, H_g), and –2.41 (s, H_a) ppm (Figure 2 and Table 1). Like the isomorphous cavitand **1·1**,¹¹ the large upfield chemical shifts of encapsulated guest **4** in capsule **2·2** are observed up to 4.50 ppm for the methyl group (H_a, 2.09 ppm) of free **4** in the ¹H NMR spectrum as a result of the shielding by aromatic ring current of capsule **2·2**. Compared to the chemical shifts (Δδ₁) for **4** in **1·1**, those (Δδ₂) in **2·2** are slightly smaller (ΔΔδ = 0.40, 0.37 for H_a and H_g, respectively), which is consistent with the molecular dimensions (17.21 vs 17.43 Å through the long C₄ axis of **1·1** and **2·2**, respectively) calculated using semiempirical AM1.

The two different chemical shifts of the N–H of urea moieties for cavitand **2** (10.40 and 10.36 ppm) confirm that

(13) Körner, S. K.; Tucci, F. C.; Rudkevich, D. M.; Heinz, T.; Rebek, J., Jr. *Chem. Eur. J.* **2000**, *6*, 187.

(14) Craig, S. L.; Lin, S.; Chen, J.; Rebek, J., Jr. *J. Am. Chem. Soc.* **2002**, *124*, 8780.

(15) Shivanyuk, A.; Rebek, J., Jr. *Angew. Chem., Int. Ed.* **2003**, *42*, 684.

(16) Shivanyuk, A.; Rebek, J., Jr. *J. Am. Chem. Soc.* **2002**, *124*, 12074.

(17) Scarso, A.; Shivanyuk, A.; Hayashida, O.; Rebek, J., Jr. *J. Am. Chem. Soc.* **2003**, *125*, 6239.

(18) Shivanyuk, A.; Scarso, A.; Rebek, J., Jr. *Chem. Commun.* **2003**, 1230.

(19) Shivanyuk, S.; Rebek, J., Jr. *Chem. Commun.* **2002**, 2326.

(20) Hayashida, O.; Shivanyuk, A.; Rebek, J., Jr. *Angew. Chem., Int. Ed.* **2002**, *41*, 3423.

(21) Chen, J.; Körner, S.; Craig, S. L.; Lin, S.; Rudkevich, D. M.; Rebek, J., Jr. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 2593.

(22) Hof, F.; Rebek, J., Jr. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 4775.

(23) Far, A. R.; Shivanyuk, A.; Rebek, J., Jr. *J. Am. Chem. Soc.* **2002**, *124*, 2854.

(24) Ebbing, M. H. K.; Villa, M.-J.; Valpuesta, J.-M.; Prados, P.; de Mendoza, J. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99*, 4962.

(25) Rudkevich, D. M.; Hilmersson, G.; Rebek, J., Jr. *J. Am. Chem. Soc.* **1997**, *119*, 9911.

(26) Rudkevich, D. M.; Hilmersson, G.; Rebek, J., Jr. *J. Am. Chem. Soc.* **1998**, *120*, 12216.

(27) Tucci, F. C.; Rudkevich, D. M.; Rebek, J., Jr. *J. Org. Chem.* **1999**, *64*, 4555.

(28) Cram, D. J.; Choi, H.-J.; Bryant, J. A.; Knobler, C. B. *J. Am. Chem. Soc.* **1992**, *114*, 7748.





Figure 2. ^1H NMR spectrum of encapsulation complex 4-methyl-*N-p*-tolylbenzamide@**2·2** in toluene- d_8 at 100 °C.

two chemically nonequivalent methyl groups of guest **4** occupy each cavitand of the dimeric capsule and the guest's head-to-tail rotation is tightly restricted on the ^1H NMR time scale even at 100 °C.

The capsular complex **4@2·2** was unusually inert in nonpolar solvents such as toluene- d_8 . No guest exchange with the solvent molecule was observed in toluene- d_8 at 100 °C for 24 h. However, in polar solvents such as DMSO- d_6 the guest **4** in capsular complex **4@2·2** escaped rapidly at room temperature.

Table 1. Comparison of ^1H NMR Chemical Shifts (ppm) of 4-Methyl-*N-p*-tolylbenzamide **4** and Encapsulation Complexes **4@1·1** and **4@2·2** in Toluene- d_8 ^a



	protons						
	H _a	H _b	H _c	H _d	H _e	H _f	H _g
free 4	2.09	6.96(d)	7.54(d)	<i>d</i>	7.40(d)	6.92(d)	2.12
4@1·1 ^b	-2.81	<i>c</i>	<i>c</i>	<i>c</i>	<i>c</i>	<i>c</i>	-2.70
$\Delta\delta_1$	4.90						4.82
4@2·2	-2.41	3.29(d)	5.41(d)	4.54	5.20(d)	3.14(d)	-2.33
$\Delta\delta_2$	4.50	3.67	2.13		2.20	3.78	4.45
$\Delta\Delta\delta$	0.40						0.37

^a $\Delta\delta_1 = \delta$ of free **4** - δ of **4@1·1**, $\Delta\delta_2 = \delta$ of free **4** - δ of **4@2·2**. $\Delta\Delta\delta = \Delta\delta_1 - \Delta\delta_2$. ^b In mesitylene- d_{12} at 295 K.¹¹ ^c Not reported. ^d Obscured with solvent peak.

Capsular complexes with smaller guests such as *p*-xylene were prepared by heating cavitand **2** in *p*-xylene. In the toluene- d_8 , *p*-xylene escaped rapidly as a result of the dynamic equilibrium with toluene- d_8 by mass law ($t_{1/2} = 37$ min at 50 °C).

It is obvious that the hydrogen bonding mode in the self-assembly **2·2** is superior to that in **1·1**, which has been already observed by de Mendoza.²⁴ For molecular capsule **2·2**, two sets of lone pair electrons in the sp^2 hybridized carbonyl oxygen complementarily hydrogen bond with two adjacent H-N protons, which consumes all of the potential hydrogen bond donors and acceptors. However, in the self-assembly **1·1** only one set of lone pair electrons of the carbonyl oxygen hydrogen bonds to the adjacent H-N proton, as shown in Figure 1.

The unusual stability of capsule **4@2·2** in toluene- d_8 without exchanging guest **4** with solvent even at 100 °C suggests that, in addition to the exceptional tight hydrogen bond between two cavitands **2**, van der Waals interaction between guest **4** and capsule **2·2** and the dipolar attraction or hydrogen bond between the amide group of guest **4** and the urea groups of the capsule **2·2** are efficiently operating.

However, in polar solvents such as DMSO- d_6 the guest **4** in capsular complex **4@2·2** escaped rapidly at room temperature. The polar DMSO- d_6 molecules compete for the hydrogen bond acceptors in the dimeric capsule **2·2**, which weakens the stability of capsule **2·2** and allows the fast escape of guest **4** and/or the fast exchange of guest **4** with solvent. The observed vase conformation of **2** in DMSO- d_6 confirms that cavitand **2** still exists as a dimeric capsule DMSO- d_6 @**2·2** because monomeric cavitand **4** could only exist as rectangular kite conformation as a result of the repulsion between adjacent urea groups.²⁸

In summary, the solvent- and/or guest-assisted molecular capsule formations **G@2·2** were observed and characterized by heating a mixture of cavitand **2** and guests in a nonpolar solvent at the elevated temperature.

Acknowledgment. This work was supported by grant R05-2001-000-00225-0 (2003) from the Basic Research Program of the Korea Science & Engineering Foundation.

Supporting Information Available: Synthetic procedures, encapsulation experiments, molecular modeling, and selected NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL048446Y