

Acyclic Stereocontrol Based on Nonchelation-controlled Ene Reactions with α -Haloaldehydes

Koichi Mikami,* Teck-Peng Loh and Takeshi Nakai*

Department of Chemical Technology, Tokyo Institute of Technology, Ookayama, Meguro-ku, Tokyo 152, Japan

Aluminium(III)-promoted ene reactions with α -haloaldehydes are shown to exhibit a high *anti*-diastereofacial (non-chelation) selection or *syn*-diastereoselection to afford an efficient method for preparing stereochemically-defined β -haloalcohols including the 22*R*-hydroxy side chain unit in steroids.

The ene reaction involving carbonyl compounds, aldehydes in particular, as enophiles (carbonyl-ene reaction) has currently emerged as a new tool for acyclic stereoselection.¹ However, the types of aldehyde enophile explored thus far have been limited.² Herein we report a new type of Lewis acid-promoted ene reaction using α -bromo- or α -chloro-aldehydes as enophiles (haloaldehyde-ene reaction) which proceeds at relatively low temperatures under effective nonchelation control with high *syn*-diastereoselectivity.

The reactions of α -bromopropanal **1** and isobutene **2a** (2 equiv.) at -78°C in dichloromethane (Scheme 1) were found to give the ene products **3a** and **4a** in good yields, using organoaluminium reagents (1 equiv.) as the Lewis acid (Table 1). The *anti* (nonchelation) stereoisomer **3a** was obtained as the major product, the ratio depending on the nature of the Lewis acid employed. Of special value is the Me_2AlCl -promoted reaction which provides **3a** in relatively high selectivity. The structural assignment of the ene products **3** and **4** was based on conversion to the epoxides **5** and **6**; the epoxide **6** derived from the minor ene product **4** showed a relatively strong NOE between the methylene and methyl

protons. The *anti*-diastereofacial selectivity thus observed is reasonably explained in terms of Felkin-Anh's or Cram's dipolar model.³ Thus, this new type of ene reaction is proved to proceed under effective nonchelation control.





Table 1 Haloaldehyde-ene reactions^a

Alkene	AlL_n	3 : 4 ^b	Total yield (%) ^c
2a	Me_2AlCl	79 : 21	90
2a	ArOAlMeCl^d	72 : 28	82
2a	$(\text{ArO})_2\text{AlCl}^d$	68 : 32	80
2b	Me_2AlCl	75 : 25	87
2c	Me_2AlCl	79 : 21	87

^a All reactions were carried out on a 1 mmol scale under argon. ^b The isomer ratio was determined by ^{13}C NMR and HPLC analyses. ^c Yield of isolated product after silica gel chromatography. ^d $\text{Ar} = 2,4,6\text{-Me}_3\text{-C}_6\text{H}_2$.

Scheme 1 Reagents and conditions: i, AlL_n , CH_2Cl_2 , -78°C ; ii, NaH , dimethylformamide



Scheme 2 Reagents and conditions: i, ClCH_2CHO **8**, Me_2AlCl , CH_2Cl_2 , -78°C ; ii, NaH ; iii, H_2 , PtO_2

Next, we examined the simple diastereoselection of the haloaldehyde–ene reaction in the context of steroid side chain synthesis.⁴ Thus, the reaction of the easily available steroidal alkene **7**⁵ and chloroacetaldehyde **8** (1 equiv. each) with Me_2AlCl was found to show an extremely high level of simple *syn*-diastereoselection. The 20*S*,22*R*-*syn* product **9** was obtained as a single stereoisomer in 73% yield. The stereochemistry was assigned on the basis of the ^{13}C and ^1H NMR analyses,⁶ after conversion to the steroidal epoxide **10**, a key intermediate of 22*R*- α -hydroxylated steroid side chains.⁶

Received, 3rd September 1990; Com. 0/03997G

References

- Reviews: K. Mikami, M. Terada, M. Shimizu and T. Nakai, *J. Synth. Org. Chem., Jpn.*, 1990, **48**, 292; B. B. Snider, *Acc. Chem. Res.*, 1980, **13**, 426.
- Chloral–ene reactions: (a) G. B. Gill and B. Wallace, *J. Chem. Soc., Chem. Commun.*, 1977, 382; J. P. Benner, G. B. Gill, S. J. Parrott, B. Wallace and M. J. Begley, *J. Chem. Soc., Perkin Trans. I*, 1984, 315; (b) K. Maruoka, Y. Hoshino, T. Shirasaka and H. Yamamoto, *Tetrahedron Lett.*, 1988, **29**, 3967. Glyoxylate–ene reactions: (c) O. Achmatowicz, Jr., and B. Szechner, *J. Org. Chem.*, 1972, **37**, 964; (d) B. B. Snider and J. W. van Straten, *J. Org. Chem.*, 1979, **44**, 3567; (e) J. K. Whitesell, *Acc. Chem. Res.*, 1985, **18**, 280; (f) K. Mikami, T.-P. Loh and T. Nakai, *Tetrahedron Lett.*, 1988, **29**, 6305; (g) K. Mikami, M. Terada and T. Nakai, *J. Am. Chem. Soc.*, 1990, **112**, 3949. Propynal–ene reactions: (h) K. Mikami, T.-P. Loh and T. Nakai, *J. Chem. Soc., Chem. Commun.*, 1988, 1430; (i) K. Mikami, T.-P. Loh and T. Nakai, *J. Am. Chem. Soc.*, 1990, **112**, 6737. Alkoxyaldehyde–ene reactions: (j) K. Mikami and T.-P. Loh, *Tetrahedron Asymmetry*, 1990, **1**, 13. Aminoaldehyde–ene reactions: (k) K. Mikami, M. Kaneko, T.-P. Loh, M. Terada and T. Nakai, *Tetrahedron Lett.*, 1990, **31**, 3909.
- N. T. Anh, *Top. Curr. Chem.*, 1980, **88**, 145.
- Reviews: D. M. Patai and J. Wicha, *Chem. Rev.*, 1978, **78**, 199; J. Redpath and F. Zeelen, *Chem. Soc. Rev.*, 1983, **12**, 75; R. Pardo and M. Santelli, *Bull. Chem. Soc. Fr.*, 1985, 99; K. Mori, *J. Synth. Org. Chem., Jpn.*, 1985, **43**, 849.
- N. R. Schmuff and B. M. Trost, *J. Org. Chem.*, 1983, **48**, 1404.
- For the synthesis of the 20*S*,22*R*-epoxide **10** and its 22*S*-epimer, and their transformations to the 22*R*-hydroxycholesterol and 22*S*-hydroxylated ecdysone side chains respectively, see: M. Koreeda and D. J. Ricca, *J. Org. Chem.*, 1986, **51**, 4090. Also see: K. Mikami, K. Kawamoto and T. Nakai, *Tetrahedron Lett.*, 1985, **26**, 5799 and ref. 2f.