

Spotlight on Pegfilgrastim in Chemotherapy-Induced Neutropenia¹

James E. Frampton and Gillian M. Keating

Adis International Inc., Yardley, Pennsylvania, USA

Contents

Abstract	405
1. Pharmacological Properties	405
2. Therapeutic Efficacy	406
3. Tolerability	406
4. Pharmacoeconomic Considerations	406

Abstract

Pegfilgrastim (Neulasta®), the sustained-duration form of filgrastim (recombinant human granulocyte colony-stimulating factor [G-CSF]), is created by the addition of a polyethylene glycol (PEG) moiety to filgrastim. Its approved indication in the US is to decrease the incidence of infection, as manifested by febrile neutropenia, in patients with non-myeloid malignancies receiving myelosuppressive chemotherapy.

A single subcutaneous injection of pegfilgrastim once per chemotherapy cycle was more effective than placebo as an adjunct to moderately myelosuppressive chemotherapy for breast cancer, no less effective than daily injections of filgrastim as an adjunct to highly myelosuppressive chemotherapy for breast cancer, and as effective as daily filgrastim as an adjunct to chemotherapy for lymphoma (predominantly non-Hodgkin lymphoma [NHL]) and acute myeloid leukemia. Pegfilgrastim has also successfully supported delivery of dose-dense chemotherapy, stem cell mobilization, and stem cell transplantation after high-dose chemotherapy in patients with non-myeloid or myeloid malignancies. By offering a convenient alternative to daily filgrastim, once-per-cycle administration of pegfilgrastim has the potential to simplify the management of chemotherapy-induced neutropenia, further improve patient health-related quality of life, and reduce total treatment costs in breast cancer and NHL, and possibly other cancer settings. Pegfilgrastim should, likewise, permit simplification of G-CSF-based stem cell mobilization and transplantation procedures.

1. Pharmacological Properties

Pegfilgrastim is produced by covalently binding a 20 kDa polyethylene glycol (PEG) molecule to the *N*-terminal methionyl residue of filgrastim.^[1] Pegfilgrastim and filgrastim share the same mechanism of action (stimulation of granulocytopoiesis) and pharmacodynamic profile (e.g. elevation of absolute neutrophil count [ANC] and reduction in duration of chemotherapy-induced neutropenia);^[2] both recombinant human granulocyte colony-stimulating factors (G-CSFs) are administered by the subcutaneous route. However, the greater weight and size of the pegfilgrastim mole-

cule results in reduced renal clearance, an increased terminal elimination half-life, and hence a sustained duration of action relative to filgrastim.^[3,4] In patients with chemotherapy-induced neutropenia after myelosuppressive chemotherapy, plasma concentrations of pegfilgrastim remained elevated throughout the ANC nadir and declined rapidly as ANC recovered.^[4-6] This is consistent with the suggestion that self-regulating, neutrophil-mediated clearance, rather than renal elimination, is the predominant route of elimination of pegfilgrastim.^[7]

¹ This spotlight is derived from abstract and summary text of an Adis Drug Evaluation originally published in full in *American Journal of Cancer* 2005; 4 (4): 255-74. Reviewers of the original full text article are listed in the Acknowledgments section.

2. Therapeutic Efficacy

The efficacy of subcutaneous pegfilgrastim once per cycle in providing neutrophil support after standard regimens of myelosuppressive chemotherapy has been demonstrated in patients, most of whom were previously untreated, with breast cancer^[6,8,9] or lymphomas (non-Hodgkin lymphoma [NHL] or Hodgkin disease),^[10-12] or acute myeloid leukemia (AML).^[13]

In large, noninferiority, pivotal, phase III trials in patients with high-risk stage II–IV breast cancer^[6,9] and a phase II trial in patients with lymphoma (predominantly NHL),^[10] a fixed (6mg) or bodyweight-based (100 µg/kg) dose of pegfilgrastim administered once per chemotherapy cycle was no less effective than daily filgrastim 5 µg/kg (mean/median ≈11 doses per cycle) in reducing chemotherapy-induced neutropenia after highly myelosuppressive chemotherapy. In the studies in patients with breast cancer,^[6,9] which were identical except for the mode of pegfilgrastim dosing, between-group differences in the mean duration of severe neutropenia (ANC <0.5 × 10⁹/L) during the first chemotherapy cycle (primary endpoint) were <1 day, thus satisfying the noninferiority criteria for pegfilgrastim versus filgrastim. A similar outcome was seen in the NHL trial,^[10] although this was not formally designed as a noninferiority study.

Notwithstanding the noninferiority design of the studies in patients with breast cancer,^[6,9] a significant reduction in the incidence of febrile neutropenia across all chemotherapy cycles was seen with pegfilgrastim versus filgrastim in the trial of pegfilgrastim 100 µg/kg per cycle (9% vs 18%)^[9] as well as in a combined analysis of both trials (11% vs 19%).^[14] Additionally, subset analyses in the studies in patients with breast cancer suggested that the effect of pegfilgrastim on the duration of severe neutropenia following one or all (four) cycles of chemotherapy was similar regardless of prior chemotherapy or radiotherapy exposure and, in patients receiving pegfilgrastim 6mg per cycle, irrespective of bodyweight.^[6,9]

Relative to placebo, pegfilgrastim 6mg once per cycle significantly reduced rates of febrile neutropenia (17% vs 1%), hospitalization (14% vs 1%) and intravenous anti-infectives (10% vs 2%) over all cycles in a large, randomized, double-blind, phase III trial in 928 patients with stage I–IV breast cancer who received moderately myelosuppressive chemotherapy.^[8]

Pegfilgrastim was also effective in supporting the delivery of dose-dense cytotoxic chemotherapy regimens for previously untreated solid tumors (breast cancer^[15,16] and small cell lung cancer^[17]) and lymphomas (NHL^[18,19] and Hodgkin disease^[20,21]).

Pegfilgrastim, alone or in combination with myelosuppressive chemotherapy, successfully mobilized CD34+ cells in patients with solid tumors, lymphomas, or multiple myeloma;^[4,22-24] it also supported stem cell transplantation after high-dose chemotherapy in patients with lymphomas or multiple myeloma.^[25]

Pegfilgrastim was, in general, at least as effective as daily filgrastim based on 'head-to-head' and/or retrospective comparisons conducted in the settings of AML,^[13] dose-dense chemotherapy,^[18] and stem cell mobilization.^[23]

3. Tolerability

Pegfilgrastim was generally well tolerated in a total of 465 patients with solid tumors or lymphomas undergoing myelosuppressive chemotherapy in clinical trials; the drug had an adverse event profile (including changes in laboratory values) similar to that of filgrastim.^[26]

Medullary bone pain was the most common treatment-related adverse event with pegfilgrastim, occurring in approximately one-quarter of patients who received the drug in clinical trials.^[26] The incidence, severity, and duration of bone pain with a single injection of pegfilgrastim 6mg or 100 µg/kg per cycle was not significantly different from that with daily doses of filgrastim 5 µg/kg in the two pivotal phase III trials in patients with breast cancer.^[6,9,27] However, if bone pain occurred, the onset was earlier after the initial and final cycles of chemotherapy in the fixed-dose trial of pegfilgrastim 6mg once per cycle.^[6,27] Bone pain was generally of mild-to-moderate intensity and usually resolved with non-narcotic analgesics.^[27]

In the large, noninferiority, phase III studies in patients with breast cancer,^[6,9] transient elevations in plasma levels of lactate dehydrogenase, alkaline phosphatase, and uric acid with pegfilgrastim were not clinically significant, and no neutralizing antibodies against the drug were detected.

4. Pharmacoeconomic Considerations

Preliminary reports of economic modelling studies based on outcomes in the two pivotal phase III trials in patients with breast cancer suggested that once-per-cycle pegfilgrastim 6mg or 100 µg/kg may reduce total chemotherapy-induced neutropenia management costs over all four cycles relative to daily filgrastim 5 µg/kg.^[28-30]

According to other preliminary reports, universal (i.e. non-targeted) primary prophylaxis with pegfilgrastim is predicted to be cost saving compared with no G-CSF treatment when the risk of developing febrile neutropenia in the first chemotherapy cycle is >23% in patients receiving cyclophosphamide, doxorubicin, vincristine, and prednisolone for NHL and >16% in patients receiving docetaxel, with or without doxorubicin, for breast cancer.^[31,32]

Acknowledgments

The full text article in *American Journal of Cancer* 2005; 4 (4): 255-74 was reviewed by: **R.A. Beveridge**, Inova Fairfax Hospital, Fairfax, Virginia, USA; **P.S. Carey**, Sunderland Royal Infirmary, Sunderland, England; **D.C. Dale**, University of Washington, Department of Medicine, Seattle, Washington, USA; **D.B. Daniel**, Duke University Medical Center, Department of

Medicine, Durham, North Carolina, USA; **M.D. Green**, Royal Melbourne Hospital, Melbourne, Victoria, Australia; **E. Kubista**, Medical University of Vienna, Vienna, Austria; and **F. Willis**, St. George's Hospital, London, England.

References

- Molineux G. The design and development of pegfilgrastim (PEG-rmetHuG-CSF, Neulasta®). *Curr Pharm Des* 2004; 10 (11): 1235-44
- Lord BI, Woolford LB, Molineux G. Kinetics of neutrophil production in normal and neutropenic animals during the response to filgrastim (r-metHu G-CSF) or filgrastim SD/01 (PEG-r-metHu G-CSF). *Clin Cancer Res* 2001 Jul; 7: 2085-90
- Molineux G. Pegylation: engineering improved biopharmaceuticals for oncology. *Pharmacotherapy* 2003 Aug; 23 (8 Pt 2): 3S-8S
- Johnston E, Crawford J, Blackwell S, et al. Randomized, dose-escalation study of SD/01 compared with daily filgrastim in patients receiving chemotherapy. *J Clin Oncol* 2000 Jul; 18 (13): 2522-8
- Holmes FA, Jones SE, O'Shaughnessy J, et al. Comparable efficacy and safety profiles of once-per-cycle pegfilgrastim and daily injection filgrastim in chemotherapy-induced neutropenia: a multicenter dose-finding study in women with breast cancer. *Ann Oncol* 2002 Jun; 13 (6): 903-9
- Green MD, Koelbl H, Baselga J, et al. A randomized double-blind multicenter phase III study of fixed-dose single-administration pegfilgrastim versus daily filgrastim in patients receiving myelosuppressive chemotherapy. *Ann Oncol* 2003 Jan; 14 (1): 29-35
- Zamboni WC. Pharmacokinetics of pegfilgrastim. *Pharmacotherapy* 2003 Aug; 23 (8 Pt 2): 9S-14S
- Vogel CL, Wojtukiewicz MZ, Carroll RR, et al. First and subsequent cycle use of pegfilgrastim prevents febrile neutropenia in patients with breast cancer: a multicenter, double-blind, placebo-controlled phase III study. *J Clin Oncol* 2005; 23 (6): 1178-84
- Holmes FA, O'Shaughnessy JA, Vukelja S, et al. Blinded, randomized, multicenter study to evaluate single administration pegfilgrastim once per cycle versus daily filgrastim as an adjunct to chemotherapy in patients with high-risk stage II or stage III/IV breast cancer. *J Clin Oncol* 2002 Feb 1; 20 (3): 727-31
- Vose JM, Crump M, Lazarus H, et al. Randomized, multicenter, open-label study of pegfilgrastim compared with daily filgrastim after chemotherapy for lymphoma. *J Clin Oncol* 2003 Feb 1; 21 (3): 514-9
- Grigg A, Solal-Celigny P, Hoskin P, et al. Open-label, randomized study of pegfilgrastim vs. daily filgrastim as an adjunct to chemotherapy in elderly patients with non-Hodgkin's lymphoma. *Leuk Lymphoma* 2003 Sep; 44 (9): 1503-8
- George S, Yunus F, Case D, et al. Fixed-dose pegfilgrastim is safe and allows neutrophil recovery in patients with non-Hodgkin's lymphoma. *Leuk Lymphoma* 2003 Oct; 44 (10): 1691-6
- Bosi A, Szer J, Kassiss J, et al. A multicentre, double-blind, randomised, phase 2 trial comparing pegfilgrastim with filgrastim as an adjunct to chemotherapy for acute myeloid leukaemia (AML) [abstract no. 866]. *Blood* 2004 Nov 16; 104 (11 Pt 1): 247a. Plus poster presented at the 46th Annual Meeting of the American Society of Hematology; 2004 Dec 4-7; San Diego (CA)
- Siena S, Piccart MJ, Holmes FA, et al. A combined analysis of two pivotal randomized trials of a single dose of pegfilgrastim per chemotherapy cycle and daily filgrastim in patients with stage II-IV breast cancer. *Oncol Rep* 2003 May; 10 (3): 715-24
- Burstein HJ, Parker LM, Doherty J, et al. Use of the long-acting hematopoietic growth factors pegfilgrastim and darbepoetin alfa in support of dose-dense adjuvant chemotherapy [abstract no. 1055 plus poster]. *San Antonio Breast Cancer Society Annual Meeting*; 2004 Dec 8-11; San Antonio (TX)
- Schwartzberg LS, Tauer KT, Barry FV, et al. Dose dense chemotherapy (DDC) supported by pegfilgrastim (peg) and an erythropoietic agent (EA) in operable breast cancer [abstract no. 6035 plus poster]. *San Antonio Breast Cancer Society Annual Meeting*; 2004 Dec 8-11; San Antonio (TX)
- Pirker R, Ulsperger E, Aigner K, et al. A phase II study of pegfilgrastim to support ACE 14 chemotherapy for the treatment of subjects with small cell lung cancer (SCLC; extensive disease) [abstract no. 7216]. *Proc Am Soc Clin Oncol* 2004 Jun; 23: 667. Plus poster presented at the 40th Annual Meeting of the American Society of Clinical Oncology; 2004 Jun 5-8; New Orleans (LA)
- Lopez A, Fernandez de Sevilla A, Castaigne S, et al. Pegfilgrastim supports delivery of CHOP-R chemotherapy administered every 14 days: a randomised phase II study [abstract no. 3311]. *Blood* 2004 Nov 16; 104 (11 Pt 1): 904a. Plus poster presented at the 46th Annual Meeting of the American Society of Hematology; 2004 Dec 4-7; San Diego (CA)
- Moore TD, Patel T, Segal ML, et al. A single pegfilgrastim dose per cycle supports dose-dense (q14d) CHOP-R in patients with non-Hodgkin's lymphoma [abstract no. 2365]. *Blood* 2003 Nov 16; 102 (11 Pt 1): 641a. Plus poster presented at the 45th Annual Meeting of the American Society of Hematology; 2003 Dec 6-9; San Diego (CA)
- Engert A, Doehner H, Ho AD, et al. Pegfilgrastim supports delivery of BEACOPP chemotherapy administered every 14 days [abstract no. 1316]. *Blood* 2004 Nov 16; 104 (11 Pt 1): 370-1a. Plus poster presented at the 46th Annual Meeting of the American Society of Hematology; 2004 Dec 4-7; San Diego (CA)
- Younes A, Fayad L, Romaguera J, et al. Long-term safety and efficacy of single administration of a fixed dose pegfilgrastim (Neulasta) in inducing neutrophil count recovery after ABVD chemotherapy in patients with Hodgkin lymphoma [abstract no. 4646]. *Blood* 2004 Nov 16; 104 (11 Pt 2): 245b
- Isidori A, Tani M, Bonifazi F, et al. Phase II study of a single pegfilgrastim injection as an adjunct to chemotherapy to mobilize stem cells into the peripheral blood of a pretreated lymphoma patients. *Haematologica* 2005; 90: 225-31
- Willis F, Pettengell R, Woll PJ, et al. Pegfilgrastim alone successfully mobilizes peripheral CD34+ cells in chemotherapy naive subjects with solid tumours: initial results of a phase 1-2 study [abstract no. 2687.5]. *Blood* 2004 Nov 16; 104 (11 Pt 1): 734a. Plus poster presented at the 46th Annual Meeting of the American Society of Hematology; 2004 Dec 4-7; San Diego (CA)
- Steidl U, Fenk R, Bruns I, et al. Successful transplantation of peripheral blood stem cells mobilized by chemotherapy and a single dose of pegylated G-CSF in patients with multiple myeloma. *Bone Marrow Transplant* 2005; 35: 33-6
- Jagasia M, Greer J, Kassim A, et al. Pegylated filgrastim after high dose chemotherapy and autologous stem cell transplant [abstract no. 6639]. *Proc Am Soc Clin Oncol* 2004 Jun; 23: 590. Plus poster presented at the 40th Annual Meeting of the American Society of Clinical Oncology; 2004 Jun 5-8; New Orleans (LA)
- Amgen Inc. Neulasta® (pegfilgrastim) prescribing information. 2005 Jan
- Kubista E, Glaspy J, Holmes FA, et al. Bone pain associated with once-per-cycle pegfilgrastim is similar to daily filgrastim in patients with breast cancer. *Clin Breast Cancer* 2003 Feb; 3 (6): 391-8
- Green MD, Lu ZJ. Once-per-cycle fixed-dose administration of pegfilgrastim reduced resource utilization and costs compared with daily filgrastim in the prevention of chemotherapy-induced neutropenia [abstract no. PCN4]. *Value Health* 2002 Nov; 5: 537-8
- Heckinger E, Lee J, Calhoun E, et al. Cost minimization analysis of filgrastim versus pegfilgrastim for stage II-IV breast cancer patients receiving chemotherapy: assessments based on third-party and societal perspectives [abstract no. 2116]. *Proc Am Soc Clin Oncol* 2003 May 31; 22: 526
- Suzcs TD, Standaert B, Lu ZJ. Analysis on cost difference between daily filgrastim and once per cycle pegfilgrastim for prophylaxis against chemotherapy-induced neutropenia in France and Germany. *Value Health* 2003 Nov; 6 (6): 732
- Lyman GH, Kuderer NM, Crawford J, et al. Economic impact of pegfilgrastim use based on the risk of febrile neutropenia in NHL patients treated with CHOP [abstract no. 2384]. *Proc Am Soc Clin Oncol* 2003 May 31; 22: 593
- Eldar-Lissai A, Cosler L, Lyman GH. Economic analysis of prophylactic pegfilgrastim in cancer patients [abstract no. 2212]. *Blood* 2004 Nov 16; 104 (11 Pt 1): 608a. Plus poster presented at the 46th Annual Meeting of the American Society of Hematology; 2004 Dec 4-7; San Diego (CA)

Correspondence: *James E. Frampton*, Adis International Inc., 770 Township Line Road, Suite 300, Yardley, PA 19067, USA.
E-mail: demail@adis.com