

CrossMark
click for updatesCite this: *RSC Adv.*, 2015, 5, 87221

Heterogeneous Cu/OMS-2 as an efficient catalyst for the synthesis of tetrasubstituted 1,4-enediones and 4*H*-pyrido[1,2-*a*]-pyrimidin-4-ones†

Jinqi Zhang,^{ac} Xu Meng,^{ab} Chaoying Yu,^a Gexin Chen^a and Peiqing Zhao^{*ab}

An efficient Cu/OMS-2-catalyzed oxidative heterogeneous protocol has been developed for the synthesis of tetrasubstituted unsymmetrical 1,4-enediones and 4*H*-pyrido[1,2-*a*]-pyrimidin-4-ones from 1,4-enediones and amines or 2-aminopyridines. The present catalytic system employs air as the terminal oxidant and tolerates a broad range of substrates using low loading copper (1.3 wt%). The Cu/OMS-2 was characterized by various methods, including XRD, XPS, BET as well as TEM/SEM, and the results of H₂-TPR shows that the redox ability of the catalyst was improved by the combination of Cu and OMS-2. Moreover, the catalyst is recyclable.

Received 27th August 2015
Accepted 5th October 2015

DOI: 10.1039/c5ra17351e

www.rsc.org/advances

Introduction

From economic and industrial points of views, heterogeneous catalysis is more attractive because of catalyst recyclability, handling and easy separation of catalyst from the products.^{1,2} In recent decades, manganese oxide octahedral molecular sieves (OMS-2) have gained attention mainly due to their unique nature in heterogeneous catalysis. OMS-2 is a microporous material with a composition of KMn₈O₁₆·*n*H₂O. The structure of it contains 4.6 × 4.6 Å tunnels that arise from the 2 × 2 arrangement of octahedron and the average oxidation state of Mn is 3.8 with the presence of Mn⁴⁺, Mn³⁺, and Mn²⁺ ions in the framework.³ Consequently, the mixed valent, crystalline and semi-conductive properties make OMS-2 have a wide range of applications.⁴

Our group has been committed to studying OMS-2-based heterogeneous catalysis and found that solid supported catalyst Cu/OMS-2 with the low-loading copper has the interaction of electron transfer between metal catalytic species and support.⁵ Based on the previous reports, it is believed that the catalytic ability of Cu/OMS-2 can be significantly improved by the electronic interaction, which might facilitate organic transformations and make them proceed under a low-energy pathway.^{6,7} Therefore, the development of efficient heterogeneous protocol for organic transformation using Cu/OMS-2 is appreciated.

In recent years, the 1,4-enedione skeleton has received great attention since it widely presents in bioactive natural products and pharmaceutical compounds (Fig. 1).⁸ More importantly, 1,4-enediones are significant synthetic drug precursors for further elaboration.⁹ Thus, numerous synthetic methods have been established to produce this unit.¹⁰ Nevertheless, to the best of our knowledge, only Wu's group reported a homogeneous method promoted by Cu(OAc)₂ for the synthesis of tetrasubstituted unsymmetrical 1,4-enediones in 2013.¹¹ Based on the above works and our previous studies,¹² we herein describe a heterogeneous oxidative method for the




Fig. 1 Natural products and pharmaceutical compounds containing 1,4-enedione motifs.

^aState Key Laboratory for Oxo Synthesis and Selective Oxidation, Lanzhou Institute of Chemical Physics (LICP), Chinese Academy of Sciences, Lanzhou 730000, China. E-mail: zhaopq@licp.cas.cn; Fax: +86 931 827700

^bState Key Laboratory for Oxo Synthesis and Selective Oxidation, Suzhou Research Institute of LICP, Lanzhou Institute of Chemical Physics (LICP), Chinese Academy of Sciences, Lanzhou 730000, China

^cUniversity of Chinese Academy of Sciences, Beijing 100049, China

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c5ra17351e

synthesis of tetrasubstituted unsymmetrical 1,4-enediones using Cu/OMS-2 as catalyst under air.

Experimental section

Preparation of OMS-2 (ref. 4a)

5.89 g of KMnO_4 in 100 mL of deionized water was added to a solution of 8.8 g of $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ in 30 mL of deionized water and 3 mL concentrated HNO_3 . The solution was refluxed at 100 °C for 24 hours, and the product was filtered, washed, and dried at 120 °C for 8 hours. Finally, the dry OMS-2 was calcined in a muffle furnace at 350 °C for 2 hours. Then, the black powder OMS-2 was obtained.

Preparation of Cu/OMS-2 (ref. 5)

Support OMS-2 (2 g) was added to a 50 mL round-bottom flask. A solution of $\text{Cu}(\text{NO}_3)_2 \cdot 3\text{H}_2\text{O}$ (0.15 g) in deionized water (10 mL) was added to OMS-2, and additional deionized water (10 mL) was added to wash down the sides of the flask. Then the flask was submerged into an ultrasound bath for 3 hours at room temperature and stirred for further 20 hours at room temperature. After that, the water was distilled under reduced pressure on a rotary evaporator at 80 °C for more than 2 hours. Finally, the black powder was dried into an oven at 110 °C for 4 hours followed by calcination at 350 °C for 2 hours. The Inductive Coupled Plasma Optical Emission Spectrum (ICP-OES) showed Cu content is 1.3 wt%.

General procedure for Cu/OMS-2-catalyzed synthesis of tetrasubstituted 1,4-enediones 3

Cu/OMS-2 (30 mg, 2 mol%), 1,4-enediones **1** (0.3 mmol), *N*-substituted nucleophiles **2** (0.6 mmol), DMSO (1.8 mL) were added to a flask with a stirring bar. The flask was stirred at 80 °C for 12 hours under air. After cooling to room temperature, the mixture was diluted with ethyl acetate and filtered. The filtrate was removed under reduced pressure to get the crude product, which was further purified by silica gel chromatography (petroleum/ethyl acetate = 4/1 as eluent) to yield corresponding product. The identity and purity of the products was confirmed by ^1H and ^{13}C NMR spectroscopic analysis.

General procedure for Cu/OMS-2-catalyzed synthesis of 4H-pyrido[1,2-*a*]-pyrimidin-4-ones 5

Cu/OMS-2 (30 mg, 2 mol%), 1,4-enediones **1** (0.3 mmol), 2-aminopyridines **4** (0.6 mmol), DMSO (1.8 mL) were added to a flask with a stirring bar. The flask was stirred at 80 °C for 12 hours under air. After cooling to room temperature, the mixture was diluted with ethyl acetate and filtered. The filtrate was removed under reduced pressure to get the crude product, which was further purified by silica gel chromatography (petroleum/ethyl acetate = 2/1 as eluent) to yield corresponding product. The identity and purity of the products was confirmed by ^1H and ^{13}C NMR spectroscopic analysis.

Catalyst durability

The Cu/OMS-2 catalyst was isolated by centrifugation after each run of reactions, then washed by ethanol and water. After drying at 110 °C for 5 hours, the catalyst was reused with fresh charge of solvent and reactant for subsequent reaction under the same conditions.

Characterization

The specific surface areas (S_{BET}) of the catalysts were measured from a multipoint Brunauer–Emmett–Teller (BET) analysis of the nitrogen adsorption isotherms at 77 K recorded on a Quantachrome Autosorb-1 apparatus. The morphologies of the samples were characterized by a TF20 transmission electron microscope (TEM) and SM-5600LV scanning electron microscope (SEM). The crystal phase and composition of catalysts were determined by power X-ray diffraction (XRD) using a X-Pert PRO X-ray diffractometer with Cu K α radiation in the 2θ ranges of 10–90°. The surface properties of the catalyst were analyzed by a Thermo Scientific ESCALAB 250Xi X-ray photoelectron spectrometer (XPS). The reducibility of the catalysts was measured by the hydrogen temperature-programmed reduction (H_2 -TPR) technique. 50 mg of catalyst was placed in a quartz reactor that was connected to a TPR apparatus and the reactor was heated from r.t. to 550 °C with a heating rate of 10 °C min^{-1} . The reducing atmosphere was the mixture of H_2 and N_2 with a total flow rate of 30 mL min^{-1} and the amount of H_2 uptake during the reduction was measured by a thermal conductivity detector (TCD).

Results and discussion

The catalyst was synthesized by wet impregnation method. The BET surface areas and porosities of Cu/OMS-2 were determined by N_2 adsorption–desorption at 77 K and the results showed that the BET surface area is 127 $\text{m}^2 \text{g}^{-1}$, pore volume is 0.48 $\text{cm}^3 \text{g}^{-1}$ and pore size is 137.2 Å (Table S1, see ESI†).

The morphology of Cu/OMS-2 was inspected by TEM and SEM (Fig. 2), the images show that the synthetic material is uniform nanorod with diameters of about 8 nm and lengths of 70–120 nm.¹³

Fig. 3 plots the XPD patterns of OMS-2 and Cu/OMS-2, the results show that no signals caused by copper oxide or copper metal (cluster) were observed and the XRD pattern of Cu/OMS-2 was identical to that of support OMS-2,¹⁴ which further illustrated copper oxide is highly dispersed on support OMS-2.




Fig. 2 TEM (a) and SEM (b) image of Cu/OMS-2.




Fig. 3 XRD patterns of (a) OMS-2, (b) Cu/OMS-2 and (c) Cu/OMS-2 after one cycle.

Besides, the structure of reused catalyst was not altered (see below for the test of catalyst recyclability).

To explore the surface properties of the catalyst, Cu/OMS-2 was characterized by XPS (Fig. S2, see ESI†). The Mn 2p spectrum shows that Mn 2p_{1/2} peak and Mn 2p_{3/2} peak are centred at 654.0 eV and 642.4 eV respectively, indicating this sample contains both Mn³⁺ and Mn⁴⁺.¹⁵ The Cu 2p spectrum shows that the binding energies of Cu 2p_{1/2} and Cu 2p_{3/2} are about 954.0 eV and 934.3 eV, and the shake-up satellite peaks of Cu 2p_{1/2} and Cu 2p_{3/2} are at 962.1 eV and at 944.7 eV, respectively, which demonstrates that Cu²⁺ species are existed on the surface of the catalyst.^{15,19a}

The redox ability of the catalyst was studied by H₂-temperature programmed reduction (H₂-TPR) and the profiles are presented in Fig. 4. For the support OMS-2, the overlapping peaks (marked as α, β) from 369 °C to 390 °C are ascribed to the reduction of MnO₂ to MnO.^{16,18} Compared with OMS-2, the reduction peaks of Cu/OMS-2 shift to low temperature, suggesting that the existence of copper species improves the reducibility of manganese oxide by a hydrogen spillover effect.^{17,19a} More importantly, the reduction peaks α' and β' of Cu/OMS-2 became obvious, which




Fig. 4 H₂-TPR profiles of OMS-2 and Cu/OMS-2.

probably is caused by the enhanced mobility of oxygen.^{15,18} These results all confirm the interaction between the catalytic metal copper and the support OMS-2. Moreover, the extra peaks γ and δ appeared in the TPR profile of the Cu/OMS-2 are related to the reduction of CuO to Cu. The γ peak observed at 210 °C corresponds to the reduction of the finely dispersed CuO particles and the higher-temperature peak δ represents the reduction of the bulk CuO particles.¹⁹

Next, we employed 1,4-enedione **1a** and pyrazole **2a** as the model substrates to optimize the reaction conditions (Table 1). In the first attempt, the reaction was run using OMS-2 as the catalyst under air at room temperature, but no desired product was observed (Table 1, entry 1). To our delight, 58% yield of the target product was obtained when Cu/OMS-2 was used as the catalyst (Table 1, entry 2). Then, Cu(OH)_x/OMS-2 was examined in the reaction and it did not better the reaction (Table 1, entry 3). Besides, various supports were tested, such as ATP (attapulgit), C (charcoal) and Al₂O₃, and it was found that OMS-2 is optimal for this reaction (Table 1, entries 4–6). Importantly, a very low yield of **3aa** was observed when the reaction temperature was increased from r.t. to 120 °C with the Cu/Al₂O₃ as the catalyst. These observations perhaps indicate that Cu/Al₂O₃ is not able to lower the redox energy barriers while high temperature

Table 1 Optimization of the reaction conditions^a

| | 1a | 2a | | 3aa | |
|-----------------|-----------------------------------|---------------------------------|------------|----------------|------------------------|
| Entry | Cat. | Solvent | Temp. (°C) | Oxidant | Yield ^b (%) |
| 1 | OMS-2 | DMSO | r.t. | Air | 0 |
| 2 ^c | Cu/OMS-2 | DMSO | r.t. | Air | 58 |
| 3 | Cu(OH) _x /OMS-2 | DMSO | r.t. | Air | 0 |
| 4 | Cu/ATP | DMSO | r.t. | Air | 0 |
| 5 | Cu/C | DMSO | r.t. | Air | 0 |
| 6 | Cu/Al ₂ O ₃ | DMSO | r.t. | Air | 0 |
| 7 | Cu/Al ₂ O ₃ | DMSO | 120 | Air | 7 |
| 8 ^c | Cu/OMS-2 | DMF | r.t. | Air | 0 |
| 9 ^c | Cu/OMS-2 | DCB | r.t. | Air | 0 |
| 10 ^c | Cu/OMS-2 | CH ₃ NO ₂ | r.t. | Air | 36 |
| 11 ^c | Cu/OMS-2 | Toluene | r.t. | Air | 43 |
| 12 ^c | Cu/OMS-2 | CH ₂ Cl ₂ | r.t. | Air | 34 |
| 13 ^c | Cu/OMS-2 | THF | r.t. | Air | 27 |
| 14 ^c | Cu/OMS-2 | DMSO | 40 | Air | 60 |
| 15 ^c | Cu/OMS-2 | DMSO | 80 | Air | 72 |
| 16 ^c | Cu/OMS-2 | DMSO | 100 | Air | 54 |
| 17 ^c | Cu/OMS-2 | DMSO | 80 | O ₂ | 75 |
| 18 ^c | Cu/OMS-2 | DMSO | 80 | N ₂ | <5 |
| 19 ^d | CuO + OMS-2 | DMSO | 80 | Air | 15 |
| 20 | CuO (1 equiv.) | DMSO | 80 | N ₂ | 56 |
| 21 | CuO (0.1 equiv.) | DMSO | 80 | Air | 41 |

^a Unless otherwise specified, all reactions were carried out using **1a** (0.3 mmol), **2a** (0.6 mmol) and catalysts (30 mg) in 1.8 mL solvent under air for 12 h. ^b Isolated yields. ^c Cu content: 1.3 wt%. ^d CuO (2 mol%), OMS-2 (30 mg).


made the reaction work to some extent by overcoming the redox energy barriers (Table 1, entry 7). Next, we tried different solvents, which shows the reaction did not proceed in DMF (dimethyl formamide) and DCB (dichlorobenzene) (Table 1, entries 8 and 9). DMSO (dimethylsulfoxide) led to the best result and other solvents, like nitromethane, toluene, dichloromethane and THF (tetrahydrofuran), could give the desired product to some extent (Table 1, entries 10–13). Subsequently, the reaction temperature was optimized, and we found that 80 °C is the best choice (Table 1, entries 14–16). More experiments demonstrated that using O₂ as the oxidant did not better the reaction obviously and only trace amount of **3aa** was observed under N₂ (Table 1, entries 17 and 18). Finally, the catalytic activity of physical mixture of bulk CuO and OMS-2 was much lower than this of Cu/OMS-2, which indicates highly dispersed Cu species played a crucial role in the reaction (Table 1, entry 19). Furthermore, without the external oxidant, equivalent unsupported CuO could also give moderate yield (Table 1, entry 20), while catalytic amount of CuO gave the poor yield of **3aa** with oxygen as the external oxidant (Table 1, entry 21). These observations indicate that the reduced copper species cannot be reoxidized by O₂ due to the high-energy barrier of direct reoxidation without the help of OMS-2.

To examine whether the observed catalytic activity is generated from the solid catalyst Cu/OMS-2 or the leached copper species, the hot filtration experiment was carried out with 1,4-enedione **1g** and pyrazole **2a** as substrates (specific details see ESI†). The catalyst was removed after the reaction was run for 3 hours, then the reaction was kept on, but the yield did not increase. At the same time, the filtrate was analyzed by inductively coupled plasma-atomic emission spectrometry (ICP-AES), and the result displayed that the

concentration of copper species in the solution is 0.45 ppm. Hence, it is concluded that the copper species leached from the catalyst is catalytically inactive and the catalysis is truly heterogeneous. Additionally, the life span of the catalyst is another aspect of the heterogeneous catalysis. The reactions between 1,4-enedione **1g** and pyrazole **2a** was used as the model reactants to test the recyclability of Cu/OMS-2. The catalyst was recovered by filtration and centrifugation, and results showed that the catalyst can recycle at least 3 times with slightly decrease of catalytic activity (Fig. 5). Furthermore, the result of TEM indicated that the used Cu/OMS-2 was intact in morphology, implying the stability of Cu/OMS-2 (Fig. S1, see ESI†). From the results of the hot filtration experiment (specific details see ESI†), we can speculate that the loss of the catalytic activity most likely results from the leaching of copper from the support.

With the optimized conditions in hand, we investigated the scope of this amination catalyzed by Cu/OMS-2 (Table 2). Generally, this catalytic system could tolerate a large range of substrates and lead to moderate to good yields of corresponding products. In terms of R¹ substituents, good yields were gained with various substituted groups, such as –Me, –OMe and halogenated substituents (Table 2, entries 1–6). Delightedly, when R³ was alkoxy groups (–OMe, –OEt), the reaction proceeded

Table 2 Scope of the reactions between 1,4-enediones **1** and pyrazole **2a**^a

|  | | | | | | |
|--|-----------|-----------------------|----------------|----------------|------------|-------------------------------------|
| Entry | 1 | R ¹ | R ² | R ³ | 3 | Yield ^b (%) |
| 1 | 1a | Ph | Ph | Ph | 3aa | 72 |
| 2 | 1b | 4-Me-Ph | Ph | Ph | 3ba | 57 |
| 3 | 1c | 4-OMe-Ph | Ph | Ph | 3ca | 68 |
| 4 | 1d | 4-Cl-Ph | Ph | Ph | 3da | 56 |
| 5 | 1e | 4-Br-Ph | Ph | Ph | 3ea | 59 |
| 6 | 1f | 4-F-Ph | Ph | Ph | 3fa | 75 |
| 7 ^c | 1g | Ph | Ph | OEt | 3ga | 80 (<i>E</i> : <i>Z</i> = 66 : 34) |
| 8 ^c | 1h | 4-Me-Ph | Ph | OEt | 3ha | 78 (<i>E</i> : <i>Z</i> = 77 : 23) |
| 9 ^c | 1i | 4-OMe-Ph | Ph | OEt | 3ia | 87 (<i>E</i> : <i>Z</i> = 67 : 33) |
| 10 ^c | 1j | 4-NO ₂ -Ph | Ph | OEt | 3ja | 67 (<i>E</i> : <i>Z</i> = 87 : 13) |
| 11 ^c | 1k | 4-CF ₃ -Ph | Ph | OEt | 3ka | 88 (<i>E</i> : <i>Z</i> = 90 : 10) |
| 12 ^c | 1l | 2-Naphthyl | Ph | OEt | 3la | 90 (<i>E</i> : <i>Z</i> = 81 : 19) |
| 13 ^c | 1m | 4-Cl-Ph | Ph | OEt | 3ma | 84 (<i>E</i> : <i>Z</i> = 88 : 12) |
| 14 ^c | 1n | 4-Br-Ph | Ph | OEt | 3na | 87 (<i>E</i> : <i>Z</i> = 89 : 11) |
| 15 ^c | 1o | 4-F-Ph | Ph | OEt | 3oa | 81 (<i>E</i> : <i>Z</i> = 88 : 12) |
| 16 ^c | 1p | Ph | 4-OMe-Ph | OEt | 3pa | 78 (<i>E</i> : <i>Z</i> = 63 : 37) |
| 17 ^c | 1q | Ph | 4-F-Ph | OEt | 3qa | 86 (<i>E</i> : <i>Z</i> = 86 : 14) |
| 18 ^c | 1r | Ph | 4-OMe-Ph | OMe | 3ra | 91 (<i>E</i> : <i>Z</i> = 74 : 26) |
| 19 ^c | 1s | Ph | 4-Cl-Ph | OMe | 3sa | 83 (<i>E</i> : <i>Z</i> = 92 : 8) |

^a Unless otherwise specified, all reactions were carried out using **1** (0.3 mmol), **2a** (0.6 mmol) and Cu/OMS-2 (30 mg, 2 mol%) in 1.8 mL solvent under air for 12 h. ^b Isolated yields. ^c A mixture of *E/Z*-isomers was obtained, *E* : *Z* ratio determined by ¹H NMR.




Fig. 5 The recyclability of Cu/OMS-2. Reaction conditions: **1g** (0.3 mmol), **2a** (0.6 mmol) and Cu/OMS-2 (30 mg, 2 mol%) in 1.8 mL DMSO under air for 12 h.

smoothly with good to excellent yields of tetrasubstituted unsymmetrical 1,4-enediones (Table 2, entries 7–19).












Next, the scope of *N*-substituted nucleophiles was tested under the standard conditions with 1,4-enedione **1g** as the model substrate (Table 3). Anilines proceeded very well in this system and moderate to good yields of corresponding products were observed with electron-neutral (**3gb**), electron-donating (**3gc**, **3gd**), electron-withdrawing (**3ge**, **3gf**, **3gg**) substituents. Besides, *N*-methylaniline (**3gh**) and 6-chloropyridin-2-amine (**3gi**) could also be used as suited substrates. To our delight, aliphatic *N*-nucleophiles were also tolerated in this reaction and gave moderate yields of desired products (**3gj**, **3gk**) with excellent *E/Z* ratio.

Surprisingly, a cyclized product 4*H*-pyrido[1,2-*a*]-pyrimidin-4-one was obtained in 91% yield when 1,4-enedione **1g** reacted with 2-aminopyridine under our standard reaction conditions. It is believed after 2-aminopyridine and 1,4-enedione undergo the aza-Michael addition and aerobic dehydrogenation, the nitrogen in pyridine ring further undergoes intramolecular amidation under the present conditions to generate the heterocycles **5**. It is an encouraging result because 4*H*-pyrido

[1,2-*a*]-pyrimidin-4-ones are a class of nitrogen-containing heterocycles with pharmacological activities, and they extensively exist in natural products and pharmaceutical drugs.²⁰ So far, a number of synthetic protocols for construction of this skeleton have been reported.²¹ However, as far as we know, this is the first example to synthesize 4*H*-pyrido[1,2-*a*]-pyrimidin-4-ones under the heterogeneous conditions. Therefore, the scope of cyclizations was examined with a series of different 1,4-enediones **1** and various substituted 2-aminopyridines **4** under the present reaction conditions (Table 4). Many 1,4-enediones with different substituted groups were employed as the substrates and good to excellent yields of desired products were obtained. We also examined the scope of 2-aminopyridines in this reaction. Specifically, 3-methylpyridin-2-amine afforded the corresponding product in good yield (**5gb**). However, 6-chloropyridin-2-amine provided the product **3gi** via C–N bond-forming instead of cyclized product **5gc** because of the steric hindrance.












Based on the reported literatures and our results, we proposed a possible mechanism for these two kinds of reactions respectively. The first amination reaction consists of aza-Michael addition reaction between amine **2a** and 1,4-enedione **1a** with assistance of Cu and a following supported copper species catalyzed aerobic oxidative dehydrogenation of intermediate **A** under air (Fig. 6).^{14,22} For the cyclization, the first two steps are same as these of the amination. After

Table 3 Scope of the reactions between 1,4-enedione **1g** and *N*-substituted nucleophiles **2**^a

|  | | |
|---|---|---|
| 1g | 2 | 3 |
|  |  |  |
| 3gb , 76% (<i>E/Z</i> =56:44) | 3gc , 53% (<i>E/Z</i> =57:43) | 3gd , 62% (<i>E/Z</i> =60:40) |
|  |  |  |
| 3ge , 70% (<i>E/Z</i> =60:40) | 3gf , 71% (<i>E/Z</i> =52:48) | 3gg , 57% (<i>E/Z</i> =47:53) |
|  |  |  |
| 3gh , 31% (<i>E/Z</i> =91:9) | 3gi , 64% (<i>E/Z</i> =56:44) | 3gj , 53% (<i>E/Z</i> >99:1) |
| |  | |
| | 3gk , 36% (<i>E/Z</i> >99:1) | |

^a Unless otherwise specified, all reactions were carried out using **1g** (0.3 mmol), **2** (0.6 mmol) and Cu/OMS-2 (30 mg, 2 mol%) in 1.8 mL DMSO under air for 12 h, isolated yields. *E/Z* ratio determined by ¹H NMR.

Table 4 Scope of the synthesis of 4*H*-pyrido[1,2-*a*]-pyrimidin-4-ones **5**^a

|  | | |
|--|---|---|
| 1 | 4 | 5 |
|  |  |  |
| 5ga , 89% | 5ha , 87% | 5ia , 86% |
|  |  |  |
| 5ka , 83% | 5la , 93% | 5ma , 81% |
|  |  |  |
| 5pa , 91% | 5qa , 77% | 5gb , 74% |
| |  | |
| | 5gc , 0% | |

^a Unless otherwise specified, all reactions were carried out using **1** (0.3 mmol), **4** (0.6 mmol) and Cu/OMS-2 (30 mg, 2 mol%) in 1.8 mL DMSO under air for 12 h, isolated yields.




Fig. 6 The proposed mechanism of the amination.




Fig. 7 The proposed mechanism of the cyclization.

that, the *E*-isomer amination intermediate (*E*)-C produces its *Z*-isomer amination intermediate (*Z*)-C through isomerization, finally the cyclized product **5ga** is formed by supported copper species catalyzed intramolecular amidation (Fig. 7).^{21f} In the two kinds of reaction mechanisms, the reduced copper species is reoxidized by OMS-2, and the reduced OMS-2 is reoxidized by O₂ of air to finish the whole redox cycle.

Conclusions

In summary, we have developed an efficient heterogeneous strategy for the synthesis of tetrasubstituted 1,4-enediones and 4*H*-pyrido[1,2-*a*]-pyrimidin-4-ones catalyzed by Cu/OMS-2. The excellent catalytic ability of Cu/OMS-2 is attributed to OMS-2

which accelerates the reoxidation of reduced copper species through rapid electron transfer. This superiority might allow it to apply to other oxidative processes and reactions. Further applications of this strategy are underway in our laboratory.

Acknowledgements

We gratefully acknowledge the National Natural Science Foundation of China (Grant No. 21403256, 21573261) and the Suzhou Industrial Technology and Innovation Project (Grant No. SYG201531).

Notes and references

- (a) M. Pagliaro, V. Pandarus, R. Ciriminna, F. Béland and P. D. Carà, *ChemCatChem*, 2012, **4**, 432; (b) M. J. Climent, A. Corma and S. Iborra, *RSC Adv.*, 2012, **2**, 16; (c) R. A. Sheldon and H. V. Bekkum, *Fine Chemicals through Heterogeneous Catalysis*, Wiley-VCH, Weinheim, 2001.
- For selected papers on heterogeneous catalysis, see: (a) X. Meng, X. Xu, T. Gao and B. Chen, *Eur. J. Org. Chem.*, 2010, 5409; (b) J. B. Bharate, S. K. Guru, S. K. Jain, S. Meena, P. P. Singh, S. Bhushan, B. Singh, S. B. Bharate and R. A. Vishwakarma, *RSC Adv.*, 2013, **3**, 20869; (c) F. Nemat, A. Elhampour, H. Farrokhi and M. B. Natanzi, *Catal. Commun.*, 2015, **66**, 15; (d) B. Yang, Z. Mao, X. Zhu and Y. Wan, *Catal. Commun.*, 2015, **60**, 92; (e) A. T. Nguyen, L. T. Pham, N. T. S. Phan and T. Truong, *Catal. Sci. Technol.*, 2014, **4**, 4281; (f) L. Zhang, P. Li, C. Liu, J. Yang, M. Wang and L. Wang, *Catal. Sci. Technol.*, 2014, **4**, 1979.
- J. R. Kona, C. K. Kingöndu, A. R. Howell and S. L. Suib, *ChemCatChem*, 2014, **6**, 749.
- (a) Y. C. Son, V. D. Makwana, A. R. Howell and S. L. Suib, *Angew. Chem.*, 2001, **113**, 4410; *Angew. Chem., Int. Ed.*, 2001, **40**, 4280; (b) V. V. Krishnan and S. L. Suib, *J. Catal.*, 1999, **184**, 305; (c) S. Sithambaram, Y. Ding, W. Li, X. Shen, F. Gaenzler and S. L. Suib, *Green Chem.*, 2008, **10**, 1029; (d) R. Kumar, S. Sithambaram and S. L. Suib, *J. Catal.*, 2009, **262**, 304; (e) K. Yamaguchi, H. Kobayashi, T. Oishi and N. Mizuno, *Angew. Chem.*, 2012, **124**, 559; *Angew. Chem., Int. Ed.*, 2012, **51**, 544; (f) K. Yamaguchi, Y. Wang and N. Mizuno, *ChemCatChem*, 2013, **5**, 2835; (g) K. Yamaguchi, H. Kobayashi, Y. Wang, T. Oishi, Y. Ogasawara and N. Mizuno, *Catal. Sci. Technol.*, 2013, **3**, 318; (h) Y. Wang, H. Kobayashi, K. Yamaguchi and N. Mizuno, *Chem. Commun.*, 2012, **48**, 2642; (i) X. Jin, K. Yamaguchi and N. Mizuno, *RSC Adv.*, 2014, **4**, 34712; (j) X. Jin, K. Yamaguchi and N. Mizuno, *Angew. Chem.*, 2014, **126**, 465; *Angew. Chem., Int. Ed.*, 2014, **53**, 455; (k) K. Yamaguchi, Y. Wang, T. Oishi, Y. Kuroda and N. Mizuno, *Angew. Chem.*, 2013, **125**, 5737; *Angew. Chem., Int. Ed.*, 2013, **52**, 5627; (l) X. Meng, J. Zhang, G. Chen, B. Chen and P. Zhao, *Catal. Commun.*, 2015, **69**, 239.
- X. Meng, C. Yu, G. Chen and P. Zhao, *Catal. Sci. Technol.*, 2015, **5**, 372.
- T. Oishi, K. Yamaguchi and N. Mizuno, *ACS Catal.*, 2011, **1**, 1351.

- 7 (a) B. P. Babu, X. Meng and J. E. Bäckvall, *Chem.–Eur. J.*, 2013, **19**, 4140; (b) Y. Endo and J. E. Bäckvall, *Chem.–Eur. J.*, 2011, **17**, 12596; (c) B. P. Babu, Y. Endo and J. E. Bäckvall, *Chem.–Eur. J.*, 2012, **18**, 11524; (d) M. Johansson, B. W. Purse, O. Terasaki and J. E. Bäckvall, *Adv. Synth. Catal.*, 2008, **350**, 1807; (e) N. Gigant and J. E. Bäckvall, *Org. Lett.*, 2014, **16**, 1664; (f) N. Gigant and J. E. Bäckvall, *Chem.–Eur. J.*, 2013, **19**, 10799; (g) X. Meng, C. Li, B. Han, T. Wang and B. Chen, *Tetrahedron*, 2010, **66**, 4029; (h) J. Piera and J. E. Bäckvall, *Angew. Chem.*, 2008, **120**, 3558; *Angew. Chem., Int. Ed.*, 2008, **47**, 3506.
- 8 (a) X. F. Niu, X. Liu, L. Pan and L. Qi, *Fitoterapia*, 2011, **82**, 960; (b) A. Aiello, E. Fattorusso, S. Magno, M. Menna and M. Pansini, *J. Nat. Prod.*, 1991, **54**, 281; (c) P. Cai, F. Kong, M. E. Ruppen, G. Glasier and G. T. Carter, *J. Nat. Prod.*, 2005, **68**, 1736; (d) H. Abou-Gazar, E. Bedir, S. Takamatsu, D. Ferreira and I. A. Khan, *Phytochemistry*, 2004, **65**, 2499; (e) F. Lv, M. Xu, Z. Deng, N. J. de Voogd, R. W. M. van Soest, P. Proksch and W. Lin, *J. Nat. Prod.*, 2008, **71**, 1738; (f) R. Ballini and P. Astolfi, *Liebigs Ann.*, 1996, **11**, 1879; (g) M. Fouad, R. A. Edrada, R. Ebel, V. Wray, W. E. G. Müller, W. H. Lin and P. Proksch, *J. Nat. Prod.*, 2006, **69**, 211.
- 9 (a) Y. Yang, M. Gao, C. Deng, D. X. Zhang, L. M. Wu, W. M. Shu and A. X. Wu, *Tetrahedron*, 2012, **68**, 6257; (b) Y. Yang, M. Gao, D. X. Zhang, L. M. Wu, W. M. Shu and A. X. Wu, *Tetrahedron*, 2012, **68**, 7338; (c) G. Yin, Z. Wang, A. Chen, M. Gao, A. Wu and Y. J. Pan, *Org. Chem.*, 2008, **73**, 3377; (d) Y. Yang, M. Gao, L. M. Wu, C. Deng, D. X. Zhang, Y. Gao, Y. P. Zhu and A. X. Wu, *Tetrahedron*, 2011, **67**, 5142; (e) M. Gao, Y. Yang, Y. D. Wu, C. Deng, W. M. Shu, D. X. Zhang, L. P. Cao, N. F. She and A. X. Wu, *Org. Lett.*, 2010, **12**, 4026; (f) W. M. Shu, Y. Yang, D. X. Zhang, L. M. Wu, Y. P. Zhu, G. D. Yin and A. X. Wu, *Org. Lett.*, 2013, **15**, 456; (g) D. Zhang, Y. Yang, M. Gao, W. Shu, L. Wu, Y. Zhu and A. Wu, *Tetrahedron*, 2013, **69**, 1849; (h) L. Wu, Y. Yang, M. Gao, D. Zhang, W. Shu, Y. Zhu and A. Wu, *Synlett*, 2012, **23**, 2137; (i) Q. Gao, Y. Zhu, M. Lian, M. Liu, J. Yuan, G. Yin and A. Wu, *J. Org. Chem.*, 2012, **77**, 9865.
- 10 (a) M. K. Eberhardt, *J. Org. Chem.*, 1993, **58**, 497; (b) J. Q. Yu and E. J. Corey, *J. Am. Chem. Soc.*, 2003, **125**, 3232; (c) M. Gao, Y. Yang, Y. D. Wu, C. Deng, L. P. Cao, X. G. Meng and A. X. Wu, *Org. Lett.*, 2010, **12**, 1856; (d) G. Yin, B. Zhou, X. Meng, A. Wu and Y. Pan, *Org. Lett.*, 2006, **8**, 2245; (e) Z. Wang, G. Yin, A. Chen, S. Hu and A. Wu, *Synth. Commun.*, 2007, **37**, 4399; (f) B. Crone and S. F. Kirsch, *Chem. Commun.*, 2006, 764; (g) O. Prakash, A. Batra, V. Chaudhri and R. Prakash, *Tetrahedron Lett.*, 2005, **46**, 2877; (h) A. M. Echavarren, M. Perez, A. M. Castano and J. M. Cuerva, *J. Org. Chem.*, 1994, **59**, 4179; (i) S. K. Kang, Y. T. Lee and S. H. Lee, *Tetrahedron Lett.*, 1999, **40**, 3573; (j) K. A. Runcie and R. J. K. Taylor, *Chem. Commun.*, 2002, 974; (k) C. Deng, Y. Yang, M. Gao, Y. P. Zhu, A. X. Wu, J. R. Ma and G. D. Yin, *Tetrahedron*, 2012, **68**, 3828; (l) P. Wasnaire, T. de Meroode and I. E. Markó, *Chem. Commun.*, 2007, 4755; (m) A. Del Zotto, W. Baratta, G. Verardo and P. Rigo, *Eur. J. Org. Chem.*, 2000, 2795; (n) C. D. S. Lisboa, V. G. Santos, B. G. Vaz, N. C. de Lucas, M. N. Eberlin and S. J. Garden, *J. Org. Chem.*, 2011, **76**, 5264.
- 11 Y. Yang, F. Ni, W. M. Shu, S. B. Yu, M. Gao and A. X. Wu, *J. Org. Chem.*, 2013, **78**, 5418.
- 12 (a) X. Meng, Y. Wang, C. Yu and P. Zhao, *RSC Adv.*, 2014, **4**, 27301; (b) X. Meng, C. Yu and P. Zhao, *RSC Adv.*, 2014, **4**, 8612; (c) J. Liu, C. Yu, P. Zhao and G. Chen, *Appl. Surf. Sci.*, 2012, **258**, 9096; (d) C. Yu, P. Zhao, G. Chen and B. Hu, *Appl. Surf. Sci.*, 2011, **257**, 7727.
- 13 S. Sithambaram, R. Kumar, Y. C. Son and S. L. Suib, *J. Catal.*, 2008, **253**, 269.
- 14 S. Sithambaram, L. Xu, C. H. Chen, Y. Ding, R. Kumar, C. Calvert and S. L. Suib, *Catal. Today*, 2009, **140**, 162.
- 15 W. Y. Hernández, M. A. Centeno, S. Ivanova, P. Eloy, E. M. Gaigneaux and J. A. Odriozola, *Appl. Catal., B*, 2012, **123–124**, 27.
- 16 V. P. Santos, M. F. R. Pereira, J. J. M. Orfao and J. L. Figueiredo, *Appl. Catal., B*, 2010, **99**, 353.
- 17 M. R. Morales, B. P. Barbero and L. E. Cadús, *Fuel*, 2008, **87**, 1177.
- 18 Y. Yang, J. Huang, S. Zhang, S. Wang, S. Deng, B. Wang and G. Yu, *Appl. Catal., B*, 2014, **150–151**, 167.
- 19 (a) X. S. Liu, Z. N. Jin, J. Q. Lu, X. X. Wang and M. F. Luo, *Chem. Eng. J.*, 2010, **162**, 151; (b) M. F. Luo, Y. J. Zhong, X. X. Yuan and X. M. Zheng, *Appl. Catal., A*, 1997, **162**, 121; (c) M. F. Luo, Y. P. Song, J. Q. Lu, X. Y. Wang and Z. Y. Pu, *J. Phys. Chem. C*, 2007, **111**, 12686; (d) M. F. Luo, J. M. Ma, J. Q. Lu, Y. P. Song and Y. J. Wang, *J. Catal.*, 2007, **246**, 52.
- 20 (a) I. Hermecz and Z. Mészáros, *Med. Res. Rev.*, 1988, **8**, 203; (b) I. Hermecz and L. Vasvari-Debreczy, *Comprehensive Heterocyclic Chemistry III*, ed. K. Jones Execut and A. R. Katritzky, 2007, pp. 77–217; (c) C. Fenton and L. J. Scott, *CNS Drugs*, 2005, **19**, 429; (d) F. Awouters, J. Vermeire, F. Smeyers, P. Vermote, R. Van Beek and C. J. E. Niemegeers, *Drug Dev. Res.*, 1986, **8**, 95; (e) L. E. J. Kennis, F. P. Bischoff, C. J. Mertens, C. J. Love, F. A. F. van den Keybus, S. Pieters, M. Braeken, A. A. H. P. Megens and J. E. Leysen, *Bioorg. Med. Chem. Lett.*, 2000, **10**, 71; (f) R. L. Smith, R. J. Barrett and E. Sanders-Bush, *J. Pharmacol. Exp. Ther.*, 1995, **275**, 1050; (g) A. Pettersson, K. Gradin, T. Hedner and B. Persson, *Naunyn-Schmiedeberg's Arch. Pharmacol.*, 1985, **329**, 394.
- 21 (a) I. Hermecz, *Adv. Heterocycl. Chem.*, 2003, **85**, 173; (b) J. Modranka and T. Janecki, *Tetrahedron*, 2011, **67**, 9595; (c) G. Bentabed-Ababsa, S. C. S. Ely, S. Hesse, E. Nassar, F. Chevallier, T. T. Nguyen, A. Derdour and F. Mongin, *J. Org. Chem.*, 2010, **75**, 839; (d) P. Čebasěk, D. Bevk, S. Pirc, B. Stanovnik and J. Svete, *J. Comb. Chem.*, 2006, **8**, 95; (e) A. Begouin, S. Hesse, M. J. R. P. Queiroz and G. Kirsch, *Synthesis*, 2006, 2794; (f) Y. Yang, W. M. Shu, S. B. Yu, F. Ni, M. Gao and A. X. Wu, *Chem. Commun.*, 2013, **49**, 1729.
- 22 (a) O. A. Attanasi, G. Favi, P. Filippone, F. Mantellini, G. Moscatelli and F. R. Perrulli, *Org. Lett.*, 2009, **12**, 468; (b) L. W. Xu, J. W. Li, C. G. Xia, S. L. Zhou and X. X. Hu, *Synlett*, 2003, 2425; (c) D. Wang, Y. Shiraishi and T. Hirai, *Chem. Commun.*, 2011, **47**, 2673.