

Radiolysis of Ethane-1,1,1-d 3

Louis J. Stief and P. Ausloos

Citation: *The Journal of Chemical Physics* **36**, 2904 (1962); doi: 10.1063/1.1732400

View online: <http://dx.doi.org/10.1063/1.1732400>

View Table of Contents: <http://scitation.aip.org/content/aip/journal/jcp/36/11?ver=pdfcov>

Published by the AIP Publishing

Articles you may be interested in

H(D)-atom yields in the quenching of Xe ($6s\ [3/2]\ 1$) by methane, ethane, ethene, ethyne, and their deuterated isotopologues

J. Chem. Phys. **125**, 034306 (2006); 10.1063/1.2213256

Ion-Molecule Reactions in the Radiolysis of Ethane

J. Chem. Phys. **54**, 2612 (1971); 10.1063/1.1675220

Near-Infrared Spectrum of Ethane-1,1,1-d 3

J. Chem. Phys. **48**, 918 (1968); 10.1063/1.1668736

Effect of Electrical Fields and Density in the Radiolysis of Ethane

J. Chem. Phys. **42**, 343 (1965); 10.1063/1.1695697

Thermal Isomerization of Isopropyl-1,1,1-d 3 Radicals

J. Chem. Phys. **36**, 2272 (1962); 10.1063/1.1732875



NEW Special Topic Sections

NOW ONLINE

Lithium Niobate Properties and Applications:
Reviews of Emerging Trends

AIP

Applied Physics
Reviews

fortunately, there is some uncertainty in making the comparison, i.e.,

$$\bar{S}_2^{E*}(p) = \bar{S}_2^{E*}(p') + \int_p^{p'} \left(\frac{\partial \bar{V}_2^*}{\partial T} \right)_p dp, \quad (29)$$

where p' is 350 atm and p is the highest pressure at which experimental values of \bar{V}_2^* are available. Because \bar{V}_2^* was only measured to 150 atm, the evaluation of the integral in Eq. (29) involves an extrapolation of equations which are linear in $1/\bar{V}_2^*$.⁶ This extrapolation

can be made with reasonable certainty at low temperatures where $(\partial \bar{V}_2^*/\partial T)_p$ is small, but the certainty gets less as the temperature is increased. Therefore, while the agreement between Eq. (28) and experiment in Table IV is probably meaningful up to 200°C, that at 230°C is fortuitous.

ACKNOWLEDGMENTS

The author thanks W. B. Kay and W. A. Junk for their advice.

Radiolysis of Ethane-1,1,1- d_3 †

LOUIS J. STIEF† AND P. AUSLOOS

National Bureau of Standards, Washington, D. C.


(Received December 26, 1961)

The radiolysis of CH_3CD_3 has been investigated with particular emphasis on the effect of pressure, temperature, and addition of xenon on the molecular hydrogen formation. The hydrogen formed by molecular detachment was determined by irradiating CH_3CD_3 in the presence of C_2H_4 , NO , O_2 , or I_2 . The formation of molecular hydrogen in the unimolecular decomposition of the parent ion and in the decomposition of the neutral excited ethane molecule are considered. Methane formed in the presence of scavengers may be attributed to an ion-molecule reaction or to an intramolecular process similar to that proposed in the direct photolysis of ethane. Evidence is presented for occurrence of a reaction producing ethylene in addition to the one in which molecular hydrogen accompanies the formation of ethylene.

INTRODUCTION

IN recent investigations^{1,2} of the radiolysis of C_2H_6 - C_2D_6 mixtures, it has been clearly demonstrated that nearly 50% of the hydrogen is formed by molecular-elimination processes. It was also shown in the course of these investigations that a considerable fraction of the methane is formed by molecular detachment from ethane.

Recently³ similar processes have been observed in the short-wavelength photolysis of ethane. Of special interest are the results obtained in the photolysis of CH_3CD_3 , which indicated that at 1470 Å, hydrogen is mostly formed by elimination of a hydrogen molecule from a single carbon atom.



The present investigation was undertaken in order to determine (a) if similar processes occur in the

radiolysis of CH_3CD_3 , (b) whether the distribution of the molecular hydrogen isotopes can be related to the mass spectral cracking pattern, (c) whether the distribution of the isotopic methanes formed in the presence of radical scavengers is consistent with the results obtained in the C_2H_6 - C_2D_6 system, and (d) whether the distribution of the isotopic ethylenes formed in the presence of scavengers is consistent with the distribution of the corresponding hydrogen isotopes.

EXPERIMENTAL

Materials

Ethane-1,1,1- d_3 was obtained from Merck and Company, Ltd. Small amounts of methane were removed by subjecting the ethane to successive cycles of freezing, pumping, and warming. The absence of ethylene in the CH_3CD_3 was checked by gas chromatographic analysis [Perkin-Elmer No. 154, silica gel (J) column, 35°]; as little as 0.001% ethylene would have been readily detected by this method. The isotopic purity of the CH_3CD_3 is discussed in the next section. Deuteroethane and ethane were obtained from Merck and Company, Ltd. and Union Carbide and Carbon Chemicals, respectively. Both were purified of noncondensables by a vacuum distillation procedure similar to that employed for CH_3CD_3 . Ethylene was removed by gas chromatography (conditions as above),

† This research was supported by a grant from the U. S. Atomic Energy Commission; presented in part at the 139th Meeting of the American Chemical Society, St. Louis, Missouri, March 1961.

‡ National Academy of Science-National Research Council Postdoctoral Research Associate, 1960-1961; present address: Chemistry Department, The University, Sheffield 10, England.

¹ L. M. Dorfman, J. Phys. Chem. **60**, 826 (1956).

² L. M. Dorfman, J. Phys. Chem. **62**, 29 (1958).

³ H. Okabe and J. R. McNesby, J. Chem. Phys. **34**, 668 (1961).

TABLE I. Mass spectral data for CH₃CD₃.

<i>m/e</i>	Ionizing voltage						
	75 ^a	70 ^b	70	25	18	14	12
12	1.19	10.07	4.8				
13	1.20	10.85	5.2				
14	3.42	22.7	11.8	0.17			
14.5	1.62	2.0	1.2				
15	6.19	24.4	15.2	3.63	1.06		
15.5	3.14	2.85	2.5				
16	4.77	14.4	11.8	1.15	0.29		
17	1.20	6.7	3.6	1.11	0.30		
18	2.5	13.0	13.3	4.38	1.31		
24	2.4	10.2	4.0				
25	5.2	20.4	7.8				
26	26.5	92.3	28.4	5.3	1.4		
27	61.2	150.7	75.4	24.2	5.7		
28	82.7	220.0	98.8	45.2	11.0	0.6	6.7
29	131.8	321.5	138.8	90.1	44.5	25.4	6.7
30	285.9	507.7	435.5	425.0	364.3	316.5	140.3
31	59.6	107.7	88.7	80.0	55.0	33.4	6.5
32	43.9	81.5	57.0	55.4	38.5	21.0	4.07
33	100.0	100.0	100.0	100.0	100.0	100.0	100.0

^a Reference 7.^b Reference 8. All other patterns were obtained using Consolidated model 21-101, trap current 50 μ a; all relative intensities have been corrected for C₁₃ abundance.

successively trapping and reintroducing the sample until the ethylene peak was reduced to less than 0.001% of the ethane. The oxygen, xenon, nitric oxide, and ethylene have been described previously.⁴ Reagent-grade iodine was obtained from the Mallinckrodt Chemical Works and used without purification.

Irradiation Procedure

The 2000 curie cobalt-60 gamma-ray source at the National Bureau of Standards was used in this investigation. The procedure has been described previously.⁵ The majority of the experiments in which scavengers were added were done at 0.2% conversion.

Analysis

The analytical system has been described.² The gases which are noncondensable at -210° were removed by a Toepler pump, measured in a gas buret, and analyzed for isotopic hydrogen and methanes with a Consolidated model 21-101 mass spectrometer. The ethane was subjected to at least two successive warmup cycles to ensure complete removal of the methane. Only about half of the methane is removed by simply freezing and pumping. The distribution of the isotopic methanes was calculated from cracking patterns reported in the literature.⁶

The ethylene was determined by gas chromatography [Perkin-Elmer No. 154, silica gel (*J*) column, 25°]. Ethane was used as an internal reference. Analysis of known mixtures of ethane and ethylene, under conditions identical to those used for analysis of the irradiated

samples, showed that to within 5% the relationship between peak area and concentration was the same for both ethane and ethylene. The ethylene was trapped from the helium stream in a glass spiral trap immersed in liquid nitrogen. The helium was removed by pumping through a solid-nitrogen trap (to retain the ethylene) and the ethylene was then pumped from a Ward still at -170° . This sample was analyzed in the mass spectrometer for isotopic ethylenes. Patterns for all the isotopic ethylenes were obtained by running authentic samples (obtained from Merck and Company, Ltd.) in the same mass spectrometer.

RESULTS AND DISCUSSION

I. Mass Spectrum of CH₃CD₃

Before discussing the radiolysis results, it is essential to establish the isotopic purity of the ethane-1,1,1-*d*₃ used in this investigation.

A. High-Energy Electron Cracking Pattern

Taking into consideration the different types of mass spectrometers used, it can be seen (Table I) that there is fair agreement between the cracking pattern of the CH₃CD₃ used in this investigation and the one obtained by Schissler *et al.*⁷ A sample of the CH₃CD₃ used in this investigation was also examined on a Nier-type instrument through the kindness of Dr. F. P. Lossing at the National Research Council of Canada. The cracking pattern obtained agreed within 5% of the one obtained by us. There is, however, considerable disagreement between the cracking pattern

⁴ L. J. Stief and P. Ausloos, *J. Phys. Chem.* **65**, 1560 (1961).⁵ L. J. Stief and P. Ausloos, *J. Phys. Chem.* **65**, 877 (1961).⁶ F. L. Mohler, V. H. Dibeler, and E. Quinn, *J. Research Natl. Bur. Standards* **61**, 171 (1958).⁷ D. O. Schissler, S. O. Thompson, and J. Turkevich, *Discussions Faraday Soc. Hydrocarbons*, 46 (1951).

TABLE II. Radiolysis of ethane-1,1,1- d_3 :Effect of scavengers.

	No scavenger	3% C_2H_4	0.4% O_2	3% NO	0.05% I_2	30% I_2
cc(NTP)/min-mole $\times 10^3$						
H_2	15.27	5.95 ^a	7.09 ^b	5.49	4.67	3.52
HD	17.93	10.42	13.07 ^b	10.81	9.09	6.60
D_2	3.88	3.42	4.47 ^b	3.65	3.10	2.28
CH_4	0.59	0.65	0.17	0.14	0.13	0.12
CH_3D	0.59	0.52	0.29	0.35	0.42	0.26
CH_2D_2	0.30	0.33	n.d.	0.13	0.16	0.03
CD_3H	1.17	1.14	0.68	0.71	n.d.	0.42
CD_4	0.32	0.12	0.26	0.13	n.d.	0.07
HD/ D_2	4.62	3.05	2.92	2.96	2.93	2.90

Pressure of CH_3CD_3 —250 mm at 25°.
 Reaction temp.—16° (165° for 30% I_2 experiment).
 Conversion—0.2% (0.05% for scavenger-free experiment).
 n.d.—not determined.

^a Corrected for contribution from C_2H_4 decomposition.

^b These yields may be too high because of the uncertainty in measuring small amounts of noncondensibles in the presence of a large quantity of oxygen.

reported by Quinn and Mohler⁸ and the one obtained by us, although the mass spectrometers used were of a very similar design. This difference is even more surprising if one considers the fact that the spectra of C_2H_6 and C_2D_6 obtained by us agreed closely with the ones reported by Quinn and Mohler. It was stated by these authors that there is a wide variation in the abundances of the parent-molecule ions of the different ethanes. By making the sums of the ions in each ethane cracking pattern equal, relative intensities, 22.1, 13.0, and 18.7 were reported for the parent ions $C_2H_6^+$, $CH_3CD_3^+$, and $C_2D_6^+$, respectively. It was thus concluded that highly asymmetrical ions gave low values for the abundance of parent-molecule ion. However, if we use the cracking pattern of CH_3CD_3 obtained by us, the distribution is as follows: 22.1, 20.0, 18.7. Because the latter distribution seems more reasonable⁹ than the one presented by Quinn and Mohler, we are led to believe that the sample used by these authors must have contained impurities contributing to the abundance of the peaks in the mass region 24–32.

B. The Low-Energy Electron Cracking Pattern

The following two conclusions may be drawn from the cracking pattern obtained at 12 v¹⁰:


- (1) The maximum amount of $C_2H_4D_2$ is about 4%.
- (2) The abundance of the fragment ion correspond-

⁸ E. I. Quinn and F. L. Mohler, J. Research Natl. Bur. Standards **65A**, 93 (1961).

⁹ A similar distribution can be deduced from the data of Schissler *et al.* Also, for other hydrocarbons, only gradual variations in the abundance of the parent ions with varying degrees of deuteration, have been reported in the literature. For instance, in the case of ethylene [V. H. Dibeler, F. L. Mohler, and M. de Hemp-trime, J. Research Natl. Bur. Standards **53**, 107 (1954)] the cracking pattern of the asymmetrical CH_2CD_2 and the symmetrical $CHDCHD$ are very similar and the abundances of the parent ions of these two isomers lie in between those of C_2H_4 and C_2D_4 .

¹⁰ The values of the electron voltages given in this paper are only nominal values, which are accurate within 0.5 v.

ing to mass 30 is more than 20 times that of any other fragment ion. This clearly demonstrates the importance of the reaction



which in turn indicates that CH_3CD_3 is the major isotopic component of the $C_2H_3D_3$ used in this investigation.

It may be noted that after making a correction¹¹ for the contribution of the $C_2H_3D_2^+$ ion to mass 31, the amounts of hydrogen (H_2 : HD: D_2) removed at 14 v are in the relation 0.058: 0.872: 0.07. Assuming there is 4% of CHD_2CH_3 in the ethane sample, these probabilities become¹² 0.065: 0.886: 0.049. Incidentally, these distributions differ considerably from the one which Schissler *et al.* deduced from their 13-v cracking pattern. Their probabilities for removal of H_2 : HD: D_2 are 0.04: 0.71: 0.25 which lead to the unlikely conclusion that the removal of D_2 is strongly favored over the removal of H_2 from the CH_3CD_3 ion.

In view of the presence of possible traces of isotopic impurities and the difficulty of obtaining accurate cracking patterns below 12 v, it is not possible to decide whether or not the 29 and 31 peaks can be attributed to a reaction such as



followed by a loss of hydrogen molecule from two adjacent carbon atoms.¹³ The distribution obtained at 12 v indicates, however, that reshuffling of hydrogen atoms is not an important process at low electron

¹¹ A weighting factor of 1.67, as deduced by Quinn and Mohler from the cracking pattern of C_2D_6H at 70 v was used for estimating probabilities of removal of an H and a D atom from the parent ion.

¹² It was approximated by a statistical argument that the probability for the loss of HD from CHD_2CH_3 is twice that for the loss of H_2 .

¹³ M. Krauss, A. L. Wahrhaftig, and H. Eyring, Ann. Rev. Nuclear Sci. **5**, 256 (1955).

energies. At high energies, on the other hand, the presence of ions of mass 17 shows that a process such as (4) may precede the split of the ethane-molecule ion along the carbon-carbon axis. The decrease in the ratio of the relative intensities of masses 17 and 18 with decrease in electron energy may be related to the diminishing importance of reaction (4) at lower electron energies.

II. Hydrogen Formation in Radiolysis of Ethane

A. Experimental Observations

The radiolysis of an equimolar mixture of C₂H₆ and C₂D₆ in presence of iodine at 16° gave the following distribution of the hydrogens, H₂, HD, D₂: 50.8, 6.8, 42.4. The latter distribution is in good agreement with the one obtained by Dorfman² in experiments using C₂H₄ as a radical scavenger, and thus confirms Dorfman's conclusion that most of the hydrogen not removed by the ethylene is formed in a molecular detach-

TABLE III. Radiolysis of ethane-1,1,1-*d*₃: effect of pressure.

Pressure (mm Hg)	3.0	50.0	150.0	250.0
cc. (NTP)/min.-mol×10 ³				
H ₂	5.40	5.93	5.46	5.96
HD	10.44	11.29	9.91	10.42
D ₂	3.10	3.51	3.20	3.42
CH ₄	0.64	0.25	0.55	0.65
CH ₃ D	0.42	0.47	0.53	0.52
CH ₂ D ₂	0.20	0.28	0.31	0.33
CD ₃ H	1.17	0.90	1.00	1.14
CD ₄	0.18	0.20	0.13	0.12
HD/D ₂	3.37	3.27	3.09	3.05
HD/H ₂	1.94	1.91	1.82	1.75

Reaction temp.—16°.

All experiments performed in presence of 3% C₂H₄.

H₂ yields corrected for contribution from C₂H₄ decomposition.

ment not involving free hydrogen atoms. It may thus be safely stated that the hydrogens produced in the radiolysis of CH₃CD₃ should arise from a similar process.

The scavenger experiments (Table II) indicate that in the radiolysis of CH₃CD₃, molecular H₂, DH and D₂ are eliminated from the ethane molecule. The distribution H₂:HD:D₂ is, within experimental error, 1.7:3.0:1.0 for all scavengers.¹⁴ It is interesting to note that although addition of iodine up to 30% does reduce the total hydrogen yield, it does not significantly affect the distribution of the three hydrogens.

The results of Table III show that, within experimental error, the total yield of molecular hydrogen is independent of pressure. The relative yield of HD as

¹⁴ The ratio HD/D₂ given in the last row of Table II is a good measure of the constancy of the hydrogen distribution. It has been shown before⁴ that some H₂ may be produced from the walls of the reaction vessel, and may cause small fluctuations of the ratio H₂/HD.

TABLE IV. Radiolysis of ethane-1,1,1-*d*₃: effect of temperature.

	16° ^a	155° ^a	221° ^a	314° ^a	320°	394°
cc(NTP)/min.-mol×10 ³						
H ₂	4.60	4.93	5.12	8.85	23.10	23.04
HD	8.16	8.82	8.96	12.27	19.51	25.84
D ₂	2.71	2.85	2.90	3.70	4.88	7.37
CH ₄	0.41	0.74	0.93	n.d.	3.21	7.34
CH ₃ D	0.30	0.52	0.71	n.d.	2.14	6.33
CH ₂ D ₂	0.36	0.30	0.38	n.d.	1.56	4.74
CD ₃ H	0.85	1.01	1.37	n.d.	3.37	9.67
CD ₄	0.19	0.11	0.33	n.d.	0.96	2.99
HD/D ₂	3.01	3.09	3.09	3.32	4.00	3.51

Pressure of CH₃CD₃—250 mm at 25°.

n.d.—not determined.

Conversion—0.2%.

^a 3% C₂H₄ added; H₂ corrected for contribution from C₂H₄ decomposition.

compared to the yields of H₂ and D₂ does undergo a slight but definite increase with decrease in pressure.

An increase in temperature from 16° to 221° (Table IV) has apparently no effect on the total yield or on the distribution of the molecular hydrogen isotopes. The increase in the yield of hydrogen at 314° can be ascribed to the inefficiency of ethylene in trapping hydrogen atoms at this temperature.

It may finally be seen (Table V) that addition of xenon considerably increases the total yield¹¹ of molecular hydrogen, but does not affect the relative yields of isotopic hydrogens.

B. Discussion

The unimolecular decomposition of the parent ion and that of the neutral excited ethane molecule are probably the two principal sources for the formation of molecular hydrogen. As pointed out before,² the formation of molecular hydrogen by an ion-molecule reaction may be tentatively excluded because, as yet, no efficient reaction leading to the formation of molecular hydrogen has been observed in mass spectrometric studies of ethane.

a. *Decomposition of the ethane ion.* If we assume that the decomposition of an ion in the radiolysis system can be deduced from its mass spectral cracking pattern, it can be concluded that a large fraction of the ethane

TABLE V. Radiolysis of ethane-1,1,1-*d*₃: effect of xenon.


Xe/CH ₃ CD ₃	0 ^a	1 ^a	2 ^a	4 ^a	1 ^b
cc(NTP)/min.-mol×10 ³					
H ₂	5.96	23.77	32.26	72.98	19.04
HD	10.42	37.20	52.88	105.70	33.18
D ₂	3.42	12.66	17.69	35.17	10.61
HD/D ₂	3.05	2.94	2.99	3.00	3.13

Reaction temperature—16°.

^a 3% C₂H₄ added; H₂ yield corrected for contribution from C₂H₄ decomposition.


^b 1% NO added.

ions produced in the radiolysis decompose according to process (5)




At 70 v, the relative intensity of the peak corresponding to mass 30 is about equal to the combined relative intensities of all other fragment ions, while at 12 v about 90% of the ethane ions break down according to reaction (5). In order to obtain a more accurate estimate of the distribution of the molecular hydrogens formed in the fragmentation of CH_3CD_3^+ at 70 v, the following decomposition steps will have to be considered:

(1) From the contribution to mass 31, it can be deduced¹¹ that





may occur with a probability which is at most one-tenth that of process (5). If one accepts the existence of an isotope effect, it may be estimated that the analogous process





will only be about half as important as process (6).







(2) C_2H_3^+ , which is a relatively important fragment ion in the C_2H_6 spectrum, can be formed in two alternative ways which are both thermodynamically possible.¹⁵



In view of the fact that in the case of CH_3CD_3 the ethylene ion is largely formed by the loss of a hydrogen molecule from two adjacent carbon atoms, process (8) may be written as follows:




The analog of reaction (9) for CH_3CD_3 consists of six different reaction steps.



¹⁵ A. Kropf, Ph.D. thesis, University of Utah (1954).

If one accepts that randomization of the hydrogen atoms in the ethyl ion is rapid,¹⁶ and neglects the isotope effect, it can be readily seen that reactions (a), (b), (c) correspond to a calculated distribution of: 0.3, 0.6, 0.1.

(3) The acetylene ion, which is the only important ion still to be considered, may be expected to be formed in the following reaction sequence



From the cracking patterns of CHDCHD and CH_2CD_2 it can be deduced¹⁷ that the ethylene- d_2 ion in reaction (10) eliminates H_2 , HD , D_2 with the probabilities: 0.23, 0.65, 0.12.

Using the reactions discussed above, it can be calculated from the cracking pattern that the molecular hydrogen produced in the fragmentation of CH_3CD_3 at 70 v consists of 78 to 88% HD ,¹⁸ at low electron energies, the percentage of molecular HD will be considerably higher.

b. Decomposition of neutral CH_3CD_3 . As mentioned in the introduction, photochemical studies on CH_3CD_3 have clearly demonstrated³ that the elimination of a hydrogen molecule from one carbon atom [Eq. (1) and (2)] is a relatively important process. At 1470 Å, processes (1) and (2) account for more than 80% of the total hydrogen yield. Although the effect of wavelength on the relative importance of primary processes (1), (2), and (11) is not known,



there are indications¹⁹ that the distribution of the molecular hydrogens at 1237 Å is not greatly different from that observed at 1470 Å. A similar independence of wavelength on the molecular hydrogen distribution has been observed in the photolysis of CH_2CD_2 .²⁰

c. Hydrogen formation in the radiolysis of CH_3CD_3 . From the foregoing discussion we may conclude that the average molecular hydrogen distribution: 0.30, 0.52, 0.18 observed in the radiolysis of CH_3CD_3 cannot

¹⁶ A. Kropf, E. M. Eyring, A. L. Wahrhaftig, and H. Eyring, J. Chem. Phys. **32**, 149 (1960). Experimental evidence for an efficient rearrangement of the hydrogen atoms in CH_3CD_2^+ has been obtained recently [P. Ausloos and S. Lias, J. Chem. Phys. (to be published)].

¹⁷ P. Ausloos and R. Gorden, Jr., J. Chem. Phys. **36**, 5 (1962).

¹⁸ The lower value has been obtained by assuming that reaction (9) accounts for the C_2H_3^+ contribution in the spectral pattern of C_2H_6 , while the higher value is based on the occurrence of reaction (8). Although the values for the lower and upper limits are only approximate, it can be estimated that they are reliable within 5%.

¹⁹ The increased importance of H- and D-atom production at the shorter wavelengths (1237 Å) complicates the interpretation of the observed hydrogen distribution. (H. Okabe, unpublished results.)

²⁰ H. Okabe and J. McNesby, J. Chem. Phys. (to be published).

be adequately explained by either a decomposition of the ion or of the neutral excited molecule. A combination of the two modes of decomposition may, however, fit the data. A similar conclusion had to be drawn from the results obtained in the radiolysis of ethylene-*d*₂.¹⁷

It is obvious that an accurate estimate of the relative importance of molecule-ion decomposition and neutral molecule decomposition cannot be obtained from the information available at present. Needed are: (a) a complete analysis of the cracking patterns, as well as the energy distribution of the ions produced in the radiolysis, (b) quantum yields of the molecular elimination processes in the photolysis of ethane as a function of wavelength, and (c) the degree of excitation of the neutral ethane molecules in the radiolysis system.

In view of the disagreement between the molecular hydrogen distribution calculated from the mass spectral pattern of CH₃CD₃ and the one observed in the radiolysis, we may conclude that any correspondence²¹⁻²³ between the *G* value for molecules H₂ calculated from the cracking pattern of C₂H₆ and the experimentally determined *G* values, does not necessarily mean that ion decompositions are solely responsible for the radiolysis products.

The gradual decrease of the ratios HD/D₂ and HD/H₂ with increase in pressure (Table III) indicates that the higher the pressure, the more the distribution of the molecular hydrogens approaches the photochemically observed distribution. A very similar effect of pressure can be deduced from the isotopic hydrogens formed in the radiolysis of CH₂CD₂.¹⁷ This effect may be due to a variation of the energy distribution of the secondary electrons with pressure. An alternative explanation could be based on a competition between reaction and decomposition of the ethane ion.²⁴

Addition of xenon (Table V) does lead to an augmentation of the total yield of molecular hydrogen which approximates the expected increase based on the absorption characteristics. The fact that in the xenon experiments the isotopic distributions of the hydrogens are very similar to those obtained in the pure ethane runs indicates that the relative importance of ion to neutral molecule decomposition is approximately the same for both cases. In contrast to this behavior, it may be pointed out that the distributions of molecular products observed in the radiolyses of CH₂CD₂¹⁷ and CH₃CD₂CH₃²⁵ differ considerably from those found in mixtures containing xenon.

III. The Formation of Methane


The relative concentrations of the methanes produced in the radiolysis of an equimolar mixture of C₂H₆ and

C₂D₆ in the presence of iodine are as follows:


CH ₄	CH ₃ D	CD ₂ H ₂	CD ₃ H	CD ₄
100	16.2	16.2	30.2	86.0.

This distribution is in close agreement with that obtained by Dorfman² in the radiolysis of C₂H₆-C₂D₆-C₂H₄ mixtures. The scavenger experiments thus clearly indicate that a considerable fraction of the methanes is formed in a molecular-detachment process.² Consequently, the methanes produced in the radiolysis of CH₃CD₃-scavenger mixtures should consist largely of CH₃D and CD₃H. The results of Table II indicate that in the presence of NO, I₂, and O₂ this is indeed the case. It can also be seen that the ratio CD₃H/CH₃D is roughly equal to 2, indicating the existence of a considerable isotope effect.²⁶

In order to explain the formation of molecular methane in the radiolysis of ethane the following reaction has been proposed⁴:




However, in view of the fact that recent photolysis studies of C₂H₆-C₂D₂ mixtures have shown that excited ethane molecules may decompose in part according to the process



reaction (10) should not be considered as the only possible mode of molecular methane formation in the ethane radiolysis.

The results of Table IV show that an increase in temperature leads to an increase in the yields of all the methanes. This may be explained by abstraction of a hydrogen atom by a methyl radical from ethane. No obvious explanation can be given for the formation of CH₂D₂. The facts that its yield can be reduced by scavengers (Table II) and increased with temperature (Table IV) suggest that CH₂D₂ is partly formed in a radical reaction. A sequence of reactions such as:



followed by abstraction reactions involving the radicals CH₂D and CD₂H would be consistent with the results on CH₃CD₃, but would not explain the formation of CH₂D₂ in the radiolysis of C₂H₆-C₂D₆ mixtures.²

²⁶ In view of the presence of CD₄, CH₂D₂, CH₄ among the methanes, it may be expected that a fraction of the total CH₃D and CD₃H is formed in bimolecular reactions. If one makes the reasonable assumption that the yields of CH₃D and CD₃H produced by these reactions are, respectively, equal to the yields of CD₄ and CH₄, the isotope effect reaches a value of 3. The latter value depends strongly on the accuracy by which CH₄ can be determined.

²¹ R. A. Back, J. Phys. Chem. **64**, 124 (1960).

²² L. M. Dorfman and M. C. Sauer, Jr., J. Chem. Phys. **32**, 1886 (1961).

²³ K. Yang and P. L. Gant, J. Phys. Chem. **65**, 1861 (1961).

²⁴ D. P. Stevenson, Radiation Research **10**, 610 (1959).



²⁵ S. Lias and P. Ausloos, J. Chem. Phys. (to be published).

TABLE VI. Radiolysis of ethane-1,1,1-*d*₃: ethylene formation.



Scavenger pressure	I ₂ 0.3 mm	NO 7.5 mm	O ₂ 7.5 mm
Relative concentrations			
C ₂ HD ₃	30.0	22.1	38.5
C ₂ H ₂ D ₃ ^a	127.5	92.4	120.0
C ₂ H ₃ D	14.7	11.5	18.3
H ₂	50.0	50.8	54.2
HD	100.0	100.0	100.0
D ₂	32.0	33.8	34.0

^a C₂H₂D₃ consists of more than 85% CH₃CD₂. Pressure of CH₃CD₃: 250 mm at 25°C.

On the other hand, a reaction sequence such as²⁷



followed by an abstraction or disproportionation reaction, would explain the formation of scavengable CH₂D₂. It is reasonable to assume that the CH₂D₂ observed in the C₂H₆-C₂D₆-I₂ system is formed by the reactions



It may be pointed out that although the formation of CH₃D and CD₃H in the radiolysis of C₂H₆-C₂D₆-I₂ can be partly due to a hot radical effect, the production of CH₂D₂ cannot be explained this way.

IV. The Formation of Ethylene



The composition of the ethylene produced in the radiolysis of an equimolar mixture of C₂H₆ and C₂D₆ in the presence of iodine is as follows:

C ₂ H ₄	C ₂ H ₃ D	C ₂ H ₂ D ₂	C ₂ D ₃ H	C ₂ D ₄
41.1	7.5	0.1	8.9	42.5.



²⁷ See for instance: R. A. Marcus, J. Chem. Phys. **20**, 352 (1952).

These results indicate that a large fraction of the ethylene may be formed by a decomposition of the ethane molecule. The data of Table VI show that in the radiolysis of CH₃CD₃, ethylene-*d*₂ is the principal ethylene formed. The fact that the ethylene-*d*₂ is mainly CH₂CD₂ indicates that randomization of the hydrogen atoms in the ethane molecule or ethyl fragment, prior to decomposition, is not important.

The decomposition of neutral excited CH₃CD₃ according to reactions (16) and (17) probably accounts for a large fraction of the CH₂CD₂. If, as



mentioned before, HD is mostly formed in the decomposition of the ethane ion, it may be concluded that process (17) is the major source of CH₂CD₂.²⁸ The latter reaction may eventually be written as a sequence of two stages



where CH₂CD₃^{*} denote a vibrationally excited ethyl radical. Evidence for a process analogous to 17 has recently been obtained in the photolysis of ethylene,²⁹ and has recently been confirmed independently.²⁰

Since, under all conditions H₂ > C₂D₃H and D₂ > C₂H₃D, we conclude that the ethylidene di-radicals, if formed, do not all rearrange to form ethylene. The extent of bimolecular mixing evidenced in the C₂H₆-C₂D₆-I₂ system is enough to account for a large portion of the C₂D₃H and C₂H₃D formed from CH₃CD₃.

²⁸ C₂H₃D₂⁺ produced in the decomposition of CH₃CD₃⁺, will not lead to the formation of a stable CH₂CD₂ molecule. Neutralization of C₂H₃D₂⁺ will always result in the formation of a highly excited species which will undergo decomposition into smaller fragments. There is also evidence¹⁷ which indicates that the hydrogen atoms in the ethylene ion rearrange rapidly.

²⁹ M. C. Sauer, Jr. and L. M. Dorfman, J. Chem. Phys. **35**, 497 (1961).