

- 10 McDevitt NB, McDonagh J, Taylor HL, Roberts HR. An acquired inhibitor to factor XIII. *Arch Intern Med* 1972; **130**: 772–7.
- 11 Lorand L, Maldonado N, Fradera J, Atencio AC, Robertson B, Urayama T. Hemorrhagic syndrome of autoimmune origin with a specific inhibitor against fibrin-stabilizing factor (factor XIII). *Br J Haematol* 1972; **23**: 17–27.
- 12 Krumdieck R, Shaw DR, Huang ST, Poon MC, Rustagi PK. Hemorrhagic disorder due to an isoniazid-associated acquired factor XIII inhibitor in a patient with Waldenström's macroglobulinemia. *Am J Med* 1991; **90**: 639–45.
- 13 Lopaciuk S, Bykowska K, McDonagh JM, McDonagh RP, Yount WJ, Fuller CR, Cooperstein L, Gray A Jr, Lorand L. Differences between type I autoimmune inhibitors of fibrin stabilization in two patients with severe hemorrhagic disorder. *J Clin Invest* 1978; **61**: 1196–203.
- 14 McDonagh J. Structure and function of factor XIII. In: Colman RW, Hirsh J, Marder VJ, Saltzman EW, eds. *Hemostasis and Thrombosis: Basic Principles and Clinical Practice*, 3th edn. Philadelphia, PA: JB Lippincott Co., 1994: 301.
- 15 Jennings I, Kitchen S, Woods TA, Preston FE. Problems relating to the laboratory diagnosis of factor XIII deficiency: a UK NEQAS study. *J Thromb Haemost* 2003; **1**: 2603–8.
- 16 Anwar R, Miloszewski KJA. Factor XIII deficiency. *Br J Haematol* 1999; **107**: 468–84.
- 17 Fickenscher K, Aab A, Stuber W. A photometric assay for blood coagulation factor XIII. *Thromb Haemost* 1991; **65**: 535–40.
- 18 Larsen PD, Wallace JW, Frankel LS, Crisp D. Factor XIII deficiency and intracranial hemorrhages in infancy. *Pediatr Neurol* 1990; **6**: 277–8.

## Independence of tumor necrosis factor- $\alpha$ -induced adhesion molecule expression from sphingosine 1-phosphate signaling in vascular endothelial cells

Y. MIURA,\* Y. YATOMI,† T. OHMORI, M. OSADA and Y. OZAKI

Department of Laboratory Medicine, University of Yamanashi Faculty of Medicine, Yamanashi, Japan; \*Department of Applied Cell Biochemistry and Cell Culture, Faculty of Agriculture, Okayama University, Okayama, Japan and †Department of Laboratory Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

**To cite this article:** Miura Y, Yatomi Y, Ohmori T, Osada M, Ozaki Y. Independence of tumor necrosis factor- $\alpha$ -induced adhesion molecule expression from sphingosine 1-phosphate signaling in vascular endothelial cells. *J Thromb Haemost* 2004; **2**: 1019–21.

Adhesion of circulating leukocytes, especially monocytes, to activated endothelial cells, leading to the resultant transendothelial migration to the subendothelium, is an initial and important step in atherosclerosis [1]. Since this process is mediated by adhesion molecules, including E-selectin and vascular cell adhesion molecule (VCAM)-1, the regulation of expression of these molecules is related to the development of atherosclerosis [1]. The inappropriate expression of these adhesive proteins is induced by various inflammatory stimuli, of which the inflammatory cytokine tumor necrosis factor (TNF)- $\alpha$  is important [1,2]. Since sphingosine 1-phosphate (Sph-1-P) is abundantly stored in platelets and released extracellularly upon stimulation and since vascular endothelial cells express the G protein-coupled Sph-1-P receptors SIPs, it is


important to study the effect of Sph-1-P on endothelial cells from the viewpoint of platelet–endothelial cell interaction [3]. In this study, we examined the effect of Sph-1-P, in comparison with the inflammatory cytokines, on adhesion molecule expression; Sph-1-P was previously reported to mediate the response elicited by TNF- $\alpha$  [4].

In human umbilical vein endothelial cells (HUVECs), it was confirmed that TNF- $\alpha$ , as well as interleukin (IL)-1 $\beta$ , potently triggers the expression of E-selectin and VCAM-1 (Fig. 1A). Sph-1-P did induce adhesion molecule expression, but only very weakly (Fig. 1A). Similar results, i.e. strong responses induced by TNF- $\alpha$  and weak responses by Sph-1-P, were obtained when HUVEC plasminogen activator inhibitor-1 release was examined (data not shown). These findings are in contrast with the strong intracellular Ca<sup>2+</sup> mobilization and migration responses induced by Sph-1-P, which have been established to be due to its extracellular actions via SIPs [3,5]. Since NF- $\kappa$ B is an essential transcriptional factor for not only the expression of adhesion molecules but also that of PAI-1, it is likely that the ability of Sph-1-P to activate this transcriptional regulatory protein is weak. Our findings are consistent with the report that Sph-1-P fails to induce E-selectin expression or NF- $\kappa$ B activation [6], but not with one describing

Correspondence: Yutaka Yatomi, Department of Laboratory Medicine, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan.

Tel.: +81 3 5800 8730; fax: +81 3 5689 0495; e-mail: yatomi-y-lab@h.u-tokyo.ac.jp

Received 24 January 2004, accepted 5 February 2004



**Fig. 1.** Expression of E-selectin and vascular cell adhesion molecule (VCAM)-1 in human umbilical vein endothelial cells (HUVECs). To examine the expression of adhesion molecules, HUVECs, after the indicated reactions, were washed with PBS and incubated with anti-VCAM-1 (Berkeley Antibody Co., Richmond, CA, USA) or anti-E-selectin (Upstate Biotechnology, Lake Placid, NY, USA) monoclonal antibody for 30 min. Cells were then incubated with fluorescein isothiocyanate-conjugated goat antimouse IgG (Nippon Becton Dickinson, Tokyo, Japan) and fixed in 3% paraformaldehyde. The number of positive cells for the expression of cell-surface adhesion molecules was measured using a FACScan flow cytometer (Nippon Becton Dickinson), and expressed as a percentage of the total cell number. The stimulation-dependent expression was calculated as (positive cells upon stimulation) – (positive cells without stimulation). (A) HUVECs were stimulated with 100 U mL<sup>-1</sup> of tumor necrosis factor (TNF)- $\alpha$  (○) or interleukin (IL)-1 $\beta$  (●), or 5  $\mu$ M sphingosine 1-phosphate (Sph-1-P) (□), for various durations, and the cell surface expression of E-selectin (upper panel) or VCAM-1 (lower panel) was measured. (B) HUVECs were pretreated with 100 ng mL<sup>-1</sup> of pertussis toxin for 16 h, and then stimulated with 100 U mL<sup>-1</sup> of TNF- $\alpha$  (T) or IL-1 $\beta$  (I), or 5  $\mu$ M Sph-1-P (S), for 4 h. Cell surface expression of E-selectin (left panel) or VCAM-1 (right panel) was measured. The results are expressed as a percentage of the control (without the toxin pretreatment). Columns and error bars represent the mean  $\pm$  SD ( $n = 3$ ).

through the generation of intracellular Sph-1-P is critically involved in mediating TNF- $\alpha$ -induced endothelial activation [4]. Xia *et al.* reported that: (i) TNF- $\alpha$  induces a rapid Sph kinase activation and the resultant Sph-1-P generation; (ii) Sph-1-P, as well as TNF- $\alpha$ , potentially stimulates NF- $\kappa$ B activation and adhesion protein expression; and (iii) the Sph kinase inhibitor dimethylsphingosine inhibits the TNF- $\alpha$ -induced responses [4]. Thus, it was postulated that intracellular Sph-1-P serves as a second messenger in mediating TNF- $\alpha$ -induced adhesion molecule expression [4]. However, it should be kept in mind that dimethylsphingosine inhibits not only Sph kinase but also protein kinase C and MAP kinase, and the data obtained using this non-specific inhibitor should be interpreted with caution [7].

In this study, we observed that: (i) the stimulatory effect of Sph-1-P on HUVEC adhesion molecule expression is much weaker than that of TNF- $\alpha$  (Fig. 1A); (ii) the Sph-1-P (but not TNF- $\alpha$ ) response was abolished by pertussis toxin (Fig. 1B); and (iii) Sph conversion into Sph-1-P, reflecting Sph kinase activation, was not affected by TNF- $\alpha$  (data not shown). These present results suggest that Sph-1-P induction of adhesion molecule expression, although weak, is mediated by cell surface receptors, possibly one or more of S1Ps, and argue against the second messenger role of Sph-1-P. Further studies are needed to clarify the mechanism by which TNF- $\alpha$  induces adhesion molecule expression in vascular endothelial cells.

#### Acknowledgements

The authors thank Mitsubishi Kagaku Iatron, Inc. (Tokyo, Japan) for the donation of LPIA/tPAI test kits. This study was supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan and by Mitsubishi Pharma Research Foundation.

that Sph-1-P can mimic the effect of TNF- $\alpha$  leading to the adhesive responses [4].

Sph-1-P has been shown to act not only as an extracellular mediator but also as an intracellular second messenger. Both types of Sph-1-P actions can even be observed within one system, and the HUVEC has been considered as one such example; Sph-1-P induces migration, proliferation, angiogenesis, and nitric oxide formation through the cell surface receptors S1P<sub>1</sub> and S1P<sub>3</sub> [3,5], while the Sph kinase pathway

## References

- Zimmerman GA, McIntyre TM, Prescott SM. Adhesion and signaling in vascular cell-cell interactions. *J Clin Invest* 1996; **98**: 1699–702.
- Madge LA, Pober JS. TNF signaling in vascular endothelial cells. *Exp Mol Pathol* 2001; **70**: 317–25.
- Yatomi Y, Ozaki Y, Ohmori T, Igarashi Y. Sphingosine 1-phosphate: synthesis and release. *Prostaglandins Other Lipid Med* 2001; **64**: 107–22.
- Xia P, Gamble JR, Rye KA, Wang L, Hii CS, Cockerill P, Khew-Goodall Y, Bert AG, Barter PJ, Vadas MA. Tumor necrosis factor- $\alpha$  induces adhesion molecule expression through the sphingosine kinase pathway. *Proc Natl Acad Sci USA* 1998; **95**: 14196–201.
- Lee MJ, Thangada S, Claffey KP, Ancellin N, Liu CH, Kluk M, Volpi M, Sha'afi RI, Hla T. Vascular endothelial cell adherens junction assembly and morphogenesis induced by sphingosine-1-phosphate. *Cell* 1999; **99**: 301–12.
- Masamune A, Igarashi Y, Hakomori S. Regulatory role of ceramide in interleukin (IL)-1 $\beta$ -induced E-selectin expression in human umbilical vein endothelial cells. Ceramide enhances IL-1 $\beta$  action, but is not sufficient for E-selectin expression. *J Biol Chem* 1996; **271**: 9368–75.
- Yang L, Yatomi Y, Satoh K, Igarashi Y, Ozaki Y. Sphingosine 1-phosphate formation and intracellular Ca<sup>2+</sup> mobilization in human platelets: evaluation with sphingosine kinase inhibitors. *J Biochem* 1999; **126**: 84–9.

## More on: unusual expression of the F9 gene in peripheral lymphocytes hinders investigation of F9 mRNA in hemophilia B patients

JACQUELINE A. CUTLER, MICHAEL J. MITCHELL and GEOFFREY F. SAVIDGE

Reference Centre for Haemostatic and Thrombotic Disorders, St. Thomas' Hospital, London, UK

**To cite this article:** Cutler JA, Mitchell MJ, Savidge GF. More on: unusual expression of the F9 gene in peripheral lymphocytes hinders investigation of F9 mRNA in hemophilia B patients. *J Thromb Haemost* 2004; **2**: 1021.

See also Green PM, Rowley G, Giannelli F. Unusual expression of the F9 gene in peripheral lymphocytes hinders investigation of F9 mRNA in hemophilia B patients. *J Thromb Haemost* 2003; **1**: 2675–6.

We read with interest the recent publication in this journal by Green *et al.*: 'Unusual expression of the F9 gene in peripheral lymphocytes hinders investigation of F9 mRNA in hemophilia B patients' [1]. The use of RNA from peripheral lymphocytes to examine exons g-h has been documented over many years [2,3]. Publication of data covering the remainder of the coding region of this eight-exon gene has, however, not been forthcoming.

Green *et al.* postulate that an incomplete transcript is produced in peripheral lymphocytes and other tissues, consisting only of exons g-h. Their results support our own findings, that F9 mRNA in lymphocytes is abnormal.

Of 54 hemophilia B families referred to this center for analysis, we have failed to identify a mutation in the genomic DNA of five index cases. RNA from whole blood from three of these five was extracted. In each case we successfully reverse transcribed RNA into cDNA and sequenced exons g-h. In 2/3 cases we have also been able to amplify and sequence exons

a-b. We have been unable to obtain amplification products specific to exons c-f from patient or control lymphocyte RNA.

These results indicate that abnormal transcripts are produced in peripheral lymphocytes, but that these are not comprised exclusively of exons g-h, as suggested by the previous authors. While it may be that discrete transcripts are produced from the 5' and 3' ends of the gene, it is possible that partially processed transcripts are being produced, comprising exons a, b, g and h.

It is unlikely that a full-length F9 transcript will be produced from peripheral lymphocytes. It is not feasible, in patients with hemophilia B, to obtain RNA from the tissues successfully utilized by Green *et al.* Thus, in cases where no genetic abnormality can be identified by exon-exon sequencing, the only remaining option is analysis of the intronic regions.

## References

- Green PM, Rowley G, Giannelli F. Unusual expression of the F9 gene in peripheral lymphocytes hinders investigation of F9 mRNA in hemophilia B patients. *J Thromb Haemost* 2003; **1**: 2675–700.
- Sarkar G, Sommer SS. Access to a messenger RNA sequence or its protein product is not limited by tissue or species specificity. *Science* 1989; **244**: 331–4.
- Chan V, Au P, Lau P, Chan TK. A novel haemophilia B defect due to partial duplication of the factor IX gene. *Br J Haematol* 1994; **86**: 601–9.

Correspondence: Jacqueline A. Cutler, Reference Centre for Haemostatic and Thrombotic Disorders, St. Thomas' Hospital, London, UK.

Tel.: +44 20 7188 2798; fax: +44 20 7620 1659; e-mail: jacqueline.cutler@gstt.sthames.nhs.uk

Received 5 January 2004, accepted 29 January 2004