

## H<sub>2</sub>O/Triton X-100 Solvent Effect in the Micellar Catalysis of the Aquation of Tris-(3,4,7,8-tetramethyl-1,10-phenanthroline)iron(II)

Jide Ige,\* John N. Lambi and Oladega O. Soriyan

*Chemistry Department, University of Ife, Ile-Ife, Nigeria*

The effect of triton X-100 micelles on the aquation of  $\text{Fe}(\text{Me}_4\text{phen})_3^{2+}$  has been investigated with triton X-100 as solvent. In liquid triton X-100, over a range of  $[\text{H}_2\text{O}]_T$  (0.2–2.0 mol dm<sup>-3</sup>), significant rate enhancement factors of 35–140 are observed. Acid inhibits the rate of aquation at fixed  $[\text{H}_2\text{O}]_T$ . A mechanism based on effective solvent participation in a chemical environment similar to that in reversed micelles is proposed in liquid triton X-100 with dispersed water pockets. This mechanism predicts direct H<sub>2</sub>O substitutions into the coordination sphere of  $\text{Fe}(\text{Me}_4\text{Phen})_3^{2+}$  in the highly polar water pockets or cavities where the Fe<sup>II</sup> complex molecules are solubilized. Changes in the tumbling rate, mobility, structure, nucleophilicity and activity of water are suggested to account for the observed changes in the rate of aquation as a function of  $[\text{H}_2\text{O}]_T$ , in the peculiar chemical environment of the water pockets. All  $k_w$ - $[\text{H}_2\text{O}]_T$  profiles are structured and exhibit maxima with  $k_w(\text{max})$  shifted to progressively higher  $[\text{H}_2\text{O}]_T$  as the fixed concentration of  $[\text{H}^+]_T$  is increased.

Recently, we reported<sup>1,2</sup> the effect of micelles on the aquation of phenanthroline complexes of iron(II). Our results reflect some of the kinetic and mechanistic features of micelle-catalysed systems and emphasise the significance of substrate partitioning between the micellar and bulk water phases.

Such substrate partitioning in pure surfactant medium must differ, as the bulk-water phase is now replaced by the bulk-surfactant phase with the H<sub>2</sub>O molecules and Fe<sup>II</sup> complex molecules dispersed in the latter. Few studies of chemical reactions in purely surfactant medium have been investigated in the literature. The effect of the pure surfactant medium on the overall kinetics and mechanism of the aquation reactions is therefore of interest. The present work is in this direction. We report the aquation of  $\text{Fe}(\text{Me}_4\text{phen})_3^{2+}$  in triton X-100, an important emulsifying agent which has 10 monomers in its aqueous micelle. Triton X-100 is a neutral surfactant for which hydrophobic interaction with the bulky and hydrophobic  $\text{Fe}(\text{Me}_4\text{phen})_3^{2+}$  will be significant.

With triton X-100 as the bulk solvent, addition of small amounts of water creates water pockets with a chemical environment which may be similar to that in reversed micellar systems.<sup>3-13</sup> The kinetics and mechanistic features of the dissociation of the Fe<sup>II</sup> complex is therefore of comparative interest with respect to observed data in ordinary aqueous system.†

### Experimental

#### Materials

$\text{Fe}(\text{Me}_4\text{phen})_3\text{SO}_4$  (G. F. Smith Chemical Company) gave a molar extinction coefficient in good agreement with the literature<sup>13</sup> value at  $\lambda_{\text{max}} = 500$  nm and was therefore used

† We have studied the aquation of this complex in aqueous triton X-100. Inhibition was observed and inhibition factors of 2.4 to 2.7 were calculated.

without further purification. Scintillation-grade triton X-100 was also used without further purification. All other chemicals were of AnalaR grade (BDH). Because of the viscous nature of triton X-100, vigorous shaking was always necessary for thorough mixing in the reaction cells. The  $\text{Fe}^{\text{II}}$  complex concentration was fixed at  $3.162 \times 10^{-5} \text{ mol dm}^{-3}$ , except where stated otherwise.  $\text{H}_2\text{SO}_4$  was used preferentially because the iron(II) complex is precipitated in almost all the other mineral acids.

### Kinetics

The aquation rate was followed by monitoring the disappearance of the iron(II) complex at  $\lambda_{\text{max}} = 500 \text{ nm}$ , using an SP 500 series 2 Pye-Unicam spectrophotometer fitted with an automatic cell changer and thermostatted cell compartments. Constant temperature was maintained by immersion in a water bath fitted with a Gallenkamp thermoregulator and a cryocool CC-60T compressor coil.  $k_w$ , the aquation rate constant was obtained from the slopes of plots of  $\ln(A_t - A_\infty)$  vs.  $t$ , which were always linear. Temperature was maintained at  $25^\circ\text{C}$ .

### Results

$k_w$  vs.  $[\text{H}_2\text{O}]_T$  profiles (fig. 1), at fixed  $[\text{H}^+]_T$  ( $0.00$ – $0.02 \text{ mol dm}^{-3}$ ), exhibit maxima with a shift in the position of the maximum to progressively higher  $[\text{H}_2\text{O}]_T$  as fixed  $[\text{H}^+]_T$  is increased. Comparison of the profiles indicates inhibition by acid. At the maxima, calculated catalysis factors ( $k_w/k_w^\dagger$ ) are in the range 35–140. This is significant.

$k_w$  decreases with increase in  $[\text{Fe}(\text{Me}_4\text{phen})]_T^{2+}$  at fixed  $[\text{H}_2\text{O}]_T$  ( $0.6 \text{ mol dm}^{-3}$ ). Fig. 2 shows a 2.5-fold decrease in  $k_w$  for a threefold increase in complex concentration.

### Mechanism and Discussion


In fig. 3, we propose that in triton X-100 solvent medium the surfactant molecules adopt an approximately reversed micellar arrangement $\ddagger$  around the limited  $\text{H}_2\text{O}$  molecules ( $< 4\% \text{ v/v H}_2\text{O}$ ) present,<sup>4–12</sup> with the polar OH headgroups pointing towards the centre of the polar cavities where the  $\text{H}_2\text{O}$  molecules are localised as in other reverse micellar systems.<sup>4–12</sup> The activity of water is highly enhanced in this environment owing to its reduced mobility/tumbling rate.<sup>4, 6–12</sup> Hydrogen bonding with the OH head group can only be such that the oxygen of  $\text{H}_2\text{O}$  points towards the centre of the cavity where, we propose, the bulky  $\text{Fe}(\text{Me}_4\text{phen})_3^{2+}$  is localised. Owing to the constraint of space in the cavity, the tumbling rate of the iron(II) complex is significantly reduced. This is enhanced by the inability of the substrate to escape into the bulk of the solvent because of the steric blockage of possible outlets by  $\text{H}_2\text{O}$  molecules hydrogen bonded to the OH headgroups. The iron(II) complex is therefore suitably oriented for direct attack by  $\text{H}_2\text{O}$  molecules.

Further support for the above proposed mechanism comes from lipid systems<sup>15</sup> (the liquids are just as viscous) where the tumbling rate of water, at low  $\text{H}_2\text{O}$  content (*ca.*  $0.20\% \text{ v/v}$ ), is negligible. Given this condition the mobility of the  $\text{H}_2\text{O}$  in the present system is influenced predominantly by the overall tumbling rate of the micelle and iron(II) complex. It is reasonable to expect that at higher water compositions additional water molecules are free with a mobility approaching that in bulk water, as has been observed for lipids and reversed micellar systems.<sup>13, 15, 16</sup>


The initial observed rate enhancement (fig. 1) and the observed maxima are well within the range of water concentration specified for the first type of  $\text{H}_2\text{O}$  molecules hydrogen bonded to the OH head groups. The rate enhancement is therefore a

$\dagger k_w$  = aquation rate constant in micelle free medium ( $2.40 \times 10^{-4} \text{ s}^{-1}$  in  $1.0 \text{ mol dm}^{-3} \text{ H}^+$ ).

$\ddagger$  In the absence of any calculations or experimental data on the size number of  $\text{H}_2\text{O}$  molecules in the water pools or cavities of pure triton X-100, the true number of substrates/monomers may be different from that illustrated in fig. 3.



**Fig. 1.** Variation of aquation rate constant with  $[\text{H}_2\text{O}]_T$  at  $\text{H}^+$  concentrations ( $\text{mol dm}^{-3}$ ) of 0.00 (○), 0.002 (●), 0.010 (○) and 0.02 (▲).  $T = 25^\circ\text{C}$  (Triton X-100 solvent medium).  $[\text{Fe}(\text{Me}_4\text{phen})_3^{2+}]_T = 3.162 \times 10^{-5} \text{ mol dm}^{-3}$ .



**Fig. 2.** Variation of aquation rate constant with iron(II) complex concentration in triton X-100 solvent.





Fig. 3. Substrate-micelle interaction.  $R = (CH_3)_3C-CH_2-C(CH_3)_2-$ .

consequence of both the increased activity of water and the suitable orientation of the cornered iron(II) complex localised in the interior of the polar cavities and this enhances direct displacement of  $Me_4phen$  ligands from the coordination sphere of the iron(II) complex. The subsequent decrease in rate results in an increase in the tumbling rate of  $H_2O$  molecules and the iron(II) complex with an accompanying decrease in the activity of water, which gradually approaches that in bulk water at  $2.0 \text{ mol dm}^{-3} H_2O$  (3.7% v/v).

These observations are consistent with the following mechanism:



which expresses the bonding of  $H_2O$ , through hydrogen bonds, to the OH headgroups.



represents the solubilisation of  $S^{2+}$  in the polar cavities.



is the aquation path at low  $H_2O$  concentration, when the mobility of  $H_2O$  and other substrates in the polar cavities are significantly reduced, and



is the aquation path at high  $\text{H}_2\text{O}$  concentration. It can be shown that eqn (1)–(4) yield

$$k_\psi = \frac{\text{rate}}{[\text{S}^{2+}]_T} = k_4 + \frac{K(k_3 - k_4)[\text{H}_2\text{O}]_T}{K[\text{S}^{2+}]_T + K[\text{H}_2\text{O}]_T + K_1 + 1} \quad (5)$$

where  $K = K_1 K_2$  and  $[\text{M}]_T$  does not appear in the expression for  $k_\psi$  because the surfactant is the solvent, and hence to a good approximation  $[\text{M}]_T$  is constant and is contained in  $K_1$ . Eqn (5) is reduced to

$$k_\psi = \frac{k_3 K[\text{H}_2\text{O}]_T}{K[\text{H}_2\text{O}]_T + K[\text{S}^{2+}]_T + K_1 + 1} \quad (6)$$


at small  $[\text{H}_2\text{O}]_T$  when reaction is predominantly through eqn (3) and predominantly through reaction path 4.  $k_\psi$  is then given by

$$k_\psi \approx k_4 - \frac{k_4 K[\text{H}_2\text{O}]_T}{K[\text{H}_2\text{O}]_T + K[\text{S}^{2+}]_T + K_1 + 1} \quad (7)$$


at higher  $[\text{H}_2\text{O}]_T$  when conditions approach that in bulk water and reaction is predominantly through reaction path 4. Eqn (6) and (7) are in agreement with the observed variation of  $k_\psi$  at high and low  $[\text{H}_2\text{O}]_T$ .

Inhibition by acid (fig. 2) is due to the protonation of the OH headgroups, making the first type of favourably oriented  $\text{H}_2\text{O}$  molecules in the water pockets or polar cavities to form hydroxonium ions. This reduces considerably the nucleophilicity of the  $\text{H}_2\text{O}$  molecules for direct substitution into the coordination sphere of the positively charged iron(II) complex [*i.e.* there is Coulombic repulsion between  $\text{H}_3\text{O}^+$  and  $\text{Fe}(\text{Me}_4\text{phen})_3^{2+}$ ]. Inhibition by acid contrasts with the observed† catalysis in micelle/micelle-free aqueous triton X-100 and strongly supports the postulate that aquation does not occur predominantly by dissociative mechanism.<sup>1</sup>


Direct substitution of  $\text{H}_2\text{O}$  into the coordination sphere of metal-ion complexes in reverse micellar medium has been proposed by Fendler and co-workers;<sup>17</sup> however, their system is quite different and hydrogen-bonding and proton transfer by  $\text{H}_2\text{O}$  molecules are central to their proposed mechanism for aquation. This is significantly complementary to the present study. Direct water substitution into the coordination sphere of  $(\text{Me}_4\text{phen})_3^{2+}$  in the polar cavities of triton X-100 is either by two consecutive steps,



† The aquation rate constant ( $k_w$ ), for the catalysis by acid in a micelle-free medium, fits the expression  $k_w = k_0 + k[\text{H}^+]$  ( $k_0$ ,  $k$  are constants). In aqueous triton X-100 medium, at fixed triton X-100 composition, a similar expression is obtained for the variation of the aquation rate constant ( $k_\psi$ ) with  $[\text{H}^+]$  at low surfactant compositions.



or by a single concerted attack by two water molecules



The present kinetic data cannot distinguish between these two modes of water substitution.

Inclusion of eqn (10) [in addition to eqn (1)–(4)]



in the derivation of the expression for  $k_\psi$  yields

$$k_\psi = k_4 + \frac{K(k_3 - k_4)[H_2O]_T}{K[H_2O]_T + K[S^{2+}]_T + K_8[H^+]_T + K_1 + 1} \quad (11)$$

which predicts inhibition by acid.

The presence of  $SO_4^{2-}$  ions, the dissociation product of  $H_2SO_4$ , in addition, inhibits aquation by forming a negative field around the iron(II) complex. This effect is, however, only significant at high acid concentrations. The shifts in the position of the maxima progressively to higher  $[H_2O]_T$  with increase in  $[H^+]_T$  is a combination of both the decreasing activity of water and the changing number of surfactant headgroups surrounding the polar cavities.

The  $k_\psi$  vs.  $[H_2O]_T$  profile in acid-free medium (uppermost curve in fig. 1) does not completely fit into the comparative trends in  $k_\psi$  at low fixed  $[H_2O]_T$ . While we cannot explain this deviation, we have observed a similar effect in  $k_\psi$  vs.  $[SDS]_T$  profiles.<sup>18</sup>

We thank Professor H. D. Burrows for supplying the Tritonx-100 and for his useful discussions.

## References

- 1 H. Burrows, J. Ige and S. A. Umoh, *J. Chem. Soc., Faraday Trans. 1*, 1982, **78**, 947.
- 2 J. Ige and O. Soriyan, *J. Chem. Soc., Faraday Trans. 1*, 1986, **82**, 2011.
- 3 J. H. Fendler and J. E. Fendler, *Catalysis in Micellar and Macromolecular Systems* (Academic Press, New York, 1975).
- 4 (i) M. Wong, J. K. Thomas and T. Nowak, *J. Am. Chem. Soc.*, 1977, **99**, 4730; (ii) *Microemulsions* (Proc. Conf. Phys. Chem. Microemulsions), ed. I. D. Robb (Plenum Press, New York, 1982), p. 103.
- 5 J. H. Fendler and L. J. Liu, *J. Am. Chem. Soc.*, 1975, **97**, 999.
- 6 *Physics Reports: Surfactant–Water–Liquid Crystal Phases* ed. G. J. T. Tiddy (North Holland, Amsterdam, 1980), vol. 57, p. 1.
- 7 J. E. Fendler, J. H. Fendler, R. T. Medary and V. A. Woods, *J. Chem. Soc., Chem. Commun.*, 1972, 1497; *J. Am. Chem. Soc.*, 1972, **94**, 7288.
- 8 J. H. Fendler and S. A. Chang, *J. Am. Chem. Soc.*, 1973, **95**, 3273.
- 9 C. J. O'Connor, J. E. Fendler and J. H. Fendler, *J. Org. Chem.*, 1973, **38**, 3371.

- 10 C. J. O'Connor, J. E. Fendler and J. H. Fendler, *J. Chem. Soc., Dalton Trans.*, 1974, 625.
- 11 C. J. O'Connor, J. E. Fendler and J. H. Fendler, *J. Am. Chem. Soc.*, 1974, **96**, 370.
- 12 J. H. Fendler, F. Nome and H. C. Van Woert, *J. Am. Chem. Soc.*, 1974, **96**, 6745.
- 13 J. H. Fendler, *Acc. Chem. Res.*, 1976, **9**, 153.
- 14 G. F. Smith, *Chemical Company Catalogue*; 1985, contains many references on this complex.
- 15 J. Simplicio and K. Schwenzer, *Biochemistry*, 1973, **12**, 1923.
- 16 W. Wong, J. K. Thomas and T. Nowak, *J. Am. Chem. Soc.*, 1973, **95**, 286.
- 17 C. J. O'Connor, E. J. Fendler and J. H. Fendler, *J. Am. Chem. Soc.*, 1973, **95**, 600.
- 18 J. Ige and O. Soriyan, to be submitted.

*Paper 6/115; Received 15th January, 1986*