



IONIC AND RADICAL MECHANISMS IN OLEFINIC SYSTEMS, WITH SPECIAL REFERENCE TO PROCESSES OF DOUBLE-BOND DISPLACEMENT, VULCANISATION AND PHOTO-GELLING.

BY ERNEST HAROLD FARMER.

Received 29th April, 1942.

The α -methylenic reactions discussed in the two preceding papers recall a series of interesting observations by Baker and Nathan¹ which indicate that a *p*-methyl substituent attached to the benzene nucleus can permit electron-release to the nucleus in a manner that appears only in lesser degree in higher alkyl groups and may be absent in some (*e.g.* Bu^y). Thus in *p*-methylbenzyl bromide the suggested function of the methyl group (dotted arrows in (I)) permits additional electron-release



at the C—Br bond, and so facilitates the anionisation of the bromine. Baker and Nathan suggest that the electrons of the duplet constituting

¹ *J. Chem. Soc.*, 1935, p. 1847; see also *ibid.*, 1942, p. 191.

the C—H bond of the methyl group are less localised than those in a similarly placed C—C bond, and hence that a methyl group attached to the necessary conjugated unsaturated system is capable of electron-release by a mechanism similar to the tautomeric effect (II).

They make clear, however, that the new mechanism functions *in addition* to the generally-accepted inductive (+ I) effect of alkyl groups, and they find experimentally that the efficiency of representative alkyl groups is in the order $\text{Me} > \text{Et} > \text{Pr}^\beta > \text{Bu}^\gamma$, and that the suggested electron-release must be of mesomeric rather than of tautomeric type. The phenomenon noted by Baker and Nathan seems to be part of a much wider phenomenon which applies to unconjugated (as well as to conjugated) polyolefines, and even to mono-olefins which contain the system $-\text{CH}_2-\text{CH}=\text{CH}-$, and it appears likely that in general the three-carbon system exists in a mesomeric state corresponding to the

electron-shifts in $-\text{CH}-\text{CH}=\text{CH}$. This conclusion has, if correct,

$$\begin{array}{c} \text{H} \\ \curvearrowright \\ -\text{CH}-\text{CH}=\text{CH} \end{array}$$

certain consequences with respect to the ionic reactivity and physical properties of olefinic substances: these consequences are exemplified in the following two sections. In addition, however, the mesomeric condition as here conceived appears to lie at the root of the well-known "cracking" rule of diene polymers (as it is illustrated in the cracking of rubber to isoprene, the reversal of the diene synthesis, etc.), whereby the links which break, *i.e.* the weakest links, are never the C—C links adjoining the double bonds $(-\text{C}=\text{C}-\text{C}-)$; also it appears to exert a marked influence on the tendency of the α -methylenic C—H bond to dissociate symmetrically. Such observations as have been made in the latter respect indicate that alkyl substituents attached to the ethylenic carbon atoms do indeed appear to facilitate the symmetrical dissociation; nevertheless it is difficult at present to disentangle the purely inductive effect of the alkyl groups from the mesomeric effect in a system such as $\text{R}-\overset{*}{\text{CH}}_2-\text{C}(\overset{*}{\text{CH}}_3)=\text{CH}-\overset{*}{\text{CH}}_2-\text{R}'$, characteristic of the polyisoprenes and rubber. The available evidence suggests that dissociative reaction does not occur principally at the methyl groups, but it is too fragmentary to distinguish quantitatively between the three C—H bonds marked *.

Ionic Double-bond Displacement.


Double-bond displacement induced by the action of alkali is well known under the name of three-carbon or prototropic change and is a common property of $\alpha\beta$ - or $\beta\gamma$ -unsaturated fatty acids, ketones, nitriles, etc. The same type of change has been observed in the case of the aryl-olefins, β -phenylpropylene, eugenol and Δ^2 -dihydronaphthalene in which the aryl group replaces $>\text{CO}$ or $-\text{CN}$ as an activating group, but until recently was unknown in olefinic chains in which the double bond was remote from all activating groups of the usual types. Moore,² however, found that linolenic acid, $\text{C}_2\text{H}_5-(\text{CH}=\text{CH}-\text{CH}_2)_3-(\text{CH}_2)_6-\text{CO}_2\text{H}$, when heated with alkali passed slowly into an isomeric conjugated form. Linoleic acid and linoleyl alcohol similarly pass into conjugated forms,³ and it is undoubtedly a

^{1a} Cf. Schmidt, *Ber. B.*, 1936, **69**, 1885; Scheller, *ibid.*, 1939, **72**, 1917.


² *Biochem. J.*, 1937, **31**, 139; 1939, **33**, 1635.

³ Kass and Burr, *J. Amer. Chem. Soc.*, 1939, **62**, 1796.

In the case of the 1 : 5-diene system as it is represented by the linked isoprene units $\text{—CH}_2\text{—CMe=CH—CH}_2\text{—}$ of squalene, dihydromyrcene and rubber, prolonged treatment with alkali under the conditions which are successful with linolenic and lineoleic acids has been found to cause no recognisable degree of bond shift.⁴ With regard to the simple monoolefinic system there is on record the early statement of Varrentrapp⁵ that oleic acid when fused with caustic alkali gave a considerable yield of palmitic and acetic acids, as though by hydrolytic fission of Δ^2 -octadecenoic acid :


$$-\text{CO}-\text{CH}_2-\text{CO}- \quad \text{and} \quad -\text{CH}=\text{CH}-\text{CH}_2-\text{CO}-,$$
$$\begin{array}{ccc} \text{H} & & \\ | & & \\ -\text{CH}_2-\text{CH}=\text{CH}-\text{CH}-\text{CH}=\text{CH}-\text{CH}_2 & & -\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2 \\ \text{a} \quad \text{a}' & & \text{a} \quad \text{a}' \\ \text{(A)} & & \text{(B)} \end{array}$$

If we consider the three-carbon system (C) in terms of Ingold's conception of prototropic mechanism, the removal of a proton from C_α by the




⁵ *Annalen*, 1840, 35, 196.

base B and the complementary and concurrent supply of a proton to C γ by some substance in the system together constitute a process with a definite activation energy (say E). This energy, equal to the difference in energy between the transition state (D) and the initial state C, may be considered to be reduced by a quantity α , due to the polarisations a and b , and representing resonance energy resulting from the electron distribution symbolised by a and b . If the process actually does occur, then the system becomes (E), *i.e.* the process necessitates a double-bond shift, which in the absence of an imposed external constraint and with an energy in the system after the change essentially the same as before it (as, *e.g.*, in a purely hydrocarbon system with only saturated carbon atoms beyond the three-carbon system) may be in either direction. But since in the case of oleic acid double-bond displacement not only succeeds, but apparently proceeds only in one direction (towards the CO $_2^-$ group), it must be concluded both that the required activation energy $E - \alpha$ is actually developed under the conditions of alkali fusion, and that the negative charge on the CO $_2^-$ ion exercises a sufficient, albeit a very feeble, inductive influence through a chain of six saturated carbon atoms (or alternatively through the medium) to determine that a hydrogen atom is detached from

C $_8$ in the system $\text{---CH}_2\text{---CH=CH---CH}_2^{\delta+} \rightarrow (\text{CH}_2)_6 \rightarrow \text{CO}_2^-$ in preference to one from C $_{11}$, *i.e.* to ensure that a transition state involving the three-carbon system C $_8$ ---C $_{10}$ rather than one involving the system C $_8$ ---C $_{11}$ becomes operative.

Turning to the diene system (F), the activation energy of the isomerisation which applies to the simple three-carbon system (*i.e.* $E - \alpha$) will here be further reduced by a quantity, γ , due to incipient conjugation resulting from the polarisation c , so that the activation energy is $E - (\alpha + \gamma)$



and since a condition of complete conjugation of the double bonds finally ensues, the resonance energy corresponding to this condition reduces the energy of the system (say by Y), so that the degree of conjugation in the transition state will be less than in the final state, *i.e.* $\gamma < Y$.† But it is important to note that the process which gives rise to Y is that which in a less complete manner gives rise to γ . The double-bond displacement then in the diene system is likely to be more easily achieved than in the mono-olefinic system, and, since the same (central) methylene group loses a proton whichever of the two three-carbon systems comes into play, the occurrence and direction of displacement are likely to be largely unaffected by polar groups terminating the carbon chain. From similar considerations it appears that double-bond displacement in rubber and the polyisoprene hydrocarbons is unlikely to occur with any facility under alkali-treatment, if at all.

It is of some interest that the polyisoprene hydrocarbon, squalene (C $_{30}$ H $_{50}$), shows a persistent exaltation of the molecular refraction of 1.6 units and the normal-chained unconjugated fish-oil acid, docosahexaenoic acid (C $_{22}$ H $_{32}$ O $_2$; F6), one of 0.8 units⁶ provided that each compound is scrupulously freed from its autoxidation products. Rubber also shows the same anomaly. Whether these exaltations are significant and refer

† The author gratefully acknowledges a discussion of this matter with Dr. L. C. Bateman.

⁶ Farmer and Sutton, *unpublished work*.

to an element of conjugation in the polyene chain analogous to that shown for the diene system in formula (G), which exists in spite of the fact that the double bonds are here separated by two methylene groups, remains to be tested by other methods.

Double-bond Displacement by Heat.

The above-mentioned optical anomalies were at first thought to be due to conjugation brought about by progressive heat-isomerisation during slow fractional distillation of the high-boiling materials; but this was not the case, since the anomaly appears in specimens which have not been heated above 100°. Prolonged heating of these latter specimens in entire absence of oxygen at temperatures a little below their decomposition points, has caused no changes in the absorption spectra which can be attributed to conjugation. Yet at high temperatures (above 270°, or possibly at somewhat lower temperatures in presence of a contact catalyst) olefinic hydrocarbons and the esters of vegetable oil acids give polymerised and cyclised products whose constitutions are such that conjugation can be fairly definitely concluded to have preceded polymerisation or cyclisation.⁷ This isomerisation must be attributed to radical dissociation of α -methylenic hydrogen atoms.

Vulcanisation.


The process of vulcanisation is usually hastened by the use of organic accelerators, some of which can themselves act as efficient vulcanisers, even when no sulphur is present. The basis of the success of these self-sufficient compounds appears to be their capacity for decomposing thermally to give free radicals which then act precisely analogously to dibenzoyl peroxide⁸ and bring about direct C—C linking of the rubber units at the α -methylenic carbon atoms; tetramethylthiuram disulphide,




one of the most efficient of these is a close sulphur-analogue of dibenzoyl peroxide, and polyalkyl-hydrazines, which are prone to decompose thermally to give radicals, have been found to have a vulcanising effect. Where sulphur alone, or sulphur together with an accelerator, forms the vulcanising agent, three processes of bridging seem to be possible, *viz.* (a) direct C—C linking at the α -methylene groups caused by radicals derived from peroxidic chain-starters contained in the rubber, or by radicals derived from the decomposing accelerator, or by atmospheric oxygen:—



(b) sulphur bridging at the α -methylene group, initiated by chain-starters:—



and (c) C—C linking brought about by sulphur functioning purely as a hydrogen-acceptor:—




With regard, however, to the mechanism by which sulphur becomes steadily and extensively incorporated in rubber during the later stages of reaction, with concurrent loss of the unsaturation of the latter, it would

⁷ Farmer, *Trans. Faraday Soc.*, 1939, **35**, 1039.

⁸ Cf. Farmer, preceding paper, p. 348.

† Or the corresponding disulphide-bridged compound.

be unprofitable without further investigation of sulphur-olefin reactions to attempt to formulate the precise sequence of events. There are, however, two possibilities of double-bond saturation in the vulcanisation reaction which do not necessitate recourse to postulation of the direct addition of sulphur in one or other of its known modifications at the double bonds. The first is addition at the double bonds of hydrogen sulphide generated by the attack of sulphur at the methylene groups (equation (c) above)—an addition which is known to be catalysed by sulphur, and the second a reaction analogous to that by which a molecule of oxygen becomes attached to an olefin at an α -methylene group



and then by secondary reaction attacks and saturates (inter- or intra-molecularly) a near-by double bond.

The Photo-gelling of Rubber.

This phenomenon is one of a group of insolubilisation reactions which are of embarrassingly frequent occurrence in connection with the manipulation of rubber and its derivatives, and for which the exact cause usually remains obscure. Solutions of rubber containing such "sensitisers" as ketones, aldehydes, carbon tetrachloride, chloroform, peroxides, etc., undergo rapid gelling in sunlight, the best sensitisers being those which break down photochemically to give free-radicals, and the poorest comparatively stable substances such as pure petroleum fractions and benzene. The very considerable extent to which fragments from the sensitisers or photochemically-decomposing solvents enter into the composition of the photo-gels, leaves little doubt as to the radical nature of the reaction, and there seems to be no difference at all in reaction-principle between the type of combined cross-linking and radical-addition here encountered and that observed with rubber containing thermally-decomposing dibenzoyl peroxide molecules. The increasing toughness of the photo-gels with increasing degree of reaction (increasing cross-linking) is impressive.

The role of oxygen in the photo-gelling reactions seems from the observations of Stevens to be a mixed one. As in the case of the substitutional chlorination of olefins, traces of oxygen appear to function catalytically, although in this case the gelling promoted is very slow; larger amounts of oxygen on the contrary hinder oxidation, and serve to cause severe oxidative degradation of the gels.

The above paper and the two preceding ones relate to subjects forming part of a programme of fundamental research undertaken by the Board of the British Rubber Producers' Research Association.

*British Rubber Producers' Research Association,
48 Tewin Road, Welwyn Garden City, Herts.*