


Keto–Enol Equilibria in Mixed Acetic Sulphuric Anhydrides †


By Enzo Montoneri,* Ezio Tempesti, Luigi Giuffrè, and Aldo Castoldi, Istituto di Chimica Industriale del Politecnico di Milano, Piazza L. da Vinci 32, 20133 Milano, Italy

Mixed acetic sulphuric anhydrides obtained from both equimolar $\text{SO}_3\text{--AcOH}$ and $\text{SO}_3\text{--Ac}_2\text{O}$ systems in liquid SO_2 are found at low temperature ($\leq -20^\circ$) in equilibrium between the keto ($\text{CH}_3\text{CO}\cdot\text{OSO}_3\text{Y}$) and enol ($\text{CH}_2\text{:COH}\cdot\text{OSO}_3\text{Y}$) forms ($\text{Y} = \text{SO}_3^-$ or Ac). At room temperature a cleavage reaction is shown to occur through the intermediacy of keten, which reacts further to yield oligomers and sulpho-derivatives [reaction (i)]. This




cleavage reaction appears specifically not to involve participation of acetylum ions, whereas previously these ions were normally considered as the principal precursor of keten intermediates.

Mixed anhydrides (RCOY ; $\text{Y} = \text{NO}_3$,¹ HCO_2 ,² CF_3SO_3 ,³ ClO_4 ,⁴ or halogen⁵) have been studied by numerous workers since they are known to be key intermediates in many acid-catalysed reactions. Under specific conditions some of these species^{3,5} have been shown to ionize to the corresponding acylium ions



[reaction (1)]; these ions, due to the ease of preparation and relatively high stability, constitute one of the best characterized classes of stable carbocations.

For their structure a resonance hybrid of the oxonium, oxocarbenium, and 'keten-like' forms is generally assumed [equilibrium (2)]. Specifically for acetylum



ions, participation of the 'keten-like' form has been reported to be negligible (crystalline salts)⁶ or substantial (in solution)⁷ according to the experimental conditions.

For pure mixed sulphuric carboxylic anhydrides ($\text{RCO}\cdot\text{OSO}_3\cdot\text{Y}$, $\text{Y} = \text{SO}_3^-$ or RCO), which have been characterized in equimolar solutions of carboxylic acids⁸ or anhydrides⁹ and SO_3 in liquid SO_2 , it has been also shown that on increasing the concentration of SO_3 ionization of these mixed anhydrides to acylium ions occurs.

We now report that specifically for $\text{Ac}_2\text{O}\text{--}\text{SO}_3$ and $\text{AcOH}\text{--}\text{SO}_3$ equimolar solutions in liquid SO_2 the formation of the corresponding mixed anhydrides is observed to involve at low temperature ($\leq -20^\circ$) the establishment of tautomeric equilibria not previously observed. We also report that depending on the experimental conditions the enol species involved in these tautomeric equilibria are cleaved at room temperature and imply the intermediacy of keten without any significant participation by acetylum ions.

† Reported in part at the 8th I.C.A.S. Conference, Cambridge, 1979.

EXPERIMENTAL

Most of the experimental details have been previously reported.^{8,9} In addition to i.r. and n.m.r. spectra, the $\text{Ac}_2\text{O}\text{--}\text{SO}_3$ and $\text{AcOH}\text{--}\text{SO}_3$ mixtures were analysed in other ways to supplement the spectroscopic data.

Classical Analyses.—*Analysis of sulphur derivatives.* Samples were quenched with ice and water and SO_2 was completely evaporated (as checked by iodometric titration on the resulting aqueous solution); free sulphates, organic sulphur, and total acidity were then determined as previously reported.

Olefinic double bonds. To obtain evidences of olefinic double bonds in unquenched samples chlorine (in slight excess relative to initial Ac_2O or AcOH) was added at -70° to the SO_2 solutions; after quenching with ice and water, the determination of the relative concentrations of chloroacetic and acetic acid was performed either by n.m.r. spectroscopy (free acids) or by g.l.c. (butyl esters).

Autocondensation products. The test to obtain evidence of the formation of aceto- and diaceto-acetic species in $\text{SO}_3\text{--}\text{SO}_2$ mixtures with AcOH or Ac_2O included quenching of samples with ethanol at -50° and determination of the corresponding esters as previously reported.⁷ Dehydroacetic acid was determined after quenching the samples with ice and water as reported above, extracting the aqueous solutions with CCl_4 , and concentrating the organic layer. The crystalline (-10°) product was identified by its i.r. and n.m.r. spectra which were identical with those of an authentic sample.

Elaboration of N.m.r. Data.—*Determination of mole ratio of diacetyl sulphate enol (II) and keto (I) forms.* From relative band areas at 1.0 mole ratio (Table 2a), a tentative quantitative determination of mole ratio of (II) to (I) at equilibrium may be obtained from equation (3) where n_{CH_3} , n_{CH_2} , and n_{OH} are the numbers of methyl, methylene,

$$(3n_{\text{CH}_3} + 2n_{\text{CH}_2})/n_{\text{OH}} = A_{\text{CH}_3}/A_{\text{OH}} = 8 \quad (3)$$

and hydroxy groups, respectively, present at equilibrium and A is the band area. The assumption that $A_{\text{OH}} = 1$ and $n_{\text{CH}_3} = n_{\text{OH}}$ [based on structure (II)] yields $n_{\text{CH}_3}/n_{\text{CH}_2} = 2$ and a (II)/(I) mole ratio of 2. Furthermore the chemical shift values for the methyl and methylene protons (δ° 2.32 and 2.57, respectively) in species (I) and (II) may be ten-

tatively obtained. As shown for the AcOH system (Table 2d), where separate n.m.r. bands are observed for AcO_2O_6^- (δ 2.50) and $\text{CH}_2\text{COH}\cdot\text{OS}_2\text{O}_6^-$ (δ 2.75), the methylene proton resonance (enol form) is 0.25 p.p.m. downfield from that of the methyl protons (keto-form). Assuming then the same $\Delta\delta$ for (I) and (II), from the (II)/(I) mole ratio and equations (4) and (5), where X = proton fraction of methyl

$$X_{\text{CH}_3}\delta^0\text{CH}_3 + X_{\text{CH}_2}\delta^0\text{CH}_2 = 2.38 \quad (4)$$

$$\delta^0\text{CH}_3 + 0.25 = \delta^0\text{CH}_2 \quad (5)$$

and methylene groups and 2.38 is the observed δ at 1.0 mole ratio, the δ^0 values are obtained.

Determination of the relative concentrations of the acetyl disulphate enol (VIII) and keto (VII) forms and of protonated acetic acid. From the band areas of acetyl disulphate [keto-form (VII) δ 2.50; enol form (VIII) δ 2.75] and of protonated acetic acid (δ 2.66) the relative concentrations of each species [(VIII) : (VII) : AcOH_2^+ = 0.4 : 0.6 : 1.0] can be obtained. Since only acetyl disulphate is involved in the tautomeric equilibrium (6) with the aid of equations (7) and (8), where n and A are the moles and band areas for each species, the mole ratios may be obtained.

$$n(\text{AcOH}_2^+) = n(\text{VII}) + n(\text{VIII}) \quad (6)$$

$$n(\text{VII}) = A(\text{VII})/3 \quad (7)$$

$$3n(\text{AcOH}_2^+) + 2n(\text{VIII}) = A(\text{AcOH}_2^+) + A(\text{VIII}) \quad (8)$$

RESULTS

The behaviour of equimolar $\text{Ac}_2\text{O}-\text{SO}_3$ and $\text{AcOH}-\text{SO}_3$ solutions in liquid SO_2 was investigated between -78° and room temperature. I.r. spectra were always recorded at -60° and n.m.r. spectra at -78° and are reported in Tables 1 and 2, respectively. Liquid SO_2 bands are omitted from the i.r. spectra. Spectra a and c * refer to freshly

TABLE 1

I.r. frequencies (cm^{-1}) of equimolar $\text{Ac}_2\text{O}-\text{SO}_3$ and $\text{AcOH}-\text{SO}_3$ systems in liquid SO_2 at -60°

System	$\text{Ac}_2\text{O}-\text{SO}_3$		$\text{AcOH}-\text{SO}_3$		
Spectrum	a	b	c	d	e
	2 950w,br	3 000s,br	2 800m,br	2 850s,br	2 850s,br
			1 875w,br	1 875w,br	
			1 805ms	1 805w	
1 820s					1 710m,br
		1 725s,br			
1 680w					
1 600m		1 580s,br	1 612s	1 612s	1 612s
1 555m		1 545s,br	1 545s	1 545s	1 545s
1 416s			1 405s	1 405w	
1 375s		1 365s,br	1 370m	1 370m	1 370s,br
1 230m					
1 218s					
1 185ms	1 170s,br		1 200s,br	1 200s,br	1 212s,br
1 045m	1 030s,br		1 045m	1 040m	1 040m
998ms			995m		
940m	925s,br		930m	940m	935s
850s	885sh		850m	892w	895sh
802s	810m,br				795s
765m			760s	775ms	705m

prepared (at -60°) mixtures; the other spectra were recorded after the mixtures had been left at -20° for 48 h (spectrum d) and at room temperature for 90 min (spectra b and e). The spectra (which are not reported in the Tables)

* The values reported in column c are taken from ref. 8. The bands at 1405 and 1370 cm^{-1} appear as shoulders of the SO_2 band at 1340 cm^{-1} , which in this system is quite broad.

of freshly prepared 0.0 and 0.5 $\text{SO}_3:\text{Ac}_2\text{O}$ mole ratio mixtures contained the following bands: δ 2.18; ν 1 821s, 1745s, 1 430m, 1 375ms, 1 230w, 1 045w, 1 000ms, 900m, 810w, and 780w cm^{-1} (0.0 mole ratio); δ 2.30 (19 H) and 15.10 (1.0 H) (0.5 mole ratio). Dehydroacetic acid formation (10 and 1% w/w yields relative to initial Ac_2O and AcOH, respectively) and loss of free sulphate (75 and 50%

TABLE 2

Proton chemical shifts (δ values, tetramethylsilane as internal standard) of $\text{Ac}_2\text{O}-\text{SO}_3$ and $\text{AcOH}-\text{SO}_3$ system in liquid SO_2


System	$\text{Ac}_2\text{O}-\text{SO}_3$		$\text{AcOH}-\text{SO}_3$		
Spectrum	a	b	c	d	e
	2.38	2.0	2.61	2.50	2.15
	(8.0 H)	(2.5 H)	(3.0 H)	(0.85 H)	(0.14 H)
		2.24		2.66	2.66
		(0.15 H)		(1.6 H) *	(0.83 H)
		2.62		2.75	3.41
		(0.15 H)			(0.10 H)
		4.10			4.40
		(1.3 H)			(0.28 H)
					5.04
					(0.03 H)
					5.38
					(0.05 H)
					5.80
					(0.05 H)
					13.49
					(1.0 H)
	15.13	12.65	14.47	14.52	
	(1.0 H)	(1.0 H)	(1.0 H)	(1.0 H)	

* The integrated intensity refers to both δ 2.66 and 2.75 bands since they were both broad and separate values could not be obtained. The latter band was however much weaker.

mol per mol relative to initial SO_3 in the Ac_2O and AcOH systems, respectively) and of organic acidity (15 and 25% relative to initial Ac_2O and AcOH, respectively) occurred only in mixtures from which spectra b and e are derived. Addition of Cl_2 to yield chloroacetic acid (MCA) occurred only in mixtures from which spectra a and d are derived with 30 and 25% mol per mol yields (relative to initial Ac_2O and AcOH), respectively. No evidence of the formation of aceto- and diaceto-acetic species was obtained.

DISCUSSION


(a) *Tautomerization of Mixed Acetic Sulphuric Anhydrides.*—(1) *The $\text{Ac}_2\text{O}-\text{SO}_3$ system at low temperature* ($\leq -20^\circ$). The establishment of tautomeric equilibria involving mixed acetic sulphuric anhydrides has been thoroughly investigated in this work starting from a mixture of Ac_2O and SO_3 in SO_2 in order to eliminate spectroscopic interferences due to the presence of acid protons (e.g. in $\text{AcOH}-\text{SO}_3$). The equimolar addition of SO_3 to Ac_2O at -60° leads quantitatively to diacetyl sulphate formation [reaction (9)] on the basis of the



following spectroscopic evidence. The absence of Ac_2O C=O bands (ν 1 745 and 1 821 cm^{-1}) in the i.r. spectrum (Table 1a) shows that the conversion of acetic anhydride is complete. The absorptions at ν 1 820 (C=O), 998 and 1 045 (C-O), 1 416 and 1 218 (S=O), and 765–850 (S-O) cm^{-1} are characteristic of diacetyl sulphate (I), and similar to those previously reported for biscyclohexylcarbonyl sulphate.⁹ Two additional bands (1 600 and 1 555 cm^{-1}) indicated however, the presence of

a new species. If we assume that a donor (carbonyl and not ether oxygen⁹)-acceptor complex, such as $\text{Ac}_2\text{O}\cdot\text{SO}_3$ is formed (see *e.g.* the donor-acceptor complexes of carboxylic anhydrides with metal halides),¹⁰ then we would expect to find additional S=O absorptions at wavelengths lower than 1390 and 1065 cm^{-1} (free SO_3),¹¹ respectively. The two bands, instead, which have been identified as additional S=O absorptions fall at 1375 and 1185 cm^{-1} . Since a broad band is also observed around 2950 cm^{-1} (which is reminiscent of a strong hydrogen-bonded OH function)¹² the formation of a new enolic species $\text{CH}_2\cdot\text{COH}\cdot\text{OSO}_2\text{Ac}^*$ in equilibrium with the keto-form (I) seems plausible.

Referring again to the new bands at 1600 and 1555 cm^{-1} , it is known that these are frequently indicative of



enolic compounds. In β -diketo-compounds, for instance, strong absorptions in this range (1640–1540 cm^{-1}) have been assigned to the C=O (chelated) and C=C stretching vibrations.¹⁴ A pair of absorptions in this range, however, does not necessarily require a β -diketo-structure; in fact it may also be due to protonated C=O groups as in (III) where intramolecular interaction between the proton formally attached to the carbonyl oxygen atom and hydroxy ($\text{R}' = \text{H}$) or ether ($\text{R}' = \text{alkyl}$) oxygen atoms may occur.¹⁵ Furthermore Hunt and Satchell¹⁶ suggested that in some cases, even if a β -diketo-chelated structure is present (*e.g.* the enolic form of the $\text{Ac}_2\text{O}\cdot\text{SnCl}_4$ donor-acceptor complex), the absorption between 1630 and 1500 cm^{-1} would rise from the cyclic chelated structure (IV) and not from C=O stretching vibrations.

Based on these findings, the 1600 and 1555 cm^{-1} absorptions in Table 1a can be due to the enolic form of diacetyl sulphate (I) where strong intramolecular hydrogen bonding of the chelate ring structure (II) occurs.[†]

* Only the keto-form was observed at low temperature [$\leq -20^\circ$] for biscyclohexylcarbonyl sulphate⁹ and propionyl sulphate.¹³

† Stable six-membered chelate ring systems have been experimentally established in carbohydrates¹⁷ and in enolized systems.¹⁸

‡ Although intense methyl absorptions may also occur at 1375 cm^{-1} (*e.g.*, that of Ac_2O), it may reasonably be assumed from the following discussion that in this case the band is due to ν_{as} ($\text{S} \cdots \text{O} \cdots \text{H}$).

Accordingly, the remaining new bands reported in Table 1a, may be assigned as follows. The broad absorption (in the 3400–2300 cm^{-1} range) centred at 2950 cm^{-1} relates to the bonded hydroxy-group in the chelated form,¹² while the 1375 \ddagger and 1185 cm^{-1} bands may be attributed to hydrogen-bonded sulphuryl groups ($\text{S} \cdots \text{O} \cdots \text{H}$). The latter frequencies, which are almost equally displaced from the free S=O [as in (I)] absorptions, exhibit absorption shifts of the same magnitude as observed from hydrogen-bonding studies with sulphones and sulfoxides.¹⁹ Finally the weak band at 1680 cm^{-1} could be tentatively assigned to the stretching vibration of the C=C bond \S in (II) by analogy with the corresponding absorption (1672 cm^{-1}) in isopropenyl acetate.²¹

Further information on structure (II) can be obtained from n.m.r. spectroscopy. The high-field resonance dependent on the SO_3 mole ratio between 0.0 and 1.0 may be attributed to the carbon protons of fast exchanging Ac_2O , (I), and (II) species. The value (δ 2.38) at 1.0 mole ratio (Table 2a) is, however, due to exchange of species (I) and (II) only since conversion of Ac_2O is shown to occur quantitatively from the i.r. spectrum.


Tentative chemical shift values for pure methyl (δ^0 2.32) and methylene (δ^0 2.57) protons in species (I) and (II) have been obtained (see Experimental section). The methyl proton shift in diacetyl sulphate is 0.14 p.p.m. deshielded relative to the value observed for pure methyl protons in Ac_2O . This deshielding is well within the range observed for other mixed carboxylic sulphuric anhydrides when they are compared with the corresponding pure carboxylic species.^{8,9}

The low-field band (δ 15.13) indicates a substantially deshielded exchangeable proton associated with at least two oxygen atoms such as those of a very strong acid or a strong intramolecularly hydrogen-bonded species. The behaviour of this band is quite peculiar; although the relative intensity of the low- to high-field proton bands increases from 0.0 to 0.12 upon changing the mole ratio from 0.0 to 1.0, indicating increasing concentration of hydroxy-protons, the hydroxy-proton chemical shift is constant. This may be assumed to be consistent²² with a strong intramolecular-hydrogen-bonded hydroxy-proton such as in (II), whose chemical shift is not dependent on concentration.

(2) *The $\text{Ac}_2\text{O}\cdot\text{SO}_3$ system at room temperature.* Starting from Ac_2O and H_2SO_4 (or AcOH and SO_3) the room temperature formation of a stable sulphur derivative, sulphoacetic acid, is known. This sulphonation reaction has been assumed²³ to proceed through the intermediacy of both acetylsulphuric acid and diacetyl sulphate. While evidence for the existence of acetylsulphuric acid has been reported^{8,24} no direct observation of diacetyl sulphate is known. We have shown here that at low

\S The presence of an olefinic double bond has been confirmed by chlorination as indicated above; the formation of MCA occurs immediately at low temperature, while it is known that, normally,²⁰ starting from AcOH , high temperatures and long induction periods are required.

The n.m.r. spectrum of an equimolar $\text{Ac}_2\text{O-SO}_3$ solution, which contains at low temperature a single low field proton band (δ 2.38), yields at room temperature two main bands at δ 2.0 and 4.1 and two weak bands at δ 2.24 and 2.62 (Table 2b). These latter were identified as the methyl proton bands of dehydroacetic acid, which was recovered from the reaction mixture with a yield in agreement with its spectroscopic concentration. This oligomerization may occur through thermal cleavage of (II) to acetylsulphuric acid and keten [reaction (10)]



In this system the participation of diacetyl sulphate in


tautomerization cannot be established unambiguously due to its low concentration. No appreciable modification is indeed observed in the weak 1875 cm^{-1} band, which in the freshly prepared mixture had been previously assigned to sulphur-oxygen protonated diacetyl sulphate.

Finally neither AcOSO_3^- nor AcOH_2^+ appear involved in tautomerization equilibria since no changes are observed in the 0.5 mole ratio spectra on standing at -20° .


The modifications observed in the i.r. spectra are in fair agreement with those observed in the n.m.r. spectra. At a SO_3/AcOH mole ratio of 1.0, the n.m.r. spectrum (Table 2c) of the freshly prepared mixture contains a single high-field proton band at $\delta\ 2.61$, which has been previously justified with the rapid exchange of equimolar $\text{AcOS}_2\text{O}_6^-$ and AcOH_2^+ species. At -20° , however, this band splits (Table 2d) and the ratio of the high-field ($\delta \leq 2.75$) band area sum to the low-field ($\delta \geq 14.47$) band area decreases from 3 to 2.45 (spectra c and d, respectively), indicating increasing concentration of acid protons. The $\delta\ 2.50$ chemical shift is in fair agreement with that previously extrapolated⁸ for the methyl protons of acetyl disulphate. The other bands may be assigned to the methyl protons of AcOH_2^+ ($\delta\ 2.66$) and to the methylene * protons of $\text{CH}_2\text{:COH}\cdot\text{OS}_2\text{O}_6^-$ ($\delta\ 2.75$).

Finally the relative equilibrium concentration of (VIII), (VII), and AcOH_2^+ can be obtained (see Experimental section).

(4) *The $\text{AcOH}\text{--}\text{SO}_3$ system at room temperature.* While no changes are observed at the SO_3/AcOH 0.5 mole ratio, thus indicating that AcOSO_3^- is stable even at room



temperature, at the SO_3/AcOH 1.0 mole ratio at the same temperature we found that $\text{AcOS}_2\text{O}_6^-$ undergoes thermal cleavage † to keten [reaction (17)] which may further react by (18) and (19).



The n.m.r. bands initially (Table 2d) assigned to (VII) ($\delta\ 2.50$) and (VIII) ($\delta\ 2.75$) are absent from the spectrum (Table 2e) of the room temperature sample, whereas a major band at $\delta\ 4.4\ \ddagger$ is observed and can be assigned to the methylene protons of sulphaacetic acid (IX). The $\delta\ 2.66$ band may be common to both AcOH_2^+ and AcOSO_3H ; the other bands at $\delta\ 2.15$, 3.41 , and $5.0\text{--}5.5$ may be due to other oligomerization products which, except for dehydroacetic acid, were not identified.

The i.r. evidence agrees with the n.m.r. data. A new broad band at 1710 cm^{-1} (Table 1e), which may be assigned to $\text{HO}_3\text{SCH}_2\text{CO}_2\text{H}$,^{24b} appears in place of the

* The presence of an olefinic double bond inferred from spectroscopic data has been further supported through chlorination experiments.

1805 cm^{-1} band of (VII) while two strong bands at 1612 and 1545 cm^{-1} can be assigned to both AcOH_2^+ and AcOSO_3H . Finally the absorptions at 795 and 705 cm^{-1} are due to HS_2O_7^- ,²⁹ which is not completely converted through equation (18) since (19) occurs too.

(b) *Effect of Excess of SO_3 on the Tautomerization of Mixed Acetic Sulphuric Anhydrides.*—Both systems (Ac_2O — or $\text{AcOH}\text{--}\text{SO}_3$) upon increasing the SO_3 mole ratio above 1.0 evolve unequivocally towards the formation of acetylum ions. The complex equilibria involved in this case will be dealt with in a subsequent paper.

(c) *Mechanistic Considerations on the Acetylating (and/or Sulphonating) Properties of Acetyl Sulphates.*—Acetylation performed with acetyl sulphates²³ are known to exhibit complications since, depending on the experimental conditions, these species may also induce intra- or inter-molecular sulphonation.²⁴ With other acylating systems (acyl and metal halides)⁵ it has been shown that, according to the system considered, acylations may occur with or without acylium ion participation; in the former case the involvement of 'keten-like' species has been considered as in (2).⁶ In agreement with this assumption, starting from acids or anhydrides in sulphuric media, due to the complications reported above, previous investigators have not considered the possibility of the involvement of keten species without the unequivocally established participation of acetylum ions.³⁰

The involvement of keten in sulphuric acid reactions³¹ may, however, occur on raising the temperature (80°). In this case it has been inferred that the active precursor of a keten intermediate^{26,27b} is a mixed carboxylic sulphonic anhydride (RCOY ; R = cyclohexyl, Y = 1-carboxycyclohexanesulphonate). It has now been found that starting from mixed carboxylic sulphuric anhydrides such as acetyl sulphates, even at lower temperatures keten oligomers may be isolated without evident participation of acetylum ions. This may be


due to the higher (relative to mixed carboxylic sulphonic anhydrides) electron-withdrawing effect of AcOSO_3 or S_2O_6^- moieties present in (I) and (VII) respectively, which through an evident enolization step facilitates the mechanism of thermal cleavage [reaction (20)].

On the basis of this finding it becomes necessary to reconsider some controversial aspects related to other acylating systems. For $\text{Ac}_2\text{O}\text{--}\text{SnCl}_4$, the formation of condensation products (oligomers) has been suggested by Hunt and Satchell¹⁶ as a consequence of preliminary



† The participation of diacetyl sulphate, however, cannot be rigorously excluded.

‡ A medium effect may be responsible for the 0.35 p.p.m. low-field-shifted value relative to that ($\delta\ 4.05$) reported for the Ac_2O system.^{24b}


enolization* of the active acetylating intermediate (the donor-acceptor complex $\text{Ac}_2\text{O}\cdot\text{SnCl}_4$; based on i.r. data, no mention is made of Ac^+ participation). On the other hand, starting from AcCl and AlCl_3 ,⁷ a plausible mechanism has recently been reported for the formation of



diacetoacetylum ions in which the rate-limiting step is assumed to be the deprotonation (at low acidities) of acetylum ions [reactions (21) and (22)]. Although



equilibrium (21) seems justified on the basis of the time dependent i.r. absorption intensity of Ac^+ ($2\,300\text{ cm}^{-1}$), no such evidence was reported for a very similar system ($\text{AcBr}\cdot\text{AlBr}_3$)³³ though diacetoacetylum ions were again shown to be present. Furthermore, referring again to equilibrium (21), while the deprotonation of the acetoacetylum ion has been observed directly in super-acid systems³⁴ through an enolization step which seems necessary in terms of charge delocalization [reaction (23)], no such observation has been reported directly for



acetylum ions. Although basicity of the system may be necessary for the establishment of equilibrium (21), other arguments also come into consideration. So far as our system is concerned, we have to admit that the involvement of 'keten-like' species mainly implies the enolization of acetyl sulphates without evident participation of acetylum ions.

[9/705 Received, 8th May, 1979]

* Somewhat confusing evidence of enolization in 1 : 1 acetyl-metal halide complexes leading to the formation of the 'Illari' compound, $\text{CH}_2\cdot\text{CClO}\cdot\text{AlCl}_2$, has been produced.³²

REFERENCES

- J. Chedin, S. Fernant, and R. Vandoni, *Mem. Serv. Chim. Etat*, 1950, **35**, 53.
- W. Stevens and A. Vas Es, *Rec. Trav. chim.*, 1964, **89**, 863.
- A. Germain, A. Commeyras, and A. Casadevall, *Bull. Soc. chim. France*, 1973, 2527.
- A. M. Avedikian and A. Commeyras, *Bull. Soc. chim. France*, 1970, 1258.
- B. Chevrier and R. Weiss, *Angew. Chem. Internat. Edn.*, 1974, **13**, 1.
- G. A. Olah, A. Germain, and A. M. White, in 'Carbonium Ions' eds. G. A. Olah and P. v. R. Schleyer, Wiley-Interscience, New York, 1976, ch. 35.
- A. Germain, A. Commeyras, and A. Casadevall, *Bull. Soc. chim. France*, 1972, 3177.
- E. Montoneri, M. Fornaroli, L. Giuffr , E. Tempesti, and G. Sioli, *J.C.S. Perkin II*, 1976, 1784.
- E. Montoneri, L. Giuffr , M. Cassago, E. Tempesti, and M. Fornaroli, *Canad. J. Chem.*, 1977, **55**, 355.
- C. Cook, *Canad. J. Chem.*, 1962, **40**, 445.
- G. E. Walrafen, *J. Chem. Phys.*, 1964, **40**, 2326.
- S. Bratoz, D. Hazdi, and G. Rossmy, *Trans. Faraday Soc.*, 1956, 464.
- E. Montoneri, unpublished data.
- H. Ogoshi and K. Nakamoto, *J. Chem. Phys.*, 1969, **45**, 3113.
- C. J. Clemett, *Chem. Comm.*, 1970, 211.
- P. Hunt and D. P. N. Satchell, *J. Chem. Soc.*, 1964, 5437.
- L. Mester, A. Stephen, and J. Parello, *Tetrahedron Letters*, 1968, 4119.
- S. Forsen and M. Nillson, *Acta. Scand.*, 1960, **14**, 1333; L. D. Reeves, *Canad. J. Chem.*, 1957, **35**, 1351; J. L. Burdett and M. T. Rogers, *J. Amer. Chem. Soc.*, 1964, **86**, 2105.
- D. Barnard, J. M. Fabian, and H. P. Koch, *J. Chem. Soc.*, 1949, 2242; 1953, 6305.
- W. Richarz and A. Mathey, *Chimia*, 1967, **21**, 388.
- G. C. Smith, *J. Amer. Chem. Soc.*, 1953, **75**, 1134.
- G. A. Olah, K. Dunne, Y. K. Mo, and P. Szilagyi, *J. Amer. Chem. Soc.*, 1972, **94**, 4200.
- E. A. Jeffery and D. P. N. Satchell, *J. Chem. Soc.*, 1962, 1887.
- A. Casadevall and A. Commeyras, *Bull. Soc. chim. France*, 1970, (a) 1850; (b) 1856.
- M. H. Karger and Y. Mazur, *J. Org. Chem.*, 1971, **36**, 528; E. Tempesti, L. Giuffr , G. Sioli, M. Fornaroli, and G. Airoidi, *J.C.S. Perkin I*, 1974, 771.
- A. Casadevall and A. Commeyras, *Bull. Soc. chim. France*, 1970, 1850.
- (a) E. Tempesti, L. Giuffr , M. Fornaroli, and G. Airoidi, *Chem. and Ind.*, 1973, 183; (b) E. Tempesti, C. Buzzi Ferraris, G. Airoidi, and E. Montoneri, *ibid.*, 1978, **60**, 401.
- L. J. Bellamy, 'The I.r. Spectra of Complex Molecules', Methuen, London, 1960.
- G. E. Walrafen, D. R. Irish, and T. F. Young, *J. Chem. Phys.*, 1962, **37**, 662.
- N. C. Deno, C. U. Pittman, jun., and M. J. Wisotsky, *J. Amer. Chem. Soc.*, 1964, **86**, 4370.
- L. Giuffr , E. Tempesti, G. Sioli, M. Fornaroli, and G. Airoidi, *Chem. and Ind.*, 1971, 1098.
- B. P. Susz and J. J. Wuhrmann, *Helv. Chim. Acta*, 1957, **40**, 971.
- A. Germain, J. L. Pascal, and J. Potier, *Canad. J. Chem.*, 1977, **55**, 3096.
- G. A. Olah, A. Germain, C. Lin, and K. Dunne, *J. Amer. Chem. Soc.*, 1975, **97**, 5477.