

PRIMER NOTE

Polymorphic microsatellite loci from the Southeast Asian cyprinid, *Barbodes gonionotus* (Bleeker)

W. KAMONRAT,* S. K. J. McCONNELL† and D. I. COOK‡

*Ornamental Fish Research Institute, Department of Fisheries, Chatuchak, Bangkok 10900, Thailand, †School of Biological Sciences, University of Wales, Swansea, Singleton Park, Swansea SA2 8PP, UK, ‡Department of Biology and Marine Gene Probe Laboratory, Dalhousie University, Halifax, Nova Scotia, Canada, B3H 4J1

Abstract

The cyprinid *Barbodes gonionotus* (Bleeker) is a commercially important fish in both capture fisheries and aquaculture in Southeast Asia. Five polymorphic microsatellite loci from *B. gonionotus* are described. Four are highly variable, with 9–30 alleles observed per locus in four populations sampled from Thailand ($H_O = 0.694–0.808$). These will be of use in studies of population genetic structure and in pedigree analyses.

Keywords: *Barbodes* spp., cyprinid, genetic markers, microsatellites, silver barb

Received 22 June 2001; revision accepted 14 September 2001

Barbodes gonionotus (Bleeker, 1850) is a cyprinid fish, common in large river systems in Thailand, Laos, Cambodia, Vietnam, Sumatra and Java. It has also been widely introduced outside this range. In Thailand, the species occurs throughout the country, but is most abundant in major rivers of the central region, such as the ChaoPhaya, MaeKlong, NakornNayok and PaSak (Smith 1945). It comprises approximately 20% of the total inland fishery (25 000 metric tons; Anonymous 1996) and since 1960, the Thai Department of Fisheries has successfully used induced spawning to produce *B. gonionotus* fingerlings for restocking and aquaculture. Here we describe five microsatellite loci developed from *B. gonionotus*.

B. gonionotus samples were obtained either directly from the rivers or from local fishermen during April to August 1994. A muscle tissue sample was collected from each fish and preserved in 100% ethanol. Nuclear DNA was extracted using a standard phenol extraction procedure. Genomic DNA from a single *B. gonionotus* (50 µg) was digested with the restriction endonucleases; *PvuII*, *RsaI*, *HincII* and *AluI* (Pharmacia). Genomic fragments of 300–700 bp were recovered from a 1% agarose gel using a phenol freeze fracture procedure (Qian & Wilkinson 1991). The size selected DNA was ligated to *SmaI*-digested and dephosphorylated pUC18 vector (Pharmacia) and trans-

formed into MAX efficiency Δ H5 α ™ competent cells (BRL). Transformed cells were plated on 2 \times YT medium (1.6% Bactotryptone, 1% yeast extract, 0.1 M NaCl) containing 100 µg/mL of ampicillin, and grown out at 37 °C overnight. Colonies were lifted onto Hybond-N nylon filters (Amersham).

A hybridization probe was prepared by end-labelling 100 pm of a synthetic oligonucleotide [(GT)₁₅]. The filters were prehybridized in [6.25 mL 20 \times SSPE pH 7.6 (174 g NaCl, 27.6 g NaH₂PO₄·H₂O, 7.4 g EDTA and ddH₂O to a total volume of 1.0 litre), 2.5 mL 50 \times Denhardt's solution (5 g Ficoll-Type 400, 5 g Polyvinyl pyrrolidone, 5 g BSA and ddH₂O to a total volume of 500 mL), 0.5% SDS (w/v), 100 µL/mL tRNA] at 65 °C for 1 h prior to addition of heat denatured (70 °C for 5 min) labelled probe. Hybridization was carried out at 60 °C for 12 h. The filters were washed three times; 2 \times SSC, 0.1% SDS for 15 min at room temperature, 1 \times SSC, 0.1% SDS for 15 min at room temperature, and 1 \times SSC, 0.1% SDS for 15 min at 60 °C and exposed to autoradiographic film.

Eighty-eight positive colonies were identified and sequenced. Thirty-nine clones contained microsatellites, five of which contained two or more microsatellite loci (separated by three or more bases). Three clones contained tetranucleotide microsatellites and two contained trinucleotide microsatellites. Thirty-six clones contained dinucleotide microsatellites, 23 of which were (AC/GT)_n repeats (26.1%). Most of the *B. gonionotus* microsatellites found in this study were perfect repeats (73.2%), four clones

Correspondence: W. Kamonrat. Fax: 662 5580177; E-mail: wongpatk@fisheries.gov.th

Table 1 Details of five microsatellite loci developed from *Barbodes gonionotus*. H_O , observed heterozygosity; H_E , expected heterozygosity; T_a , annealing temperature

Locus	Accession number	Motif	Primers 5'–3'	Allele size range (bp)/ (sequenced allele size)	No. of alleles	H_O/H_E	T_a (°C)
<i>Bgon-17</i>	AJ291684	(AC) ₃₄	TTACAAGGGGTTACATACTG CAGTCTCATATTTGAAAGCAG	108–182 (152)	30	0.808/0.885	49
<i>Bgon-22</i>	AJ291680	(TCC) ₆	TCTTGTTGATCACACGGACG GTGACTGTATCAATGAGTCTG	73–99 (89)	2	—	49
<i>Bgon-69</i>	AJ291681	(TG) ₁₂	GCAAAGGTTCTGTCAAGG GTATCCAGAAACATGTTTCAG	87–103 (95)	9	0.694/0.75	49
<i>Bgon-75</i>	AJ291682	(AC) ₁₀	CTGGTAAAGACTTCAGATGC GCATGCCAAATGAGAAAGGCT	96–118 (104)	11	0.729/0.759	53
<i>Bgon-79</i>	AJ291683	(CA) ₁₂	GCCAGACTGGAGCGAGG GTTCCGTGAAGCCATGAGG	101–137 (109)	17	0.789/0.783	53

contained imperfect repeats (9.8%), and seven clones contained compound repeats (17%). The motif copy-number varied from four to 35 repeats, the most common size of dinucleotide motif was 12, but tri- and tetranucleotide microsatellite loci were comprised of a smaller number of repeats. The largest motif copy-number of tri- and tetranucleotides microsatellite loci found was 10, and the averages were eight and seven repeats, respectively. The polymerase chain reaction (PCR) primers were designed from the unique flanking sequences of five microsatellites (Table 1) and tested using four unrelated populations from Thailand: Chainat, Khonkaen, Phitsanulok and Surin. One primer of each pair was end-labelled with [γ^{32}]-PATP. Five microlitre PCR reactions contained 1× PCR buffer (10 mM Tris pH 8.3, 50 mM KCl, 1 mM MgCl₂, 0.01% gelatin), 300 µM each dNTP, 0.6 µM each of primer, 0.025 µM [γ^{32}]-PATP labelled primer, 0.25 U *Taq* polymerase and 0.05% Tween20, overlaid with mineral oil. PCR reactions with *Bgon-17* and *Bgon-69* included: 0.05 mM [γ^{32}]-PATP labelled primer, 0.6 mM forward primer and 0.075 µM reverse primer. Programme parameters were: six cycles of 1 min at 94 °C, 30 s at a primer specific annealing temperature (T_a) and 30 s at 72 °C; 39 cycles of 30 s at 90 °C, 30 s at T_a and 30 s at 72 °C. Amplification products were denatured in stop dye (99% formamide, 0.1% bromophenol blue, 0.1% xylene cyanol green, 10 mM NaOH; 95 °C for 15 min), and 2.5 µL was run on an 8% denaturing polyacrylamide gel for 2 h at 55 mV. Gels were dried and exposed overnight to an autoradiography film (Kodak XAR5) at –80 °C. Microsatellite alleles were sized relative to a sequencing ladder generated from single-stranded M13 DNA.

Four loci were highly polymorphic in the population samples surveyed with heterozygosities ranging from 0.62 to 0.9. Thirty alleles were observed at locus *Bgon-17* with sizes ranging from 108 to 182 bp, nine alleles at locus *Bgon-69* with sizes ranging from 87 to 103 bp, 11 alleles at locus *Bgon-75* with sizes ranging from 96 to 118 bp, and 17 alleles

at locus *Bgon-79* with sizes ranging from 101 to 137 bp. At *Bgon-22* only two alleles were observed in the populations surveyed. All loci exhibited consecutive alleles that differed by multiple numbers of the microsatellite motif except for *Bgon-69* where one allele, *allele*92*, differed by only 1 bp (2% and 6% frequency in Chainat and Phitsanulok, respectively).

Genotypic frequencies at the four microsatellite loci; *Bgon-17*, *Bgon-69*, *Bgon-75* and *Bgon-79*, conformed to Hardy–Weinberg expectations (with sequential Bonferroni correction for multiple testing; results not shown). Allelic segregation was tested in a full-sib family consisting of 20 progeny (results not shown). The expected genotypes were observed in F₁ individuals. A significant deviation of gametic segregation from a 1:1 expectation was observed for the female parent at *Bgon-69* ($P < 0.05$) possibly due to a sampling error because only 20 offspring were studied. These polymorphic microsatellite loci will be of use both for pedigree analysis and for population genetic studies in this species.

Acknowledgements

This work was supported by the International Development Research Centre (IDRC, Canada) under the Fish Genetics (Thailand Project 89–0219) and the Aquatic Biodiversity (Project 92–0032), and by an NSERC research grant to Roger W. Doyle of Dalhousie University, Halifax, Canada.

References

- Anonymous (1996) *Fisheries Statistics Sub-Division*. Department of Fisheries, Bangkok.
- Qian L, Wilkinson M (1991) DNA fragment purification: removal of agarose 10 minutes after electrophoresis. *Biotechniques*, **10**, 736–738.
- Smith HM (1945) *The Fresh-Water Fishes of Siam or Thailand*. Smithsonian Institution, Washington D. C.