



An unusual, but not unexpected, evolutionary step taken by syrphid flies: the first record of true primary parasitoidism of ants by Microdontinae

GABRIELA PÉREZ-LACHAUD^{1*}, MARK A. JERVIS², MENNO REEMER³ and JEAN-PAUL LACHAUD^{1,4}

¹*El Colegio de la Frontera Sur, Conservación de la Biodiversidad, Avenida Centenario Km 5.5, Chetumal 77014, Quintana Roo, Mexico*

²*Cardiff School of Biosciences, Cardiff University, Cardiff CF10 3AX, UK*

³*Naturalis Biodiversity Center, European Invertebrate Survey, PO Box 9517, 2300 RA Leiden, The Netherlands*

⁴*Centre de Recherches sur la Cognition Animale, CNRS-UMR 5169, Université de Toulouse UPS, 118 route de Narbonne, 31062 Toulouse Cedex 09, France*

Received 7 September 2013; revised 18 October 2013; accepted for publication 18 October 2013

Parasitoidism of ants by flies is known only for Phoridae and Tachinidae. We report the first record of a hoverfly (Syrphidae, Microdontinae, *Hypselosyrphus*) acting as a true primary parasitoid of ants. Previously, larvae of Microdontinae were known only as obligate predators of ant brood. This is also the first host record for any *Hypselosyrphus* species, the first reliable record of an association between a syrphid fly and a ponerine ant, and the first record of a dipteran parasitoid developing upon the immature stages of ants. We reared adults of *Hypselosyrphus trigonus* from cocoons of the arboreal ponerine ant, *Pachycondyla villosa*, nesting in *Aechmea bracteata* bromeliads in southern Quintana Roo, Mexico, and we succeeded in determining various aspects of the parasitoid's life history. The findings obtained in the present study provide novel insights into the evolutionary transformation and diversification of the feeding habits of microdontine syrphid larvae, from an obligatory, specific predatory association with the ant brood, to parasitoidism of ant prepupae. We also highlight the need for more detailed studies of the interactions of arboreal ants and their parasites. We conclude with an overview of the evolutionary transitions and diversification of larval feeding habits that have taken place within the family Syrphidae. © 2013 The Linnean Society of London, *Biological Journal of the Linnean Society*, 2014, **111**, 462–472.

ADDITIONAL KEYWORDS: bromeliad – Diptera – Formicidae – *Hypselosyrphus* – Mexico – *Pachycondyla villosa* – parasitoid – ponerine.

INTRODUCTION

To date, insect parasitoids attacking ant immature stages or adults have been assumed to comprise members of nine families of Hymenoptera (Lachaud & Pérez-Lachaud, 2012; Pérez-Lachaud, Noyes & Lachaud, 2012), and two families of Diptera (Gösswald, 1950; Feener & Brown, 1997; Brown, 2000). Known dipteran/ant associations include scavenging within the ant nest, cleptoparasitism, predation, and

parasitoidism, and involve more than 20 families of flies (Bequaert, 1922; Donisthorpe, 1927; Duffield, 1981; Kistner, 1982; Hölldobler & Wilson, 1990; Rotheray, Barr & Hewitt, 1996; Feener & Brown, 1997; Schmid-Hempel, 1998; Reemer, 2012). Primary parasitoidism of ants by Diptera has so far only been recorded for members of the Phoridae (Feener & Brown, 1997), the so-called ant-decapitating flies, and for one tachinid species, *Strongygaster globula* (Meigen), which is endoparasitic in *Lasius* spp. colony-founding queens (Gösswald, 1950).

The Syrphidae (hoverflies or flower flies) is one of the most diverse dipteran families, comprising almost

*Corresponding author. E-mail: igperez@ecosur.mx

6000 described species and having an almost world-wide distribution (Rotheray & Gilbert, 2011). Its members have been intensively studied as a result of: (1) the important role that the adults of most species play as pollinators (Ssymank *et al.*, 2008); (2) their potential usefulness as biological control agents (the aphidophagous predatory species) (Ankersmit *et al.*, 1986; Laubertie, Wratten & Hemptinne, 2012); and (3) their postulated Batesian mimicry of noxious Hymenoptera (Chittka & Osorio, 2007). The current classification of Syrphidae recognizes three subfamilies: Microdontinae, Eristalinae, and Syrphinae (Thompson & Rotheray, 1998; Ståhls *et al.*, 2003). Each of these includes species with myrmecophilous larvae (Hölldobler, 1929; Pontin, 1959; Dixon, 1960; Thompson, 1972; Speight, 1976, 1988, 2011), although the Microdontinae is certainly the best-known myrmecophile-containing group (Reemer, 2013a; Reemer & Ståhls, 2013a). Microdontine larvae resemble slugs or coccids, and have been described as molluscs in several instances (see Wheeler, 1908; Reemer, 2012). All of the species whose life histories are known have been found within ant nests or in their immediate vicinity (Cheng & Thompson, 2008; Reemer, 2012, 2013a) and some species are highly host-specific (Elmes *et al.*, 1999; Witek *et al.*, 2013). The nature of the feeding habits of the larvae of Microdontinae has long remained unclear. Early studies suggested that immatures lived as scavengers (Wheeler, 1908; Donisthorpe, 1927; Hartley, 1961; Wilson, 1971), although accumulated evidence showed that larvae of at least some species of *Microdon* Meigen and *Omegasyrphus* Giglio-Tos are obligate predators that feed on eggs, larvae, and pupae of particular ant taxa (Van Pelt & Van Pelt, 1972; Duffield, 1981; Garnett, Akre & Sehlke, 1985; Barr, 1995; Schönrogge *et al.*, 2006; Wolton, 2011; Witek *et al.*, 2012). By contrast, larvae of non-microdontine myrmecophilous syrphid genera [namely *Chrysotoxum* Meigen (Syrphinae), *Pipizella* Rondani (Eristalinae), and *Xanthogramma* Schiner (Syrphinae)] are predatory on ant-attended root aphids within the host nest (Rotheray *et al.*, 1996; Speight, 2011). Until now, no reliable case of true primary parasitoidism on ants has ever been reported for syrphid flies.

The Microdontinae is mainly a tropical group. According to Reemer & Ståhls (2013a), 43 genera with 454 valid species are recognized worldwide, with almost half of the species known to occur in the Neotropics. Data on the life histories of microdontines are scarce, especially for the Neotropical fauna. In the present study, we report the Neotropical arboreal ant *Pachycondyla villosa* (Fabricius) as the first known primary host for a species of the microdontine genus *Hypselosyrphus* Hull. This constitutes the first

recorded instance of a syrphid fly developing as a parasitoid upon ants, and the first reliable record of a ponerine ant association with a microdontine. We conclude by reviewing the available information on microdontine-ant interactions and discussing both the evolutionary transitions and diversification of larval feeding habits that have taken place within the Syrphidae.

MATERIAL AND METHODS

STUDY SITE

The present study was conducted in the lowland tropical forest of southern Quintana Roo, Mexico. A plot in a secondary growth, lowland forest enclosing a population of *Aechmea bracteata* (Swartz) Grisebach (Monocotyledones, Bromeliaceae) was situated near an almost perennial small pond ('aguada'), 7 km along the road to Kohunlich archaeological site (18°25'36.23"N, 88°48'16.39"W, 120 m a.s.l.). Within the plot, the tank bromeliads *A. bracteata*, were examined for the presence of ants. *Aechmea bracteata* occurs frequently in the low inundated forest of the Yucatan Peninsula, Mexico (Dejean & Olmsted, 1997). Organic matter decomposes between the basal portions of the leaves that are arranged in compact rosettes, creating soil-like microsites (Picado, 1913). The plants frequently serve as nesting sites for various ant species, such as the ponerine *P. villosa*, the dolichoderine *Dolichoderus bispinosus* (Olivier), and various small myrmicine ants (Dejean, Olmsted & Snelling, 1995; Dejean & Olmsted, 1997; Durou *et al.*, 2002).

ANT HOST SPECIES

Pachycondyla villosa is an opportunistic cavity-breeder that nests in dead wood, in cavities at the junctions of live branches, in bromeliad bases and in abandoned or peripheral hollow stems of various species of myrmecophytic *Cecropia* Loeffling (Lachaud, Fresneau & García-Pérez, 1984; Pérez-Bautista, Lachaud & Fresneau, 1985; Dejean & Olmsted, 1997; MacKay & MacKay, 2010). Colonies are usually founded by several queens (pleometrosis) (Trunzer, Heinze & Hölldobler, 1998). This ant species is found from southern Texas to northern Argentina, and is widespread both in wet and dry forests. Workers are generalized arboreal predators of other arthropods; and collect extrafloral nectar (Lachaud *et al.*, 1984; Dejean & Corbara, 1990). Most of the foraging activity occurs at twilight and during the night (Lachaud *et al.*, 1984).

SAMPLING METHOD

A plot of approximately 8000 m² around the 'aguada' was delimited and all the *A. bracteata* bromeliads

present were examined for the presence of ants. A portion of each epiphyte was cut off from the supporting branch, dismantled leaf-by-leaf in plastic bins coated with Fluon (Whitford GmbH), and all *P. villosa* ants and their brood were collected. Ten colonies (or parts of colonies) of *P. villosa* were obtained between 10 January and 12 January 2012. Ants were counted and their sex, caste, and developmental stage recorded. Larvae were examined for the presence of any external sign of parasitism (i.e. scars, visible external changes or respiratory funnels) under a stereomicroscope. Cocoons from each nest were kept separately in glass vials at room temperature for several days ($28\text{ }^{\circ}\text{C} \pm 2\text{ }^{\circ}\text{C}$ and $75\% \pm 5\%$ relative humidity), and then a sample of the cocoons was dissected under a stereomicroscope. This process yielded several fly puparia together with a complete series of developmental stages of the flies. Puparia were further incubated until adult fly emergence. A sample of ants and fly parasitoids was preserved in 70% ethanol, and voucher specimens were deposited at the Arthropod Collection of El Colegio de la Frontera Sur-Chetumal, Quintana Roo, Mexico (ECO-CH-AR), at the Smithsonian Institution, Washington, DC, USA, and at Naturalis Biodiversity Center (RMNH), Leiden, The Netherlands.

RESULTS

ANT COLONY COMPOSITION

Seven complete *P. villosa* colonies were collected. Three others were incomplete (no or almost no brood, and no queen present) and were not taken into account for further analysis (Table 1). Five colonies contained at least one egg-laying female (1, 2, 2, 5, and 8 dealate females present). Two further colonies contained no queen but workers and brood were abundant. Colonies contained between 62 and 365 workers (mean \pm SEM: 188 ± 45.2 , $N = 7$), two and 111 pupae (54.9 ± 17.4) and seven and 141 larvae (53.9 ± 18.6); two colonies were producing winged females at the moment of collection but males were absent from all colonies.

PARASITISM LEVEL

Five out of the seven complete colonies collected (71.4%) were parasitized by immature stages of a microdontine syrphid fly species (Table 1). Parasitized colonies each contained between two and six parasitized cocoons; the mean parasitism level for all ant cocoons was 4.7%. Syrphid adults (Fig. 1) were reared from puparia found inside ant cocoons. In the taxonomic key of Reemer (2013b), specimens run to *Hypselosyrphus trigonus* Hull, 1937, a Neotropical microdontine species that phenotypically resembles

Table 1. Summary of colony composition of *Pachycondyla villosa* ants nesting in *Aechmea bracteata* bromeliads, percent parasitism rate by *Hypselosyrphus trigonus*, and number and stage of development of retrieved parasitoids in parenthesis

Nest number	Cocoon dissection										Percentage parasitism
	Queen	Alate females	Workers	Cocoons (pupae)	Larvae	Eggs	♀Pupae	♂Pupae	Worker pupae	Prepupae	Parasitized cocoons
1	1	0	65	16	8	No	0	0	6	7	3 (2♂, 1P)
2	0	3	99	99	12	No	4	0	73	17	5 (2♂, 1P, 2L)
3	0	0	34	1	0	No	0	0	1	0	0
4	0	0	2	0	0	No	0	0			
5	0	0	11	0	0	No					
6	5	0	175	51	78	Yes	2	1	11	31	6 (1♀, 1♂, 2L ₁ , 2L ₂)
7	2	0	282	111	60	Yes	0	0	76	33	2 (1♀, 1L ₂)
8	0	0	62	2	7	Yes	0	0	2	0	0
9	8	7	365	93	141	Yes	2	0	36	53	2 (2L ₁ , 1egg)
10	2	0	268	12	71	Yes	0	0	0	12	0

P, puparium; L, larva. Rows in italics correspond to incomplete colonies that were not taken into account for further analysis.




Figure 1. *Hypselosyrphus trigonus*. A, female, lateral habitus. B, male, lateral habitus. C, female head, frontal view. D, male head, frontal view. E, male genitalia.

stingless bees (Apidae: Apinae: Meliponini). Specimens were also compared with pictures of the holotype (Museum of Comparative Zoology, Cambridge, MA, USA), and with an additional specimen from Belize in the RMNH collection (Leiden, The Netherlands), with which the Mexican specimens agree well. Although Hull (1937) stated that the type of *H. trigonus* is a male, examination of the holotype revealed that it is a female. Hull based his description on only one specimen, and the label data of the type specimen agree with those mentioned in the description, which suggests that there cannot have been a confusion of type specimens. Instead, Hull appears to have confused the sexes. Thus, the present record is the first for the male of *H. trigonus*. The male is quite similar to the female, except for usual sexual differences (e.g. compound eyes more closely approaching in the male, which can even be considered holoptic, a highly unusual condition in Microdontinae, Fig. 1D). Its genitalia are shown in Figure 1E.

LIFE-HISTORY VARIABLES OF *H. TRIGONUS*

Through dissections of ant cocoons, the life history of this parasitoid could be partly reconstructed. The egg of *H. trigonus* was found attached to the outer surface of the *P. villosa* prepupa, within the host's protective silk cocoon ($N = 1$) (Fig. 2A, inset). On one occasion, two first-instar larvae were found attached to a single ant prepupa, although their development could not be monitored. Larvae of *H. trigonus* are ectoparasitic on




Figure 2. *Hypselosyrphus trigonus*. A, egg upon a *Pachycondyla villosa* prepupa (arrow), inset: close-up of egg. B, first-instar larva. C, first-second-instar larva. The ant cocoon has been cut open.

the ant prepupa (Figs 2B, C, 3A) and pass through three larval instars as is typical of Aschiza (the Diptera section to which Syrphidae belong) (Rotheray & Gilbert, 1999). The host could still be observed




Figure 3. *Hypselosyrphus trigonus*. A, second–third-instar larva feeding on a *Pachycondyla villosa* prepupa. B, last-instar larva, lateral view.

moving during the initial phase of parasitism (see Supporting information, Video S1), which is unsurprising because only parasitoid wasp females are known to inject venom, although it did not have time to pupate as a result of the rapid growth of the syrphid larva. Both the third larval instar (Fig. 3B) and the puparium (Fig. 4A) were always found in the apical, wider portion of the ant cocoon that corresponds to the ant's head (at the opposite extremity to the meconium). Almost all of the contents of the ant prepupa were consumed by the *H. trigonus* larva that pupated (and eclosed as an adult from the puparium) within the ant cocoon.

As for most aschizans, the puparium of *H. trigonus* has lines of weakness through which the adult fly can easily emerge. However, it is likely that adult flies cannot emerge from the host cocoons without the help of the ants, which usually assist nestmates during emergence (see Discussion). On one occasion, we found a young imago that successfully emerged from its puparium but failed to emerge from the ant cocoon and died (Fig. 4B). Adults (two females and five




Figure 4. *Hypselosyrphus trigonus*. A, puparium inside ant cocoon. B, young imago found dead inside the host cocoon.

males) emerged between 24 January and 5 February, with males emerging earlier than females; two of the males did not expand their wings.

DISCUSSION

HOST LOCATION AND PARASITOID–ANT INTERACTIONS

We determined most of the life-cycle of *H. trigonus*. However, several uncertainties remain concerning the host microhabitat selection, the host location process, and the nature of the behavioural interactions between the adult flies and the ants.

Both hymenopteran and dipteran parasitoids use a diverse array of environmental cues to locate their hosts and/or their local habitat. Phorid parasitoids of ants, for example, have evolved the means to exploit the communication systems of their hosts (Feener & Brown, 1997; Mathis & Philpott, 2012): the females are attracted by the alarm pheromones produced by the mandibular glands of workers, which are released when workers fight, are injured or are otherwise stressed (Feener, Jacobs & Schmidt, 1996). One of the compounds that elicits host location behaviour in phorids, 4-methyl-3-heptanone (Feener *et al.*, 1996), is also present in *P. villosa* (Duffield & Blum, 1973), and workers of this species that were experimentally

injured were attractive to various species of *Apocephalus* Coquillett (Brown & Feener, 1991a, b; Brown, 2000). This or other ant-derived compounds might be used by syrphid fly females to locate ant colonies. For example, in laboratory bioassays, *Microdon mutabilis* (L.) females responded to specific volatile compounds from host ant nests (Schönrogge *et al.*, 2008).

The method of host location by dipteran parasitoids commonly involves two separate phases. First, adult females locate the microhabitat of potential hosts and lay eggs nearby. Thereafter, it is up to the first-instar larva to make contact with the host (Hagen, 1964; Feener & Brown, 1997). This general two-phase strategy is also found in some hymenopterous parasitoids, such as the Eucharitidae, which are specific parasitoids of the ant brood (Heraty, 2002; Lachaud & Pérez-Lachaud, 2012). Most females of non-parasitoid microdontines presumably lay eggs close to the ant nest entrance (Wheeler, 1908; Garnett *et al.*, 1985; Elmes *et al.*, 1999), and do not interact with the ants. Occasionally, oviposition within the ant nest has been recorded (Verhoeff, 1890 in Wheeler, 1910: 385; Wolton, 2011). Although phorid and tachinid ant parasitoids are known to parasitize adult ants during foraging, *H. trigonus* is the first dipteran parasitoid to be found attacking the immature stages of ants. From our observations, it is clear that females of *H. trigonus* lay their eggs directly on the prepupae of ant hosts (Fig. 2A); however, it is unclear at what point in the host's life history they do so, as well as how they cope with ant aggression. Because *H. trigonus* females have no piercing ovipositor, they probably lay an egg upon the ant host larva, within the ant nest, before the larva completes the spinning of its cocoon. Furthermore, we never detected puncture marks on parasitized cocoons. Nest moving by *P. villosa* has never been recorded and, if it does occur, it is certainly infrequent; therefore, ant larvae and cocoons are almost never exposed outside the nest, and the *H. trigonus* female must need to enter it and find the target host stage. Consequently, she will encounter workers in the process of parasitizing the brood, and must circumvent nest defensive behaviours somehow. For example, the female of *Ichneumon eumerus* Wesmäl (Hymenoptera: Ichneumonidae), a parasitoid that attacks larvae of *Maculinea rebeli* Hirschke (Lepidoptera: Lycaenidae) in nests of *Myrmica schencki* Emery, releases semiochemicals that induce in-fighting between worker ants, so enabling her to oviposit unhindered (Thomas *et al.*, 2002).

Other aspects that deserve further attention are the means by which *H. trigonus* adults manage to emerge from the host cocoon and their destination after emer-

gence. Adult microdontines typically mate and reproduce close to their own natal sites (Schönrogge *et al.*, 2006). Therefore, we can assume that adult *H. trigonus* have to escape from host nests to mate. Because adult syrphid mouthparts are of the suctorial type (Speight, 1987; Gilbert & Jervis, 1998), *H. trigonus* adults lack the means to pierce, cut or tear the ant cocoon. The fact that recently emerged adult flies cannot make an exit-hole in the host cocoon suggests that worker ants must help parasitoid flies to emerge, just as they commonly assist with the eclosion of their own callow nestmates (Crosland, Crozier & Jefferson, 1988; Roces & Protomastro, 1988). To our knowledge, there is only one other case of a dipteran larva pupating within the cocoon of an ant: the cleptoparasitic phorid *Cataclinusa pachycondylae* (Brues), which parasitizes a close relative of *P. villosa*, *Pachycondyla harpax* (Fabricius) (Wheeler, 1901). The phorid larva consumes part of the food supplied to the ant larva by the ant workers; in this case, however, the adult fly emerges after the surviving ant, through the latter's exit-hole (Wheeler, 1901).

Once emerged, the *H. trigonus* flies somehow avoid being killed by their host; presumably, they achieve this by either being furtive and quickly escaping from ant nests (e.g. predatory *Microdon*; Forti *et al.*, 2007) or by emitting chemical appeasement substances (e.g. aphidiid parasitoid wasps: Takada & Hashimoto, 1985; slave maker ants: Mori *et al.*, 2000) or repellent allomones (e.g. social parasitic ants: Lloyd, Schmuff & Hefetz, 1986; Ruano *et al.*, 2005). As far as is currently known, newly-eclosed adults of obligate myrmecophiles are not chemically protected from ant attacks (unlike their larvae), and are sometimes killed by their hosts (e.g. Lycaenidae: Fiedler & Maschwitz, 1989). However, protection from ant aggression may be obtained by other means. For example, in some ant-parasitic lycaenid butterflies, the newly-emerged adult butterfly is thickly clothed in loosely attached scales that slough off when grasped by an ant, enabling it escape, relatively unharmed, from ant workers (Pierce *et al.*, 2002).

Other ant guests apparently take advantage of the hygienic behaviour of ants. For example, newly-emerged adult eucharitid parasitoid wasps are not treated aggressively by ants (Wheeler, 1907; Vander Meer, Jouvenaz & Wojcik, 1989) and, under laboratory conditions, the wasps have been observed to be transported to the nest's exterior by the workers, unharmed (Howard, Pérez-Lachaud & Lachaud, 2001). In the field, parasitized ant cocoons are, at least occasionally, discarded by the ants (Pérez-Lachaud *et al.*, 2006), and the adult wasps cut themselves emergence holes in the host cocoons.

A GENERAL PERSPECTIVE ON MICRODONTINE–ANT INTERACTIONS

So far, 108 records of associations of Microdontinae with ants have been documented (summarized in Reemer, 2013a). These concern larvae of 13 microdentine genera (out of 43) but mostly refer to species of *Microdon* s.s. (as presently understood; Reemer & Ståhls, 2013a) and ants from four subfamilies (Dolichoderinae, Pseudomyrmecinae, Formicinae, and Myrmicinae). Weber (1946) reported the presence of ‘*Microdon*’ type larvae in a nest of the ectatommine ant, *Ectatomma ruidum* (Roger), although this record was recently questioned (Reemer, 2013a). Moreover, in all known previous records, microdentine larvae are stated to prey upon the ant brood. Thus, our report of the ponerine ant *P. villosa* as the true primary host of *H. trigonus* is the first reliable record of a poneroid associated with a syrphid fly. This finding has two important implications. First, because parasitism would presumably take longer than predation to evolve given the necessary physiological and behavioural intimacy between host and parasite, our report of a microdentine species with a parasitoid lifestyle suggests the association between microdentine flies and ants evolved earlier than has previously been thought (Reemer, 2013a). Second, our finding of an unusual feeding mode for a microdentine suggests that the parasitoid lifestyle has evolved at least twice independently in the Syrphidae (see below).

Poneroid larvae have large, heavily sclerotized, mandibles and flexible necks enabling them to reach prey items placed near them and to feed by themselves without previous malaxation by the adults (Wheeler & Wheeler, 1976), and they may constitute risky prey for microdentine larvae. Indeed, associations between predatory microdintines and ants are confined to that subset of ant subfamilies in which liquid food exchange by regurgitation (stomodeal trophallaxis) is strongly developed (Hölldobler & Wilson, 1990). The development among certain clades of ants of trophallaxis towards larvae, which are no longer able to feed on solid food, may even have created the opportunity for Microdontinae to prey on these less dangerous stages, and thus may have triggered the evolution of the group. Possibly, parasitoidism by *H. trigonus* of nonfeeding late larval instars is an adaptation to counter the defensive capability of *Pachycondyla* larvae.

Finally, several considerations suggest that our finding might not be unique: (1) Microdontinae are mainly Neotropical (Duffield, 1981; Reemer & Ståhls, 2013a) and, presently, only 18 reported associations with ants (including ours) are from this biogeographical region. These records are for seven microdentine genera (*Ceratophya* Wiedemann, *Hypselosyrphus*,

Masarygus Brèthes, *Microdon*, *Mixogaster* Macquart, *Pseudomicrodon* Hull, and *Stipomorpha* Hull) and ten unidentified microdentine species (reported as *Microdon* sp.) (Reemer, 2013a; present study). Apart from the present results, in only one instance (*Microdon tigrinus* Curran; Forti *et al.*, 2007) are the feeding habits of the larvae known (predation); (2) most of the *P. villosa* colonies collected from our study site were parasitized, although the number of parasitized hosts per colony was relatively low; and (3) parasitoidism of late larval instars of relatively large ants as in *H. trigonus* (only one host is needed to complete development) is a very efficient strategy compared to predatory microdintines. An individual predatory *M. mutabilis* larva needs to consume approximately 1300 ant brood items (eggs, small larvae) to complete its development (Schönrogge *et al.*, 2006).

Ponerinae are extensively distributed within the Neotropics, and further studies focusing on the immature stages of species of the subfamily might increase substantially the list of associations involving syrphid flies, particularly in the case of arboreal species.

DIVERSITY OF LIFESTYLES IN SYRPHIDS

According to Eggleton & Belshaw (1992), it is likely that parasitoidism has evolved over 100 times independently in the Diptera. However, parasitoidism of ants has evolved only in three dipteran families; this includes our own record of Syrphidae. Almost all syrphid adults feed on nectar and pollen from flowers (Rotheray & Gilbert, 2011). By contrast, syrphid larvae are very variable in structure, habits, and feeding modes (Roberts, 1970). All of the microdentine species whose larval stages have previously been described are predators on ant brood (see Introduction). By contrast, the subfamily Eristalinae comprises species with larval life-history traits that include saprophagy, phytophagy, and predation (Rotheray & Gilbert, 1999), and at least one species, *Volucella inanis* (L.), is an ectoparasitoid on larvae of social wasps (Rupp, 1989). Syrphine larvae are predatory mostly on soft-bodied Hemiptera, although some taxa prey on aquatic larvae in the water tanks of bromeliads (Rotheray *et al.*, 2000) and several species are leaf miners (Nishida, Rotheray & Thompson, 2002; Mengual, Ståhls & Rojo, 2008) or feed on pollen (Reemer & Rotheray, 2009). In this context, our finding of an unusual feeding mode for a microdentine is surprising but not unexpected, and raises the question of which evolutionary route was taken: did ectoparasitoidism evolve from larval predators, or did the two life-history strategies evolve independently?

A comprehensive phylogenetic analysis of Syrphidae based on both morphological and molecular characters supported entomophagy (predation) as the plesiomorphic larval feeding mode, and placed Microdontinae as a sister-group of all other Syrphidae (Ståhls *et al.*, 2003). Such placement was later confirmed by Rotheray & Gilbert (2008) using larval morphology. Apart for the single record of parasitoidism in the Eristalinae already noted (Rupp, 1989), there is, to our knowledge, no other record of a syrphid with this lifestyle. Notably, our record is the first evidence of flies succeeding in parasitizing the immature stages of ants, and suggests that a major evolutionary shift occurred in microdontine biology leading to the most advanced evolutionary step in the syrphid/ant association. Such adaptive evolution, by means of natural selection or other process (Kull, 2013), is likely to have been derived from predation, although presently only the feeding modes of species of *Microdon* and *Omegasyrphus* are known. Combining the new evidence with the most recent phylogenetic hypothesis of Microdontinae (Reemer, 2013a; Reemer & Ståhls, 2013b) does not provide insights into the evolutionary pathway taken.

Very similar to what has been hypothesized to have occurred several times among hymenopteran parasitoids associated with ants (Huggert & Masner, 1983; Hanson, West-Eberhard & Gauld, 1995; Hansson, Lachaud & Pérez-Lachaud, 2011; Pérez-Lachaud *et al.*, 2012), a possible evolutionary path to the parasitism of ants by parasitoid microdontine syrphid flies would have been a shift from predation on other arthropods present in ant nests to predation on the brood of the host ant and then, through a gradual process of close association and life-history integration with the ant hosts, to primary parasitoidism on the ant brood. Research that encompasses other associations between ants and microdontine flies might provide more examples of such a shift. The time is ripe for a reassessment of the ecology and phylogeny of the subfamily Microdontinae.

ACKNOWLEDGEMENTS

We are indebted to Alain Dejean, Bruno Corbara, Laurent Pélozuelo, and Yann Hénaut, who collected the bromeliads; to Jacques H. C. Delabie (Laboratório de Mirmecologia CEPEC/CEPLAC, Itabuna-BA, Brazil) who confirmed the identity of the ant; to F. Christian Thompson (Smithsonian) who initially identified the larva as a syrphid; and to three anonymous reviewers for their helpful comments. Holger Weissenberger (ECOSUR) kindly helped with the German literature.

REFERENCES

- Ankersmit GW, Dijkman H, Keuning NJ, Mertens H, Sins A, Tacoma HM. 1986. *Episyrphus balteatus* as a predator of the aphid *Sitobion avenae* on winter wheat. *Entomologia Experimentalis et Applicata* **42**: 271–277.
- Barr B. 1995. Feeding behaviour and mouthpart structure of larvae of *Microdon eggeri* and *Microdon mutabilis* (Diptera, Syrphidae). *Dipterist Digest* **2**: 31–36.
- Bequaert JC. 1922. The predaceous enemies of ants. *Bulletin of the American Museum of Natural History* **45**: 271–331.
- Brown BV. 2000. Revision of the 'Apocephalus miricauda-group' of ant-parasitizing flies (Diptera: Phoridae). *Contributions in Science* **482**: 1–62.
- Brown BV, Feener DH Jr. 1991a. Behavior and host location cues of *Apocephalus paraponerae* (Diptera: Phoridae), a parasitoid of the giant tropical ant, *Paraponera clavata* (Hymenoptera: Formicidae). *Biotropica* **23**: 182–187.
- Brown BV, Feener DH Jr. 1991b. Life history parameters and description of the larva of *Apocephalus paraponerae* (Diptera: Phoridae), a parasitoid of the giant tropical ant *Paraponera clavata* (Hymenoptera: Formicidae). *Journal of Natural History* **25**: 221–231.
- Cheng X-Y, Thompson FC. 2008. A generic conspectus of the Microdontinae (Diptera: Syrphidae) with the description of two new genera from Africa and China. *Zootaxa* **1879**: 21–48.
- Chittka L, Osorio D. 2007. Cognitive dimensions of predator responses to imperfect mimicry? *PLoS Biology* **5**: e339.
- Crosland MWJ, Crozier RH, Jefferson E. 1988. Aspects of the biology of the primitive ant genus *Myrmecia* F. (Hymenoptera: Formicidae). *Journal of the Australian Entomological Society* **27**: 305–309.
- Dejean A, Corbara B. 1990. Predatory behavior of a Neotropical arboricolous ant: *Pachycondyla villosa* (Formicidae: Ponerinae). *Sociobiology* **17**: 271–286.
- Dejean A, Olmsted I. 1997. Ecological studies on *Aechmea bracteata* (Swartz) (Bromeliaceae). *Journal of Natural History* **31**: 1313–1334.
- Dejean A, Olmsted I, Snelling RR. 1995. Tree-epiphyte-ant relationships in the low inundated forest of Sian Ka'an Biosphere Reserve, Quintana Roo, Mexico. *Biotropica* **27**: 57–70.
- Dixon TJ. 1960. Key to and descriptions of the third instar larvae of some species of Syrphidae (Diptera) occurring in Britain. *Transactions of the Royal Entomological Society of London* **112**: 345–379.
- Donisthorpe HStJK. 1927. *The guests of British ants, their habits and life histories*. London: Routledge & Sons Ltd.
- Duffield RM. 1981. Biology of *Microdon fuscipennis* (Diptera: Syrphidae) with interpretations of the reproductive strategies of *Microdon* species found North of Mexico. *Proceedings of the Entomological Society of Washington* **83**: 716–724.
- Duffield RM, Blum MS. 1973. 4-Methyl-3-heptanone: identification and function in *Neoponera villosa* (Hymenoptera: Formicidae). *Annals of the Entomological Society of America* **66**: 1357.

- Durou S, Dejean A, Olmsted I, Snelling RR. 2002. Ant diversity in coastal zones of Quintana Roo, Mexico, with special reference to army ants. *Sociobiology* **40**: 385–402.
- Eggleton P, Belshaw R. 1992. Insect parasitoids: an evolutionary overview. *Proceedings of the Royal Society of London Series B, Biological Sciences* **337**: 1–20.
- Elmes GW, Barr B, Thomas JA, Clarke RT. 1999. Extreme host specificity by *Microdon mutabilis* (Diptera: Syrphidae), a social parasite of ants. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences* **266**: 447–453.
- Feener DH Jr, Brown BV. 1997. Diptera as parasitoids. *Annual Review of Entomology* **42**: 73–97.
- Feener DH Jr, Jacobs LF, Schmidt JO. 1996. Specialized parasitoid attracted to a pheromone of ants. *Animal Behaviour* **51**: 61–66.
- Fiedler K, Maschwitz U. 1989. The symbiosis between the weaver ant, *Oecophylla smaragdina*, and *Anthene emolus*, an obligate myrmecophilous lycaenid butterfly. *Journal of Natural History* **23**: 833–846.
- Forti LC, Camargo RS, Verza SS, Andrade APP, Fujihara RT, Lopes JFS. 2007. *Microdon tigrinus* Curran, 1940 (Diptera, Syrphidae): populational fluctuation and specificity to the nest of *Acromyrmex coronatus* (Hymenoptera: Formicidae). *Sociobiology* **50**: 909–919.
- Garnett WB, Akre RD, Sehlike G. 1985. Cocoon mimicry and predation by myrmecophilous diptera (Diptera: Syrphidae). *The Florida Entomologist* **68**: 615–621.
- Gilbert F, Jervis M. 1998. Functional, evolutionary and ecological aspects of feeding-related mouthpart specializations in parasitoid flies. *Biological Journal of the Linnean Society* **63**: 495–535.
- Gösswald K. 1950. Pflege des Ameisenparasiten *Tamiclea globula* Meig. (Diptera) durch den Wirt mit Bemerkungen über den Stoffwechsel in der parasitierten Ameise. *Verhandlungen der Deutschen Zoologischen* **1949**: 256–264.
- Hagen KS. 1964. Developmental stages of parasites. In: DeBach P, ed. *Biological control of insect pests and weeds*. New York, NY: Reinhold, 168–246.
- Hanson PE, West-Eberhard MJ, Gauld ID. 1995. Interspecific interactions of nesting Hymenoptera. In: Hanson PE, Gauld ID, eds. *The hymenoptera of Costa Rica*. Oxford: Oxford University Press, 76–88.
- Hansson C, Lachaud J-P, Pérez-Lachaud G. 2011. Entedoninae wasps (Hymenoptera, Chalcidoidea, Eulophidae) associated with ants (Hymenoptera, Formicidae) in tropical America, with new species and notes on their biology. *ZooKeys* **134**: 65–82.
- Hartley JC. 1961. A taxonomic account of the larvae of some British Syrphidae. *Proceedings of the Zoological Society of London* **136**: 505–573.
- Heraty JM. 2002. A revision of the genera of Eucharitidae (Hymenoptera: Chalcidoidea) of the world. *Memoirs of the American Entomological Institute* **68**: 1–367.
- Hölldobler B, Wilson EO. 1990. *The ants*. Berlin: Springer-Verlag.
- Hölldobler K. 1929. Über die Entwicklung der Schwirrflye *Xanthogramma citrofasciatum* im Neste von *Lasius alienus* und *niger*. *Zoologischer Anzeiger* **82**: 171–176.
- Howard RW, Pérez-Lachaud G, Lachaud J-P. 2001. Cuticular hydrocarbons of *Kapala sulcifacies* (Hymenoptera: Eucharitidae) and its host, the ponerine ant *Ectatomma ruidum* (Hymenoptera: Formicidae). *Annals of the Entomological Society of America* **94**: 707–716.
- Huggert L, Masner L. 1983. A review of myrmecophilic-symphilic diapiiid wasps in the Holarctic realm, with descriptions of new taxa and a key to genera (Hymenoptera: Proctotrupoidea: Diapriidae). *Contributions of the American Entomological Institute* **20**: 63–89.
- Hull FM. 1937. New species of exotic syrphid flies. *Psyche* **44**: 12–32.
- Kistner DH. 1982. The social insects' bestiary. In: Hermann HR, ed. *Social insects*, Vol. 3. New York, NY: Academic Press, 1–244.
- Kull K. 2013. Adaptive evolution without natural selection. *Biological Journal of the Linnean Society* doi: 10.1111/bij.12124.
- Lachaud J-P, Fresneau D, García-Pérez J. 1984. Étude des stratégies d'approvisionnement chez 3 espèces de fourmis ponérines. *Folia Entomologica Mexicana* **61**: 159–177.
- Lachaud J-P, Pérez-Lachaud G. 2012. Diversity of species and behavior of hymenopteran parasitoids of ants: a review. *Psyche* **2012**: Article ID 134746.
- Laubertie EA, Wratten SD, Hemptinne J-L. 2012. The contribution of potential beneficial insectary plant species to adult hoverfly (Diptera: Syrphidae) fitness. *Biological Control* **61**: 1–6.
- Lloyd HA, Schmuff NR, Hefetz A. 1986. Chemistry of the anal glands of *Bothriomyrmex syrius* Forel. Olfactory mimetism and temporary social parasitism. *Comparative Biochemistry and Physiology* **83**: 71–73.
- Mackay WP, Mackay EE. 2010. *The systematics and biology of the New World ants of the genus Pachycondyla* (Hymenoptera: Formicidae). New York, NY: The Edwin Mellen Press.
- Mathis KA, Philpott SM. 2012. Current understanding and future prospects of host selection, acceptance, discrimination, and regulation of phorid fly parasitoids that attack ants. *Psyche* **2012**: Article ID 895424.
- Mengual X, Ståhls G, Rojo S. 2008. Molecular phylogeny of *Allograpta* (Diptera, Syrphidae) reveals diversity of lineages and non-monophyly of phytophagous taxa. *Molecular Phylogenetics and Evolution* **49**: 715–727.
- Mori A, Visicchio R, Sledge MF, Grasso DA, Le Moli F, Turillazzi S, Spencer S, Jones GR. 2000. Behavioral assays testing the appeasement allomone of *Polyergus rufescens* queens during host-colony usurpation. *Ethology Ecology and Evolution* **12**: 315–322.
- Nishida K, Rotheray G, Thompson FC. 2002. First non-predaceous syrphine flower fly (Diptera: Syrphidae): a new leaf-mining *Allograpta* from Costa Rica. *Studia Dipterologica* **9**: 421–436.
- Pérez-Bautista M, Lachaud J-P, Fresneau D. 1985. La división del trabajo en la hormiga primitiva *Neoponera*

- villosa* (Hymenoptera: Formicidae). *Folia Entomologica Mexicana* **65**: 119–130.
- Pérez-Lachaud G, Heraty JM, Carmichael A, Lachaud J-P. 2006.** Biology and behavior of *Kapala* (Hymenoptera: Eucharitidae) attacking *Ectatomma*, *Gnamptogenys*, and *Pachycondyla* (Formicidae: Ectatomminae and Ponerinae) in Chiapas, Mexico. *Annals of the Entomological Society of America* **99**: 567–576.
- Pérez-Lachaud G, Noyes J, Lachaud J-P. 2012.** First record of an encyrtid wasp (Hymenoptera: Chalcidoidea) as a true primary parasitoid of ants (Hymenoptera: Formicidae). *The Florida Entomologist* **95**: 1066–1076.
- Picado C. 1913.** Les broméliacées épiphytes considérées comme milieu biologique. *Bulletin Scientifique de la France et de la Belgique* **47**: 215–360.
- Pierce NE, Braby MF, Heath A, Lohman DJ, Mathew J, Rand DB, Travassos MA. 2002.** The ecology and evolution of ant association in the Lycaenidae (Lepidoptera). *Annual Review of Entomology* **47**: 733–771.
- Pontin AJ. 1959.** Some records of predators and parasites adapted to attack aphids attended by ants. *The Entomologist's monthly magazine* **95**: 154–155.
- Reemer M. 2012.** Unravelling a hotchpotch: phylogeny and classification of the Microdontinae (Diptera: Syrphidae). DPhil Thesis, Leiden University.
- Reemer M. 2013a.** Review and phylogenetic evaluation of associations between Microdontinae (Diptera: Syrphidae) and ants (Hymenoptera: Formicidae). *Psyche* **2013**: Article ID 538316.
- Reemer M. 2013b.** Taxonomic exploration of Neotropical Microdontinae (Diptera: Syrphidae) mimicking stingless bees. *Zootaxa* **3697**: 1–88.
- Reemer M, Rotheray GE. 2009.** Pollen feeding larvae in the presumed predatory syrphine genus *Toxomerus* Macquart (Diptera, Syrphidae). *Journal of Natural History* **43**: 939–949.
- Reemer M, Ståhls G. 2013a.** Generic revision and species classification of the Microdontinae (Diptera, Syrphidae). *ZooKeys* **288**: 1–213.
- Reemer M, Ståhls G. 2013b.** Phylogenetic relationships of Microdontinae (Diptera: Syrphidae) based on molecular and morphological characters. *Systematic Entomology* **38**: 661–688.
- Roberts MJ. 1970.** The structure of the mouthparts of syrphid larvae (Diptera) in relation to feeding habits. *Acta Zoologica* **51**: 43–65.
- Roces F, Protomastro JJ. 1988.** Prey availability and eclosion-help of callow workers in the formicine ant *Camponotus mus*. *Oecologia* **77**: 387–389.
- Rotheray G, Gilbert F. 1999.** Phylogeny of Palaearctic Syrphidae (Diptera): evidence from larval stages. *Zoological Journal of the Linnean Society* **127**: 1–112.
- Rotheray GE, Barr B, Hewitt SM. 1996.** The myrmecophilous larvae of *Chrysotoxum arcuatum*, *Pipizella varipes* and *Xanthogramma pedissequum* from Europe and *Platycheirus milleri* from New Zealand (Dip.: Syrphidae). *The Entomologist's Record and Journal of Variation* **108**: 257–265.
- Rotheray GE, Gilbert F. 2008.** Phylogenetic relationships and the larval head of the lower Cyclorrhapha (Diptera). *Zoological Journal of the Linnean Society* **153**: 287–323.
- Rotheray GE, Gilbert F. 2011.** *The natural history of hoverflies*. Ceredigion: Forrest Text.
- Rotheray GE, Zumbado M, Hancock EG, Thompson FC. 2000.** Remarkable aquatic predators in the genus *Ocyptamus* (Diptera, Syrphidae). *Studia Dipterologica* **7**: 385–398.
- Ruano F, Hefetz A, Lenoir A, Francke W, Tinaut A. 2005.** Dufour's gland secretion as a repellent used during usurpation by the slave-maker ant *Rossomyrmex minuchae*. *Journal of Insect Physiology* **51**: 1158–1164.
- Rupp L. 1989.** Die mitteleuropäischen Arten der Gattung *Volucella* (Diptera, Syrphidae) als Kommensalen und parasitoide in den Nestern von Hummeln und sozialen wespen. Unpublished dissertation, Albert-Ludwigs-University.
- Schmid-Hempel P. 1998.** *Parasites in social insects*. Princeton, NJ: Princeton University Press.
- Schönrogge K, Gardner MG, Elmes GW, Napper EKV, Simcox DJ, Wardlaw JC, Breen J, Barr B, Knapp JJ, Pickett JA, Thomas JA. 2006.** Host propagation permits extreme local adaptation in a social parasite of ants. *Ecology Letters* **9**: 1032–1040.
- Schönrogge K, Napper EKV, Birkett MA, Woodcock CM, Pickett JA, Wadhams LJ, Thomas JA. 2008.** Host recognition by the specialist hoverfly *Microdon mutabilis*, a social parasite of the ant *Formica lemani*. *Journal of Chemical Ecology* **34**: 168–178.
- Speight MCD. 1976.** The puparium of *Chrysotoxum festivum* (L.) (Diptera, Syrphidae). *The Entomologist's Record and Journal of Variation* **88**: 51–52.
- Speight MCD. 1987.** External morphology of adult Syrphidae (Diptera). *Tijdschrift voor Entomologie* **130**: 141–175.
- Speight MCD. 1988.** *Doros destillatorius*, espèce nouvelle pour la France, avec désignation des types des deux espèces européennes du genre *Doros*, description de leurs pupes et clés de détermination des adultes et des pupes (Dipt. Syrphidae). *Bulletin de la Société Entomologique de France*. **92**: 193–200.
- Speight MCD. 2011.** *Species accounts of European Syrphidae (Diptera), Glasgow 2011. Syrph the Net, the database of European Syrphidae*, Vol. 65. Dublin: Syrph the Net publications.
- Ssymanck A, Kearns CA, Pape T, Thompson FC. 2008.** Pollinating flies (Diptera): a major contribution to plant diversity and agricultural production. *Biodiversity* **9**: 86–89.
- Ståhls G, Hippa H, Rotheray G, Muona J, Gilbert F. 2003.** Phylogeny of Syrphidae (Diptera) inferred from combined analysis of molecular and morphological characters. *Systematic Entomology* **28**: 433–450.
- Takada H, Hashimoto Y. 1985.** Association of the root aphid parasitoids *Aclitus sappaphis* and *Paralipsis eikoeae* (Hymenoptera, Aphidiidae) with the aphid-attending ants *Pheidole fervida* and *Lasius niger* (Hymenoptera, Formicidae). *Kontyû* **53**: 150–160.

- Thomas JA, Knapp JJ, Akino T, Gerty S, Wakamura S, Simcox DJ, Wardlaw JC, Elmes GW. 2002.** Parasitoid secretions provoke ant warfare. *Nature* **417**: 505–506.
- Thompson FC. 1972.** A new *Platycheirus* from New Zealand: first record of a melanostomine syrphid fly associated with ants. *New Zealand Journal of Science* **15**: 77–84.
- Thompson FC, Rotheray GE. 1998.** Family Syrphidae. In: Papp L, Darvas B, eds. *Contributions to a manual of Palaearctic Diptera (with special reference to flies of economic importance)*, vol. 3, higher Brachycera. Budapest: Science Herald, 81–139.
- Trunzer B, Heinze J, Hölldobler B. 1998.** Cooperative colony founding and experimental primary polygyny in the ponerine ant *Pachycondyla villosa*. *Insectes Sociaux* **45**: 267–276.
- Van Pelt AF, Van Pelt SA. 1972.** *Microdon* (Diptera: Syrphidae) in nests of *Monomorium* (Hymenoptera: Formicidae) in Texas. *Annals of the Entomological Society of America* **65**: 977–978.
- Vander Meer RK, Jouvenaz DP, Wojcik DP. 1989.** Chemical mimicry in a parasitoid (Hymenoptera: Eucharitidae) of fire ants (Hymenoptera: Formicidae). *Journal of Chemical Ecology* **15**: 2247–2261.
- Weber NA. 1946.** Two common ponerine ants of possible economic significance, *Ectatomma tuberculatum* (Olivier) and *E. ruidum* Roger. *Proceedings of the Entomological Society of Washington* **48**: 1–16.
- Wheeler GC, Wheeler J. 1976.** Ant larvae: review and synthesis. *Memoirs of the Entomological Society of Washington* **7**: 1–108.
- Wheeler WM. 1901.** An extraordinary ant-guest. *American Naturalist* **35**: 1007–1016.
- Wheeler WM. 1907.** The polymorphism of ants, with an account of some singular abnormalities due to parasitism. *Bulletin of the American Museum of Natural History* **23**: 1–93.
- Wheeler WM. 1908.** Studies of myrmecophiles. III. *Microdon*. *Journal of the New York Entomological Society* **16**: 202–213.
- Wheeler WM. 1910.** *Ants; their structure, development and behavior*. New York, NY: Columbia University Press.
- Wilson EO. 1971.** *The insect societies*. Cambridge, MA: Harvard University Press.
- Witek M, Canterino S, Balletto E, Bonelli S. 2012.** Life cycle and growth pattern of the endangered myrmecophilous *Microdon myrmicae* (Diptera: Syrphidae). *European Journal of Entomology* **109**: 457–461.
- Witek M, Casacci LP, Barbero F, Patricelli D, Sala M, Bossi S, Maffe M, Woyciechowski M, Balletto E, Bonelli S. 2013.** Interspecific relationships in co-occurring populations of social parasites and their host ants. *Biological Journal of the Linnean Society* **109**: 699–709.
- Wolton R. 2011.** Observations on the ecology and behaviour of *Microdon myrmicae* Schönrogge *et al.* (Diptera, Syrphidae), with a description of egg and early instar morphology. *Dipterists Digest* **18**: 55–67.

SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Video S1. First–second-instar *Hypselosyrphus trigonus* larva feeding upon an ant prepupa. Note the movements of the prepupa that is still alive.