

The Caudal Ventrolateral Medulla as an Important Inhibitory Modulator of Pain Transmission in the Spinal Cord

Isaura Tavares and Deolinda Lima

Abstract: The caudal ventrolateral medulla (VLM) has emerged during the last decade as one of the main components of the endogenous pain control system. Profound and long-lasting analgesia is produced by mild stimulation of the VLM. The VLMLat, the reticular formation located between the spinal trigeminal nucleus and the lateral reticular nucleus (LRt), appears to play a major role in that antinociceptive action. The projections to spinal cord laminae involved in nociceptive transmission originate exclusively in the VLMLat. The VLMLat participates in a disynaptic pathway involving spinally projecting pontine A5 noradrenergic neurons, which appears to convey α_2 -adrenoreceptor-mediated analgesia produced from the VLM. Neurons in the VLMLat and in lamina I are reciprocally connected by a closed loop that is likely to mediate feedback control of supraspinal nociceptive transmission. On the other hand, the LRt, which is targeted by ventral (lamina VII) and deep dorsal (laminae IV to V) horn inputs, projects to the premotor lamina VII. Nociceptive input ascending from the cord and increases in blood pressure are discussed as possible physiologic triggers of the analgesia produced by the VLM. The overall role of the VLM as a center for integration of nociceptive, cardiovascular, and motor functions is discussed. The putative therapeutic benefits of manipulating the VLM for the control of chronic pain are envisaged.

© 2002 by the American Pain Society

Key words: Pain control, supraspinal centers, stimulation-produced analgesia, nociceptive and cardiovascular integration.

Nociceptive transmission from the spinal cord is subject to modulatory influences originating from widespread brain areas, which we will collectively refer to as the endogenous pain control system.^{11,20,34,43,61} The existence of a system that selectively modulates pain was predicted by the Gate Control Theory of Melzack and Wall⁸¹ and received support from studies showing that electrical or chemical stimulation of the periaqueductal gray (PAG) or nucleus raphe magnus and adjacent reticular formation (collectively known as the rostral ventromedial medulla, RVM) inhibited behavioral responses to noxious stimulation without affecting reactions to other environmental stimuli.^{11,34,57,64,101} This so-called stimulation-produced analgesia (SPA) was later found to be also elicited from several other brain

regions, providing further evidence for the wide extent and heterogeneity of the endogenous pain control system.^{20,34,43,49,61,83,94,97,99}

Several studies have shown that the SPA produced from most components of the endogenous pain control system results from inhibition of nociceptive activity at the level of the spinal cord. Tract tracing studies showed direct or indirect connections with the spinal dorsal horn, whereas electrophysiologic recordings showed that local electrical or chemical stimulation in various brain areas selectively inhibited nociceptive responses of dorsal horn neurons.^{61,105} Moreover, the magnitude of the analgesia obtained by manipulating the system correlated with the decrease of noxious-evoked spinal induction of the *c-fos* proto-oncogene, a reliable marker of nociceptive neuronal activation in the spinal cord.^{44,51,53,98,99,128} As to the spinal dorsal horn circuitry targeted by the descending modulatory system, direct inhibition of neurons projecting in the spinothalamic and spinomesencephalic tracts has been clearly established,⁶¹ and actions on spinal interneurons, namely, inhibition of excitatory interneurons⁶⁵ and activation of inhibitory interneurons,⁸² have also been inferred.¹²³ With respect to their neurochemical nature, 3 main types of descending mod-

From the Institute of Histology and Embryology, Faculty of Medicine and IBMC, University of Oporto, Porto, Portugal.

Supported by BIOTEC project no. BIO4-CT98-0076 (EU) and Gulbenkian Pain Program.

Address reprint requests to Isaura Tavares, PhD, Institute of Histology and Embryology, Faculty of Medicine of Oporto, Alameda Prof Hernâni Monteiro, 4200-319 Porto, Portugal. E-mail: isatav@med.up.pt

© 2002 by the American Pain Society

1526-5900/2002 \$35.00/0

doi:10.1054/jpai.2002.127775




Figure 1. (A) Drawing depicting the components (dashed regions) of the endogenous pain modulatory system at a coronal section of the medulla oblongata obtained 5.3 mm caudal to the interaural line. Adapted from references 61 and 91. (B) Photomicrograph of the VLM. 10, Dorsal motor nucleus of the vagus; 12, hypoglossal nucleus; cc, central canal; Cu, nucleus cuneatus; Gr, nucleus gracilis; IO, inferior olive; LRtm, magnocellular part of the LRt; LRtp, parvocellular part of the LRt; pyx, pyramidal decussation; nA, nucleus ambiguus. Scale bar in (B) represents 0.20 mm.

ulatory pathways have been demonstrated by anatomic and pharmacologic studies: opioidergic, serotonergic, and noradrenergic.^{34,94} The intrathecal administration of suitable agonists and antagonists has shown that the SPA frequently involves the participation of the 3 neurotransmitters, which interact in a complex manner, specific for each component of the pain control system.⁶¹

The existence of a similar pain modulating system in humans has been postulated on the basis of the observation that patients with chronic pain experienced analgesia after electrical stimulation of some of the pain control sites identified in animals, namely, the thalamus, the PAG, and periventricular gray areas.^{12,14,104} The invasive nature of deep brain stimulation in humans has, however, precluded a complete demonstration of the phylogenetic homology of the endogenous pain modulatory system between rodents and humans. However, the distribution of the neurotransmitters involved in pain modulation is highly conserved across mammalian species, including humans.^{30,92} That the opioidergic component of the system is engaged in pain control by morphine and its derivatives is evidenced by the profound hyperalgesia produced in humans on administration of the opioid an-

tagonist naloxone.^{13,46,92} Furthermore, the interactions between the opioidergic and noradrenergic components of the pain control system, demonstrated at the experimental level, are clinically relevant. The coadministration of opioid derivatives and α_2 -adrenoreceptor agonists results in improved analgesia without some of the adverse effects of opioid treatment alone.^{24,26,47,90}

In recent years a multitude of data has demonstrated the complexity of the endogenous pain control system. The role played by the medulla oblongata in the endogenous pain modulatory system deserves special attention. The medulla oblongata is the brain region with the higher density of areas involved in pain modulation, and it presents some peculiarities in the way it controls nociception. Besides the RVM, the medullary regions belonging to the endogenous pain modulatory system include the nucleus tractus solitarii (NTS), dorsal reticular nucleus (DRt), ventral reticular nucleus (VRt), and caudal ventrolateral medulla (VLM) (Fig 1A). Electrical or chemical stimulation of the RVM, NTS, VRt, or VLM inhibits the behavioral responses to pain and the nociceptive activity of dorsal horn neurons.⁶¹ Contrary to the SPA produced from most areas of the endogenous pain control system, opioids do not appear to be significantly involved in descending modulation from the medulla oblongata with the exception of the RVM. Another characteristic of pain modulation from the medulla oblongata is that it is the only part of the system from which facilitation of pain transmission, ie, pronociception, can be produced along with antinociception. Neurons in the RVM facilitate nociceptive transmission in the spinal cord,^{32-34,36,37} and enhanced nociceptive behavioral responses are observed after stimulation of the DRt.^{5,7} Finally, it has been claimed that pain-modulating areas also control other functions that affect nociceptive transmission and are influenced by noxious events.^{72,73} The medullary regions controlling autonomic functions, which are key components of the pain control system, are also likely to be responsible for the interaction between cardiovascular changes and pain.^{95,127}

Located at the ventrolateral quadrant of the medulla oblongata, the VLM emerged recently as an important component of the endogenous pain modulatory system. The efficacy of analgesia originated from the VLM, as assessed by the electrical thresholds for inhibition of nociceptive dorsal horn neurons and by the magnitude of duration of suppression of nociceptive reflexes,^{42,54,55} is greater than that of most other areas including PAG,^{19,57} the RVM,^{87,102} and the locus coeruleus.^{59,60} Moreover, the VLM appears to be responsible for tonic descending inhibition of spinal nociceptive neurons.^{40,48} During the last decade, a large amount of anatomic and functional data on the mechanisms underlying pain modulation from the VLM has accrued, prompting the present review of the specific role of the VLM in the endogenous pain control system in relation to previously studied areas such as the PAG and RVM.^{11,38,48} The anatomic data recently compiled point to the existence of subregions within the VLM in what concerns the participation in pain modulatory circuits, which could not be predicted

by physiologic and pharmacologic studies because of the characteristics of such functional studies.

Direct VLM-Spinal Pathways

Tract tracing studies have shown that the VLM is a heterogeneous region in terms of its efferent projections to the spinal cord. The projections to spinal areas containing nociceptive neurons, namely, laminae I, IV to V, and X,³⁵ originate in the region of the reticular formation designated as VLMLat^{109,111,112} (Fig 1B) located between the spinal trigeminal nucleus, pars caudalis (Sp₅C), and the lateral reticular nucleus (LRt). On the other hand, lamina VII, which contains mainly premotor interneurons, is targeted exclusively by the LRt, a region involved in motor control.⁵² That the VLMLat is the region within the VLM responsible for pain modulation is supported by the observation that the dorsolateral funiculus, which is the spinal pathway mediating the descending antinociception from the VLM,^{55,56} contains fibers originating in the VLMLat, whereas the LRt projects through the ventrolateral funiculus.¹¹⁴ Furthermore, although inhibition of nociceptive spinal neurons can be produced by electrical⁸⁴ or glutamate⁵⁵ stimulation of the LRt, the hypoalgesia induced from the VLMLat is more profound and long-lasting,⁴² suggesting that the LRt antinociception may be due to coactivation of the VLMLat.

That the spinal laminae targeted by the VLMLat contain nociceptive neurons that project in the main ascending pathways⁷³ suggests that the VLMLat-spinal pathway is specifically involved in the modulation of supraspinal transmission of nociceptive information. Supporting this hypothesis, potent inhibition of nociceptive spinothalamic neurons was produced by electrical stimulation of sites encompassing the VLMLat.⁵⁵ In this regard, it is noteworthy that neurons in lamina I project back to the VLMLat,⁷⁰ forming a closed reciprocal loop that is likely to be specifically involved in pain modulation¹¹⁴ (Fig 2). Terminal boutons labeled from the superficial dorsal horn (laminae I to II) contain round vesicles and establish asymmetrical synapses with VLMLat neurons projecting to those spinal layers. In lamina I, boutons labeled from the VLMLat contain either round or flattened vesicles and establish, respectively, asymmetrical or symmetrical synapses with VLMLat-projecting neurons. Considering the putative excitatory or inhibitory nature of, respectively, asymmetrical and symmetrical synapses,^{23,45,103,118,120} the arrival of nociceptive input from lamina I to the VLMLat appears to trigger descending modulation of pain transmission through both excitatory and inhibitory mechanisms. Interestingly, this reciprocal loop appears to differentially involve distinct morphologic classes of VLMLat-projecting lamina I cells. Although fusiform neurons are the main source of lamina I projections to the VLMLat,⁷⁰ pyramidal cells in lamina I are contacted in much higher numbers by fibers descending from the VLMLat.¹¹⁴ Incidentally, it must be noted that pyramidal neurons have a more widespread distribution of projections to supraspinal pain control centers than the other lamina I cell types and that spinofugal lamina I neurons




Figure 2. Diagram showing the reciprocal connections between the VLM and the spinal cord. The VLMLat–lamina I loop is closed at both the medullary and spinal levels. The LRt is reciprocally connected with lamina VII without the formation of any closed loop. The VLM projects to laminae IV–V, and this region projects to the LRt. Abbreviations as in Figure 1.

frequently have collaterals in the brain.^{6,31,66–68,70,73} It is therefore possible that the pyramidal neurons may be involved in adjusting and integrating the nociceptive responses of lamina I neurons to the overall functional status of the endogenous pain control system.

Despite being targeted by the VLMLat, deep dorsal horn neurons send ascending projections only to the LRt.⁷⁰ The participation of the LRt in motor functions is well-established on the basis of its connections with the cerebellum and inferior olive.^{9,22,29} Rather than being involved in feedback modulation at the spinal cord level, noxious input conveyed by deep dorsal horn neurons to the LRt⁸⁰ is more likely to trigger integrated nociceptive motor responses through connections with both motor control brain areas and spinal lamina VII.¹¹⁴

Indirect VLM-Spinal Pathways

Spinal α_2 -adrenoreceptors and serotonin (5-HT) receptors mediate the antinociception produced from the VLM.^{42–44,75} However, neurons of the A1 noradrenergic cell group located in VLM do not project to the cord,^{111,112,121} and serotonergic cells are absent from the VLM. The widespread connections of the VLM with other components of the endogenous pain control system^{61,79,124} point to the involvement of polysynaptic pathways in both noradrenergic and serotonergic antinociceptive effects from VLM. In the brainstem, VLM fibers are directly apposed to spinally projecting neurons only in the pontine A5 noradrenergic cell group and the




Figure 3. Effect of the administration of SAP-anti-DBH in the VLMIlat on the time course of pain-related activity (A) and spinal *c-fos* expression in the dorsal horn (B) after formalin injection in the ipsilateral hindpaw. In (A), focused pain-related activity is the time (in seconds) spent in motor activity directed toward the injected paw, including licking, biting and shaking of the paw. Filled squares, control group; open circles: group with lesions. In (B), numbers of nuclei immunoreactive for the Fos protein were counted in the dorsal horn (laminae I–VI) ipsilateral to the injected paw. Filled bar, control group; dashed bar, group with lesions. In (A) and (B), numbers represent means \pm standard error of mean. Asterisk represents statistical differences ($P < .05$) determined by 2-way analysis of variance with repeated measures (A) or by the paired Student *t* test (B). Photomicrographs depict DBH-immunoreactive neurons at the A5 cell group in a control animal (C) and in 1 animal with SAP-anti-DBH lesion (D). 7, Facial nucleus; 7n, facial nerve. Magnification bars, 42 μ m.

RVM.¹¹¹ The VLM-A5 pathway originates exclusively from the VLMIlat and establishes asymmetrical, putative excitatory, synaptic contacts with spinally projecting A5 neurons.^{111,112} According to these data, the α_2 -adrenoceptor antinociception produced from the VLM is triggered in the VLMIlat and mediated by the A5 noradrenergic cell group.^{17,94,124} Notably, the large majority of A5 noradrenergic neurons that participate in the disynaptic pathway to the cord also have collateral projections back to the VLMIlat.¹¹² Because noradrenaline and the α_2 -adrenoceptor agonist clonidine inhibit VLM neurons¹⁸ and produce hyperalgesia,⁸⁹ it is possible that the A5-VLMIlat projection represents the anatomic substrate for a negative feedback circuit mediating autoinhibition of the VLMIlat.

Despite the lack of ultrastructural studies, it is most likely that the direct appositions between VLM fibers and spinally projecting serotonergic neurons of the RVM¹¹¹ represent synaptic contacts in a manner similar to those observed in the VLM-A5 connection.¹¹¹ The well-established involvement of 5-HT in descending antinociception from the RVM^{3,15,50} suggests that the di-

synaptic VLM-RVM-spinal pathway conveys the serotonergic antinociceptive effects elicited from the VLM.^{42,54}

Similar anatomic approaches have been used to demonstrate the existence of parallel disynaptic pathways for other components of the endogenous analgesic system. For example, terminals arising from the PAG synapse on RVM neurons that project to the cord,⁶² and, in accordance, the analgesia produced by PAG stimulation are prevented by RVM lesions.^{2,41,93} As to the role of the disynaptic VLMIlat-A5-spinal circuit in pain modulation, ongoing experiments show that selective lesion of the A5 noradrenergic cell group by retrograde transport from the VLMIlat of the neurotoxin saporin-anti-dopamine- β -hydroxylase (SAP-anti-DBH) induced moderate hyperalgesia in the second phase of the formalin test (Fig 3A; unpublished results), increased noxious-evoked spinal *c-fos* expression (Fig 3B; unpublished results), and loss of the noradrenergic neuronal population of the A5 (Fig 3C and D). The data suggest that the VLMIlat input to the A5 and RVM is aimed at triggering, respectively, noradrenergic and serotonergic descending inputs. It is, however, unlikely that the VLMIlat merely regulates the an-

tinociception produced from those regions. The analgesia produced from the VLMIat presents specific characteristics distinct from the antinociception elicited from the A5 noradrenergic cell group and RVM as to its intensity and duration and to the spinal receptors involved.^{3,15,42,50,54,61,80} Given the degree of interconnectivity among the various components of the endogenous pain control system, it is likely that other areas besides A5 and RVM take part in the antinociception elicited from the VLMIat through more complex polysynaptic routes.

An interesting question is the relative roles of those polysynaptic connections and of the direct VLMIat-spinal pathway in pain modulation. According to pharmacologic data, antagonists of α_2 -adrenoreceptors and 5-HT receptors do not completely inhibit the antinociception produced from the VLMIat.^{42,54} The direct pathway may, therefore, produce the additional inhibition through neurotransmitters that remain to be established. That this direct VLMIat-spinal pathway is part of a reciprocal loop with lamina I is a further indication for its effective role in pain modulation. Similar spinomedullary loops have been shown to control pain reactions through modulation of spinal nociceptive transmission.^{4,5,7,8,28} The VLMIat is thus likely to modulate pain transmission at the spinal cord through direct and indirect pathways, and the former are probably specially suited to rapid control mechanisms.

Physiologic Signals Triggering Pain Modulation from the VLMIat

Despite accumulating data suggesting a role for the VLMIat in the descending modulation of nociceptive transmission, little is known about the physiologic signals that trigger those actions. Extracellular electrophysiologic recordings and *c-fos* studies have shown that VLMIat neurons are activated by noxious stimuli^{16,18,39,63,74,76,106} as well as by increases in blood pressure.^{58,85,115} Neurons in both laminae I and II, the main source of spinal projections to the VLMIat,^{69,70,108} are likely to convey nociceptive input. Lamina I neurons responding to peripheral noxious stimulation were antidromically activated from brainstem regions encompassing the VLMIat,⁸⁰ and noxious-evoked *c-fos* induction was observed in laminae I and II neurons that were retrogradely labeled from the VLMIat.¹⁰⁸ Curiously, the degree of nociceptive activation of lamina I neurons projecting to antinociceptive components of the endogenous pain control system was similar but much lower than for areas involved in pronociception. The percentages of lamina I neurons that project to the VLMIat, NTS, or caudal mesencephalon and express *c-fos* in response to noxious cutaneous stimulation are similar (23%, 21%, and 16%, respectively)^{5,7,54,71,73,108} and much lower than for the pronociceptive DRt (42%).⁶ Collectively, these data suggest that the nociceptive responses of lamina I neurons may be controlled by their supraspinal target.⁷² In accordance, nociceptive lamina I neurons projecting to the VLMIat have recently been shown to be directly targeted by the 2 neurotransmitters




Figure 4. Proportions of spinal neurons retrogradely labeled from the VLMIat and receiving appositions from fibers immunoreactive for DBH (hatched bars) or 5-HT (filled bars). The percentages of laminae I-III receiving appositions from fibers immunoreactive for DBH were higher than those receiving appositions from 5-HT (A). In lamina I, fusiform and flattened neurons also received higher number of appositions from fibers immunoreactive for DBH (B). Asterisks represent statistical differences ($P < .05$) between percentages determined by 1-way analysis of variance with repeated measures.

involved in the descending antinociception elicited from the VLMIat, namely, noradrenaline and 5-HT.^{42,54} Moreover, the noradrenergic input to superficial dorsal horn neurons, as measured by the numbers of cells receiving contacts from immunostained boutons, is much higher than the serotonergic (Fig 4A; unpublished results), which matches the stronger involvement of spinal α_2 -adrenoreceptors in the analgesia elicited from the VLM.^{42,54} The descending modulation of lamina I neurons appears to obey to a refined control mechanism




Figure 5. Effect of lesion in VLMLat on *c-fos* expression in spinal lamina I neurons retrogradely labeled with cholera toxin subunit b and with nuclei immunoreactive for the Fos protein (double-labeled neurons) from the same medullary region. The lesion was produced by injection of 0.3 μ L of quinolinic acid (180 nmol/ μ L) after injection of the tracer. The activation of the *c-fos* proto-oncogene was induced by noxious stimulation of the internal proximal skin of the hindpaw. The % on the y-axis represent the percentages of double-labeled neurons in the total of retrogradely labeled cells. In (A), mechanical stimulation was performed by pinching the skin with a rat-tooth forceps every 2 minutes during 2 hours. In (B), chemical stimulation consisted of subcutaneous injection of 50 μ L of 5% formalin and perfusion after 2 hours. Asterisks represent statistical differences ($P < .05$) in the percentages of retrogradely labeled neurons expressing *c-fos* between control animals (filled bars) and rats with lesions (hatched bars). Statistical differences were determined by 2-way analysis of variance for repeated measures.

that is likely to discriminate the morphologic cell type. The percentages of VLM-projecting fusiform and flattened neurons in laminae I that are apposed by noradrenergic boutons were much higher than for 5-HT, whereas for pyramidal cells no differences were detected (Fig 4B; unpublished results). Ongoing experiments further indicate that the supraspinal target may

exert an even more refined control over the nociceptive responses of lamina I neurons, depending on the nature of the noxious stimulus and the morphologic cell class. The increase in the numbers of VLMLat-projecting lamina I neurons that express *c-fos* after VLMLat lesion is more pronounced in neurons of the fusiform type after noxious mechanical cutaneous stimulation (Fig 5A; unpublished results) and of the pyramidal and flattened classes on chemical stimulation (Fig 5B; unpublished results).

The spinal input arriving at the VLMLat differs from all the other spinofugal pathways by the high incidence of lamina II cells,^{70,108} which are classically considered to be local circuit neurons.¹⁰⁷ These neurons appear to be highly involved in the transmission of nociceptive information to the VLMLat because the proportions of VLM-lat-projecting lamina II neurons that are *c-fos* activated after noxious stimulation are even higher than those of lamina I.⁸⁵ In spite of the prevalence of inhibitory neurotransmitters in lamina II, namely, gamma-aminobutyric acid (GABA), glycine, and dynorphin,^{21,117-119} it is unlikely that the lamina II-VLMLat projection is inhibitory. Lamina II neurons projecting to the VLMLat do not contain GABA, and glycine occurs exclusively in neurons containing GABA.^{118,119} On the other hand, glutamate and aspartate are present in lamina II,^{10,100} and the ultrastructural features of VLMLat terminals labeled from lamina I-II are consistent with excitatory synaptic transmission.

Increases in blood pressure have been shown to produce hypoalgesia.^{27,77,78,95,125,126} This hypertension-induced analgesia results from inhibition of nociceptive responses of spinal dorsal horn neurons^{96,110} through descending noradrenergic mechanisms.¹¹⁶ The supraspinal origin of this noradrenergic inhibition was recently ascribed to the VLMLat presumably through the noradrenergic A5 relay circuit described above. The decrease in noxious-evoked spinal *c-fos* induction observed in hypertensive animals is prevented by lesions of the VLMLat.¹¹³

Stimulation of the VLMLat produces transient decreases in blood pressure and heart rate, whereas the opposite effect is observed after local lesion,¹¹³ indicating that this specific region participates in the vasodepressor actions ascribed to the VLM.^{25,86,122} Vasodepression has been shown to result from monosynaptic, GABA-mediated inhibition of the vasopressor rostral ventrolateral medulla,¹ whereas noradrenergic antinociception is due to monosynaptic activation of the A5 noradrenergic neurons, as described above. The VLMLat activation by increases in blood pressure is thus likely to promote integrated responses of distinct neuronal populations aimed at both decreasing blood pressure through inhibition of the vasopressor nucleus and decreasing pain sensitivity through activation of the VLM-lat-A5 antinociceptive circuit. According to these data, the antinociception produced from the VLMLat is not a result of the autonomic effects. On VLMLat stimulation, the analgesia was maintained for longer than cardiovascular changes, and the latter consisted of blood pressure decreases, the opposite to what should occur if this parameter was affecting pain perception.

Conclusions and Perspectives

The VLM is an important antinociceptive center that appears to integrate nociceptive, cardiovascular, and motor functions. Although data on its performance during chronic pain are still missing, potent analgesic effects have been demonstrated for acute pain. The increase in VLM metabolic activity observed during monoarthritis⁸⁸ is, however, indicative of a strong engagement of those medullary neurons during chronic pain states. The possibility of achieving therapeutic benefits from VLM manipulation by sustained and invasive stimulation strategies is precluded because of its parallel involvement in auto-

nomic and motor regulation. However, according to the data compiled in this review, antinociception is probably promoted in a region (VLMlat) distinct from that involved in motor control (LRT). Moreover, there are strong indications that within VLMlat, the neurons involved in analgesia are distinct from those involved in autonomic control. On this basis, the possibility to selectively manipulate the VLMlat neurons involved in antinociception by recent techniques of gene therapy should be evaluated. To achieve this purpose, we recommend further detailed investigations of the anatomy and neurochemistry of the neuronal VLMlat populations involved in pain modulation or autonomic control.

References

- Agarwal SK, Calaresu FR: Monosynaptic connection from caudal to rostral ventrolateral medulla in the baroreceptor reflex pathway. *Brain Res* 555:70-74, 1991
- Aimone LD, Gebhart GF: Stimulation-produced spinal inhibition from the midbrain in the rat is mediated by an excitatory amino acid neurotransmitter in the medial medulla. *J Neurosci* 6:1803-1813, 1986
- Aimone LD, Jones SL, Gebhart GF: Stimulation-produced descending inhibition from the periaqueductal gray and nucleus raphe magnus in the rat: Mediation by spinal monoamines but not opioids. *Pain* 31:123-136, 1987
- Almeida A, Tavares I, Lima D, Coimbra A: Descending projections from the medullary dorsal reticular nucleus make synaptic contacts with spinal cord lamina I cells projecting to that nucleus: An electron microscopic tracer study in the rat. *Neuroscience* 55:1093-1106, 1993
- Almeida A, Tjølsen A, Lima D, Coimbra A, Hole K: The medullary dorsal reticular nucleus facilitates acute nociception in the rat. *Brain Res Bull* 39:7-15, 1996
- Almeida A, Lima D: Activation by cutaneous or visceral noxious stimulation of spinal neurons projecting to the medullary dorsal reticular nucleus in the rat. *Eur J Neurosci* 9:686-695, 1997
- Almeida A, Størkson R, Lima D, Hole K, Tjølsen A: The medullary dorsal reticular nucleus facilitates pain behaviour induced by formalin in the rat. *Eur J Neurosci* 11:110-122, 1999
- Almeida A, Tavares I, Lima D: Reciprocal connections between the medullary dorsal reticular nucleus and the spinal dorsal horn in the rat. *Eur J Pain* 4:373-387, 2000
- Alstermark B, Lundberg A, Norsell U, Sybirska E: Integration in descending motor pathways controlling the forelimb in the cat: Differential behavioral defects after spinal cord lesions interrupting defined pathways from higher motor centers to motoneurons. *Exp Brain Res* 42:299-318, 1981
- Antal M, Polgar E, Chalmers J, Minson JB, Llewellyn-Smith IP, Heizman CW, Somogy P: Differential populations of parvalbumin- and calbindin-D28K-immunoreactive neurons contain GABA and accumulate ³H-D-aspartate in the dorsal horn of the rat spinal cord. *J Comp Neurol* 314:114-124, 1991
- Basbaum AI, Fields HL: Endogenous pain control systems: Brainstem spinal pathways and endorphin circuitry. *Ann Rev Neurosci* 7:309-338, 1984
- Baskin DS, Mehler WR, Hosobuchi Y: Autopsy analysis of the safety, efficacy and cartography of electrical stimulation of the central gray in humans. *Brain Res* 371:231-236, 1986
- Benedetti F, Amanzio M: The neurobiology of placebo analgesia: From endogenous opioids to cholecystokinin. *Prog Neurobiol* 52:109-125, 1997
- Boivie J, Meyerson BA: A correlative anatomical and clinical study of pain suppression by deep brain stimulation. *Pain* 13:113-126, 1982
- Bowker RM, Westlund KN, Sullivan MC, Coulter JD: Organization of descending serotonergic projections to the spinal cord. *Prog Brain Res* 57:239-265, 1982
- Bullit E: Expression of C-fos protein as a marker for neuronal activity following noxious stimulation in the rat. *J Comp Neurol* 296:517-530, 1990
- Burnett A, Gebhart GF: Characterization of descending modulation of nociception from the A₅ cell group. *Brain Res* 546:271-281, 1991
- Cahusac PMB, Hill RG: Alpha-2 adrenergic receptors on neurones in the region of the lateral reticular nucleus in the rat. *Neurosci Lett* 42:279-284, 1983
- Carstens EE, Watkins LR: Inhibition of neurons in the rat spinal cord to noxious skin heating by stimulation in mid-brain periaqueductal gray or lateral reticular formation. *Brain Res* 382:266-277, 1986
- Carstens EE: Endogenous pain suppression mechanisms. *J Am Vet Med Assoc* 191:1203-1206, 1987
- Cho HJ, Basbaum AI: Ultrastructural analysis of dynorphin B-immunoreactive cells and terminals in the superficial dorsal horn of the deafferented spinal cord of the rat. *J Comp Neurol* 281:189-205, 1989
- Cledenin M, Ekerot CF, Oscarsson O, Rosén I: The lateral reticular nucleus in the cat. II. Organization of component activated from bilateral ventral flexor reflex tract (bVFRT). *Exp Brain Res* 21:487-500, 1974
- Collonier M: Synaptic patterns on different cell types in the different laminae of the cat visual cortex: An electron microscope study. *Brain Res* 9:268-287, 1968
- D'Angelo R, Evans E, Dean LA, Gaver R, Eisenach JC: Spinal clonidine prolongs labor analgesia from spinal sufentanil and bupivacaine. *Anesth Analg* 3:573-576, 1999

25. Day TA, Ro A, Renaud LP: Depressor area within the caudal ventrolateral medulla of the rat does not correspond to the A1 catecholamine cell group. *Brain Res* 279:299-302, 1983
26. De Kock MF, Pichon G, Scholtes JL: Intraoperative clonidine enhances postoperative morphine patient-controlled analgesia. *Can J Anesth* 39:537-544, 1992
27. Ditto B, Edwards MC, Miller S, D'Antono B, Blum S: The effects of sodium loading on blood pressure and pain responses to the cold pressor test. *J Psychom Res* 37:771-780, 1993
28. Dugast C, Almeida A, Lima D: The medullary dorsal reticular nucleus (DRt) increase the response of lumbar nociceptive neurons to electrical stimulation. *Soc Neurosci Abst* 26:652, 2002
29. Ekerot CF: The lateral reticular nucleus in the cat. VII. Excitatory and inhibitory projection to the ipsilateral forelimb tract (IF tract). *Exp Brain Res* 79:109-119, 1989
30. Emson PC, Corder R, Ratter SJ, Tomlin S, Lowry PJ, Ress LH, Arregui A, Rosser MN: Regional distribution of pro-opiomelanocortin-derived peptides in the human brain. *Neuroendocrinology* 38:45-50, 1984
31. Esteves F, Lima D, Coimbra A: Structural types of spinal cord marginal (lamina I) neurons projecting to the nucleus of the tractus solitarius in the rat. *Somatosens Mot Res* 10:203-216, 1993
32. Fields HL, Heinricher MM: Anatomy and physiology of a nociceptive modulatory system. *Phil Trans R Soc* 308:361-374, 1985
33. Fields HL, Barbaro NM, Heinricher MM: Brain stem neuronal circuitry underlying the antinociceptive action of opiates. *Prog Brain Res* 77:245-257, 1988
34. Fields HL, Basbaum AI: Endogenous pain control mechanisms, in Wall PD, Melzak R (eds): *Textbook of Pain*. Edinburgh, Scotland, Churchill Livingstone, 1989, pp 206-217
35. Fields HL, Gebhart GF, Randich A: Brainstem modulation of nociception, in Klemm WR, Vertes RP (eds): *Brainstem Mechanisms of Behavior*. New York, NY, John Wiley & Sons, Inc 1990, pp 315-352
36. Fields HL, Heinricher MM, Mason P: Neurotransmitters in nociceptive modulatory circuits. *Ann Rev Neurosci* 14:219-245, 1991
37. Fields HL, Malick A, Burstein R: Dorsal horn projection targets of ON and OFF cells in the rostral ventromedial medulla. *J Neurophysiol* 74:742-759, 1995
38. Fields HL: Pain modulation: Expectation, opioid analgesia and virtual pain. *Prog Brain Res* 122:245-253, 2000
39. Fleetwood-Walker SM, Coote JH, Gilbey MP: Identification of spinally projecting neurones in the A₁ catecholamine cell group of the ventrolateral medulla. *Brain Res* 273:25-33, 1983
40. Foong FW, Duggan AW: Brain-stem areas tonically inhibiting dorsal horn neurones: Studies with microinjection of the GABA analogue piperidine-4-sulphonic acid. *Pain* 27:361-371, 1986
41. Gebhart GF, Sandkuhler J, Thalhammer JG, Zimmermann M: Inhibition of spinal nociceptive information by stimulation in midbrain of the cat is blocked by lidocaine microinjected in nucleus raphe magnus and medullary reticular formation. *J Neurophysiol* 50:1446-1459, 1983
42. Gebhart GF, Ossipov MH: Characterization of inhibition of the spinal nociceptive tail-flick reflex in the rat from the medullary lateral reticular nucleus. *J Neurosci* 6:701-713, 1986
43. Gebhart GF, Randich A: Brainstem modulation of nociception, in Klemm WR, Vertes RP (eds): *Brainstem Mechanisms of Behavior*, New York, NY, John Wiley & Sons, Inc, pp 315-352, 1990
44. Gogas KR, Presley RW, Levine JD, Basbaum AI: The antinociceptive action of supraspinal opioids results from an increase in descending inhibitory control: Correlation of nociceptive behavior and c-fos expression. *Neuroscience* 41:617-628, 1991
45. Gray EG: Electron microscopy of excitatory and inhibitory synapses: A brief review. *Prog Brain Res* 31:141-155, 1969
46. Grevert P, Goldstein A: Placebo analgesia, naloxone and the role of endogenous opioids, in White L, Tursky B, Schwartz GE (eds): *Placebo: Theory, Research and Mechanisms*. New York, The Guilford Press, New York, 1985, pp 332-350
47. Gordon NC, Heller PH, Levine JD: Enhancement of pentazocine analgesia by clonidine. *Pain* 48:167-169, 1992
48. Hall JG, Duggan, AW, Morton CR, Johnson SM: The location of brainstem neurones tonically inhibiting dorsal horn neurones of the cat. *Brain Res* 244:215-222, 1982
49. Hammond DL: Control systems for nociceptive afferent processing: The descending inhibitory pathways, in Yaksh T (ed): *Spinal Afferent Processing*. New York, Plenum Press, 1986, pp 363-390
50. Hammond DL, Yaksh TL: Antagonism of stimulation-produced antinociception by intrathecal administration of methysergide or phentolamine. *Brain Res* 298:329-337, 1984
51. Hammond DL, Presley R, Gogas KR, Basbaum AI: Morphine or U-50,488 suppresses Fos protein-like immunoreactivity in the spinal cord and nucleus tractus solitarius evoked by noxious visceral stimulation in the rat. *J Comp Neurol* 315:244-253, 1992
52. Holstege G: The basic, somatic and emotional components of the motor system in mammals, in Paxinos G (ed): *The Rat Nervous System*, New York, Academic Press, 1995, pp 137-154
53. Hunt SP, Pini A, Evan G: Induction of c-fos-like protein in spinal neurons following sensory stimulation. *Nature* 328:632-634, 1987
54. Janss AJ, Gebhart GF: Spinal monoaminergic receptors mediate the antinociception produced by glutamate in the medullary lateral reticular nucleus. *J Neurosci* 7:2862-2873, 1987
55. Janss AJ, Gebhart GF: Quantitative characterization and spinal pathway mediating inhibition of spinal nociceptive transmission from the lateral reticular nucleus in the rat. *J Neurophysiol* 59:226-247, 1988
56. Janss AJ, Gebhart GF: Brainstem and spinal pathways mediating descending inhibition from the medullary lateral reticular nucleus in the rat. *Brain Res* 440:109-122, 1988
57. Jensen TS, Yaksh TL: Spinal monoamine and opiate systems partly mediate the antinociceptive effects produced by glutamate at brainstem sites. *Brain Res* 321:287-297, 1984
58. Jeske I, Morrison SF, Cravo SL, Reis DJ: Identification of baroreceptor reflex interneurons in the caudal ventrolateral medulla. *Am J Physiol* 264:R169-R178, 1993
59. Jones SL, Gebhart GF: Characterization of coeruleospinal inhibition of the nociceptive tail-flick reflex in the rat: Me-

- diation by spinal α_2 -adrenoreceptors. *Brain Res* 364:315-330, 1986
60. Jones SL, Gebhart GF: Quantitative characterization of ceruleospinal inhibition of nociceptive transmission in the rat. *J Neurophysiol* 56:1397-1410, 1986
 61. Jones SL: Descending control of nociception, in Gildenberg P (ed): *The Initial Processing of Pain and Its Descending Control: Spinal and Trigeminal Systems*. Basel, Switzerland, Karger, 1992, pp 203-295
 62. Lakos S, Basbaum AI: An ultrastructural study of the projections from the midbrain periaqueductal gray to spinally projecting, serotonin-immunoreactive neurons of the medullary nucleus raphe magnus in the rat. *Brain Res* 443:383-388, 1988
 63. Lantéri-Minet M, Fugazza JW, de Pommery J, Menétrey D: Hindbrain structures involved in pain processing as revealed by the expression of *c-fos* and other immediate early genes. *Neuroscience* 58:287-298, 1994
 64. Liebeskind JC, Guilbaud G, Besson JM, Oliveras J: Analgesia from electrical stimulation of the periaqueductal gray matter in the cat: Behavioral observations and inhibitory effects on spinal interneurons. *Brain Res* 50:441-446, 1973
 65. Light AR, Kavookjian AM: Morphology and ultrastructure of physiologically identified substantia gelatinosa (lamina II) neurons with axons that terminate in deeper dorsal horn laminae (III-V). *J Comp Neurol* 267:172-189, 1988
 66. Lima D, Coimbra A: The spinothalamic system of the rat: Structural types of retrogradely labelled neurons in the marginal zone (lamina I). *Neuroscience* 27:215-230, 1988
 67. Lima D, Coimbra A: Morphological types of spinomesencephalic neurons in the marginal zone (lamina I) of the rat spinal cord, as shown after retrograde labelling with cholera toxin subunit B. *J Comp Neurol* 279:327-339, 1989
 68. Lima D, Coimbra A: Structural types of marginal (lamina I) neurons projecting to the dorsal reticular nucleus of the medulla oblongata. *Neuroscience* 34:591-606, 1990
 69. Lima D, Coimbra A: Neurons in the substantia gelatinosa rolandi (lamina II) project to the caudal ventrolateral reticular formation of the medulla oblongata in the rat. *Neurosci Lett* 132:16-18, 1991
 70. Lima D, Mendes-Ribeiro JA, Coimbra A: The spino-latero-reticular system of the rat: Projections from the superficial dorsal horn and structural characterization of marginal neurons involved. *Neuroscience* 45:137-152, 1991
 71. Lima D, Esteves F, Coimbra A: *C-fos* activation by noxious input of spinal neurons projecting to the nucleus of the tractus solitarius in the rat, in Gebhart GF, Hammond D, Jensen TS (eds): *Proceedings of the 7th World Congress on Pain*, Seattle, WA, IASP Press, 1994, pp 423-434
 72. Lima D: Endogenous pain modulatory system in the light of the gate control theory. *Pain Forum* 5:31-39, 1996
 73. Lima D: Functional anatomy of spinofugal nociceptive pathways. *Pain Rev* 4:1-19, 1997
 74. Liu RH, Tang JS, Hou ZL, Jia H: Responses of reticulospinal neurons in the lateral reticular nucleus area of the rat to cutaneous noxious and non-noxious stimulation. *Neurosci Lett* 109:92-96, 1990
 75. Liu RH, Zhao ZQ: Selective blockade by yohimbine of descending spinal inhibition from lateral reticular nucleus but not from locus coeruleus in rats. *Brain Res* 142:65-68, 1992
 76. Liu RH, Tang JS, Hou ZL: Effect of systemic morphine on neurons in the lateral reticular nucleus area of the rat. *Brain Res Bull* 32:179-184, 1993
 77. Maixner W, Touw KB, Brody MJ, Gebhart GF, Long JP: Factors influencing the altered pain perception in the spontaneously hypertensive rat. *Brain Res* 237:137-145, 1982
 78. Maixner W: Interactions between cardiovascular and pain modulatory systems: Physiological and pathophysiological implications. *J Cardiovasc Electrophysiol* 2:S3-S12, 1991
 79. McKellar S, Loewy AD: Efferent projections of the A_1 catecholaminergic cell group in the rat: An autoradiographic study. *Brain Res* 241:11-29, 1982
 80. Menétrey D, De Pommery J, Besson JM: Electrophysiological characteristics of lumbar spinal neurons backfired from the lateral reticular nucleus in the rat. *J Neurophysiol* 52:595-609, 1984
 81. Melzak R, Wall PD: Pain mechanisms: A new theory. *Science* 150:971-979, 1965
 82. Millar J, Williams GV: Effects of iontophoresis of nor-adrenaline and stimulation of the periaqueductal gray on single-unit activity in the rat superficial dorsal horn. *J Comp Neurol* 287:119-133, 1989
 83. Miller JF, Proudfit HK: Antagonism of stimulation-produced antinociception from ventrolateral pontine sites by intrathecal administration of α -adrenergic antagonists and naloxone. *Brain Res* 530:20-34, 1990
 84. Morton CR, Johnson SM, Duggan AW: Lateral reticular regions and the descending control of dorsal horn neurones of the cat: Selective inhibition by electrical stimulation. *Brain Res* 275:13-21, 1983
 85. Murphy AZ, Ennis M, Siple MT, Behbehani MM: Directionally specific changes in arterial pressure induce differential patterns of Fos expression in discrete areas of the rat brainstem: A double-labelling study for Fos and catecholamines. *J Comp Neurol* 349:36-50, 1994
 86. Murugaian J, Sundaram K, Krieger A, Sapru H: Electrolytic lesions in the depressor area of the ventrolateral medulla abolish depressor responses to the aortic nerve stimulation. *Brain Res* 499:371-377, 1989
 87. Ness TJ, Gebhart GF: Quantitative comparison of inhibition of visceral and cutaneous spinal nociceptive transmission from the midbrain and medulla in the rat. *J Neurophysiol* 58:850-865, 1987
 88. Neto FL, Schadrack A, Ableitner A, Castro-Lopes JM, Barntenstein P, Zieglgänsberger TR, Tölle TR: Supraspinal metabolic activity changes in the rat during adjuvant monoarthritis. *Neuroscience* 94:607-621, 1999
 89. Ossipov MH, Gebhart GF: Opioid, cholinergic and α -adrenergic influences on the modulation of nociception from the lateral reticular nucleus. *Brain Res* 384:282-293, 1986
 90. Ota K, Namiki A, Ujike Y, Takahashi I: Prolongation of tetracaine spinal anesthesia by oral clonidine. *Anesth Analg* 75:262-264, 1992
 91. Paxinos G, Watson C: *The Rat Brain in Stereotaxic Coordinates*. New York, Academic Press, 1998
 92. Pitius CW, Seizinger BR, Pasi A, Mehraaen P, Herz A: Distribution and characterization of opioid peptides derived from proenkephalin A in human and rat central nervous system. *Brain Res* 304:127-136, 1984
 93. Prieto GJ, Cannon JT, Liebeskind JC: N raphe magnus

lesions disrupt stimulation-produced analgesia from ventral but not dorsal midbrain areas in the rat. *Brain Res* 261:53-57, 1983

94. Proudfit HK: Pharmacological evidence for the modulation of nociception by noradrenergic neurons. *Prog Brain Res* 77:357-370, 1988

95. Randich A, Maixner W: Interactions between cardiovascular and pain regulatory systems. *Neurosci Biobehav Rev* 8:343-367, 1984

96. Randich A, Robertson JD: Spinal nociceptive transmission in the spontaneously hypertensive and Wistar-Kyoto normotensive rat. *Pain* 58:169-183, 1994

97. Rees H, Roberts MHT: Anterior pretectal stimulation alters the responses of spinal dorsal horn neurones to cutaneous stimulation in the rat. *J Neurophysiol* 385:415-436, 1987

98. Ren K, Ruda MA: Descending modulation of Fos expression after persistent peripheral inflammation. *Neuroreport* 7:2186-2190, 1996

99. Roberts MHT, Rees H: The antinociceptive effects of stimulating the pretectal nucleus in the rat. *Pain* 25:83-93, 1986

100. Rustioni A, Cuenod M: Selective retrograde transport of D-aspartate in spinal interneurons and cortical neurons of rats. *Brain Res* 236:143-155, 1982

101. Satoh M, Akaike A, Nakazawa T, Takagi H: Evidence for involvement of separate mechanisms in the production of analgesia by electrical stimulation of the nucleus reticularis paragigantocellularis and nucleus raphe magnus in the rat. *Brain Res* 194:525-529, 1980

102. Satoh M, Oku R, Akaike A: Analgesia produced by microinjection of L-glutamate into the rostral ventromedial bulbar nuclei of the rat and its inhibition by intrathecal α -adrenergic blocking agents. *Brain Res* 261:361-364, 1983

103. Shepherd G: The synaptic organization of the brain. Oxford, Oxford University Press, 1990

104. Simpson BA: Spinal cord and brain stimulation, in Wall PD, Melzak R (eds): *Textbook of Pain*. Edinburgh, Churchill Livingstone, 1999, pp 1353-1381

105. Stamford JA: Descending control of pain. *Br J Anaesth* 75:217-227, 1995

106. Strassman AM, Vos BP: Somatotopic and laminar organization of fos-like immunoreactivity in the medullary and upper cervical dorsal horn induced by noxious facial stimulation in the rat. *J Comp Neurol* 331:495-516, 1993

107. Szentágothai J: Neuronal and synaptic arrangement in the substantia gelatinosa. *J Comp Neurol* 122:219-239, 1964

108. Tavares I, Lima D, Coimbra A: Neurons in the superficial dorsal horn of the rat spinal cord projecting to the medullary reticular formation express *c-fos* after noxious stimulation of the skin. *Brain Res* 623:278-282, 1993

109. Tavares I, Lima D: Descending projections from the caudal medulla oblongata to the superficial or deep dorsal horn of the rat spinal cord. *Exp Brain Res* 99:455-463, 1994

110. Tavares I, Albino-Teixeira A, Lima D: Hypertension inhibits noxious evoked *c-fos* expression in the rat spinal cord. *Neuroreport* 6:1664-1668, 1995

111. Tavares I, Lima D, Coimbra A: The ventrolateral medulla of the rat is connected with the spinal cord dorsal horn by an indirect descending pathway relayed in the A₅ noradrenergic cell group. *J Comp Neurol* 374:84-95, 1996

112. Tavares I, Lima D, Coimbra A: The pontine A₅ noradrenergic cells which project to the spinal cord dorsal horn are reciprocally connected with the caudal ventrolateral medulla in the rat. *Eur J Neurosci* 9:2452-2461, 1997

113. Tavares I, Almeida A, Albino-Teixeira A, Lima D: Lesions of the caudal ventrolateral medulla block the hypertension-induced inhibition of noxious-evoked *c-fos* expression in the rat spinal cord. *Eur J Pain* 1:149-160, 1997

114. Tavares I, Almeida A, Esteves F, Lima D, Coimbra A: The caudal ventrolateral medullary reticular formation is reciprocally connected with the spinal cord. *Soc Neurosci Abstr* 24:1132, 1998

115. Terui P, Masuda Y, Saeki Y, Kumada M: Activity of barosensitive neurons in the caudal ventrolateral medulla that send axonal projections to the rostral ventrolateral medulla in rabbits. *Neurosci Lett* 118:211-214, 1990

116. Thurston CL, Randich A: Acute increases in arterial blood pressure produced by occlusion of the abdominal aorta induces antinociception: Peripheral and central substrates. *Brain Res* 519:12-22, 1990

117. Todd AJ, Sullivan AC: Light microscope study of the coexistence of GABA-like and glycine-like immunoreactivities in the spinal cord of the rat. *J Comp Neurol* 296:496-505, 1990

118. Todd AJ: Evidence that acetylcholine and glycine exist in different populations of GABAergic neurons in lamina III of rat spinal cord. *Neuroscience* 44:741-746, 1991

119. Todd AJ, Spike RC: Localization of classical neurotransmitters and neuropeptides within neurons in laminae I-III of the mammalian spinal dorsal horn. *Prog Neurobiol* 41:609-645, 1993

120. Uchizono K: Characteristics of excitatory and inhibitory synapses in the central nervous system of the cat. *Nature* 207:642-643, 1965

121. Westlund KN, Bowker RM, Ziegler MG, Coulter JD: Noradrenergic projections to the spinal cord of the rat. *Brain Res* 263:15-31, 1983

122. Willete RN, Barcas PP, Krieger AJ, Sapru HN: Vasodepressor areas in the rat medulla: Identification by injection of L-glutamate. *Neuropharmacology* 22:1071-1079, 1983

123. Willis WD, Coggeshall RE: Sensory mechanisms of the spinal cord. New York, Plenum Press, 1991

124. Woulfe JM, Flumerfelt BA, Hryciyshyn AW: Efferent connections of the A₁ noradrenergic cell group: A DBH immunohistochemical and PHA-L anterograde tracing study. *Exp Neurol* 109:308-322, 1990

125. Zamir N, Shuber E: Altered pain perception in hypertensive humans. *Brain Res* 201:471-474, 1980

126. Zamir N, Simantov R, Segal M: Pain sensitivity and opioid activity in genetically hypertensive rats. *Brain Res* 184:299-310, 1980

127. Zamir N, Maixner W: The relationship between cardiovascular and pain regulatory systems. *Ann N Y Acad Sci* 467:371-389, 1986

128. Zhang RX, Wang R, Chen JY, Qiao JT: Effects of descending inhibitory systems on the *c-Fos* expression in the rat spinal cord during formalin-induced noxious stimulation. *Neuroscience* 58:299-304, 1994