

Heart rate recovery: what now?

Physicians have known for hundreds, perhaps thousands, of years that the pulse and its rate are simple, yet powerful, measures of human health [1]. In the early 17th century, Santorio Sanctorius (1561–1636) invented the ‘pulsilogy’, a clever pendulum-like device by which physicians could quantitatively assess pulse rate. About 100 years later, John Floyer (1649–1734) became the first physician to formally measure pulse as number of beats per minute. His work stimulated others to correlate pulse with other measures like age and fever [1]. In 1843, a physician correspondent to the *Lancet* commented on changes in pulse with stress, writing ‘The pulse in rising and falling from accidental and temporary excitement, rises and falls through a series of duodecimal degrees; when within the first few minutes of an interview the pulse of a patient rapidly subsides from 120 to 108, 96, 84, a knowledge is at once afforded of the highly excitable and therefore susceptible constitution of the patient’ [2]. This may be one of the earliest recordings of the clinical implications of heart rate changes occurring during and after stress.

Over the past 50 years, scientists have come to better appreciate links between heart rate and health, links that may reflect fundamental pathophysiological processes [3]. Amongst vertebrates, there is a strong inverse association between resting heart rate and life expectancy. In 1971, Coburn *et al.* [4] showed that low-dose digoxin, a parasympathomimetic agent that slows heart rate, could prolong life in mice. In 2008, Custodis *et al.* [5] used a knockout mouse model to show that ivabradine, an agent that slows heart rate at the level of the sinus node, can slow development and progression of atherosclerosis. In humans, there are extensive epidemiological data showing that higher resting heart rates predict greater mortality [3, 6]. Amongst patients randomized to beta-blockers in postmyocardial infarction trials, the magnitude of survival benefit correlates most strongly with the drug-induced reduction in heart rate [7]. And just last year, Swedberg *et al.* [8] showed in a randomized trial of over 6500 patients that ivabradine reduces the risk

of heart failure hospitalization and death, particularly amongst patients with higher heart rates at baseline [9]. Although there is still much to learn, resting heart rate has evolved as a measure going from a pathophysiological phenomenon to an independent epidemiological risk factor all the way to a viable treatment target.

Now there is increasing interest in heart rate changes with stress [3]. With the onset of physical or mental stress, there is an abrupt withdrawal of central nervous system vagal output, along with a gradual increase in sympathetic tone. During stress, heart rate is largely driven by increasing sympathetic tone, which is modulated by both central and hormonal factors. The increase in heart rate is known as the ‘chronotropic response’ [3]. Investigators studying healthy adults in the Framingham Heart Study showed that an impaired chronotropic response, or ‘chronotropic incompetence’, predicts a higher risk of death or major cardiac events [10]. These findings were later confirmed in clinical cohorts [11].

In 1994, Imai *et al.* [12] showed that vagal tone is reactivated rapidly within the first 30–40 s following exercise, but this reactivation is blunted in heart failure. Cole *et al.* [13] took advantage of these observations to evaluate the link between the decline in heart rate in the first few minutes after exercise, or ‘heart rate recovery’, and mortality. In a cohort of 2428 patients who were candidates for first-time coronary angiography and who were referred for treadmill myocardial perfusion imaging, heart rate recovery emerged as an independent predictor of death, even after accounting for exercise capacity, chronotropic response and evidence of myocardial ischaemia.

Since Cole *et al.*’s report, other groups have confirmed the association between heart rate recovery, its link to parasympathetic reactivation [6, 14] and adverse cardiovascular outcomes. An impaired heart rate recovery predicts in men and women all-cause mortality [15], cardiovascular mortality [16], sudden cardiac death [17], heart failure mortality [18] and even risk of developing atrial fibrillation [19]. Heart rate recovery has also been linked to other disease pathways, in particular insulin resistance [20]. In an animal model, heart rate recovery predicts susceptibility

Disclaimer: The views expressed herein are those of the author and do not necessarily reflect the views of NHLBI or the US Department of Health and Human Services.

to developing ventricular fibrillation during the first few minutes of an experimental myocardial infarction [21]; this observation reflects a finding in humans that heart rate recovery is a better predictor of sudden death than other modes of cardiac death [17]. It is conceivable that heart rate recovery reflects the ability of the parasympathetic nervous system to protect the heart from ischaemia-induced arrhythmias.

In the current issue of the *Journal of Internal Medicine*, Savonen *et al.* [22] followed the outcomes of 1102 healthy middle-aged men who underwent cycle ergometry as part of a standard evaluation within a population-based East Finland cohort study. During 18 years of follow-up, 238 men died. In their carefully performed study, the investigators found that heart rate recovery predicted death, even after accounting for standard risk factors. However, the association was markedly attenuated by chronotropic response and exercise capacity. It is not clear why the authors' findings are at odds with previously published reports. One likely reason is the absence of a uniform recovery protocol. Admittedly, the absence of internationally accepted and adopted exercise recovery protocols represents a major limitation in comparing reports and in implementing clinical recommendations. The authors used cycle ergometry; some investigators have found that cycle ergometry, as compared to treadmill exercise, may lead to altered measures of functional capacity and heart rate recovery (particularly during the first minute after exercise) [23].

What now for heart rate recovery? The time has come to redirect focus towards therapeutic implications. A number of tantalizing, though small, reports suggest that heart rate recovery can be modified with exercise training [24], weight loss [25], bariatric surgery [26], angiotensin-converting enzyme inhibitors [27], pyridostigmine (a cholinesterase inhibitor) [28] and even statins [29]. One group found that heart rate recovery may predict which heart failure patients are most likely to respond to biventricular pacing [30]. Should otherwise healthy adults with abnormal heart rate recovery be targeted for primary prevention interventions, including rigorous exercise programs? Should heart rate recovery be used to identify patients who might be most likely to benefit from heart rate modulating and other therapies? Could heart rate recovery itself function as a viable treatable biomarker, similar to resting heart rate in patients with established disease? Should our new insights into the central role of heart rate and autonomic dysfunction stimulate us to develop and test new wholly new approaches,

including modulators like ivabradine and pyridostigmine? By thinking about questions like these, the scientific community can further leverage medicine's longstanding fascination with heart rate into new paradigms for improving health.

Conflict of interest

No conflict of interest to declare.

Funding

None.

M. S. Lauer

From the Office of the Director, Division of Cardiovascular Sciences of the National Heart, Lung, and Blood Institute (NHLBI), Bethesda, MD, USA

References

- 1 Ghasemzadeh N, Zafari AM. A brief journey into the history of the arterial pulse. *Cardiol Res Pract* 2011; **2011**: 164832.
- 2 Anthropos. Diagnosis by the pulse. *Lancet* 1843; **40**: 87–88.
- 3 Lauer MS. Autonomic function and prognosis. *Cleve Clin J Med* 2009; **76**(Suppl 2): S18–S22.
- 4 Coburn AF, Grey RM, Rivera SM. Observations on the relation of heart rate, life span, weight and mineralization in the digoxin-treated A-J mouse. *Johns Hopkins Med J* 1971; **128**: 169–193.
- 5 Custodis F, Baumhakel M, Schlimmer N *et al.* Heart rate reduction by ivabradine reduces oxidative stress, improves endothelial function, and prevents atherosclerosis in apolipoprotein E-deficient mice. *Circulation* 2008; **117**: 2377–2387.
- 6 Lahiri MK, Kannankeril PJ, Goldberger JJ. Assessment of autonomic function in cardiovascular disease: physiological basis and prognostic implications. *J Am Coll Cardiol* 2008; **51**: 1725–1733.
- 7 Reil JC, Bohm M. The role of heart rate in the development of cardiovascular disease. *Clin Res Cardiol* 2007; **96**: 585–592.
- 8 Swedberg K, Komajda M, Bohm M *et al.* Ivabradine and outcomes in chronic heart failure (SHIFT): a randomised placebo-controlled study. *Lancet* 2010; **376**: 875–885.
- 9 Bohm M, Swedberg K, Komajda M *et al.* Heart rate as a risk factor in chronic heart failure (SHIFT): the association between heart rate and outcomes in a randomised placebo-controlled trial. *Lancet* 2010; **376**: 886–894.
- 10 Lauer MS, Okin PM, Larson MG, Evans JC, Levy D. Impaired heart rate response to graded exercise. Prognostic implications of chronotropic incompetence in the Framingham Heart Study. *Circulation* 1996; **93**: 1520–1526.
- 11 Lauer MS, Francis GS, Okin PM, Pashkow FJ, Snader CE, Marwick TH. Impaired chronotropic response to exercise stress testing as a predictor of mortality. *JAMA* 1999; **281**: 524–529.
- 12 Imai K, Sato H, Hori M *et al.* Vagally mediated heart rate recovery after exercise is accelerated in athletes but blunted in patients

- with chronic heart failure. *J Am Coll Cardiol* 1994; **24**: 1529–1535.
- 13 Cole CR, Blackstone EH, Pashkow FJ, Pothier CE, Lauer MS. Heart-rate recovery immediately after exercise as a predictor of mortality. *N Engl J Med* 1999; **341**: 1351–1357.
 - 14 Goldberger JJ, Le FK, Lahiri M, Kannankeril PJ, Ng J, Kadish AH. Assessment of parasympathetic reactivation after exercise. *Am J Physiol Heart Circ Physiol* 2006; **290**: H2446–H2452.
 - 15 Shetler K, Marcus R, Froelicher VF *et al*. Heart rate recovery: validation and methodologic issues. *J Am Coll Cardiol* 2001; **38**: 1980–1987.
 - 16 Cole CR, Foody JM, Blackstone EH, Lauer MS. Heart rate recovery after submaximal exercise testing as a predictor of mortality in a cardiovascularly healthy cohort. *Ann Intern Med* 2000; **132**: 552–555.
 - 17 Jouven X, Empana JP, Schwartz PJ, Desnos M, Courbon D, Ducimetiere P. Heart-rate profile during exercise as a predictor of sudden death. *N Engl J Med* 2005; **352**: 1951–1958.
 - 18 Tang YD, Dewland TA, Wencker D, Katz SD. Post-exercise heart rate recovery independently predicts mortality risk in patients with chronic heart failure. *J Card Fail* 2009; **15**: 850–855.
 - 19 Maddox TM, Ross C, Ho PM, Magid D, Rumsfeld JS. Impaired heart rate recovery is associated with new-onset atrial fibrillation: a prospective cohort study. *BMC Cardiovasc Dis* 2009; **9**: 11.
 - 20 Lind L, Andren BC. Heart rate recovery after exercise is related to the insulin resistance syndrome and heart rate variability in elderly men. *Am Heart J* 2002; **144**: 666–672.
 - 21 Smith LL, Kukielka M, Billman GE. Heart rate recovery after exercise: a predictor of ventricular fibrillation susceptibility after myocardial infarction. *Am J Physiol Heart Circ Physiol* 2005; **288**: H1763–H1769.
 - 22 Savonen KP, Kiviniemi V, Laaksonen DE *et al*. Two-minute heart rate recovery after cycle ergometer exercise and all-cause mortality in middle-aged men. *J Intern Med*. Jul 30. DOI: 10.1111/j.1365-2796.2011.02434.x. [Epub ahead of print].
 - 23 Maeder MT, Ammann P, Rickli H, Brunner-La Rocca HP. Impact of the exercise mode on heart rate recovery after maximal exercise. *Eur J Appl Physiol* 2009; **105**: 247–255.
 - 24 Adams BJ, Carr JG, Ozonoff A, Lauer MS, Balady GJ. Effect of exercise training in supervised cardiac rehabilitation programs on prognostic variables from the exercise tolerance test. *Am J Cardiol* 2008; **101**: 1403–1407.
 - 25 Brinkworth GD, Noakes M, Buckley JD, Clifton PM. Weight loss improves heart rate recovery in overweight and obese men with features of the metabolic syndrome. *Am Heart J* 2006; **152**: 693.e691–696.
 - 26 Wasmund SL, Owan T, Yanowitz FG *et al*. Improved heart rate recovery after marked weight loss induced by gastric bypass surgery: two-year follow up in the Utah Obesity Study. *Heart Rhythm* 2011; **8**: 84–90.
 - 27 Sipahi I, Tekin G, Yigit Z, Guzelsoy D, Guven O. Effect of quinapril on the attenuated heart rate recovery of type 2 diabetic subjects without known coronary artery disease. *Clin Cardiol* 2004; **27**: 480–484.
 - 28 Serra SM, Costa RV, Teixeira De Castro RR, Xavier SS, Nobrega AC. Cholinergic stimulation improves autonomic and hemodynamic profile during dynamic exercise in patients with heart failure. *J Card Fail* 2009; **15**: 124–129.
 - 29 Tekin G, Tekin A, Canatar T *et al*. Simvastatin improves the attenuated heart rate recovery of type 2 diabetics. *Pharmacol Res* 2006; **54**: 442–446.
 - 30 Thomas DE, Exton SA, Yousef ZR. Heart rate deceleration after exercise predicts patients most likely to respond to cardiac resynchronisation therapy. *Heart* 2010; **96**: 1385–1389.

Correspondence: Michael S. Lauer MD, 6701 Rockledge Drive, Room 8128, Bethesda, MD 20892, USA.
(fax: 301-480-1864; e-mail: lauerm@nhbli.nih.gov).