

**Response to Elia *et al.*****'Tauroursodeoxycholic acid in the treatment of patients with amyotrophic lateral sclerosis'**

M. Michaud<sup>a,b</sup>, C. Gauchet<sup>c</sup> and C. Dray<sup>d</sup>

<sup>a</sup>CHU Toulouse, Service de Médecine Interne et Gériatrie, <sup>b</sup>Université de Toulouse, Faculté de Médecine, <sup>c</sup>CHU Toulouse, Direction de la Recherche et de l'Innovation, and <sup>d</sup>Inserm, U1048, Toulouse, France

Correspondence: M. Michaud, Post Urgence Gériatrique, Pavillon Sénac, Hôpital Purpan, Place Dr Baylac, 31049 Toulouse Cedex, France (tel.: +33 561 77 92 88; fax: +33 561 77 99 27; e-mail: martin.michaud85@gmail.com).

**Keywords:** amyotrophic lateral sclerosis, endoplasmic reticulum stress, tauroursodeoxycholic acid, unfolded protein response

doi:10.1111/ene.12729

Received: 20 February 2015

Accepted: 26 February 2015

Dear Editor,

We have read with attention the original paper by Elia *et al.* [1] about tauroursodeoxycholic acid (TUDCA) in the treatment of patients with amyotrophic lateral sclerosis (ALS).

The choice of TUDCA as a potential treatment of ALS is justified by potential neuroprotective and anti-cytotoxic effects of TUDCA. Regarding this point we do think that the role of endoplasmic reticulum (ER) stress in the pathophysiology of ALS could have been developed in this paper.

ER is responsible for the folding, post-translational modification and trafficking of many transmembrane and secretory proteins [2]. Accumulation of unfolded and misfolded proteins leads to ER stress, also called unfolded protein response. Unfolded protein response is initially a physiological response to restore ER homeostasis. In contrast, when the ER stress is not resolved apoptosis pathways are triggered.

In animal models of ALS, an excessive ER stress activation has been observed [3]. Moreover, ER stress has been observed in spinal cord tissues of patients suffering from sporadic ALS [4–6]. As ER stress can induce oxidative stress, apoptosis and autophagy, targeting ER stress is a potential therapeutic direction in several diseases, including ALS [7,8]. Indeed, there is a growing interest in TUDCA and its potential therapeutic effects. Elia *et al.* suggest that the beneficial effect of TUDCA is mediated by mitochondrial dysfunction. We advocate that a main point with TUDCA is its ability to reduce ER stress. Studies report that TUDCA could be an anti-apoptotic agent for a number of other neurodegenerative diseases (Alzheimer's disease, Parkinson's disease and Huntington's disease) and metabolic diseases (diabetes, obesity, stroke, acute myocardial infarction) [9].

Finally, a measurement of ER stress levels in peripheral blood mononuclear cells of patients in both groups (TUDCA and placebo) would have been interesting [10].

**Disclosure of conflicts of interest**

The authors declare no financial or other conflicts of interest.

**References**

1. Elia AE, Lalli S, Monsurro MR, *et al.* Tauroursodeoxycholic acid in the treatment of patients with amyotrophic lateral sclerosis. *Eur J Neurol* 2015; doi:10.1111/ene.12664. [Epub ahead of print].
2. Schröder M. Endoplasmic reticulum stress responses. *Cell Mol Life Sci* 2008; **65**: 862–894.
3. Walker AK, Atkin JD. Stress signaling from the endoplasmic reticulum: a central player in the pathogenesis of amyotrophic lateral sclerosis. *IUBMB Life* 2011; **63**: 754–763.
4. Ilieva EV, Ayala V, Jové M, *et al.* Oxidative and endoplasmic reticulum stress interplay in sporadic amyotrophic lateral sclerosis. *Brain J Neurol* 2007; **130**: 3111–3123.
5. Atkin JD, Farg MA, Walker AK, McLean C, Tomas D, Horne MK. Endoplasmic reticulum stress and induction of the unfolded protein response in human sporadic amyotrophic lateral sclerosis. *Neurobiol Dis* 2008; **30**: 400–407.
6. Ito Y, Yamada M, Tanaka H, *et al.* Involvement of CHOP, an ER-stress apoptotic mediator, in both human sporadic ALS and ALS model mice. *Neurobiol Dis* 2009; **36**: 470–476.
7. Maly DJ, Papa FR. Druggable sensors of the unfolded protein response. *Nat Chem Biol* 2014; **10**: 892–901.
8. Dufey E, Sepúlveda D, Rojas-Rivera D, Hetz C. Cellular mechanisms of endoplasmic reticulum stress signaling in health and disease. 1. An overview. *Am J Physiol Cell Physiol* 2014; **307**: C582–C594.
9. Vang S, Longley K, Steer CJ, Low WC. The unexpected uses of urso- and tauroursodeoxycholic acid in the treatment of non-liver diseases. *Glob Adv Health Med* 2014; **3**: 58–69.
10. Mozzini C, Fratta Pasini A, Garbin U, *et al.* Increased endoplasmic reticulum stress and Nrf2 repression in peripheral blood mononuclear cells of patients with stable coronary artery disease. *Free Radic Biol Med* 2014; **68**: 178–185.