


Tandem 5-*exo-trig* allyl and 3-*exo-trig* radical cyclisation and rearrangement to copa and ylanga type sesquiterpene skeleton

K. Kaliappan and G. S. R. Subba Rao*

Department of Organic Chemistry, Indian Institute of Science, Bangalore 560 012, India


A novel tandem 5-*exo-trig* allyl and 3-*exo-trig* radical cyclisation and rearrangement to copa and ylanga type sesquiterpene skeleton is reported.

The use of radicals in organic synthesis has been given increased attention during the last two decades.¹ Although allyl radicals have been known for almost a decade,² they have rarely been used in organic synthesis^{3,4} as they are less reactive and more stable when compared to their saturated and vinylic counterparts. In continuation of our interest in the synthesis of sesquiterpenes using radical cyclisation,⁵ herein we describe the 5-*exo-trig* allyl radical cyclisation route to isotwistane **1**, which underwent further cyclisation and rearrangement to a copa and ylanga sesquiterpene skeleton **2**.



Our synthetic sequence, starting from the known⁶ bicyclo-octenone **3** having a bridgehead methoxy group, is depicted in Scheme 1. Although Grignard addition to the bicyclo-octenone is not selective, it is known⁷ that alkylation of the bicyclo-octenone proceeds at low temperature stereoselectively to afford the *endo* alkylated product. Thus, alkylation of the lithium enolate of **3** with methyl iodide gave the ketone **4** in 95% yield having the methyl group in the *endo* position. Further alkylation of the lithium enolate generated from **4** at -78°C with 1,4-dibromo-2-methylbut-2-ene⁸ proceeded stereoselectively and regioselectively to give the *endo* bromide **5**.



Radical cyclisation of **5** under standard conditions² [0.005 M benzene solution of **5** with 1.1 equiv. of tributyltin hydride (TBTH) and 0.1 equiv. of AIBN, reflux, 1–2 h] afforded a



Scheme 1 Reagents and conditions: i, LDA, THF, MeI, -78°C ; ii, LDA, THF, HMPA, 1,4-dibromo-2-methylbut-2-ene, -78°C ; iii, AIBN, TBTH, benzene, reflux, 1–2 h; iv, PTSA, benzene, reflux, 0.5 h

mixture containing the reduced product **6**[†] (5%) and a new compound **7** (71%), whose IR spectrum showed an absorption band at 1740 cm^{-1} . The ^{13}C NMR spectrum of **7** showed a methine carbon at δ 78.2 indicating that **7** is different from the 5-*exo-trig* allyl radical cyclised product **9**. On treatment with toluene-*p*-sulfonic acid (PTSA), compound **7** was quantitatively converted into a new isomer **8**,[†] whose IR spectrum showed the presence of a carbonyl absorption at 1740 cm^{-1} . The off-resonance ^{13}C NMR spectrum of **8** showed the presence of four singlets, four doublets, three triplets and four quartets. A doublet at δ 78.53 clearly showed that the OMe group is attached to a carbon atom bearing a hydrogen. This data clearly established the structure of the cyclised and isomerised products as **7** and **8**, and that the isopropenyl substituent present in **7** is isomerised to the isopropylidene group under acidic conditions to give **8**.

A probable mechanism for the formation of the compounds **7** and **8** is indicated in Scheme 2. As expected, the initial 5-*exo-trig* allyl radical cyclisation gave the radical **11** which underwent a 3-*exo-trig* radical cyclisation onto the carbonyl group resulting in the cyclopropyloxyl radical **12** which



rearranged to give **7**. Formation of a radical adjacent to the methoxy group appears to be the driving force for this rearrangement.[‡]

A number of natural products possess this skeleton, *e.g.* copacamphor **13**,⁹ sinularene **14**¹⁰ and sativene **15**,¹¹ and the above strategy might be contemplated for their total synthesis.

We thank the CSIR for the award of a fellowship to K. K.

Footnotes

[†] All the compounds exhibited spectral data consistent with their structures. *Selected spectral data for 5*: $\nu_{\max}/\text{cm}^{-1}$ 3020, 2920, 1716; δ_{H} (90 MHz, CDCl_3) 6.51 (1 H, m), 6.21 (1 H, dd, J 6.4, 1.8 Hz), 5.73 (1 H, t, J 6.8 Hz), 3.99 (2 H, s), 3.52 (3 H, s), 2.62 (1 H, m), 1.2–2.36 (6 H, m), 1.73 (3 H, s), 1.11 (3 H, s); δ_{C} (22.5 MHz, CDCl_3) 212.4(s), 136.47(d), 134.4(s), 127.6(d), 125.5(d), 84.1(s), 52.1(q), 46.9(s), 40.8(t), 39.9(d), 36.8(t), 26.4(q), 21.8(t), 21.7(t), 14.6(q). *For 8*: $\nu_{\max}/\text{cm}^{-1}$ 3010, 2920, 1740; δ_{H} (200 MHz, CDCl_3) 3.42 (1 H, m), 3.35 (3 H, s), 2.68 (1 H, d, J 1.6 Hz), 2.42 (1 H, br s), 1.2–2.22 (7 H, m), 1.66 (3 H, s), 1.50 (3 H, s), 1.09 (3 H, s); δ_{C} (22.5 MHz, CDCl_3) 218.1(s), 130.7(s), 121.7(s), 78.5(s), 55.4(q), 55.0(s), 53.2(d), 49.5(d), 48.0(d), 41.2(t), 25.2(t), 20.9(t), 19.7(q), 19.7(q), 10.48(q). *For 6* $\nu_{\max}/\text{cm}^{-1}$ 3010, 2915, 1720; δ_{H} (90 MHz, CDCl_3) 6.45 (1 H, m), 6.17 (1 H, dd, J 6.7, 1.7 Hz), 5.12 (1 H, t, J 7 Hz), 3.52 (3 H, s), 2.61 (1 H, m), 1.25–2.18 (6 H, m), 1.73 (3 H, s), 1.59 (3 H, s), 1.08 (3 H, s); δ_{C} (22.5 MHz, CDCl_3) 213.1(s), 136.5(d), 134(s), 127.4(d), 118.7(d), 84.2(s), 52.7(q), 47.8(s), 39.5(d), 36.5(t), 26.2(t), 25.7(t), 21.1(q), 21.0(q), 17.6(q).

[‡] A bicyclooctenone analogous to **5** having a bridgehead methyl group underwent a smooth 5-*exo-trig* allyl radical cyclisation exclusively to give a compound analogous to **9**.

References

- 1 For a recent review on radical cyclisation route to natural product synthesis, see C. P. Jasperse, D. P. Curran and T. L. Fevig, *Chem. Rev.*, 1991, **91**, 1237.
- 2 G. Stork and M. E. Reynolds, *J. Am. Chem. Soc.*, 1988, **110**, 6911.
- 3 N. J. G. Cox, G. Pattenden and S. D. Mills, *Tetrahedron Lett.*, 1989, **30**, 621.
- 4 S. A. Hitchcock and G. Pattenden, *Tetrahedron Lett.*, 1990, **31**, 3641.
- 5 G. S. R. Subba Rao and K. Vijaya Bhaskar, *J. Chem. Soc., Perkin Trans. I*, 1993, 2813.
- 6 D. A. Evans, W. L. Scott and L. K. Truesdale, *Tetrahedron Lett.*, 1972, 121.
- 7 G. Stork and N. H. Baine, *Tetrahedron Lett.*, 1985, **26**, 5927.
- 8 H. Staudinger, O. Muntwyler and O. Kupfer, *Helv. Chim. Acta*, 1922, **5**, 756.
- 9 F. Bohlmann, H. Suding, J. Cuatrecasas, H. Robinson and R. M. King, *Phytochemistry*, 1980, **19**, 2399.
- 10 C. M. Beecham, C. Djerassi, J. S. Finer and J. Clardy, *Tetrahedron Lett.*, 1977, 2395.
- 11 P. de Mayo and R. M. Williams, *J. Am. Chem. Soc.*, 1965, **87**, 3275.

Received, 22nd July 1996; Com. 6/05064F