

ISSN 0300-9246

JOURNAL OF THE CHEMICAL SOCIETY

Dalton Transactions

A journal of inorganic chemistry

1989



ROYAL SOCIETY OF CHEMISTRY

© The Royal Society of Chemistry, 1989. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form, or by any means, electronic, mechanical, photographic, recording, or otherwise, without the prior permission of the publishers.

Printed in Great Britain by
Richard Clay Ltd.,
Bungay, Suffolk

AUTHOR INDEX

- Abad, J. Antonio, 2241
 Abel, Edward W., 701, 711, 885, 985, 2315
 Abid, Khalid Y., 1697
 Adams, Harry, 1831, 2415
 Adatia, Trushar, 5, 1455, 1555
 Adlington, Neil K., 457
 Agarwala, U. C., 1139
 Agh-Atabay, Naz M., 1027
 Ahmed, Etmira, 645
 Ahsan, Mohammad Q., 93
 Aihara, Yukiko, 631
 Aime, Silvio, 1277
 Ainscough, Eric W., 39, 2167
 Akagi, Hisako, 179
 Akita, Munetaka, 1083, 1627
 Akitt, J. W., 1889, 1897
 Akui, Toshifumi, 943
 Al-Juaid, Salih S., 447
 Al-Salim, Najih I., 1697
 Al Obaidi, Najat, 127
 Albano, Vincenzo G., 879, 2343
 Albers, Michael O., 2303
 Albertin, Gabriele, 2353
 Alcock, Nathaniel W., 139, 471, 1589, 2069
 Alibrandi, Giuseppe, 375
 Allen, Christopher W., 2423
 Allen, Frank H., S1
 Allen, Stephen R., 885
 Almaraz, Alejandra E., 1973
 Almeida, Sílvia S. P. R., 2381
 Amadelli, Rossano, 1197
 Amigó, Josep M., 1681
 Anderson, David G., 779, 859
 Anderson, Oren P., 185
 Andersson, Ingegård, 1915
 Annan, Theodore A., 439
 Annibale, Giuliano, 1265
 Antoniutti, Stefano, 2353
 Apreda, Maria-Carmen, 1133, 1249, 1987
 Ara, Irene, 1011
 Arena, Giuseppe, 773
 Arif, Muhammad, 1273
 Armstrong, Elaine M., 295
 Armstrong, Robert S., 1771
 Arnaud-Neu, Françoise, 367
 Arnold, John, 2149
 Arriortua, María, 53
 Arsenault, Gilles J., 761
 Aspinall, Helen C., 623
 Atkinson, Alan, 2371
 Atria, Ana María, 1435
 Attwood, Martin D., 385, 391
 Bailey, Neil A., 1727, 1831, 2415
 Bakar, W. A. Wan Abu, 991
 Baker, Edward N., 39, 2167
 Baker, Mohamad Abu, 1417
 Baker, Paul K., 295, 1321, 1903
 Bal, Wojciech, 2247
 Balázspári, Lajos, 1925
 Balcazar, José Luis, 79
 Baldus, Hans Peter, 977
 Banbery, Hilary J., 1405
 Bandoli, Giuliano, 757
 Bandy, Judith A., 331
 Banerjee, Rupendranath, 1645
 Banister, Arthur J., 1705, 2229
 Bannister, Robin M., 1609
 Barbe, Jean-Michel, 1531
 Barbieri, Renato, 519
 Barker, Geoffrey K., 1321
 Barker, James, 837
 Barnfield, Elizabeth A., 545
 Barron, Andrew R., 1625
 Barthomeuf, Denise, 1361
 Bartocci, Carlo, 1197
 Basak, Arup K., 197
 Bates, Paul A., 581, 985, 1227, 1273, 2315
 Battaglia, Luigi P., 671, 1345
 Bayón, J. Carles, 1579
 Beagley, Brian, 907
 Bear, John L., 93
 Beattie, James K., 1771
 Begley, Michael G., 2443
 Begum, Noor Shahina, 1507
 Belal, Arafa A., 931
 Belforte, Antonio, 1007
 Belletti, Daniele, 1093
 Bellitto, Carlo, 719
 Beltrán-Porter, Aurelio, 53
 Beltrán-Porter, Daniel, 53, 237
 Beltrán, Daniel, 863
 Bencheikh, Abderrahim, 2193
 Bentley, Gary W., 1831
 Bergamini, Paola, 2017, 2367
 Beringhelli, Tiziana, 1143
 Bernhardt, Paul V., 1059
 Berno, Pietro, 551
 Bertolaso, Valerio, 1265
 Best, Stephen P., 581, 1771
 Beydoun, Nohma, 1715
 Bhatia, Paramjit K., 749
 Bhattacharjee, Gobinda P., 1963, 2461
 Bhattacharya, Pabitra K., 567
 Bhattacharyya, Ramgopal, 1963, 2461
 Bhusate, Ravindra, 5
 Bibby, David M., 895
 Bingham, Alistair G., 39, 2167
 Blachnik, Roger, 977
 Blake, Alexander J., 223, 965, 1675
 Blaxill, C. Philip, 2039
 Boeyens, Jan C. A., 2407
 Bombi, G. Giorgio, 1513
 Bonamico, Mario, 719
 Bonner, Francis T., 527
 Bordignon, Emilio, 2353
 Botta, Mauro, 1277
 Bottrell, Martin, 511
 Bowmaker, Graham A., 1639
 Boyar, Esther, 5
 Boyer, Jacqueline, 43
 Bradley, Donald C., 623, 1537
 Braga, Dario, 155, 879, 959, 1455, 1721, 2343
 Braithwaite, Gerard P., 1771
 Brammer, Lee, S1
 Branca, Mario, 1283, 1289
 Brauns, Thomas, 2049
 Braunstein, Pierre, 379, 1477
 Bray, Joanne M., 589
 Brelière, Claire, 43
 Brianese, Nicola, 677
 Briend, Marguerite, 1361
 Brint, Paul, 2253
 Brisdon, Alan K., 313
 Brodie, Andrew M., 39, 2167
 Brodie, Nancy M. J., 1933
 Brown, Paul L., 397
 Brown, Scott S. D., 1227, 2039
 Byrne, Lindsay T., 105
 Cabeza, Javier A., 1093
 Cabral, Joao, 1351
 Cabral, M. Fernanda, 1351
 Calderazzo, Fausto, 1007
 Cameron, James H., 1599
 Cameron, T. Stanley, 701
 Campbell, Nicholas J., 1203
 Camus, Annamaria, 753
 Cano, Félix H., 1133, 1249, 1987
 Carassiti, Vittorio, 1197
 Carini, Clara, 289
 Carlton, Laurence, 2407
 Carmona, Daniel, 1427, 1987
 Carmona, Ernesto, 2003
 Carosiglio, Tommaso, 1957
 Carré, Francis, 43
 Carriedo, Carmen, 745, 2049
 Carriedo, Gabino A., 639
 Carvalho, M. Fernanda N. N., 1209
 Carvill, Andrew, 2435
 Casabó, Jaume, 1381, 1401, 1739
 Casas, José M., 169
 Casas, José S., 1409, 1787
 Casassas, Enric, 569
 Casella, Luigi, 1161, 1979
 Casellato, Umberto, 677, 757, 1993
 Castagno, Fabrizio, 1477
 Castaño, María V., 1409
 Castan, Paule, 403
 Castellano, Eduardo E., 1409
 Castiglioni, Mario, 1477
 Castro, Isabel, 729
 Cattalini, Lucio, 1265
 Cecconi, Franco, 211
 Cederbaum, Lorenz S., 33
 Cesariotti, Edoardo, 161
 Chabanel, Martial, 2193
 Chadha, Raj K., 439
 Chamberlain, Barbara, 465
 Chandley, Peter, 581
 Chatterjee, Chinmay, 645
 Chau, Lai K., 93
 Chavan, Madhay, 93
 Che, Chi-Ming, 2011, 2065
 Cheetham, Anthony K., 1765
 Chen, Jiabi, 635
 Chiesi-Villa, Angiola, 145, 551, 953, 1793, 1957
 Chijimatsu, Tamiko, 2359
 Choy, Jin-Ho, 2335
 Chu, C. K., 1951
 Ciani, Gianfranco, 1143
 Cini, Renzo, 575
 Ciriano, Miguel A., 25, 689
 Claire, Preet P. K., 57
 Clark, David L., 1037
 Clark, Robin J. H., 581, 1689
 Claver, Carmen, 1579
 Clayden, Nigel J., 843
 Clearfield, Abraham, 1617
 Clegg, William, 203, 1127, 1705, 1887, 2229
 Clemenson, Peter I., 61, 67, 1101
 Cockayne, John S., 2049
 Coddington, Jan M., 2223
 Colacio, Enrique, 2323
 Colborn, Robert E., 1799
 Colbran, Stephen B., 2023, 2033
 Cole-Hamilton, David J., 489
 Collison, David, 1, 127, 413
 Compton, Neville A., 1887
 Connelly, Neil G., 283, 683, 689, 745, 1751, 2049
 Contreras, Leopoldo, 2003
 Cooksey, Christopher J., 645
 Cooper, Mervyn K., 1067
 Corain, Benedetto, 1513
 Coronado, Eugenio, 237, 863
 Corradi, Anna Bonamartini, 671, 1345
 Corriu, Robert J. P., 43
 Costes, Jean-Pierre, 1017
 Coston, Timothy P. J., 701, 711
 Coville, Neil J., 2407
 Crennell, Susan J., 1363
 Crespo, Margarita, 1089
 Crook, Joseph R., 465
 Crossley, Neil S., 2139
 Cruz-Garritz, Diana, 419
 Cunningham, Desmond, 267
 Cusumano, Matteo, 375
 D'Alfonso, Giuseppe, 1143
 D'Errico, John J., 1879
 Dahan, Françoise, 1017
 Daniele, Pier G., 1745
 Das, Amitava, 1645
 Dasgupta, Subrata, 1645
 Datta, Dipankar, 115
 Davidson, Jack L., 497, 991, 1027
 Davies, David L., 1799
 Davies, Simon J., 785, 1855, 1865, 2269, 2289
 Davis, Martin J., 223
 De Robertis, Alessandro, 1745
 De Stefano, Concetta, 1745
 de V. Steyn, Margot M., 2303
 Deb, Alok Kumar, 1635
 Debaerdemaeker, Tony, 1681
 Decock-Le Reverend, Brigitte, 1471
 Deeming, Antony J., 5, 2211
 Del Zotto, Alessandro, 607, 1045
 Delafosse, Denise, 1361
 Delfs, Christopher D., 1779
 Delgado, Rita, 133
 Demartin, Francesco, 1109
 Dengel, Andrew C., 1203
 Dens, Annick J. J., 539
 Dent, Andrew J., 1227
 Dessim's, Alessandro, 1283, 1289
 Devore, David D., 303, 797, 1363
 Dhillon, Shivcharan Singh, 1341
 Dhubhghaill, Orla Ni, 1657
 di Bugno, Cristina, 155
 Dillen, Jan L. M., 2199
 Dixon, Kim W., 937
 Dobson, Christopher M., 843
 Dolcetti, Giuliano, 607
 Donald, Caroline E., 527
 Dossett, Stephen J., 1871
 Douglas, Graeme, 497
 Downs, Anthony J., 545
 Drake, Simon R., 243, 505, 1455
 Drew, Michael G. B., 57, 1903

- Driessen, Willem L., 2309
 du Preez, Jan G. H., 253
 Duckworth, Paul A., 1067
 Duer, Melinda J., 2109
 Dukat, Wolfgang, 739
 Dupont, Jaïrton, 1715
 Dyke, Andrew F., 1799
 Dyson, David B., 907
 Eaborn, Colin, 447
 Ebsworth, E. A. V., 223
 Edelmann, Frank T., 1815
 Edwards, Anthony J., 127, 1393
 Edwards, Catherine J., 1203
 Edwards, Peter G., 1253
 El-Hendawy, Ahmed M., 901
 El-Kheil, Mohamed N. A., 447
 Elders, Julie M., 1889, 1897
 Ellis, S. Richard, 413
 Elsevier, Cornelis J., 317
 Emmerich, Christiane, 797
 Emsley, John, 1273
 Engelhardt, Lutz M., 105, 595
 Englert, Ullrich, 1787
 Ercolani, Claudio, 193
 Erdbrügger, Cristina Freire, 73
 Ernsting, Jan Meine, 317
 Errington, R. John, 1887
 Escrivé, Lluís, 1381, 1739
 Escrivá, Emilio, 863
 Espinet, Pablo, 1583
 Esteban, Pedro, 1579
 Etches, Sheena J., 2281
 Etienne, Michael, 2187
 Ettorre, Renato, 927
 Evans, David G., 985, 2315
 Fabbrizzi, Luigi, 229
 Fadnis, Anand G., 1237
 Fanizzi, Francesco P., 947, 1689
 Fares, Vincenzo, 719
 Faridoon, 383, 1657
 Farina, América, 849
 Farkas, Etelka, 1053, 2247
 Farrar, David H., 557
 Farrugia, Louis J., 931, 1375
 Farwaha, Rajeev, 761
 Faus, Juan, 729, 1681
 Fava, Giovanna Gasparri, 361
 Favarato, Mosé, 1513
 Felix, Mark S. B., 5
 Fenton, David E., 1727
 Ferguson, George, 85, 383, 915,
 1657, 2183
 Fern, Adrian, 843
 Fernández, Eduardo, 2127
 Fernández, María J., 683, 689,
 1249, 2073
 Ferrari, Marisa Belicchi, 361
 Ferrer, Montserrat, 379
 Ferretti, Valeria, 1265
 Ferry, Joseph, 767
 Fielding, Lee, 873
 Fields, Roy, 907
 Figgis, Brian N., 1779
 Flanagan, John, 1651
 Florencio, Feliciana, 79
 Floriani, Carlo, 33, 145, 551, 953,
 1793, 1957
 Flower, Kevin R., 1903
 Foces-Foces, Concepción, 1249
 Foces-Foces, Concha, 1133, 1987
 Folgado, José V., 53, 237
 Fontaine, Xavier L. R., 383,
 1465, 1657, 1889, 1897
 Fornies, Juan, 169, 1011, 1583
 Forsyth, George, 1903
 Fortuño, Consuelo, 1583
 Fowler, Patrick W., 2253
 Fox, David, 151
 Fraile, M. Nieves, 73
 Francois, M. Angélica, 1105
 Frankland, Julie C., 2039
 Franzoni, Doriano, 247
 Fraser, Iain, 2183
 Fraústo da Silva, João J. R., 133
 Freni, Maria, 1143
 Fülöp, Vilmos, 2331
 Fuertes, Amparo, 237, 863
 Fukui-Yasuba, Yuko, 815
 Fumagalli, Alessandro, 879, 1109,
 2343
 Furphy, Brenda M., 595, 2217
 Gallardo, Sergio, 1105
 Gallazzi, Maria C., 1819
 Gallop, Mark A., 481
 Gamasa, M. Pilar, 1499
 Gambino, Olímpia, 1277
 García, José R., 829, 1825
 Garcíá-Blanco, Severino, 79
 Gardini, Marcello, 193
 Gargano, Michele, 921
 Garland, María Teresa, 1435
 Garlaschelli, Luigi, 879, 1109
 Garner, C. David, 413
 Gasyna, Zbigniew, 2397
 Gavioli, Giovanna Battistuzzi,
 1345
 Geloso, Corrado, 1759
 Gentil, Luis A., 1973
 Gerloch, Malcolm, 2109
 Ghedini, Mauro, 1075, 1217
 Ghilardi, Carlo A., 211
 Ghosh, Nabanita, 1963, 2461
 Ghosh, Saktiprovad, 555
 Gianguzza, Antonio, 773
 Giannetto, Antonino, 375
 Gibson, Dan, 1911
 Gill, Daljit S., 1321
 Gilli, Gastone, 1265
 Gimeno, José, 1499
 Gimeno, M^a. Concepción, 1883
 Giorgi, Gianluca, 575
 Girard, Anne-Marie, 101
 Gisselbrecht, Jean-Paul, 367
 Giudice, M. Teresa Lo, 519
 Gleria, Mario, 757
 Glidewell, Christopher, 2061
 Gobetto, Roberto, 1277
 Goedken, Virgil L., 193
 Gogolashvili, Edward L., 1293
 Goh, Lai Yoong, 431, 1951
 González-Duarte, Pilar, 13
 González, Angeles Sánchez, 927
 González, Eliana, 829, 1825
 Goodgame, David M. L., 1439
 Gorrell, Ian B., 2229
 Goswami, Sreebrata, 1635
 Gotoh, Mikio, 1613, 2089
 Gouteron, Jacqueline, 1117
 Gradella, Cecilia, 1721
 Graham, Stephen, 1599
 Grassi, Antonio, 757
 Grassi, Maria, 615
 Grassi, Mariella, 161, 1819
 Graziani, Rodolfo, 677, 1993
 Greatrex, Robert, 385, 391
 Greaves, Martin, 1697
 Green, Jennifer C., 1037, 2139
 Green, Malcolm L. H., 331, 2133
 Green, Michael, 511, 1321, 1333
 Greenwood, Norman N., 385,
 391, 1465
 Grepioni, Fabrizia, 879, 1721,
 2343
 Griffith, William P., 901, 1203,
 1651
 Griffiths, Trevor R., 325
 Gross, Maurice, 367
 Gutiérrez-Puebla, Enrique, 2003
 Guastini, Carlo, 145, 551, 953,
 1793, 1957
 Gudat, Dietrich, 693
 Guerchais, Jacques E., 1799, 2187
 Guilard, Roger, 1531
 Guilhem, Jean, 1449
 Gukathasan, Ravi R., 557
 Gullotti, Michele, 1161, 1979
 Gustems, L'ó Luis, 569
 Gutiérrez-Zorrilla, Juan M.,
 2323
 Gutierrez-Jugo, Juan F., 2241
 Gutowsky, Herbert S., 275
 Györ, Miklós, 1925
 Hämäläinen, Reijo, 2309
 Haanstra, Willem G., 2309
 Haga, Masa-aki, 371
 Hague, David N., 1171
 Hambley, Trevor W., 1059, 1067,
 1951, 2205
 Hamor, Thomas A., 1405, 1697
 Hanabusa, Kenji, 1543
 Hansford, Michael I., 1705
 Hardcastle, Kenneth I., 2211
 Harjula, Risto, 101
 Harman, Mary E., 295
 Harris, Robin K., 809
 Harrison, Philip G., 2443
 Harrowfield, Jack McB., 595,
 2217
 Hart, Ian J., 2261, 2281
 Hattingh, Johannes T. Z., 1565
 Hauptman, Zdenek V., 1705
 Hay, Robert W., 85, 197, 2183
 Hayakawa, Tadao, 1543
 Hayashi, Mariko, 1947
 Hayashi, Yoshihito, 1519
 Healy, Peter C., 971, 1639
 Hearne, John A., 2371
 Helps, Ian M., 915
 Henderson, Richard A., 425
 Henderson, Sandra J. B., 1363
 Henderson, William, 259
 Henkel, Thomas, 2173
 Henrick, Kim, 1067, 1455
 Herbossa, Gabriel Garcia, 2049
 Hernández, Sergio, 849
 Hettrich, Ralph, 745
 Hidai, Masanobu, 1241
 Higgins, Kevin M., 701
 Higgins, Paula, 2435
 Higgins, Simon J., 1527
 Higgins, Tim, 267
 Hill, Anthony F., 1871, 2261
 Hill, M. N. Stuart, 203
 Hills, Adrian, 419, 1417
 Himeno, Sadayuki, 1491
 Hitchcock, Peter B., 447
 Hong, Seung-Tae, 2335
 Hope, Eric G., 313
 Hori, Toshitaka, 1491
 Hossain, M. Altaf, 2347
 Hough, Edward, 2155
 Hounslow, Andrea M., 233
 House, Donald A., 853
 Howard, Judith A. K., 303, 797,
 1363, 1855, 1871, 2269, 2289
 Howarth, Laura, 1385
 Howarth, Oliver W., 1915
 Hudson, Robert H. E., 2235
 Hughes, David L., 419, 1413,
 1417, 2389, 2453
 Hughes, Martin N., 527, 533
 Humanes, Madalena, 645
 Hummel, Hans-Ulrich, 627
 Hundal, Maninder Singh, 1341
 Hunt, Paul A., 1127
 Hunter, Geoffrey, 2115
 Huq, Rokeya, 1933
- Hursthouse, Michael B., 61, 67,
 295, 581, 623, 985, 1101, 1227,
 1273, 1537, 2061, 2149, 2315,
 2347
 Hussain, Bilquis, 623, 2149
 Hyde, Timothy L., 965
 Hynes, Michael J., 767
 Ibáñez, Fernando, 1105
 Ibrahim, Ali, 2347
 Ichiba, Sumio, 631
 Imperatori, Patrizia, 719
 Institórisz, László, 1925
 Intini, Francesca P., 947
 Iqbal, Javed, 461
 Irele, Patricia T., 2023, 2033
 Ishida, Katsuaki, 1241
 Ishiguro, Shin-ichi, 655
 Ishimori, Tomitaro, 2083
 Isobe, Kiyoshi, 1519
 Itoh, Hiroshi, 1543
 Itose, Shoichi, 2083
 Jackson, Graham E., 2429
 Jackson, Logan A., 2423
 Jacobsen, Geraldine E., 105
 Jalón, Félix A., 79
 Jarauta, M. Pilar, 1987
 Jeannin, Susanne, 1117
 Jeannin, Yves, 1117
 Jeffery, John C., 303, 1845, 2381
 Jenkins, Julian J., 453
 Jimenez-Tenorio, Manuel, 2389,
 2453
 Jørgensen, Karl Anker, 2099,
 2229
 Johnson, Brian F. G., 243, 481,
 505, 1421, 1455, 2023, 2033
 Johnson, Neil P., 403
 Johnson, Owen, 821
 Jolliffe, Jennifer M., 2179
 Jones, Christopher J., 127, 1393,
 1405
 Jones, Peter G., 73, 639, 2127,
 2241
 Jubb, Jayne, 1631
 Judd, Robert J., 2205
 Julve, Miguel, 729, 1681
 Kabir, Shariff E., 5
 Kadish, Karl M., 93, 1531
 Kafarski, Paweł, 1053
 Kahn, Olivier, 1117
 Kaitner, Branko, 1657
 Kaizaki, Sumio, 1947
 Kalil, Mutasim I., 2443
 Kamiusuki, Toshiro, 2077
 Kannadaguili, Nagesha, 1003
 Karaman, Rafik, 1911
 Katsoulis, Dimitris E., 1483
 Kawaji, Tetsuro, 943
 Kaye, Perry T., 2033
 Kelly, Mark J., 2429
 Kelly, Paul F., 2179
 Kemmitt, Raymond D. W., 259,
 345
 Kemp, Terence J., 1237, 2017
 Kendrick, Aidan, 1311
 Kennard, Olga S.
 Kennedy, John D., 383, 1465,
 1657
 Kepert, David L., 1639
 Khalil, Mutasim I., 2443
 Khan, M. Ishaque, 1139
 Khandelwal, Bandana, 885
 Khattar, Rajesh, 1421
 Kida, Sigeo, 179, 943, 1613, 2077,
 2083, 2089, 2095
 Kikabhai, Thakor, 2443
 Kilullen, Neil, 2115
 Kildea, John D., 1639
 Kilner, Melvyn, 837

- Kimura, Masaru, 357
 Kinoshita, Isamu, 815
 Kinoshita, Setsuko, 2457
 Kirkpatrick, R. James, 275
 Kirschenbaum, Louis J., 217
 Kiss, Tamás, 1053, 2247
 Kitaura, Eiji, 2077
 Kitchen, Gillian F. M., 1227
 Knight, Christopher T. G., 275
 Knights, Clive F., 2371
 Knox, Selby A. R., 1799, 1807
 Kobayashi, Nagao, 2397
 Kodera, Masahito, 943
 Koe, Julian R., 985, 2315
 Kohzuma, Takamitsu, 1613, 2089
 Koikawa, Masayuki, 1613, 2077, 2089
 Koley, Aditya P., 555
 Kondoh, Atsuo, 1083, 1627
 Korp, James D., 93
 Korybut-Daszkiewicz, Bohdan, 85
 Koyama, Toshiki, 1543
 Kozlowski, Henryk, 173, 1053, 1283, 1471, 2247
 Kpoton, Agapit, 43
 Krijnen, Bert, 317
 Kruger, Gert J., 1565
 Kucharski, Edward S., 1779
 Kundu, Amit K., 1889, 1897
 Kunwar, Ajit C., 275
 Kupryszewski, Gotfryd, 173
 L'Helgouach, Hervé, 1435
 Laguna, Antonio, 73, 1883, 2127
 Laguna, Mariano, 73, 1883
 Lahoz, Fernando J., 25, 1075, 2023, 2073
 Lai, Ting-Fong, 2011
 Lambert, Ronald J., 2061
 Lammek, Bernard, 173
 Landázuri, Cayetana, 1093
 Lanfranchi, Maurizio, 1499, 1689
 Lanfredi, Anna Maria Manotti, 753, 1217
 Larkworthy, Leslie F., 1631
 Laschi, Franco, 575
 Lastra, Elena, 1499
 Latorre, Julio, 729
 Laurent, Jean-Pierre, 1017
 Lautner, Jürgen, 639, 2127
 Lavery, Aidan J., 965
 Lawrence, Geoffrey A., 1059
 Lay, Peter A., 2205
 Lee, Seunghee, 145
 Lee, Wai-On, 2011
 Lehn, Jean-Marie, 1449
 Lehto, Jukka, 101
 Lei, Guixin, 635
 Leigh, G. Jeffery, 425, 1413, 1417, 2389, 2453
 Lejczak, Barbara, 1053
 Leonard, Gordon A., 1631
 Leoni, Piero, 155, 959
 Leporati, Enrico, 383, 1299
 Leung, Wing-Por, 105
 Levason, William, 313, 453
 Lewis, Jack, 243, 481, 505, 1421, 1455, 2023, 2033
 Lickiss, Paul D., 447
 Lifsey, Robert S., 93
 Lightfoot, Philip, 1765
 Liman, Ferid, 1471
 Lin, Guangda, 635
 Lincoln, Stephen F., 233
 Lindsell, W. Edward, 991
 Linford, Lorna, 1565
 Livera, Cynara, 173
 Llavona, Ricardo, 829, 1825
 Lloret, Francesc, 729
 Lobana, Tarlok S., 749, 2339
 Lockhart, Joyce C., 203
 Lombard, Anthoine, 1565
 Longeri, Marcello, 1217
 Loonat, Mohamed S., 2407
 Lopez-Poveda, Manuel, 489
 Lotz, Simon, 2199
 Loynes, Andrew C., 683, 689
 Mabbs, Frank E., 127
 Mabrouk, Hassan E., 1759
 Macias, Alejandro, 1409, 1787
 MacLachlan, Dugald J., 895
 Macpherson, Kirsty A., 1807
 Maeda, Yonezo, 943
 Mahmoud, Mouayed M., 837
 Maitlis, Peter M., 2415
 Maiya, G. Bhaskar, 1531
 Maldotti, Andrea, 1197
 Malito, John, 1933
 Mallah, Talal, 1117
 Manassero, Mario, 1109
 Mangani, Stefano, 1979
 Manjula, Vadaparti, 567
 Mann, Brian E., 889, 1831
 Manners, Ian, 283
 Manohar, Hattikudur, 1507
 Manoharan, Periakaruppan T., 555
 Marais, Charles F., 1565
 Markiewicz, Stan, 1933
 Marolleau, Isabelle, 367
 Marsich, Nazario, 753
 Martinez-Ripoll, Martin, 2323
 Martinez, Benjamin, 1381
 Martinengo, Secondo, 2343
 Martinez, Manuel, 1669
 Marzilli, Patricia A., 1941
 Masciocchi, Norberto, 1109
 Masuda, Etsuko, 1543
 Matsumoto, Keiji, 815, 1519
 Matsumoto, Naohide, 179, 943, 2077, 2083, 2089, 2095
 Matsumura-Inoue, Takeko, 371
 Matthes, Karen E., 915
 Mawby, Roger J., 465, 589
 Mays, Martin J., 1555
 Mazid, Muhammed, 2347
 Mazumdar, Shyamalava, 1003
 Mazzanti, Marinella, 953, 1793
 McAlees, Alan J., 761
 McArdle, Patrick, 267, 767
 McAuliffe, Charles A., 907
 McCann, G. Malachy, 2435
 McCarthy, Simon, 581
 McCleverty, Jon A., 127, 1405
 McCrindle, Robert, 761
 McCullough, Kevin J., 991
 McDonnell, Martin B., 203
 McGarvey, Bruce R., 439
 McGrady, G. Sean, 545
 McKee, Vickie, 19
 McKenna, Peter, 345
 McKillop, Kieran P., 1351
 McPartlin, Mary, 5, 1067, 1455, 1555
 McQuillan, Ferida, 1405
 McWhinnie, William R., 1697
 Mead, Kevin A., 1799
 Meah, M. Nafees, 2211
 Medhi, Okhil K., 1003
 Meille, Stefano V., 615
 Menabue, Ledi, 1345
 Menanteau, Pamela del C., 2415
 Mendiola, Maria Antonia, 1127
 Mesa, José L., 53
 Meske, Hermann, 627
 Mezzetti, Antonio, 607, 1045
 Micera, Giovanni, 1283, 1289
 Michael, David J., 1689
 Michelini, Rino A., 1149
 Micheloni, Mauro, 133
 Midollini, Stefano, 211
 Miki, Shigeru, 2359
 Miller, J. David, 457
 Mingos, D. Michael P., 539
 Minniti, Domenico, 375
 Miravitles, Carles, 1381, 1401
 Miravitles, Carlos, 863, 1739
 Mitra, Samaresh, 1003
 Modrego, Javier, 683, 1249
 Molins, Elies, 1381, 1401, 1739
 Molla, M. Elias, 447
 Mollar, Miquel, 729
 Molloy, Kieran C., 267, 289
 Moneti, Simonetta, 211
 Monge, Angeles, 2003
 Montoneri, Enzo, 757, 1819
 Moore, Laila S., 539
 Moore, Peter, 873
 Moret, Massimo, 1143
 Moreton, Anthony D., 1171
 Morgenstern-Badarau, Irène, 729
 Moro-oka, Yoshihiko, 1083, 1627
 Morris, Michael J., 1555
 Morton, David, 489
 Moss, Michael A. J., 1393
 Motlevalli, Majid, 2061
 Moussa, Mahmoud N., 901
 Muir, Kenneth W., 497, 991
 Muller, Guillermo, 1669
 Mura, Pasquale, 2367
 Murray, Martin, 821
 Musco, Alfredo, 615
 Musgrave, Rupert J., 2269
 Musmeci, M. Teresa, 519
 Nagy, Attila, 2139
 Nagy, László, 655
 Nakai, Hiroshi, 815
 Nakamura, Yukio, 1519
 Nasir, Bashir A., 1871
 Nassimbeni, Luigi R., 119
 Natile, Giovanna, 947, 1689
 Naumann, Dieter, 739
 Navarro, Rafael, 169
 Neaves, Bryan D., 127
 Nelson, Jane, 1351
 Nelson, S. Martin, 1351
 Nelson, William W. J. H., 1455
 Nestor, Karl, 1465
 Neve, Francesco, 1075, 1217
 Newman, Paul A., 331
 Nicholson, David G., 2155
 Niecke, Edgar, 693
 Nieger, Martin, 693
 Nishida, Suzuko, 357
 Niven, Margaret L., 119
 Nöth, Heinrich, 1311
 Noltmeyer, Mathias, 1815
 Norman, Nicholas C., 1887
 O'Mahoney, Caroline A., 1179
 Odiaka, Timothy I., 561
 Ogden, J. Steven, 313
 Ogden, Mark I., 2217
 Ohtaki, Hitoshi, 655
 Ohyoshi, Akira, 943
 Okawa, Hisashi, 179, 943, 1613, 2077, 2083, 2089, 2095
 Okuda, Tsutomu, 631
 Olabe, Jose A., 1973
 Oldfield, Eric, 275
 Ooi, Shun'ichiro, 815
 Organ, Gregory J., 1067
 Orlandini, Annabella, 211
 Oro, Luis A., 25, 683, 689, 1093, 1133, 1249, 1427, 1987, 2073
 Orpen, A. Guy, 511, 745, 1227, 1321, 1333, 1751, 1799, 1807, 2049, S1
 Orrell, Keith G., 701, 711
 Ortiz-Avila, C. Yolanda, 1617
 Osella, Domenico, 1277
 Oshio, Hiroki, 943, 2077
 Ossola, Franco, 677
 Otero, Antonio, 79
 Otte, Ronald, 1565
 Ozarowski, Andrzej, 439
 Ozutsumi, Kazuhiko, 655
 Padel, Lilyane, 863
 Pahor, Nevenka Bresciani, 607, 1045, 1941
 Pakkanen, Tapani A., 2449
 Panda, Rama K., 217
 Paneque-Sosa, Marguerita, 489
 Paoletti, Piero, 133
 Papasergio, Rocco I., 105
 Pappalardo, Giuseppe C., 757
 Parekh, Anjali, 1067
 Parish, R. V., 539, 907
 Parker, David, 915
 Parkin, Ivan P., 1179
 Pascard, Claudine, 1449
 Pasquali, Marco, 155, 959
 Pasquinelli, Gianfranco, 601
 Patrick, Jennifer M., 595
 Patrizio, Salvatore, 719
 Pauer, Frank, 2173
 Pérez, Pedro L., 1427
 Peacock, Robert D., 931
 Peintler, Gábor, 1925
 Pelizzi, Corrado, 247, 289, 361, 671
 Pelizzi, Giancarlo, 247, 289
 Pellerito, Lorenzo, 773
 Pelosi, Giorgio, 671
 Peltre, Marie-Jeanne, 1361
 Peña, Octavio, 1435
 Pennesi, Giovanna, 193
 Pennington, Mark, 471
 Perez-Torreto, Jesus J., 25
 Pergola, Roberto Della, 879, 1109
 Perotti, Angelo, 197
 Perry, Richard J., 453
 Pettersson, Lage, 1915
 Pettit, Leslie D., 173, 1471
 Pfesser, Michel, 1715
 Phillips, Nicolas J., 325
 Pickett, Christopher J., 425
 Pickup, Barry T., 889
 Pilotti, Massimino U., 303, 1855, 2289
 Pinciroli, Fiorella, 1161
 Pinillos, M. Teresa, 1133, 1987
 Pintar, Alessandro, 1161, 1979
 Plasencia, María M., 1409
 Platt, Andrew W. G., 139, 1193, 2069
 Plenio, Herbert, 1815
 Poë, Anthony J., 1933, 2235
 Poirier, Monique, 43
 Pombeiro, Armando J. L., 1209, 2381
 Pons, Josefina, 1401
 Pontellini, Roberto, 615
 Poojary, M. Damodara, 1507
 Pope, Michael T., 1483
 Porchia, Marina, 677
 Povey, David C., 1631, 2339
 Powell, Harold R., 5, 1537
 Powell, Richard D., 1651
 Prasad, Lakanaju
 Seeramachandra, 555
 Prati, Laura, 161
 Predieri, Giovanni, 247

- Preston, Peter N., 497
 Pringle, Paul G., 139, 1193, 2017,
 2069
 Pritchard, Robin G., 907
 Proserpio, Davide M., 229
 Prouse, Lesley J. S., 259, 345
 Puchalska, Danuta, 2193
 Puddephatt, Richard J., 2121
 Pujari, Mahesh P., 85, 197
 Purohit, Sanchita, 555
 Purrello, Roberto, 773
 Pursiainen, Jouni, 2449
 Pyburn, Simon, 173, 1471
 Quill, Kieran, 267
 Radomska, Barbara, 173
 Raithby, Paul R., 1421, 1555,
 2023, 2033
 Ramdas, Vijayalakshmi, 2339
 Randaccio, Lucio, 1941
 Randle, Neil P., 2211
 Rankin, David W. H., 223, 545,
 779, 859
 Raston, Colin L., 105
 Raubenheimer, Helgard G., 1565
 Ravasio, Nicoletta, 921
 Real, Julio, 1579
 Redfern, Catherine M., 1037
 Reedijk, Jan, 2309
 Reid, Gillian, 1675
 Rendle, Melvyn C., 1807
 Reynolds, Philip A., 1779
 Richards, Raymond L., 419, 2381
 Riera, Victor, 639
 Rigo, Pierluigi, 607, 1045
 Rius, Jordi, 1381
 Rizzarelli, Enrico, 773
 Robb, John, 931
 Robertson, Heather E., 859
 Rockenbauer, Antal, 1925
 Rodríguez-Argüelles, María C.,
 1787
 Rodríguez, Julio, 829, 1825
 Rodríguez, M. L., 639
 Rodríguez, Isabel, 1381
 Rodriguez, Maria J., 2073
 Roe, S. Mark, 1589
 Roesky, Herbert W., 1815, 2173
 Rojo, Teófilo, 53, 237
 Román, Enrique, 849
 Román, Pascual, 2323
 Romerosa, Antonio, 2323
 Ros, Renzo, 1149
 Rosé, Jacky, 1477
 Rosi, Marzio, 33
 Rossell, Oriol, 379
 Rosset, Jean-Marie, 953
 Rossetto, Gilberto, 677
 Rossi, Gentilina, 193
 Rossi, Michele, 921
 Roué, Jean, 1799
 Roy, Sujit, 2121
 Royo, Gérard, 43
 Royo, Pascual, 79
 Ruhl, Barbara L., 85
 Ruisi, Giuseppe, 519
 Ruiz-Romero, M. Elena, 2127
 Ruiz, Aurora, 1579
 Ruiz, Jose, 2323
 Ruiz, Miguel A., 1845
 Russell, David R., 259, 345
 Ryan, Helen, 2435
 Rzepa, Henry S., 1609
 Sánchez, Agustin, 1787
 Sánchez, Luis J., 2003
 Sabat, Michal, 575
 Sabatini, Antonio, 575
 Sabatino, Piera, 155, 959
 Sakamoto, Masatomi, 2083
 Sakurai, Tosio, 2359
 Saladini, Monica, 1345
 Sales, Joaquim, 1089
 Sales, Keith D., 623
 Salsini, Liana, 155
 Salter, Ian D., 1227, 2039
 Sammartano, Silvio, 1745
 Sampson, Clifford N., 2235
 Sandhu, Maninderjeet K., 2339
 Sandhu, Sarjit Singh, 1341
 Sangchakr, Benchang, 2253
 Sansoni, Mirella, 1109
 Santos, José G., 1105
 Sanz, Marta, 1739
 Sappa, Enrico, 601, 1477
 Satō, Gen P., 371
 Satoh, Keiichi, 1259
 Saunders, David R., 511
 Sawada, Koyoshi, 1259
 Scapacci, Giancarlo, 211
 Schauer, Cynthia K., 185
 Schaverien, Colin J., 1333
 Schiøtt, Birgit, 2099
 Schröder, Martin, 965, 1675
 Schwing-Weil, Marie-José, 367
 Scott, Ian D., 1375
 Scott, John D., 2121
 Seco, Miquel, 379
 Segre, Annalaura, 2367
 Sekhar, Vasu C., 1933
 Sgamellotti, Antonio, 33
 Sharma, G. Tomba, 115
 Sharp, David W. A., 461
 Sharp, Kenneth G., 1879
 Shaw, Bernard L., 1527
 Sheldrick, George M., 2173
 Sherman, Darren J., 539
 Sherwood, Paul, 1845, 1871
 Shiels, Anthony, 2435
 Shikholeslami, Afshine, 1361
 Shimizu, Kunio, 371
 Shirai, Hirofusa, 1543
 Shiro, Motoo, 815
 Short, Richard L., 61, 67, 1101
 Shtyrlin, Valery G., 1293
 Siatecki, Zdzislaw, 1053
 Siddiqi, Khwaja S., 1401
 Siegel, Andrew, 2235
 Šík, Vladimír, 701, 711, 985,
 2039, 2315
 Silva, M. Fátima C. G., 2381
 Silvestri, Arturo, 519
 Singleton, Eric, 2303
 Sinn, Ekk, 1187
 Sironi, Angelo, 615, 1143, 1455
 Skelton, Brian W., 105, 595, 971,
 1639, 2217
 Smith, Anthony K., 889
 Smith, Clyde A., 2167
 Smith, David O., 1375
 Smith, Gallienus W., 2339
 Smith, J. David, 447
 Sneddon, David W., 761
 Sola, Marco, 1345
 Solans, Xavier, 729, 1089
 Sood, Geeta, 1341
 Sordo, José, 1409, 1787
 Sosa, Plinio, 419
 Sostero, Silvana, 2017, 2367
 Spalding, Trevor R., 383, 1657
 Spaniol, Thomas P., 1871
 Spankie, Sally A. R., 497
 Speier, Gábor, 2331
 Spicer, Mark D., 453
 Spodine, Evgenia, 1435
 Stalke, Dietmar, 2173
 Stalla, Beate, 1311
 Stam, Casper H., 317
 Stedman, Geoffrey, 533
 Steel, Andrew T., 2443
 Stella, Sergio, 1957
 Stillman, Martin J., 2397
 Stone, F. Gordon A., 303, 785,
 797, 1363, 1845, 1855, 1865,
 1871, 2261, 2269, 2281, 2289
 Storch, Wolfgang, 1311
 Straughan, Brian P., 1127
 Stringer, Graham, 2139
 Suárez, María I., 1787
 Suardi, Edoardo, 1979
 Suga, Masanobu, 661
 Sullivan, Alice C., 2347
 Sunley, Glenn J., 2415
 Suzuki, Toshio, 1259
 Szabó-Plánka, Terézia, 1925
 Szóke, József, 2247
 Taha, Fatma I., 901
 Tahir, Tahir A., 457
 Takahashi, Yokino, 1259
 Tamada, Osamu, 1491
 Tamburini, Sergio, 1993
 Tanaka, Koji, 1547
 Tanaka, Nobuhiko, 631
 Tanaka, Toshio, 1547
 Tandon, Santokh S., 19
 Tang, Kaluo, 2297
 Tang, Wai-Tong, 2011
 Tang, Youqi, 2297
 Tarantelli, Francesco, 33
 Tarasconi, Pieralberto, 247, 289,
 361, 671
 Tattershall, Bruce W., 977
 Tauler, Romà, 569
 Taylor, Michael J., 2223
 Taylor, Michael W., 119
 Taylor, Robin, S1
 Teixidor, Francesc, 1381, 1401,
 1739
 Tejel, Cristina, 1133
 Thanyasiri, Thitinant, 1187
 Thompson, Arthur R., 275
 Thornton-Pett, Mark, 1465
 Tiekkink, Edward R. T., 233, 749
 Tiripicchio-Camellini, Marisa,
 25, 601, 921, 1093, 1427, 1477
 Tiripicchio, Antonio, 25, 601,
 753, 921, 1075, 1093, 1217,
 1427, 1477, 1499, 1689
 Tobe, Martin L., 645, 853, 1265
 Todd, Martin, 203
 Tomás, Milagros, 169, 1011,
 1583
 Tomita, Munetoshi, 179
 Torrens, Hugo, 419
 Traverso, Orazio, 2017, 2367
 Trocha-Grimshaw, Jadwiga,
 1351
 Tsubayama, Kaoru, 2359
 Tsubayama, Sei, 2359
 Tuck, Dennis G., 439, 1759
 Tucker, Andrew J., 1887
 Tucker, B. Joly, 1631
 Turpeinen, Urho, 2309
 Twiss, Paul, 105
 Uezumi, Satoshi, 1547
 Umakoshi, Keisuke, 815
 Underhill, Allan E., 61, 67, 1101
 Usón, Rafael, 73, 169, 1011,
 1583, 1883, 2127
 Utuk, Imelda D., 489
 Vacca, Alberto, 575
 Valle, Giovanni C., 927, 1513
 van Brecht, Bernardus J. A. M.,
 253
 van der Zeijden, Adolphus A. H.,
 317
 van Dyk, Marthie N., 2199
 van Koten, Gerard, 317
 Varga-Fábián, Mária, 1925
 Vargas, Maria D., 1455
 Vasudevan, Asha K., 2155
 Vaz, Maria Cândida T. A., 133
 Vettori, Umberto, 927
 Vicente, Jose, 2241
 Viganò, Roberto, 1161
 Vigato, Pietro Alessandro, 1993
 Viguri, Fernando, 25
 Villarroya, B. Eva, 689
 Villeneuve, Gerard, 53
 Vitali, Francesca, 247
 Vives, Josep, 13
 Wait, Simon T., 1705
 Wakita, Hisanobu, 2457
 Walker, David G., 1413, 2389
 Walker, Nigel P. C., 623
 Wallwork, Stephen C., 837
 Wang, Yongxin, 635
 Waterfield, Philip C., 289
 Watson, David G., S1
 Welch, Alan J., 223, 1321, 1583
 Weldon, Vincent J., 2253
 Wells, Cecil F., 151
 Went, Michael J., 1375
 West, Anthony P., 1651
 West, Antony A., 1697
 White, Allan H., 105, 595, 971,
 1639, 2217
 White, Jonathan M., 745
 Whiteley, Mark W., 1751
 Wilkes, Philip, 809
 Wilkinson, Geoffrey, 2149
 Willey, Gerald R., 57
 Williams, David J., 1179, 1439
 Williams, Glyn, 645
 Williams, Ian D., 511, 1321
 Williams, Mark G., 1727
 Wills, A. Richard, 1253
 Wilner, Faye R., 2217
 Wimbledon, Peter E., 533
 Wimmer, Franz L., 403
 Wimmer, Smita, 403
 Winfield, John M., 461, 937
 Winpenny, Richard E. P., 1439
 Winter, Derek J., 1727
 Winter, Mark J., 1887
 Witt, Michael, 2173
 Wong, Kwok-Yin, 2065
 Wong, Luet-Lok, 1385, 2133
 Wong, Richard C. S., 1951
 Wong, Wing-Tak, 2011
 Wood, Paul T., 809
 Woodward, Gary, 821
 Woodward, Peter, 1751
 Woollins, J. Derek, 809, 859,
 1179, 2179
 Wright, Anthony H., 481
 Xu, Quan Yun Y., 1531
 Yamada, Koji, 631
 Yamamoto, Yasuhiro, 2161
 Yamase, Toshihiro, 661
 Yamashita, Masahiro, 2457
 Yamazaki, Hiroshi, 2161
 Yang, Jianping, 2297
 Yang, Qingchuan, 2297
 Yano, Shigenobu, 1241
 Yashiro, Morio, 1241
 Yin, Jianguo, 635
 Yoshikawa, Sadao, 1241
 Young, J. Colin, 43
 Zakharov, Alexey V., 1293
 Zanella, Pierino, 677
 Zanello, Piero, 1161, 1979
 Zangrandi, Ennio, 1941
 Zatta, Paolo, 1513
 Zerbiniati, Orfeo, 1745
 Zhong, Zhuang Jin, 2095
 Ziessel, Raymond, 1449
 Zora, Jalal A., 447

SUBJECT INDEX, 1989

ACETAMIDATE

Syntheses, electrochemistry, and spectroscopy of dirhodium(II) tetra-acetamidate and tetrakis(trifluoroacetamide) complexes with axial Group 15 substituents. The *X*-ray crystal structures of $[\text{Rh}_2(\text{CH}_3\text{CONH})_4(\text{AsPh}_3)_2]$ and $[\text{Rh}_2(\text{CH}_3\text{CONH})_{4-n}(\text{CH}_3\text{CO}_2)_n(\text{MPh}_3)_2]$, M = As or Sb, $n \approx 1$, 581–8

ACETATE

Syntheses, electrochemistry, and spectroscopy of dirhodium(II) tetra-acetamidate and tetrakis(trifluoroacetamide) complexes with axial Group 15 substituents. The *X*-ray crystal structures of $[\text{Rh}_2(\text{CH}_3\text{CONH})_4(\text{AsPh}_3)_2]$ and $[\text{Rh}_2(\text{CH}_3\text{CONH})_{4-n}(\text{CH}_3\text{CO}_2)_n(\text{MPh}_3)_2]$, M = As or Sb, $n \approx 1$, 581–8

Multinuclear magnetic resonance studies of the hydrolysis of aluminium(III). Part 10. Proton, carbon-13, and aluminium-27 spectra of aluminium acetate at very high magnetic field, 1897–1902

Binuclear copper(II) complexes of new dinucleating ligands with a pyrazolate group as an endogenous bridge. Effects of exogenous azide and acetate bridges on magnetic properties, 2077–82

A convenient route to binuclear osmium(I) complexes containing diphosphine and arsinophosphine ligands. *X*-Ray crystal structure of $[\text{Os}_2(\text{MeCO}_2)_2(\text{CO})_4(\text{dppm})_2]$, 2303–8

ACETONITRILE

Synthesis and characterization of a family of nitrile-substituted triangular hydridocarbonyl rhenium clusters, $[\text{Re}_3(\mu-\text{H})_3(\text{CO})_{12-n}(\text{NCMe})_n]$ ($n = 1$ –3). *X*-Ray crystal structures of $[\text{Re}_3(\mu-\text{H})_3(\text{CO})_{11}(\text{NCMe})]$ and $[\text{Re}_3(\mu-\text{H})_3(\text{CO})_9(\text{NCMe})_3]$, 1143–8

ACETYLACETONATE

Oxovanadium(IV) and dioxomolybdenum(VI) complexes: synthesis from the corresponding acetylacetonato complexes and *X*-Ray structures, 145–50

Heteroleptic poly(pyrazolyl)borate derivatives of the lanthanide ions. The synthesis of acetylacetonato complexes and the molecular structures of $[\text{Ln}\{\text{HB}(\text{C}_3\text{N}_2\text{H}_3)_3\}_2(\text{MeC(O)CHC(O)Me})]$ ($\text{Ln} = \text{Ce}$ or Yb), 1393–1400

ACETYLACETONEIMINATE

N,N'-Ethylenebis(acetylacetoneiminato)titanium complexes: synthesis, crystal structures, and properties of titanium(III) and titanium(IV) derivatives, 953–8

ACID

Reactions of co-ordinated ligands. Part 44. The reactivity of the carbyne complex $[\text{Mo}(\equiv\text{CCH}_2\text{Bu}')\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$ towards protonic acids, diphenylphosphine, and the π -acid tetracyanoethylene; formation and crystal structures of $[\text{MoH}(\equiv\text{CCH}_2\text{Bu}')\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]\text{[BF}_4]$, $[\text{Mo}(\sigma\text{-OC(O)CF}_3)\{\eta^2\text{-HOC(O)CF}_3\}\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, and $[\text{Mo}(\equiv\text{CCH}_2\text{Bu}')\{\eta^2\text{-C}_2(\text{CN})_4\}\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, 511–8

ACTINIDE

Actinide structural studies. Part 17. The crystal and molecular structures of four uranyl(VI) complexes with *N,N*-dialkyldithiocarbamate ligands, 471–80

Tri(η -cyclopentadienyl)uranium(IV) silyl and siloxide compounds. Crystal structure of $[\text{U}(\eta^5\text{-C}_5\text{H}_5)_3(\text{OSiPh}_3)]$. Insertion of isocyanide into a uranium–silicon bond, 677–82

ACYL

Studies on olefin-co-ordinating transition-metal carbene complexes. Part 14. An unusual reaction of tricarbonyl(η^4 -cyclopenta-1,3-diene)iron: a new approach to acyldicarbonyl(η -cyclopentadienyl)iron complexes, and the *X*-ray crystal structure of $[\text{Fe}(\text{COC}_6\text{H}_4\text{CF}_3-p)(\eta\text{-C}_5\text{H}_5)(\text{CO})_2]$, 635–8

ADAMANTYL

Steric effect of the adamantyl ligand on crystal structure and dissociation rates in organocobalt B_{12} models, 1945–6

ADENOSINE

Potentiometric and electron spin resonance investigation on the interactions between adenosine 5'-triphosphate (ATP) and

triphosphate ligands with vanadyl ion. Synthesis and characterization of solid vanadyl-ATP compounds from acidic aqueous solution, 575–80

ALCOHOL

Studies on transition-metal oxo and nitrido complexes. Part 10. New oxo-ruthenium and oxo-osmium pyridine complexes, and use of the former as catalysts for oxidation of alcohols, 901–6

Chelate-stabilised alcohol- and alkoxo-palladium(II) complexes derived from the ligands $\text{Ph}_2\text{PCH}_2\text{CR}_2\text{OH}$ ($\text{R} = \text{Me}$ or H), 1193–6

Photo-oxidation of alcohols by hexachlorometalate(IV) ions ($\text{M} = \text{Pt}$, Pd , or Ir): spin trapping and matrix isolation electron spin resonances studies, 1237–40

ALKALINE EARTH

Thermodynamics of the formation of metal complexes of 1,4,10-trioxa-7,13-diazacyclopentadecane-*N,N'*-diacetic acid and of 1,4,10,13-tetraoxa-7,16-diazacyclo-octadecane-*N,N'*-diacetic acid, 133–8

ALKENE

Platinum metal complexes of potentially chelating alkene-thioether and alkene-selenoether ligands: synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{MeE}(\text{CH}_2)_n\text{CH}=\text{CH}_2\}]$ ($\text{M} = \text{Pt}$ or Pd ; $\text{X} = \text{Cl}$, Br , or I ; $\text{E} = \text{S}$ or Se ; $n = 2$ or 3) and the *X*-ray structure of *cis*-dibromo(2-thia-6-heptene)platinum(II), $[\text{PtBr}_2\{\text{MeS}(\text{CH}_2)_3\text{CH}=\text{CH}_2\}]$, 985–90

Cationic dinuclear rhodium complexes as catalyst precursors for the hydroformylation of alkenes, 1579–82

ALKENYL

Organoborazines. Part 2. Synthesis of alkenyltriborazines, 2423–8

ALKENYLIDENE

Transition metal-cyanocarbon chemistry. Part 8. Cyano-substituted buta-1,3-dienylidene bridged di-iron complexes from unprecedented cyanoalkyne insertion into the carbon–hydrogen bond of a bridging alkenylidene ligand, 2187–92

ALKOXIDE

Chelate-stabilised alcohol- and alkoxo-palladium(II) complexes derived from the ligands $\text{Ph}_2\text{PCH}_2\text{CR}_2\text{OH}$ ($\text{R} = \text{Me}$ or H), 1193–6

ALKOXYCARBONYL

Carbonyl, hydridocarbonyl, and hydridoalkoxycarbonyl complexes of iridium with phosphorus–sulphur hybrid ligands, and the crystal and molecular structure of $[\text{IrH}(\text{CO}_2\text{Me})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{SEt})_2]\text{BPh}_4$, 607–14

ALKYL

Relative importance of the inductive and steric effects of alkyl groups (R) in understanding the nature of the Co^{III} –C bond in alkyl cobaloximes, and comments on related alkyl cobalamins, 115–8

A reinvestigation of the structures of organotin sulphates and chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74

Synthesis, infrared, and tin-119 Mössbauer spectroscopic characterization of seven-co-ordinate diorganotin(IV) adducts with 2,6-diacylpyridine acylhydrazones, including the *X*-ray crystal structure of $\text{SnEt}_2(\text{dapt})$ [$\text{H}_2\text{dapt} = 2,6$ -diacetylpyridine bis(2-thienylhydrazone)], 289–94

Novel five-co-ordinate diorganorhodium compounds. *X*-Ray structure of bis-*ortho*-chelated $[\text{Rh}^{III}(\text{Me})\text{I}\{\text{C}_6\text{H}_3(\text{CH}_2\text{NMe}_2)_2\text{-}2,6\}]$, 317–24

The binding of trialkyltin(IV) moieties to rat haemoglobin, and the structure of model systems, studied by tin-119 Mössbauer spectroscopy, 519–26

Thermodynamics of hydroxy complex formation of dialkyltin(IV) ions in aqueous solution, 773–8

Reactions of cyclopalladated compounds. Part 21. Various examples of sulphur-assisted intramolecular palladation of aryl and alkyl groups, 1715–20

ALKYLATION

Formation of alkylurethanes from carbon dioxide by regioselective *O*-alkylation of alkali-metal *N,N*-diethylcarbamates in the presence of complexing agents, 1007–10

ALKYLIDENE

Organic chemistry of dinuclear metal centres. Part 12. Synthesis, X-ray crystal structure, and reactivity of the di- μ -alkylidene complex $[\text{Ru}_2(\text{CO})_2(\mu\text{-CHMe})(\mu\text{-CMe}_2)(\eta\text{-C}_5\text{H}_5)_2]$; alkylidene linking, 1799–806

ALKYLDYNE

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 84. Carbaborane tungsten–platinum complexes having a $\mu\text{-CC}_6\text{H}_3\text{Me}_2$ –2,6 ligand; crystal structures of $[\text{WPt}(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_n(\text{PEt}_3)_2(\mu\text{-}\sigma^*\text{-}\eta^5\text{-C}_2\text{B}_9\text{H}_8\text{Me}_2)]$ ($n = 2$ or 3), 303–12

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 85. Synthesis of chain and ring compounds containing molybdenum, 785–96

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 86. Alkyldyne(carborane)–molybdenum–gold, –rhodium and –iron complexes; crystal structure of $[\text{NEt}_4][\text{MoFe}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me-4})(\mu\text{-}\sigma^*\text{-}\eta^5\text{-C}_2\text{B}_9\text{H}_7\text{Me}_2)(\text{CO})_8]$, 797–808

Reactions of co-ordinated ligands. Part 45. Ligand displacement and oxidative reactions of the carbyne complex $[\text{Mo}(\equiv\text{CCH}_2\text{Bu})^2\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$; crystal and molecular structures of $[\text{Mo}(\eta^3\text{-RN=C-C}(\text{CH}_2\text{Bu})\text{-C=NR})(\text{CNR})_2(\eta\text{-C}_5\text{H}_5)]$ ($R = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), $[\text{Mo}(\text{C=CHBu})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, and $[\text{Mo}=\text{C}(\text{CH}_2\text{Bu})\text{P}(\text{O})(\text{OMe})_2]\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)$, 1321–32

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 87. Dicosahedral carbaborane(alkyldyne)tungsten complexes as reagents for the synthesis of compounds with heteronuclear metal–metal bonds; crystal structures of $[\text{NEt}_4][\text{W}(\equiv\text{CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_2(\eta^6\text{-C}_2\text{B}_{10}\text{H}_{10}\text{Me}_2)]$ and $[\text{NEt}_4][\text{WFe}(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_4(\eta^6\text{-C}_2\text{B}_{10}\text{H}_{10}\text{Me}_2)]$, 1363–74

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 88. Carbaboran tungsteniridium compounds; crystal structure of the complex $[\text{WIr}(\mu\text{-CC}_6\text{H}_4\text{Me-4})(\text{CO})_2(\text{PEt}_3)_2(\eta^5\text{-C}_2\text{B}_9\text{H}_9\text{Me}_2)]$, 1845–54

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 89. Tetra- and penta-nuclear tungsten–rhodium complexes; crystal structures of $[\text{W}_3\text{Rh}_2(\mu\text{-CO})_2(\mu\text{-C}(\text{Me})\text{C}(\text{O}))\{\mu\text{-PPPh}_2\}_2(\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3]$ and $[\text{W}_3\text{Rh}_2(\mu\text{-CO})_3(\mu\text{-CMe})\{\mu\text{-C}(\text{Me})\text{PPPh}_2\}_2(\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3]$, 1855–64

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 90. Synthesis of the complexes $[\text{W}_2\text{M}_2(\mu\text{-CO})(\mu\text{-CMe})(\mu_3\text{-CMe})(\mu\text{-PPPh}_2)_2(\text{CO})_3(\mu\text{-C}_5\text{H}_5)_2]$ ($M = \text{Rh}$ or Ir) and related tetra- and penta-nuclear metal compounds; interconversion of ketenyl and λ^5 -phospho-alkyne ligands bridging tungsten–rhodium bonds, 1865–70

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 91. Steric implications in the reactions of the complexes $[\text{M}(\equiv\text{CR})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ ($M = \text{Mo}$ or W , $R = \text{C}_6\text{H}_4\text{Me-4}$, $\text{C}_6\text{H}_4\text{OMe-2}$, or $\text{C}_6\text{H}_3\text{Me}_2\text{-2,6}$) with the compounds $[\text{Mo}(\text{CO})_3(\text{NCMe})_3]$ and $[\text{Cu}(\text{thf})(\eta\text{-C}_5\text{Me}_5)]$ ($\text{thf} = \text{tetrahydrofuran}$); crystal structure of $[\text{Mo}_3(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2]$, 1871–8

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 92. Synthesis of heteropolynuclear metal compounds with chains of seven to eleven metal atoms; crystal structure of $[\text{Mo}_3\text{W}_3\text{Pt}_6(\mu_3\text{-CMe})_3(\mu_3\text{-CC}_6\text{H}_4\text{Me-4})_4(\text{CO})_{10}(\text{PMe}_2\text{Ph})_4(\eta\text{-C}_5\text{H}_5)_5]$, 2269–80

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 95. Synthesis and crystal structures of $[\text{W}_2\text{RuPt}(\mu\text{-CC}_6\text{H}_4\text{Me-4})(\mu_3\text{-CC}_6\text{H}_4\text{Me-4})(\text{CO})_6(\text{PMe}_2\text{Ph})(\eta\text{-C}_5\text{H}_5)_2]$ and $[\text{W}_2\text{Ru}_2\text{Pt}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me-4})_2(\mu\text{-CO})_2(\text{CO})_9(\eta\text{-C}_5\text{H}_5)_2]$, 2289–96

ALKYNE

Reactions of the bis(alkyne) complexes $[\text{WI}_2(\text{CO})(\text{NCMe})(\eta^2\text{-RC}_2\text{R}_2)]$ ($R = \text{Me}$ or Ph) with mono- and bi-dentate phosphine-donor ligands and the X-ray crystal structure of $[\text{WI}_2(\text{CO})(\text{Ph}_2\text{P}(\text{CH}_2)\text{PPPh}_2)(\eta^2\text{-MeC}_2\text{Me})]$, 295–302

Alkyne insertion into ruthenium–carbon bonds: formation and ring-closing reactions of vinyl complexes, 465–70

Reactions of co-ordinated ligands. Part 44. The reactivity of the carbyne complex $[\text{Mo}(\equiv\text{CCH}_2\text{Bu})^2\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$ towards protonic acids, diphenylphosphine, and the π -acid tetracyanoethylene; formation and crystal structures of $[\text{MoH}(\equiv\text{CCH}_2\text{Bu})^2\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)][\text{BF}_4]$, $[\text{Mo}\{\sigma$

$\text{OC(O)CF}_3\}$ $\{\eta^2\text{-HOC(O)CF}_3\}$ $\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, and $[\text{Mo}(\equiv\text{CCH}_2\text{Bu})^2\{\eta^2\text{-C}_2(\text{CN})_4\}$ $\{\text{P}(\text{OMe})_3\}(\eta\text{-C}_5\text{H}_5)][\text{BF}_4]$, 511–8

Reactions of hydride complexes of ruthenium(ii) with alkynes, 589–94

Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten $[\text{Mn}(\eta^3\text{SC}(\text{CF}_3)=\text{C}(\text{CF}_3)\text{H})(\text{CF}_3\text{C}=\text{C}\text{CF}_3)(\eta^3\text{-C}_5\text{H}_5)]$ ($M = \text{Mo}$ or W). Their relevance to the mechanism of metal-promoted isomerisation of carbon–carbon double bonds, 1027–36

Heteronuclear transition metal–alkyne clusters. Part 1. Reactions of tris(alkyne)monocarbonyltungsten complexes with octacarbonyldicobalt. Synthesis, X-ray crystal structure, and two-dimensional nuclear magnetic resonance studies of $[\text{WCo}_2(\mu\text{-C}_2\text{Et}_2)(\mu\text{-C}_4\text{Et}_4)(\text{CO})_8]$, 1375–80

Preparation, characterisation, and reactions with phosphite donor ligands of the new cationic monoalkyne complexes $[\text{WI}(\text{CO})(\text{NCMe})(\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPPh}_2)(\eta^2\text{-RC}_2\text{R})][\text{BF}_4]$ ($n = 1$ –6; $R = \text{Me}$ or Ph). X-Ray crystal structure of $[\text{WI}(\text{CO})\{\text{P}(\text{OPr}')_3\}(\text{Ph}_2\text{P}(\text{CH}_2)\text{PPPh}_2)(\eta^2\text{-MeC}_2\text{Me})][\text{BF}_4]_{0.5}[\text{OH}]_{0.5}\cdot 0.5\text{H}_2\text{O}$, 1903–10

Transition metal–cyanocarbon chemistry. Part 8. Cyano-substituted buta-1,3-dienylidene bridged di-iron complexes from unprecedented cyanoalkyne insertion into the carbon–hydrogen bond of a bridging alkenylidene ligand, 2187–92

Reactions of 1-alkynes with *trans*- $[\text{ReCl}(\text{N}_2)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{-PPh}_2)_2]$: preparation of the vinylidene compounds *trans*- $[\text{ReCl}(\equiv\text{C}=\text{CH})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$ ($R = \text{alkyl}$ or aryl) and X-ray structure of *trans*- $[\text{ReCl}(\equiv\text{C}=\text{CHPh})-(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$, 2381–8

ALKYNYLALKYLIDYNE

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 94. Synthesis of di- and tri-metal compounds via the reagents $[\text{M}(\equiv\text{C}-\text{C}\equiv\text{CBu}^i)(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ ($M = \text{Mo}$ or W), 2281–8

ALKYNYL METHYLIDYNE

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 92. Synthesis of alkynylmethylidyne-molybdenum and -tungsten complexes and their reactions with $[\text{Co}_2(\text{CO})_8]$ and $[\text{Mo}_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2]$, 2261–8

ALLYL

Synthesis and stereochemical studies of $\eta^3\text{-allyl palladium(ii)}$ complexes containing a chiral chelating ligand, 161–8

Synthesis, characterization, and thermolytic behaviour of $\eta^3\text{-allyl(methyl)(tertiary phosphine)palladium(ii)}$, and X-ray structure of the cluster $[\text{Pd}_4(\mu\text{-C}_3\text{H}_5)_2(\mu\text{-Cl})_2(\mu\text{-dppm})_2]4\text{C}_4\text{H}_8\text{O}$ (dppm = $\text{Ph}_2\text{PCH}_2\text{PPPh}_2$), 1519–26

ALUM

Vibrational spectroscopic studies of tervalent hexa-aqua cations: oriented single-crystal Raman spectra between 275 and 1200 cm^{-1} of the caesium rhodium alums, $\text{CsRh}(\text{SO}_4)_2\cdot 12\text{H}_2\text{O}$, $\text{CsRh}(\text{SeO}_4)_2\cdot 12\text{H}_2\text{O}$, and $\text{CsRh}(\text{SO}_4)_2\cdot 12\text{D}_2\text{O}$, 1771–8

ALUMINIUM

Crystall and molecular structures of diaqua(nitrilotriacetato)aluminium(iii) and di- μ -hydroxobis(nitrilotriacetato)-dialuminate(iii) dianion, 1513–8

Multinuclear magnetic resonance studies of the hydrolysis of aluminium(iii). Part 9. Prolonged hydrolysis with aluminium metal monitored at very high magnetic field, 1889–96

Multinuclear magnetic resonance studies of the hydrolysis of aluminium(iii). Part 10. Proton, carbon-13, and aluminium-27 spectra of aluminium acetate at very high magnetic field, 1897–902

ALUMINIUM-27

Multinuclear magnetic resonance studies of the hydrolysis of aluminium(iii). Part 9. Prolonged hydrolysis with aluminium metal monitored at very high magnetic field, 1889–96

Multinuclear magnetic resonance studies of the hydrolysis of aluminium(iii). Part 10. Proton, carbon-13, and aluminium-27 spectra of aluminium acetate at very high magnetic field, 1897–902

AMIDE

Preparation of the bis(trimethylsilyl)amido lanthanide chlorides $[\text{Ln}[\text{N}(\text{SiMe}_3)_2]_2(\mu\text{-Cl})(\text{thf})_2]$ ($\text{thf} = \text{tetrahydrofuran}$), and the crystal and molecular structures of the gadolinium and ytterbium complexes, 623–6

The chemical and electrochemical one-electron oxidation of diamidonaphthalene-bridged complexes: paramagnetic $[\text{Rh}_2]^{3+}$ -containing A-frames, 683–8

AMIDE (contd)

- Tri-aza analogue of an acetylacetone-complex formed by nucleophilic attack of a co-ordinated nitrile by amidines. *X*-Ray crystal structure of $[\text{Pt}\{\text{HNC(Ph)NC(Ph)NH}\}_2] \cdot \frac{2}{3} \text{PhMe}$, 837–42
- Synthesis of a new P_2N_2 ligand N,N' -bis[2-(diphenylphosphino)phenyl]propane-1,3-diamine, H_2L^2 , and some of its complexes with elements of the nickel triad and rhodium: *X*-ray structure analyses of the neutral complex $[\text{NiL}^2]$ and the *trans*-spanned $[\text{Rh}(\text{CO})\text{Cl}(\text{H}_2\text{L}^2)]$, 1067–74
- Reaction of $[\text{Rh}_2\{\mu\text{-1,8-(NH)}_2\text{C}_{10}\text{H}_6\}(\text{CO})_2(\text{PPPh}_3)_2]$ with $[\text{M}(\text{PPPh}_3)]^+$ ($\text{M} = \text{Au, Ag, or Cu}$): formation of unusual 44-electron Rh_2M clusters. Crystal structure of $[\text{Rh}_2\{\mu\text{-1,8-(NH)}_2\text{C}_{10}\text{H}_6\}(\text{CO})_2(\text{PPPh}_3)_2]\text{ClO}_4 \cdot 2\text{CH}_2\text{Cl}_2$, 1249–52
- New dialkylamido complexes of transition metals in groups 4 and 5 stabilised by terdentate ligands, 1253–8
- Thermodynamic and spectrophotometric investigation of complex formation between hydrogen, cobalt(II), nickel(II), and copper(II) ions and 2-amino-*N*-hydroxypropanamide in aqueous solution, 1299–310
- Tertiary phosphine complexes of triply bonded ditungsten bis(t-butylamide) tetrachloride. The *X*-ray crystal structures of $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMMe}_3)_2]$, $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PEt}_3)_2]$, and *cis*- $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMMe}_2\text{Ph})_2]$, 1537–42
- Copper(II)-assisted hydrolysis of 2,4,6-tris(2-pyridyl)-1,3,5-triazine. Part 2. Crystal structures of [bis(2-pyridylcarbonyl)amido](pyridine-2-carboxamide)copper(II) trifluoromethanesulphonate and [bis(2-pyridylcarbonyl)amido][2,4,6-tris(2-pyridyl)-1,3,5-triazine]copper(II) trifluoromethanesulphonate, 1681–8
- Bi-, tetra-, and poly-nuclear complexes of gold(I) and silver(I) with bis(diphenylphosphino)-methane or -amide acting as six-electron donors, 1883–6
- Complex-forming properties of L- α -alaninehydroxamic acid (2-amino-*N*-hydroxypropanamide), 2247–52

AMIDINE

- Tri-aza analogue of an acetylacetone-complex formed by nucleophilic attack of a co-ordinated nitrile by amidines. *X*-Ray crystal structure of $[\text{Pt}\{\text{HNC(Ph)NC(Ph)NH}\}_2] \cdot \frac{2}{3} \text{PhMe}$, 837–42

AMINE

- Palladium complexes with the tripodal phosphine tris(2-diphenylphosphinoethyl)amine. Synthesis and structure of trigonal, tetrahedral, trigonal bipyramidal, and square planar complexes, 211–6
- The reaction of hydroxo-species of palladium(II) with bidentate nucleophiles, 375–8
- Isopropylidisilylamine and disilyl-t-butylamine: preparation, spectroscopic properties, and molecular structure in the gas phase, determined by electron diffraction, 779–84
- Reactions of amine hydrochlorides with phosphorus trichloride oxide: preparation and *X*-ray crystal structure of 2,4,6-trichloro-1,3,5-triphenyl-1,3,5,2 λ^5 4 λ^5 6A 5 -triazatriporphorinane 2,4,6-trioxide (PhNPOCl_3), 821–8
- Synthesis of a new P_2N_2 ligand N,N' -bis[2-(diphenylphosphino)phenyl]propane-1,3-diamine, H_2L^2 , and some of its complexes with elements of the nickel triad and rhodium: *X*-ray structure analyses of the neutral complex $[\text{NiL}^2]$ and the *trans*-spanned $[\text{Rh}(\text{CO})\text{Cl}(\text{H}_2\text{L}^2)]$, 1067–74
- Zinc(II) complexes of 4,7-diazadecane-1,10-diamine, 4,8-diazzaundecane-1,11-diamine, and tetraethylpenepentamine: a carbon-13 nuclear magnetic resonance study, 1171–8
- Nucleophilic (Michael) additions to $(\text{Ph}_2\text{P})_2\text{C}=\text{CH}_2$ when chelated to platinum(II) dichloride or di-iodide, 1527–30
- Syntheses, crystal structures, and Raman and resonance Raman spectroscopy of platinum(II), platinum(IV), and halogen-bridged chain platinum(II,IV) complexes involving the terdentate ligand 3-methyl-3-azapentane-1,5-diamine (4Me-dien), 1689–96

AMINO ACID

- The unusual co-ordination ability of vasopressin-like peptides; potentiometric and spectroscopic studies of some copper(II) and nickel(II) complexes, 173–8
- Possible biological activity in proton, metal, and amino hydroxamic acid equilibria. Protonation and complex formation reactions between *N*-hydroxy-D-asparagine and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution (1988, 953), 383
- Interligand interaction in mixed amino acid complexes, 567–8
- Complexes of aminophosphonates. Part 3. Copper(II) complexes of several monophosphono and diphosphono dipeptides, 1053–8

Composition, stability, and lability of copper(II) dipeptide complexes, 1293–8

Synthesis, spectroscopic, and magnetic studies of copper(II) with *N*-protected amino acids: crystal and molecular structure of diaquatetrakis(μ -*N*-benzoyl- α -alaninato)-dicopper(II), 1341–4

Sulphonamide nitrogen-containing, nitrogen protected amino acids interacting with the Cd^{2+} ion. A cadmium-113 nuclear magnetic resonance and potentiometric study. *X*-Ray diffraction of $[\text{Cd}(\text{ts}-\beta\text{-alaO}_2)\text{H}_2\text{O}]$, 1345–50

Co-ordination of nickel(II) ions by angiotensin II and its peptide fragments. A potentiometric, proton nuclear magnetic resonance and circular dichroism spectroscopic study, 1471–6

Mixed-metal complexes in solution. A potentiometric and spectrophotometric study of copper(II)-nickel(II)- δ -hydroxylysine complexes in aqueous solution, 1745–50

Electron spin resonance study of copper(II) complexes of X-glycine and glycyl-X type dipeptides, and related tripeptides. Variation of co-ordination modes with ligand excess and pH in fluid and frozen aqueous solutions, 1925–32

Complex-forming properties of L- α -alaninehydroxamic acid (2-amino-*N*-hydroxypropanamide), 2247–52

Functionalized amino acid complexation to a cobalt(III) complex of (2,5,8,11-tetraethyl-1,4,7,10-tetraazacyclododecane. Crystal structures of complexes containing (R)- or (S)-serine and (R)- α -methylserine, and release of amino acid from the complexes, 2359–66

AMMONIA

New metal-sulphur-nitrogen compounds from reactions in liquid ammonia. The *X*-ray structures of *trans*-bis(acetophenone dimethylhydrazone- N^3)dichloropalladium(II) and [di(azathien)-1-yl-S 3 N 4]2-hydrazonoethyl]phenyl]palladium(II), 1179–86

AMMONIUM

Synthesis of ammonium ions and nitrosylation reactions using nitrosyl chloride and alkyl nitrites, 1139–42

ANGIOTENSIN

Co-ordination of nickel(II) ions by angiotensin II and its peptide fragments. A potentiometric, proton nuclear magnetic resonance and circular dichroism spectroscopic study, 1471–6

ANILINE

Addition of anilines to the $[\text{Fe}(\text{CO})_3(1-5-\eta\text{-C}_7\text{H}_9)]\text{BF}_4$ complex, and the 'Ordered Transition State Mechanism', 561–6

ANTIMONY

Correlations between nuclear magnetic resonance spectra and crystal structures. A carbon-13 nuclear magnetic resonance study in the solid state of the Group 15 xanthates $[\text{A}(\text{S}_2\text{COR})_3]$ ($\text{A} = \text{As}^{III}, \text{Sb}^{III}, \text{or Bi}^{III}; \text{R} = \text{Me, Et, or Pr}'$), 233–6

Structural and motional study of pyridinium tetrahalogenoantimonates(III) by means of nuclear quadrupole resonance spectroscopy, 631–4

AQUATION

Base-catalysed aquation of $\alpha\beta$ -syn- and $\alpha\beta$ -anti-chloro- and bromo-[1,9-bis(2-pyridyl)-2,5,8-triazanonane]- and [1,11-bis(2-pyridyl)-2,6,10-triazaundecane]-cobalt(III) cations. An unusually base-sensitive halogenopentamine cobalt(III) system, 645–54

Kinetics of aquation and base hydrolysis of some *trans*- $[\text{Co}(\text{tn})_2(\text{B})\text{Cl}]^{2+}$ ($\text{tn} = 1,3$ -diaminopropane; $\text{B} = \text{OH, NCS, N}_3, \text{NO}_2, \text{NH}_3, \text{or CN}$) complexes. The effect of ring size on reactivity, 853–8

ARENÉ

Metal cluster-induced electrophilicity of arenes, 481–8

ARENEDIAZONIUM

Reactivity of new osmium dihydrides with arenediazonium cations: preparation and properties of bis(aryldiazene) and mono(aryldiazenido) complexes. A comparison with analogous iron and ruthenium derivatives, 2353–8

ARSENIC

An investigation of halide abstraction by arsenic(III): reactions of AsCl_3 with MCl_2 ($\text{M} = \text{Mg or Zn}$) and TiCl_4 in donor solvents (MeCN or POCl_3). Formation and crystal structure of the dimeric titanium(IV) compound $[\text{Ti}_2\text{Cl}_6(\text{POCl}_3)_2(\text{PO}_2\text{Cl}_2)_2]$, 57–60

Correlations between nuclear magnetic resonance spectra and crystal structures. A carbon-13 nuclear magnetic resonance study in the solid state of the Group 15 xanthates $[\text{A}(\text{S}_2\text{COR})_3]$ ($\text{A} = \text{As}^{III}, \text{Sb}^{III}, \text{or Bi}^{III}; \text{R} = \text{Me, Et, or Pr}'$), 233–6

Kinetic studies of the oxidation reaction of arsenic(III) to arsenic(V) by peroxodisulphate ion in aqueous alkaline media, 357–60

Some transition-metal complexes of 1,3,2-dithiarsolane, 885–8

ARSINE

A kinetic study of the transmission of ligand effects across the quadruple bond in $[\text{Re}_2\text{Cl}_8]^{2-}$, 767–72

ARYL

Novel five-co-ordinate diorganorhodium compounds. X-Ray structure of bis-*ortho*-chelated $[\text{Rh}^{\text{III}}(\text{Me})\text{I}\{\text{C}_6\text{H}_3(\text{CH}_2\text{NMe}_2)_2\}_{2-2,6}]$, 317–24

A kinetic and mechanistic study of the solvolysis of aryltellurium trihalides in organic solvents, 457–60

Alkyne insertion into ruthenium–carbon bonds: formation and ring-closing reactions of vinyl complexes, 465–70

Substitution reactions on sterically hindered square-planar *trans*- $[\text{NiBr}(\text{R})(\text{PR}'_3)_2]$ (R = aryl) complexes. Effects of the substituents of the aryl ligand, 1669–74

Reactions of cyclopalladated compounds. Part 21. Various examples of sulphur-assisted intramolecular palladation of aryl and alkyl groups, 1715–20

Synthesis of arylthallium complexes using organomercury compounds. Crystal and molecular structures of $[\text{NMe}_4]\text{[Tl(C}_6\text{H}_4\text{N=NP}-2)\text{Cl}_3]$ and $[\text{Tl}(\text{C}_6\text{H}_4\text{CH}_2\text{NMe}_2-2)\text{Cl}_2]$, 2241–6

ARYLAZOPYRIDINE

Synthesis, structure, and electrochemical reduction of 2-(arylaquo)pyridine complexes of rhodium(III), 1635–8

ASPARAGINE

Possible biological activity in proton, metal, and aminohydroxamic acid equilibria. Protonation and complex formation reactions between *N*-hydroxy-D-asparagine and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution (1988, 953), 383

ASPIRINATE

Synthesis, spectroscopic, electrochemical, and magnetic properties of dimolybdenum(II,II), diruthenium-(II,III) and -(II,II) complexes containing bridging aspirinate (2-acetoxybenzoate) ligands, 2435–42

AZATHIENYL

New metal–sulphur–nitrogen compounds from reactions in liquid ammonia. The X-ray structures of *trans*-bis(acetophenone dimethylhydrazone- N^{α})dichloropalladium(II) and [di(azathien-1-yl- S^1N^2)[2-(hydrazoneoethyl)phenyl]palladium(II), 1179–86

AZIDE

Reactivity of cationic molybdenum(II) complexes. Part 4. Isolation and crystal structure determination of $R_s\text{[Mo}(\text{CO})_2(\eta-\text{C}_5\text{Me}_5)(\text{PPh}_3)(\text{N}_3)]$ and $R_s\text{[Mo}(\text{CO})_2(\eta-\text{C}_5\text{Me}_5)(\text{PPh}_3)(\text{NCO})]$, 959–64

Binuclear copper(II) complexes of new dinucleating ligands with a pyrazolate group as an endogenous bridge. Effects of exogenous azide and acetate bridges on magnetic properties, 2077–82

AZIDOETHANOL

Comparative study on the multielectron reduction of $\text{HOC}_2\text{H}_4\text{N}_3$ catalyzed by $[\text{NBu}^n_4]_4\text{[MoFe}_3\text{S}_4(\text{SPh})_3(o-\text{O}_2\text{C}_6\text{Cl}_4)]_{2-}$ and $[\text{NBu}^n_4]_2\text{[Fe}_4\text{S}_4(\text{SPh})_4]$ -modified glassy carbon electrodes, 1547–54

AZINE

Iridium(III) hydrido complexes with 2-pyridinaldazines. Crystal structure of $[\text{IrH}_2(\text{mpaa})(\text{PPh}_3)_2]\text{SbF}_6$. Synthesis and characterization of homo- and hetero-nuclear complexes, 1075–82

Reactivity of the iridium nitrosyl complex $[\text{Ir}(\text{NO})(\text{paa})(\text{PPh}_3)_2]\text{[PF}_6\text{]}_2$ (paa = 2-pyridinecarbaldehyde azine) with nucleophiles: insertion of NO into a methinic C–H bond; crystal structure of $[\text{Ir}\{(2-\text{C}_5\text{H}_4\text{N})\text{CH=N-N=C}(\text{NOH})(2-\text{C}_5\text{H}_4\text{N})\}\{\text{Me}(\text{EtO})\text{C=NH}\}(\text{PPh}_3)_2]\text{[PF}_6\text{]}_2$, 1217–26

Rearrangements of $>\text{C=N-}$ functionalities promoted by cobalt(I): reactivity of cyclopentadienylbis(ethylene)cobalt(I) with dibenzophenone azine, benzophenone oxime, benzylidene(phenyl)amine and crystal structure determinations of the products, 1957–62

AZOLE

Synthesis, electrochemistry, and electron spin resonance spectra of some seventeen-electron complexes containing the $\{\text{Mo}(\text{NO})\}^{2+}$ core and the molecular structure of $[\text{Mo}(\text{NO})(\text{HB}(\text{Me}_2\text{pz}))_3](\text{pyr})_2$, 127–32

BARIUM

Lead(II) and barium(II) complexes of a potentially octadentate macrocyclic ligand (H_6L), capable of providing endogenous alkoxy and phenoxy bridges. The X-ray crystal structures of $[\text{Pb}(\text{H}_6\text{L})]\text{[ClO}_4\text{]}_2$, $[\text{Pb}_2(\text{H}_6\text{L})]\text{[ClO}_4\text{]}_2$, and $[(\text{H}_6\text{L})(\text{H}_2\text{O})_2]\text{[ClO}_4\text{]}_2$, 19–24

A barium dithiocyanate complex with an unusual macrocycle containing pendant arms, the amine precursor, and related copper(II) diperchlorate complexes: synthesis and structure, 1993–2002

BASE CATALYSIS

Base-catalysed aquation of $\alpha\beta$ -syn- and $\alpha\beta$ -anti-chloro- and bromo-[1,9-bis(2'-pyridyl)-2,5,8-triazaonane]- and -[1,11-bis(2'-pyridyl)-2,6,10-triazaundecane]-cobalt(III) cations. An unusually base-sensitive halogenopentamine cobalt(III) system, 645–54

BENZENE

Substitution reactions of the zerovalent iron complexes $[\text{Fe}(\eta-\text{C}_6\text{H}_6)(\text{PMe}_3)_2]$ and $[\text{Fe}(\eta-\text{C}_6\text{H}_6)(\text{dmpe})]$ ($\text{dmpe} = \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$), 1385–92

BENZIMIDAZOLE

Blue copper models. Spectroscopic and electrochemical studies of copper(II) complexes with new ligand systems containing sulphur and nitrogen donor atoms, 1161–70

X-Ray structure of a ternary complex, copper(II)–inosine 5'-monophosphate–benzimidazole. The first example for the absence of direct metal–nucleotide interaction, 1507–12

BENZOATE

Oxidative addition of dibenzoyl peroxide on metallic copper. The crystal structure of $[\text{Cu}(\text{C}_6\text{H}_5\text{CO}_2)_2(\text{py})_2]$, 2331–4

BENZOTHIAZOLETHIOLATE

Interaction of benzo-1,3-thiazoline-2-thione and related ligands with copper(II) salts and the single-crystal X-ray structure of μ -[1,2-bis(diphenylphosphino)ethane]-bis{(benzo-1,3-thiazole-2-thiolato-S)[1,2-bis(diphenylphosphino)ethane]copper(I)}, 2167–72

BENZYL

Synthesis and characterization of benzylindium compounds, 1625–6

BIGUANIDE

Kinetics of the uncatalysed and silver(I)-catalysed oxidation of hydrogen peroxide by [ethylenebis(biguanide)]silver(III) ion in acid perchlorate media, 1645–50

BIPYRIDINE

Kinetics of the oxidation of chloride ions by tris(2,2'-bipyridine)nickel(III) ions in aqueous perchlorate media, 151–4

Ligand additivity in the oxidation potentials of bidentate mixed-ligand ruthenium(II) complexes, 371–4

The reaction of hydroxo-species of palladium(II) with bidentate nucleophiles, 375–8

Preparation and interconversion of dimeric di- μ -hydroxo and tri- μ -hydroxo complexes of platinum(II) and palladium(II) with 2,2'-bipyridine and 1,10-phenanthroline, 403–12

Spectrophotometric and calorimetric studies on the formation of binary (2,2'-bipyridine)nickel(II) and ternary (2,2'-bipyridine)chloronickel(II) complexes in *N,N*-dimethylformamide, 655–60

Crystal and molecular structures of μ -hydrido-bis[$\text{faR}(2,2'\text{-bipyridine})$]tricarbonylrhenium(I) chloride, $\text{faR}(2,2'\text{-bipyridine})$ tricarbonylformatorhenium(I), and $\text{faR}(2,2'\text{-bipyridine})$ tricarbonyl(cyanoethylboraborate)rhenium(I), 1449–54

Complexes of lead(II) bis(O,O' -dialkyl dithiophosphates) with nitrogen donor ligands. The crystal structures of $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\cdot\text{en}$ (en = ethylenediamine), $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\cdot\text{bipy}$ (bipy = 2,2'-bipyridine), and $\{\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\}\cdot\text{en}$, 2443–8

BIPYRIDYL

Mode of binding of dimethyl sulphoxide in platinum(II) and palladium(II) cationic complexes containing a chelated diamine. X-Ray crystal structures of (2,2'-bipyridyl)chloro(dimethyl sulphoxide- O)palladium(II) tetrafluoroborate and chloro(dimethyl sulphoxide- S)(ethylenediamine)palladium(II) perchlorate, 1265–72

Structure of μ -fluoro-bis[bis(2,2'-bipyridyl)fluoronickel(II)] fluoride–ethanol (1/2) trihydrate, revealing several unique features including strong hydrogen bonds between fluoride ions and ethanol molecules, 1273–6

Reduction–oxidation properties of organotransition-metal complexes. Part 30. Triazenido-bridged carbonylrhodium bipyridyl complexes with $[\text{Rh}_2]^{2+}$, $[\text{Rh}_2]^{3+}$, and $[\text{Rh}_2]^{4+}$ cores, and the X-ray structures of $[\text{Rh}_2(\text{CO})_2(\text{bipy})(\mu-\text{RNRR}')_2]\text{[BF}_4\text{]}_2\text{CH}_2\text{Cl}_2$ and $[(\text{Rh}_2\text{I}(\text{CO})(\text{bipy})(\mu-\text{RNRR}')_2)_2]\text{[PF}_6\text{]}_2\cdot2.5\text{CH}_2\text{Cl}_2$ (R = *p*-tolyl), 2049–60

BISMUTH

Correlations between nuclear magnetic resonance spectra and crystal structures. A carbon-13 nuclear magnetic resonance study in the solid state of the Group 15 xanthates $[\text{A}(\text{S}_2\text{COR})_3]$ ($\text{A} = \text{As}^{\text{III}}, \text{Sb}^{\text{V}}$, or Bi^{III} ; $\text{R} = \text{Me}$, Et, or Pr i), 233–6

- Synthesis, characterization, and crystal structures of bismuth complexes with 2,6-diacetylpyridine bis-(2-thienylhydrazone) (H_2dapt): $[Bi(Hdapt)Cl_2] \cdot dmso \cdot H_2O$ and $[Bi(dapt)Cl] \cdot dmso$, containing a quinquedentate ligand, 671–6
- Synthetic and structural studies on organomolybdenum and organotungsten complexes containing bismuth. X-Ray crystal structures of $[BiCl\{Mo(CO)_3(\eta-C_5H_5)\}_2]$ and $[BiCl\{Mo(CO)_2(CNBu^t)(\eta-C_5H_5)\}_2]$ (1988, 2941), 1887
- BORANE**
- A kinetic study of the gas-phase thermolysis of pentaborane(11), 385–90
- Influence of added hydrogen on the kinetics and mechanism of thermal decomposition of tetraborane(10) and of pentaborane(11) in the gas phase, 391–8
- Tris(trimethylsilyl)methyl and tris(dimethylphenylsilyl)methyl derivatives of boron. Crystal structures of dihydroxy[tris(trimethylsilyl)methyl]borane and of the lithium–boron complex $[(MeOH)_2Li(\mu-OMe)_2B(OMe)_2]$, 447–52
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 87. Dicosahedral carbaborane(alkylidyne)tungsten complexes as reagents for the synthesis of compounds with heteronuclear metal–metal bonds: crystal structures of $[NEt_4][W(\equiv CC_6H_3Me_2-2,6)(CO)_2(\eta^6-C_2B_{10}H_{10}Me_2)]$ and $[NEt_4][WFe(\mu-CC_6H_3Me_2-2,6)(CO)_4(\eta^6-C_2B_{10}H_{10}Me_2)]$, 1363–74
- Polyhedral organoiridaborane chemistry. The reaction of MeNC with $[6-(\eta^5-C_5Me_5)-nido-6-IrB_9H_{13}]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[4,4-(\eta^5-C_5Me_5)(MeNC)-arachno-4-IrB_8H_{12}]$, 1465–70
- Metalla-heteroborane chemistry. Part 5. Synthesis of *creso*-[2,2-(PR_3)₂-1,2-SePtB₁₀H₁₀] ($R_3 = Et_3, Bu^n_3$, or Me_2Ph) and *arachno*-[9,9-(PM_2Ph)₂-6,9-SePtB₈H₁₀] and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of *creso*-[2,2-(PEt_3)₂-1,2-SePtB₁₀H₁₀] and *arachno*-[9,9-(PM_2Ph)₂-6,9-SePtB₈H₁₀], 1657–68
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 88. Carbaborane-tungsten-ruthenium compounds; crystal structure of the complex $[WIr(\mu-CC_6H_4Me-4)(CO)_2(PEt_3)_2(\eta^5-C_2B_9H_9Me_2)]$, 1845–54
- BORATE**
- Heteroleptic poly(pyrazolyl)borate derivatives of the lanthanide ions. The synthesis of acetylacetone complexes and the molecular structures of $[\text{Ln}\{\text{HB}(\text{C}_3\text{N}_2\text{H}_3)_3\}_2\{\text{MeC(O)CHC(O)Me}\}]$ ($\text{Ln} = Ce$ or Yb), 1393–400
- Nature of peroxyborate species in aqueous solution: a study by boron-11 nuclear magnetic resonance and Raman spectroscopy, 1651–6
- Bonding in clusters. Part 10. The nature of the BH groups in *creso*-borane anions. A study by vibrational spectroscopy, electronic spectroscopy for chemical analysis, and *ab initio* computation, 2253–60
- BORAZINE**
- Organoborazines. Part 2. Synthesis of alkenyltriborazines, 2423–8
- BORON**
- A kinetic study of the gas-phase thermolysis of pentaborane(11), 385–90
- Influence of added hydrogen on the kinetics and mechanism of thermal decomposition of tetraborane(10) and of pentaborane(11) in the gas phase, 391–8
- Tris(trimethylsilyl)methyl and tris(dimethylphenylsilyl)methyl derivatives of boron. Crystal structures of dihydroxy[tris(trimethylsilyl)methyl]borane and of the lithium–boron complex $[(MeOH)_2Li(\mu-OMe)_2B(OMe)_2]$, 447–52
- Contributions to the chemistry of boron. Part 194. Synthesis and characterization of mono, bi-, and tri-cyclic boron–nitrogen–sulphur ring systems. Crystal structure of 2,7,9,14-tetrakis(diethylamino)-3,5,6,10,12,13-hexathia-1,8-diaza-2,4,7,9,11,14-hexaboratricyclo[9.3.0.0^{4,8}]tetradecane, 1311–20
- Polyhedral organoiridaborane chemistry. The reaction of MeNC with $[6-(\eta^5-C_5Me_5)-nido-6-IrB_9H_{13}]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[4,4-(\eta^5-C_5Me_5)(MeNC)-arachno-4-IrB_8H_{12}]$, 1465–70
- Nature of peroxyborate species in aqueous solution: a study by boron-11 nuclear magnetic resonance and Raman spectroscopy, 1651–6
- Metalla-heteroborane chemistry. Part 5. Synthesis of *creso*-[2,2-(PR_3)₂-1,2-SePtB₁₀H₁₀] ($R_3 = Et_3, Bu^n_3$, or Me_2Ph) and *arachno*-[9,9-(PM_2Ph)₂-6,9-SePtB₈H₁₀] and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of *creso*-[2,2-(PEt_3)₂-1,2-SePtB₁₀H₁₀] and *arachno*-[9,9-(PM_2Ph)₂-6,9-SePtB₈H₁₀], 1657–68
- Bonding in clusters. Part 10. The nature of the BH groups in *creso*-borane anions. A study by vibrational spectroscopy, electronic spectroscopy for chemical analysis, and *ab initio* computation, 2253–60
- BORON-11**
- Nature of peroxyborate species in aqueous solution: a study by boron-11 nuclear magnetic resonance and Raman spectroscopy, 1651–6
- BROMIDE**
- The nickel(II)–bromide system in dimethyl sulphoxide: a detailed study of the influences of temperature and mole ratio, 325–30
- CADMUM**
- Polymeric thiolate complexes in solution. Stability constants of the cadmium(II) complexes of 2-aminopropane-1-thiol, 13–8
- Solvent influence on trifluoromethylation reactions with Cd(CF₃)₂: one-pot synthesis of trifluoromethyl iron complexes, 739–44
- Formation of mixed-ligand complexes of cadmium(II) halide with tri-n-octylphosphine oxide and pyridine or substituted pyridines in 1,2-dichloroethane, 1259–64
- Sulphonamide nitrogen-containing, nitrogen protected amino acids interacting with the Cd²⁺ ion. A cadmium-113 nuclear magnetic resonance and potentiometric study. X-Ray diffraction of [Cd(ts-β-al₂O)₂(H₂O)₄], 1345–50
- Metal complexes of α -hydroxyiminophosphonic acid derivatives. Preparation and crystal structures of calcium and cadmium complexes of methyl (*E*)- α -hydroxyiminobenzylphosphonate, 1911–4
- CADMUM-113**
- Sulphonamide nitrogen-containing, nitrogen protected amino acids interacting with the Cd²⁺ ion. A cadmium-113 nuclear magnetic resonance and potentiometric study. X-Ray diffraction of [Cd(ts-β-al₂O)₂(H₂O)₄], 1345–50
- CAESIUM**
- Radiotracers in fluorine chemistry. Part 12. Reactions between caesium fluoride and arsenic pentafluoride, boron trifluoride, sulphur tetrafluoride, carbonyl fluoride, and carbon dioxide under heterogeneous conditions at room temperature. Radiotracer studies using fluorine-18, sulphur-35, and carbon-14, 937–42
- Vibrational spectroscopic studies of tervalent hexa-aqua cations: oriented single-crystal Raman spectra between 275 and 1 200 cm⁻¹ of the caesium rhodium alums, CsRh(SO₄)₂·12H₂O, CsRh(SeO₄)₂·12H₂O, and CsRh(SO₄)₂·12D₂O, 1771–8
- CALCIUM**
- Metal complexes of α -hydroxyiminophosphonic acid derivatives. Preparation and crystal structures of calcium and cadmium complexes of methyl (*E*)- α -hydroxyiminobenzylphosphonate, 1911–4
- Aqueous chemistry and thermodynamic modelling of CaO-SiO₂-H₂O gels, 2371–80
- CALIXARENÉ**
- Lanthanide ion complexes of the calixarenes. Part 4. Double inclusion by *p*-t-butylcalix[4]arene (H₄L). Crystal structures of [Eu₂(HL)₂(dmf)₄]₇dmf (dmf = dimethylformamide) and H₄L·dmso (dmso = dimethyl sulphoxide), 2217–22
- CALORIMETRY**
- Spectrophotometric and calorimetric studies on the formation of binary (2,2'-bipyridine)nickel(II) and ternary (2,2'-bipyridine)chloronickel(II) complexes in *N,N*-dimethylformamide, 655–60
- CARBENE**
- Insertion of phenylethyne-thiolate or -selenolate into metal–carbene bonds: synths for co-ordinated thioacyl or selenoacetyl anions and for anionic thiocarbene or selenocarbene complexes, 1565–78
- Iron carbene complexes. Part 3. Aminolysis of di-iron alkoxycarbene complexes with ammonia and dimethylamine. X-Ray crystal structure of [Fe₂(μ-SPh)₂(CO)₅{C(NMe₂)Ph}], 2199–204
- CARBIDE**
- Redox activation of the hexanuclear ruthenium cluster [Ru₆C(CO)₁₆]²⁺ using oxidative addition, 243–6
- Polyhedral rearrangements in undeca- and dodeca-nuclear Group 1B metal osmium clusters, 505–10

CARBIDE (contd)

New carbide clusters in the cobalt subgroup. Part 17. Preparation and structural characterization of the mixed-metal octahedral dianion $[\text{Co}_2\text{Rh}_4\text{C}(\text{CO})_{13}]^{2-}$ as its $[\text{PPh}_3]^+$ salt, 879–84
 Synthesis of the cluster dianion $[\text{Os}_{11}\text{C}(\text{CO})_{27}]^{2-}$ by pyrolysis and its reactions with electrophiles; X-ray structure analysis of the mixed-metal derivative $[\text{PMePh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\{\text{Cu}(\text{NCMe})\}]\cdot\text{CH}_2\text{Cl}_2$ and the hydrido derivative $[\text{PMePh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455–64
 Mixed-metal carbido carbonyl clusters. Part 4. Synthesis and structural characterization of $[\text{Rh}_6\text{C}(\text{CO})_{13}\{\text{Au}_2(\text{PPh}_3)_2\}]$, 2343–6

CARBON-13

Zinc(ii) complexes of 4,7-diazadecane-1,10-diamine, 4,8-diazaundecane-1,11-diamine, and tetraethylenepentamine: a carbon-13 nuclear magnetic resonance study, 1171–8
 Isotopic multiplets in the carbon-13 nuclear magnetic resonance spectra due to partial deuteration of co-ordinated NH_2 and NH groups of cobalt(iii) complexes containing an N-glycoside. Importance of sugar ring oxygen atoms in the long-range isotope shifts, 1241–8
 Multinuclear magnetic resonance studies of the hydrolysis of aluminium(iii). Part 10. Proton, carbon-13, and aluminium-27 spectra of aluminium acetate at very high magnetic field, 1897–902
 Solid-state high-resolution carbon-13 magnetic resonance spectra of dithioacetate complexes of platinum, palladium, and nickel, 2457–60

CARBON DIOXIDE

Formation of alkylurethanes from carbon dioxide by regioselective O-alkylation of alkali-metal *N,N*-diethylcarbamates in the presence of complexing agents, 1007–10

CARBON DISULPHIDE

Insertion of CS_2 into Ag–S bonds. Preparations and crystal structures of $[(\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}^i_3\text{-2,4,6}))_4\cdot\text{CHCl}_3]_n$ and its insertion product with CS_2 , $[(\text{Ag}(\text{S}_2\text{CSC}_6\text{H}_2\text{Pr}^i_3\text{-2,4,6}))_2\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}^i_3\text{-2,4,6})_6]\cdot 8\text{CHCl}_3$, 2297–302

CARBON TETRACHLORIDE

Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(iii)–ethanol–carbon tetrachloride system, 1197–202

CARBONYL

Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene ($\text{C}=\text{CHOEt}$) and ethoxyethyne ($\text{CH}=\text{COEt}$) ligands respectively, 5–12

Redox activation of the hexanuclear ruthenium cluster $[\text{Ru}_6\text{C}(\text{CO})_{16}]^{2-}$ using oxidative addition, 243–6

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 84. Carbaborane tungsten–platinum complexes having a $\mu\text{-CC}_6\text{H}_5\text{Me}_2\text{-2,6}$ ligand; crystal structures of $[\text{WPt}(\mu\text{-CC}_6\text{H}_5\text{Me}_2\text{-2,6})(\text{CO})_n(\text{P}(\text{Et})_3)(\mu\text{-}\sigma\text{-}\eta^5\text{-C}_2\text{B}_9\text{H}_8\text{Me}_2)]$ ($n = 2$ or 3), 303–12

Syntheses and dynamic stereochemistry of methyl- and phenyl-carbonyl heterobinuclear metal–metal bonded complexes, 379–82

Thiolate complexes of osmium(iii) with tertiary phosphines and their dicarbonyl derivatives of osmium(ii): X-ray crystal structure of *trans,trans,trans*- $[\text{Os}(\text{SC}_6\text{F}_5)_2(\text{CO})_2(\text{P}(\text{Et})_2\text{Ph})_2]$, 419–24

Chromium chalcogen complexes. Insertion of elemental sulphur into $[\text{Cr}(\text{cp})(\text{CO})_2]_2\text{Se}$ and $[\text{Cr}(\text{cp})_2(\text{CO})_4\text{Se}_2]$ ($\text{cp} = \eta\text{-C}_5\text{H}_5$), 431–8

Metal cluster-induced electrophilicity of arenes, 481–8

Polyhedral rearrangements in undeca- and dodeca-nuclear Group 1B metal osmium clusters, 505–10

Addition of anilines to the $[\text{Fe}(\text{CO})_3(1\text{-5}-\eta\text{-C}_7\text{H}_9)]\text{BF}_4^-$ complex, and the ‘Ordered Transition State Mechanism’, 561–6

Reactions of hydride complexes of ruthenium(ii) with alkynes, 589–94

Chemistry of multi-site bound unsaturated ligands on metal carbonyl clusters. Intramolecular reductive coupling in the conversion of $[\text{Ru}_3(\mu\text{-H})(\text{CO})_8(\mu_3,\eta^2\text{-C}\equiv\text{CBu}^i)]$ to the butadienyl cluster $[\text{Ru}_3(\text{CO})_8(\mu_3,\eta^4\text{-C}(\text{Bu}^i)\text{CC}(\text{H})\text{C}(\text{Ph})(\text{PPh}_2))]$, and its crystal structure. An example of ligand isomerism on metal clusters, 601–6

Carbonyl, hydridocarbonyl, and hydridoalkoxycarbonyl complexes of iridium with phosphorus–sulphur hybrid ligands, and the crystal and molecular structure of $[\text{IrH}(\text{CO}_2\text{Me})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{SEt})_2]\text{BPh}_4^-$, 607–14

Studies on olefin-co-ordinating transition-metal carbene complexes.

Part 14. An unusual reaction of tricarbonyl(η^4 -cyclopenta-1,3-diene)iron: a new approach to acyldicarbonyl(η -cyclopentadienyl)iron complexes, and the X-ray crystal structure of $[\text{Fe}(\text{COC}_6\text{H}_4\text{CF}_3\text{-}p)(\eta\text{-C}_5\text{H}_5)(\text{CO})_2]$, 635–8

Synthesis of diphenylphosphido-bridged manganese–gold complexes. Crystal structures of $[\text{N}(\text{PPh}_3)_2][\text{Au}_2\text{Mn}(\mu\text{-PPh}_2)_2(\text{CO})_4(\text{C}_6\text{F}_5)_2]$ and $[\text{Au}_2\text{Mn}_2(\mu\text{-PPh}_2)_4(\text{CO})_8]$, 639–44
 Trinuclear mixed-valence complexes with the linear $[\text{Rh}_3]^{4+}$ core, 689–92

Solvent influence on trifluoromethylation reactions with $\text{Cd}(\text{CF}_3)_2$: one-pot synthesis of trifluoromethyl iron complexes, 739–44

New carbide clusters in the cobalt subgroup. Part 17. Preparation and structural characterization of the mixed-metal octahedral dianion $[\text{Co}_2\text{Rh}_4\text{C}(\text{CO})_{13}]^{2-}$ as its $[\text{PPh}_3]^+$ salt, 879–84

Some transition-metal complexes of 1,3,2-dithiarsolane, 885–8

The mechanisms of fluxionality of $[\text{Ir}_4(\text{CO})_{11}(\text{PEt}_3)]$: a reappraisal of the mechanism of carbonyl fluxionality on clusters, 889–94
 Reactivity of cationic molybdenum(ii) complexes. Part 4. Isolation and crystal structure determination of *cis*- $[\text{Mo}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)(\text{PPh}_3)(\text{N}_3)]$ and *cis*- $[\text{Mo}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)(\text{PPh}_3)(\text{NCO})]$, 959–64

Crystal structures of di-iron $\mu\text{-}\eta^2\text{-}(\text{C},\text{C})$ -ketene complexes, $[(\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2)_2(\mu\text{-CH}_2\text{CO})]$ and $[(\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2)_2(\mu\text{-CH}_2\text{CO})]$. Synergistic metal-to-ligand back donation as revealed by comparison with $[(\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2)_2(\mu\text{-CH}_2\text{CH}_2\text{CO})]$, 1083–8

Synthesis and reactivity of binuclear pyrazolate-bridged ruthenium(i) complexes. Crystal structures of $\text{bis}[\mu\text{-}(3,5\text{-dimethylpyrazolato-}NN')\text{tricarbonylruthenium(i)}](\text{Ru}-\text{Ru})$ and $\text{bis}[\mu\text{-}(3,5\text{-dimethylpyrazolato-}NN')]\mu\text{-iodobis}[\text{tricarbonylruthenium(i)}]\mu\text{-iodide}$, 1093–100

Synthesis and characterization of pentanuclear Fe–Ir carbonyl clusters. Solid-state structures of $[\text{N}(\text{PPh}_3)_2][\text{FeIr}_4(\text{CO})_{15}]$ and $[\text{PPh}_4][\text{Fe}_2\text{Ir}_3(\text{CO})_{14}]$, 1109–16

Synthesis and characterization of a family of nitrile-substituted triangular hydridocarbonyl rhenium clusters, $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{12-n}(\text{NCMe})_n]$ ($n = 1\text{---}3$). X-Ray crystal structures of $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{11}(\text{NCMe})]$ and $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_9(\text{NCMe})_3]$, 1143–8

The heteronuclear cluster chemistry of the Group 1B metals. Part 11. Effect of the nature of the bidentate diphosphine ligand on the metal framework structures of the gold heteronuclear cluster compounds $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})(\mu\text{-H})\{\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}(\text{CO})_{12}]$ ($n = 1\text{---}6$). X-Ray crystal structures of $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})(\mu\text{-H})\{\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}(\text{CO})_{12}]$ ($n = 1$ or 2), 1227–36

Solution structure and dynamic behaviour of two isomers of $[\text{Fe}_3(\text{CO})_9(\text{POR}_3)_3]$ ($\text{R} = \text{Me}$, Et , or Ph) derivatives, 1277–82

Reactions of co-ordinated ligands. Part 46. Formation of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}=\text{CMe}_2)(\text{CO})_{10}]$ by carbon–carbon σ -bond and carbon–hydrogen activation of 3,3-dimethylcyclopropene; crystal and molecular structure of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and evidence for its facile rearrangement to $[\text{Os}_3(\mu\text{-H})(\mu\text{-CH}=\text{CHCHMe}_2)(\text{CO})_{10}]$, 1333–40

Heteronuclear transition metal–alkyne clusters. Part 1. Reactions of tris(alkyne)monocarbonyltungsten complexes with octacarbonyldicobalt. Synthesis, X-ray crystal structure, and two-dimensional nuclear magnetic resonance studies of $[\text{WCo}_2(\mu\text{-C}_2\text{Et}_2)(\mu\text{-C}_4\text{Et}_4)(\text{CO})_8]$, 1375–80

Synthesis, structure, and reactivity of the heteronuclear clusters $[\text{Os}_5\text{H}(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}]$ and $[\text{Os}_5(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}]$ ($\text{M} = \text{Cu}$, Ag , or Au ; $\text{R} = \text{Ph}$ or Me); the X-ray crystal structure of $[\text{Os}_5\text{H}(\text{CO})_{15}\{\text{Au}(\text{PPh}_3)\}]$, 1421–6

Crystal and molecular structures of μ -hydrido-bis[*fac*-(2,2'-bipyridine)tricarbonylrhenium(i)] chloride, *fac*-(2,2'-bipyridine)tricarbonylformatorhenium(i), and *fac*-(2,2'-bipyridine)tricarbonyl(cyanotrityrhodoborato)rhenium(i), 1449–54

Synthesis of the cluster dianion $[\text{Os}_{11}\text{C}(\text{CO})_{27}]^{2-}$ by pyrolysis and its reactions with electrophiles; X-ray structure analysis of the mixed-metal derivative $[\text{PMePh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\{\text{Cu}(\text{NCMe})\}]\cdot\text{CH}_2\text{Cl}_2$ and the hydrido derivative $[\text{PMePh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455–64

Synthesis and crystal structure of $[\text{NiOs}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2\{\mu\text{-C}(\text{PPh}_3)\}(\text{CO})_9]$. Hydrogenation–dehydrogenation reactions of dienes and alcohols with heterogeneous catalysts derived from $[\text{NiM}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-X})(\text{CO})_9]$ ($\text{M} = \text{Ru}$, $\text{X} = \text{H}$; $\text{M} = \text{Os}$, $\text{X} = \text{Cu}(\text{PPh}_3)$], 1477–82

Chemistry of phosphido-bridged dimolybdenum complexes. Part 3. Reinvestigation of the reaction between $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_6]$ and P_2Ph_4 . X-ray structures of $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\text{CO})_2]$,

CARBONYL (contd)

- [$\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPPh}_2)_2(\mu\text{-CO})$] and *trans*-[$\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPPh}_2)_2\text{O}(\text{CO})_2$], 1555–64
 Insertion of phenylethyne-thiolate or -selenolate into metal–carbene bonds: synthons for co-ordinated thioacyl or selenoacyl anions and for anionic thiocarbene or selenocarbene complexes, 1565–78
 Molecular structure of heterobimetallic $\mu\text{-}\eta^2\text{-(C,C)-ketene}$ complexes, 1627–30
 Organic chemistry of dinuclear metal centres. Part 12. Synthesis, X-ray crystal structure, and reactivity of the di- μ -alkylidene complex [$\text{Ru}_2(\text{CO})_2(\mu\text{-CHMe})(\mu\text{-CMe}_2)(\eta\text{-C}_5\text{H}_5)_2$]: alkylidene linking, 1799–806
 Organic chemistry of dinuclear metal centres. Part 13. Synthesis, structure, and reactivity of [$\text{Ru}_2(\text{CO})_4(\eta^5\text{:}\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{C}_5\text{H}_4)_2$], 1807–14
 The mechanisms of fluxionality of [$\text{Fe}_3(\text{CO})_{12-n}(\text{P}(\text{OR})_3)_n$] ($\text{R} = \text{Me}$, $n = 0\text{--}3$; $\text{R} = \text{Pr}^\text{i}$, $n = 2$ or 3) and the X-ray structure of [$\text{Fe}_3(\text{CO})_{10}(\text{P}(\text{OMe})_3)_2$], 1831–44
 Synthetic and structural studies on organomolybdenum and organotungsten complexes containing bismuth. X-Ray crystal structures of [$\text{BiCl}\{\text{Mo}(\text{CO})_3(\eta\text{-C}_5\text{H}_5)\}_2$] and [$\text{BiCl}\{\text{Mo}(\text{CO})_2(\text{CNBu}^\text{i})(\eta\text{-C}_5\text{H}_5)\}_2$] (1988, 2941), 1887
 Photofragmentation kinetics of some triruthenium carbonyl clusters, 1933–40
 Chemistry of [$\text{Cr}(\text{cp})(\text{CO})_3)_2$] ($\text{cp} = \eta^5\text{-C}_5\text{H}_5$). Reaction with elemental P_4 . A high yield synthesis and crystal structures of [$\text{Cr}(\text{cp})(\text{CO})_2(\mu\text{-}\eta^2\text{-P}_2)$] and [$\text{Cr}(\text{cp})(\text{CO})_2(\eta^3\text{-P}_3)$], 1951–6
 Cluster activation *via* cyclometallation. Formation of the cyclometallated triosmium carbonyl cluster [$\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4\text{Ph})\}$] and its reactions with phosphorus donor ligands. The X-ray crystal structures of [$\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4\text{Ph})\}$] and [$\text{Os}_3\text{H}(\text{CO})_9\{\text{P}(\text{OMe})_3\}(\text{PPPh}_2)$], 2023–32
 Reactions of the cyclometallated triosmium cluster [$\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4\text{Ph})\}$] with Brønsted acids and dihydrogen: synthesis and X-ray crystal structure of [$\text{Os}_3\text{H}_2(\text{CF}_3\text{CO}_2)(\text{CO})(\mu\text{-PPPh}_2)$], 2033–8
 The heteronuclear cluster chemistry of the Group 1B metals. Part 12. Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds [$\text{M}_2\text{Ru}_4(\mu_3\text{-H}_2)(\mu\text{-L})(\text{CO})_{12}$] [$\text{M} = \text{Cu}$ or Ag , $\text{L} = \text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPPh}_2$ ($n = 1\text{--}6$) or *cis*- $\text{Ph}_2\text{PCH}=\text{CHPPPh}_2$; $\text{M} = \text{Cu}$, $\text{L} = \text{Ph}_2\text{As}(\text{CH}_2)_n\text{PPPh}_2$ ($n = 1$ or 2)] in solution, 2039–48
 Reduction–oxidation properties of organotransition-metal complexes. Part 30. Triazenido-bridged carbonylrhodium bipyridyl complexes with [$\text{Rh}_2]^{2+}$, [$\text{Rh}_2]^{3+}$, and [$\text{Rh}_2]^{4+}$ cores, and the X-ray structures of [$\text{Rh}_2(\text{CO})_2(\text{bipy})(\mu\text{-RNNNR})_2[\text{BF}_4^-]\text{CH}_2\text{Cl}_2$ and [$\text{Rh}_2(\text{CO})(\text{bipy})(\mu\text{-RNNNR})_2][\text{PF}_6^-]_2\cdot2.5\text{CH}_2\text{Cl}_2$ ($\text{R} = p\text{-tolyl}$), 2049–60
 Characterisation of tridequate macrocyclic complexes of molybdenum(0) with tetra-aza macrocycles. Crystal structure of [$\text{Mo}(\text{CO})_4\text{L}'$] ($\text{L}' = \text{C-meso-5,5,7,12,12,14-hexamethyl-1,4,8,11-tetra-azacyclotetradecane}$), 2183–6
 Transition metal–cyanocarbon chemistry. Part 8. Cyano-substituted buta-1,3-dienylidene bridged di-iron complexes from unprecedented cyanoalkyne insertion into the carbon–hydrogen bond of a bridging alkenylidene ligand, 2187–92
 Iron carbene complexes. Part 3. Aminolysis of di-iron alkoxy-carbene complexes with ammonia and dimethylamine. X-Ray crystal structure of [$\text{Fe}_2(\mu\text{-SPh})_2(\text{CO})_5(\text{C}(\text{NM}_2)\text{Ph})$], 2199–204
 Stepwise conversion of [$\text{Os}(\text{CF}_3\text{CO}_2)_2(\text{CO})_4$] into [$\text{Os}(\text{pyS})_2(\text{CO})_2$]: X-ray crystal structures of the complexes [$\text{Os}(\text{pyS})_2(\text{CO})_x$] where $x = 2$ or 3 and pyS is the pyridine-2-thionato ligand, 2211–6
 The 4-phenyl-1,2,3,5-dithiadiazole complex [$\text{Fe}_2(\text{CO})_6(\text{PhCN}_2\text{S}_2)$]: its preparation, crystal structure, and an extended-Hückel molecular-orbital study of the bonding, 2229–34
 Kinetics of reactions of hydrogen with [$\text{Os}_3(\text{CO})_{11}(\text{NCMe})$] and [$\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2$], 2235–40
 Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 92. Synthesis of alkynylmethylidyne-molybdenum and –tungsten complexes and their reactions with [$\text{Co}_2(\text{CO})_8$] and [$\text{Mo}_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2$], 2261–8
 Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 93. Synthesis of heteropolynuclear metal compounds with chains of seven to eleven metal atoms; crystal structure of [$\text{Mo}_2\text{W}_3\text{Pt}_6(\mu_3\text{-CMe})_3(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_3(\text{CO})_{10}(\text{PMc}_2\text{Ph})_4(\eta\text{-C}_5\text{H}_5)_5$], 2269–80
 Chemistry of polynuclear metal complexes with bridging carbene

or carbyne ligands. Part 94. Synthesis of di- and tri-metal compounds *via* the reagents [$\text{M}(\equiv\text{C}-\text{C}\equiv\text{CBu}^\text{i})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)$] ($\text{M} = \text{Mo}$ or W), 2281–8

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 95. Synthesis and crystal structures of [$\text{W}_2\text{RuPt}(\mu\text{-CC}_6\text{H}_4\text{Me}-4)(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)(\text{CO})_6(\text{PMe}_2\text{Ph})(\eta\text{-C}_5\text{H}_5)_2$] and [$\text{W}_2\text{Ru}_2\text{Pt}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\mu\text{-CO})_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2$], 2289–96

A convenient route to binuclear osmium(i) complexes containing diphosphine and arsinophosphine ligands. X-Ray crystal structure of [$\text{Os}_2(\text{MeCO}_2)_2(\text{CO})_4(\text{dppm})_2$], 2303–8

Mixed-metal carbido carbonyl clusters. Part 4. Synthesis and structural characterization of [$\text{Rh}_6\text{C}(\text{CO})_{13}\{\text{Au}_2(\text{PPPh}_3)_2\}$], 2343–6

Preparation and X-ray crystal structure of [$\text{Ru}_3\text{RhH}_3(\text{CO})_{12}$], 2449–52

CARBOXAMIDE

Copper(ii)-assisted hydrolysis of 2,4,6-tris(2-pyridyl)-1,3,5-triazine. Part 2. Crystal structures of [bis(2-pyridylcarbonyl)amido]pyridine-2-carboxamide)copper(ii) trifluoromethanesulphonate and [bis(2-pyridylcarbonyl)amido][2,4,6-tris(2-pyridyl)-1,3,5-triazine]copper(iii) trifluoromethanesulphonate, 1681–8

CARBOXYLATE

Highly polydentate ligands. Part 4. Crystal structures of neodymium(iii) and erbium(iii) complexes of 3,12-bis(carboxymethyl)-6,9-dioxa-3,12-diazatetradecanoate(4–), 185–92

Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN'N'*-tetra-acetate (cdta). Part 3. Structural and magnetic characterization of the dinuclear [$\text{M}(\text{OH}_2)_5\text{[M}'(\text{cdta})]\text{H}_2\text{O}$] ($\text{M}, \text{M}' = \text{Ni}, \text{Ni}; \text{Mn}, \text{Ni}; \text{Mn}, \text{Cu}; \text{Co}, \text{Ni}$; or Co, Cu) and the tetrานuclear [$\text{M}(\text{OH}_2)_4\text{[M}'(\text{cdta})(\text{OH}_2)]\cdot4\text{H}_2\text{O}$] ($\text{M}, \text{M}' = \text{Zn}, \text{Zn}; \text{Zn}, \text{Co}; \text{Co}, \text{Co}$; or Mn, Co) complexes, 863–72

An investigation of the bonding in [$\text{Ru}_2(\text{O}_2\text{CCF}_3)_4$] and [$\text{Ru}_2(\text{O}_2\text{CCF}_3)_4(\text{NO})_2$], 1037–44

Preparation and characterisation of linear mixed-metal trinuclear carboxylate complexes of general formula [$\text{M}_2\text{M}'(\text{O}_2\text{CR})_6\text{-}(\text{C}_9\text{H}_7\text{N})_2$]; crystal structure of [$\text{Zn}_2\text{Ba}(\text{O}_2\text{CCMe}_3)_6(\text{C}_9\text{H}_7\text{N})_2$], 1127–32

Phenyltin trichloroacetate complexes: preparations and structures, 1589–98

Relationship between the torsion angles of acetate chelate rings and the deuteron nuclear magnetic resonance chemical shifts for polyamine-N-carboxylatochromate(iii) complexes, 1947–50

Electrochemistry of the quasi-reversible bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate(–v) and (–iv) and bis[2-hydroxy-2-methylbutanoato(2–)]oxochromate(–v) and (–iv) redox couples and the crystal and molecular structure of sodium bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate(v) sesquihydrate, 2205–10

Stepwise conversion of [$\text{Os}(\text{CF}_3\text{CO}_2)_2(\text{CO})_4$] into [$\text{Os}(\text{pyS})_2(\text{CO})_2$]: X-ray crystal structures of the complexes [$\text{Os}(\text{pyS})_2(\text{CO})_x$] where $x = 2$ or 3 and pyS is the pyridine-2-thionato ligand, 2211–6

A convenient route to binuclear osmium(i) complexes containing diphosphine and arsinophosphine ligands. X-Ray crystal structure of [$\text{Os}_2(\text{MeCO}_2)_2(\text{CO})_4(\text{dppm})_2$], 2303–8

Oxidative addition of dibenzoyl peroxide on metallic copper. The crystal structure of [$\text{Cu}(\text{C}_6\text{H}_5\text{CO}_2)_2(\text{py})_2$], 2331–4

Copper anti-inflammatory drugs in rheumatoid arthritis. Part 2. A potentiometric and spectroscopic study of copper(ii) polyaminodicarboxylate complexes, 2429–34

Synthesis, spectroscopic, electrochemical, and magnetic properties of dimolybdenum(II,III) diruthenium-(II,III) and -(II,II) complexes containing bridging aspirinate (2-acetoxybenzoate) ligands, 2435–42

CARBOXYLIC ACID

Thermodynamics of the formation of metal complexes of 1,4,10-trioxa-7,13-diazacyclopentadecane-*N,N'*-diacetic acid and of 1,4,10,13-tetraoxa-7,16-diazacyclo-octadecane-*N,N'*-diacetic acid, 133–8

Cyclometallation of unsaturated carboxylic acids by pentamethylcyclopentadienyl-rhodium and -iridium complexes.

Crystal structures of [$(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh}=\text{CHCO}-\text{O})(\text{Me}_2\text{SO})$] and [$(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh}=\text{CHCO}-\text{OMe})\text{I}$]⁺, 2415–22

CATALYSIS

Hydrogen production from ethanol catalysed by Group 8 metal complexes, 489–96

CATALYSIS (contd)

- Catalytic oxidation of water mediated by rhenium oxides, 849–52
 Studies on transition–metal oxo and nitrido complexes. Part 10. New oxo–ruthenium and oxo–osmium pyridine complexes, and use of the former as catalysts for oxidation of alcohols, 901–6
 Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(III)–ethanol–carbon tetrachloride system, 1197–202
 Synthesis and crystal structure of $[\text{NiOs}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2\{\mu\text{-Cu}(\text{PPPh}_3)\}(\text{CO})_9]$. Hydrogenation–dehydrogenation reactions of dienes and alcohols with heterogeneous catalysts derived from $[\text{NiM}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-X})(\text{CO})_9]$ [$\text{M} = \text{Ru}, \text{X} = \text{H}; \text{M} = \text{Os}, \text{X} = \text{Cu}(\text{PPPh}_3)$], 1477–82
 Comparative study on the multielectron reduction of $\text{HOCH}_2\text{H}_2\text{N}_3$ catalyzed by $[\text{NBu}^n]_4[\text{MoFe}_3\text{S}_4(\text{SPh})_3(o\text{-O}_2\text{C}_6\text{Cl}_4)]_2$ - and $[\text{NBu}^n]_4[\text{Fe}_4\text{S}_4(\text{SPh})_4]$ -modified glassy carbon electrodes, 1547–54

Cationic dinuclear rhodium complexes as catalyst precursors for the hydroformylation of alkenes, 1579–82

Kinetics of the uncatalysed and silver(I)-catalysed oxidation of hydrogen peroxide by [ethylenebis(biguanide)]silver(III) ion in acid perchlorate media, 1645–50

CATECHOLATE

One-electron transfer processes in the reaction of tin(II) halides with substituted *o*-quinones; crystal structure of bis(3,5-di-t-butylcatecholate)tin-1,10-phenanthroline-dimethylformamide (1/1/2), 439–46

CHAIN

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 85. Synthesis of chain and ring compounds containing molybdenum, 785–96

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 93. Synthesis of heteropolynuclear metal compounds with chains of seven to eleven metal atoms; crystal structure of $[\text{Mo}_2\text{W}_3\text{Pt}_6(\mu_3\text{-CMe})_3(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\text{CO})_{10}(\text{PMe}_2\text{Ph})_4(\eta\text{-C}_5\text{H}_5)_5]$, 2269–80

CHARGE DENSITY

Charge density in the $[\text{Mo}(\text{NCS})_6]^{3-}$ ion in $\text{Cs}_3\text{Mo}(\text{NCS})_6$, 1779–86

CHLORIDE

A further investigation of the polarized single-crystal electronic absorption spectrum of $[\text{AsPh}_4][\text{CrOCl}_4]$, 1–4

An investigation of halide abstraction by arsenic(III); reactions of AsCl_3 with MCl_2 ($\text{M} = \text{Mg}$ or Zn) and TiCl_4 in donor solvents (MeCN or POCl_3). Formation and crystal structure of the dimeric titanium(IV) compound $[\text{Ti}_2\text{Cl}_6(\text{POCl}_3)_2(\text{PO}_2\text{Cl}_2)_2]$, 57–60

Kinetics of the oxidation of chloride ions by tris(2,2'-bipyridine)nickel(III) ions in aqueous perchlorate media, 151–4

Tertiary phosphine complexes of triply bonded ditungsten bis(t-butylamide) tetrachloride. The X-ray crystal structures of $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMe}_3)_2]$, $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PEt}_3)_2]$, and *cis*- $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMe}_2\text{Ph})_2]$, 1537–42

On the existence of the anion trichloro(η -cyclopentadienyl)titanate(1–). Crystal structure of the salt $[\text{Co}(\text{C}_5\text{H}_5)_2][\text{Ti}(\text{C}_5\text{H}_5)\text{Cl}_3]$, 2453–6

CHLOROMETHYL

Preparation of (chloromethyl)palladium(II) derivatives from complexes of palladium dichloride by reaction with diazomethane or bis(chloromethyl)mercury, 761–6

CHROMATE

A reinvestigation of the structures of organotin sulphates and chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74

CHROMIUM

A further investigation of the polarized single-crystal electronic absorption spectrum of $[\text{AsPh}_4][\text{CrOCl}_4]$, 1–4

Chromium chalcogen complexes. Insertion of elemental sulphur into $[\text{Cr}(\text{cp})(\text{CO})_2\text{Se}]$ and $[\text{Cr}(\text{cp})_2(\text{CO})_4\text{Se}_2]$ ($\text{cp} = \eta\text{-C}_5\text{H}_5$), 431–8

Formation in solution, synthesis, and electrochemical study of oxalato complexes of *N,N'*-ethylenebis(salicylideneiminato)-chromium(III) and -iron(III); crystal structures of piperidinium [*N,N'*-ethylenebis(salicylideneiminato)](oxalato-O¹O²)-chromate(III) and -ferrate(III), 729–38

Formation of chromium interlayers in α -zirconium phosphate, 895–900

Stacked planar molecules in antiferromagnetic di-

isothiocyanatobis(thiourea)chromium(II), a new structural type for chromium(II), 1631–4

Relation between the torsion angles of acetate chelate rings and the deuteron nuclear magnetic resonance chemical shifts for polyamine-N-carboxylatochromate(III) complexes, 1947–50

Chemistry of $[\text{Cr}(\text{cp})(\text{CO})_2\text{P}_4]$ ($\text{cp} = \eta^5\text{C}_5\text{H}_5$). Reaction with elemental P₄. A high yield synthesis and crystal structures of $[\text{Cr}(\text{cp})(\text{CO})_2\text{P}_4(\mu\text{-n}^2\text{-P}_2)]$ and $[\text{Cr}(\text{cp})(\text{CO})_2(\eta^3\text{-P}_3)]$, 1951–6

Electrochemistry of the quasi-reversible bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate-(v) and -(iv) and bis[2-hydroxy-2-methylbutanoato(2–)]oxochromate-(v) and -(iv) redox couples and the crystal and molecular structure of sodium bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate(v) sesquihydrate, 2205–10

CIRCULAR DICHROISM

Synthesis and spectroscopic studies of cobalt(III) complexes of the chiral pendant-arm macrocycle *N,N',N''*-tris[(2S)-2-hydroxypropyl]-1,4,7-triazacyclonane (H_3L); crystal structure of the hydrogen-bridged dimer $[\text{CoL}_3\text{LCo}][\text{PF}_6]_3$, 931–6

Co-ordination of nickel(II) ions by angiotensin II and its peptide fragments. A potentiometric, proton nuclear magnetic resonance and circular dichroism spectroscopic study, 1471–6

Optical absorption and magnetic circular dichroism studies of hydrogen, copper(II), zinc(II), nickel(II), and cobalt(II) crown ether-substituted monomeric and dimeric phthalocyanines, 2397–406

CLATHRATE

Studies in Werner clathrates. Part 9. Structures with bis(isothiocyanato)tetrakis(4-phenylpyridine)nickel(II) as host, 119–26

CLUSTER

Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene (C=CHOEt) and ethoxyethyne ($\text{CH}\equiv\text{COEt}$) ligands respectively, 5–12

Redox activation of the hexanuclear ruthenium cluster $[\text{Ru}_6\text{C}(\text{CO})_{16}]^{2-}$ using oxidative addition, 243–6

Metal cluster-induced electrophilicity of arenes, 481–8

Polyhedral rearrangements in undeca- and dodeca-nuclear Group 1B metal osmium clusters, 505–10

Gold-197 Mössbauer spectra and the bonding of some gold–gold and gold–platinum clusters, 539–44

Chemistry of multi-site bound unsaturated ligands on metal carbonyl clusters. Intramolecular reductive coupling in the conversion of $[\text{Ru}_3(\mu\text{-H})(\text{CO})_9(\mu_3\text{-n}^2\text{-C}\equiv\text{CBu}')]^+$ to the butadienyl cluster $[\text{Ru}_3(\text{CO})_8(\mu_3\text{-n}^4\text{-C}(\text{Bu}')\text{CC}(\text{H})(\text{C}(\text{Ph})(\text{PPh}_2))}]^+$, and its crystal structure. An example of ligand isomerism on metal clusters, 601–6

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 85. Synthesis of chain and ring compounds containing molybdenum, 785–96

New carbide clusters in the cobalt subgroup. Part 17. Preparation and structural characterization of the mixed-metal octahedral dianion $[\text{Co}_2\text{Rh}_4\text{C}(\text{CO})_{14}]^{2-}$ as its $[\text{PPh}_4]^+$ salt, 879–84

The mechanisms of fluxionality of $[\text{Ir}_4(\text{CO})_{11}(\text{PET}_3)]$; a reappraisal of the mechanism of carbonyl fluxionality on clusters, 889–94

Lewis base adducts of Group 11 metal(I) complexes. Part 48.

Heterobimetallic tris(dithiocarbamato)cobalt(III)–copper(I) halide adducts as a vehicle for novel copper(I) halide clusters (structurally characterized), 971–6

Synthesis and characterization of pentanuclear Fe–Ir carbonyl clusters. Solid-state structures of $[\text{N}(\text{PPh}_3)_2][\text{FeIr}_4(\text{CO})_{15}]$ and $[\text{PPh}_4][\text{Fe}_2\text{Ir}_3(\text{CO})_{14}]$, 1109–16

Synthesis and characterization of a family of nitrile-substituted triangular hydridocarbonyl rhenium clusters, $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{12-n}(\text{NCMe})_n]$ ($n = 1\text{--}3$). X-Ray crystal structures of $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{11}(\text{NCMe})]$ and $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_9(\text{NCMe})_3]$, 1143–8

The heteronuclear cluster chemistry of the Group 1B metals. Part 11. Effect of the nature of the bidentate diphosphine ligand on the metal framework structures of the gold heteronuclear cluster compounds $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})\{\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}(\text{CO})_{12}]$ ($n = 1\text{--}6$). X-Ray crystal structures of $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})\{\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2\}(\text{CO})_{12}]$ ($n = 1\text{ or }2$), 1227–36

Solution structure and dynamic behaviour of two isomers of $[\text{Fe}_3(\text{CO})_9(\text{P}(\text{OR}_3))_3]$ ($\text{R} = \text{Me, Et, or Ph}$) derivatives, 1277–82

Reactions of co-ordinated ligands. Part 46. Formation of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}=\text{CMe}_2)(\text{CO})_{10}]$ by carbon–carbon σ -bond and carbon–hydrogen activation of 3,3-dimethylcyclopropene; crystal and

CLUSTER (contd)

molecular structure of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and evidence for its facile rearrangement to $[\text{Os}_3(\mu\text{-H})(\mu\text{-CH}=\text{CHCHMe}_2)(\text{CO})_{10}]$, 1333–40

Synthesis, structure, and reactivity of the heteronuclear clusters $[\text{Os}_3\text{H}(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}]$ and $[\text{Os}_5(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}_2]$ ($\text{M} = \text{Cu}, \text{Ag}, \text{or Au}; \text{R} = \text{Ph or Me}$); the X -ray crystal structure of $[\text{Os}_3\text{H}(\text{CO})_{15}\{\text{Au}(\text{PPh}_3)\}]$, 1421–6

Synthesis of the cluster dianion $[\text{Os}_{11}\text{C}(\text{CO})_{27}]^{2-}$ by pyrolysis and its reactions with electrophiles; X -ray structure analysis of the mixed-metal derivative

$[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\{\text{Cu}(\text{NCMe})\}]\cdot\text{CH}_2\text{Cl}_2$ and the hydrido derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455–64

Comparative study on the multielectron reduction of $\text{HOC}_2\text{H}_4\text{N}_3$ catalyzed by $[\text{NBu}^n]_4[\text{MoFe}_3\text{S}_4(\text{SPh})_3(o\text{-O}_2\text{C}_6\text{Cl}_4)]_2$ and $[\text{NBu}^n]_2[\text{Fe}_4\text{S}_4(\text{SPh})_4]$ -modified glassy carbon electrodes, 1547–54

The mechanisms of fluxionality of $[\text{Fe}_3(\text{CO})_{12-n}\{\text{P}(\text{OR})_3\}_n]$ ($\text{R} = \text{Me}, n = 0\text{--}3; \text{R}' = \text{Pr}^i, n = 2\text{ or }3$) and the X -ray structure of $[\text{Fe}_2(\text{CO})_{10}\{\text{P}(\text{OMe})_3\}_2]$, 1831–44

Photofragmentation kinetics of some triruthenium carbonyl clusters, 1933–40

Cluster activation *via* cyclometallation. Formation of the cyclometallated triosmium carbonyl cluster

$[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ and its reactions with phosphorus donor ligands. The X -ray crystal structures of $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ and $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{OMe})_3\}(\text{PPh}_2)]$, 2023–32

Reactions of the cyclometallated triosmium cluster

$[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ with Brønsted acids and dihydrogen; synthesis and X -ray crystal structure of $[\text{Os}_3\text{H}_2(\text{CF}_3\text{CO}_2)(\text{CO})_9(\mu\text{-PPh}_2)]$, 2033–8

The heteronuclear cluster chemistry of the Group 1B metals. Part 12. Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds $[\text{M}_3\text{Ru}_4(\mu_3\text{-H})_2(\mu\text{-L})(\text{CO})_{12}]$ [$\text{M} = \text{Cu}$ or Ag , $\text{L} = \text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2$ ($n = 1\text{--}6$) or *cis*- $\text{Ph}_2\text{PCH}=\text{CHPPPh}_2$; $\text{M} = \text{Cu}$, $\text{L} = \text{Ph}_2\text{As}(\text{CH}_2)_n\text{PPh}_2$ ($n = 1$ or 2)] in solution, 2039–48

Low-valent isocyanide complexes and clusters of palladium and platinum. Crystal structure of $[\text{Pt}\{\text{C}(=\text{NR})\text{N}(\text{R})\text{C}(=\text{NR})\}(\text{RN}\bar{\text{C}})_2]$ ($\text{R} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), 2161–6

Kinetics of reactions of hydrogen with $[\text{Os}_3(\text{CO})_{11}(\text{NCMe})]$ and $[\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2]$, 2235–40

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 94. Synthesis of di- and tri-metal compounds *via* the reagents $[\text{M}(\equiv\text{C}-\text{C}\equiv\text{CBu}^i)(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ ($\text{M} = \text{Mo}$ or W), 2281–8

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 95. Synthesis and crystal structures of $[\text{W}_2\text{RuPt}(\mu\text{-CC}_6\text{H}_4\text{Me-4})(\mu_3\text{-CC}_6\text{H}_4\text{Me-4})(\text{CO})_6(\text{PMe}_2\text{Ph})(\eta\text{-C}_5\text{H}_5)_2]$ and $[\text{W}_2\text{Ru}_2\text{Pt}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me-4})_2(\mu\text{-CO})_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2]$, 2289–96

Mixed-metal carbido carbonyl clusters. Part 4. Synthesis and structural characterization of $[\text{Rh}_6\text{C}(\text{CO})_{13}\{\text{Au}_2(\text{PPh}_3)_2\}]$, 2343–6

Preparation and X -ray crystal structure of $[\text{Ru}_3\text{RhH}_3(\text{CO})_{12}]$, 2449–52

COBALAMIN

Relative importance of the inductive and steric effects of alkyl groups (R) in understanding the nature of the $\text{Co}^{\text{III}}\text{-C}$ bond in alkyl cobaloximes, and comments on related alkyl cobalamins, 115–8

COBALOXIME

Relative importance of the inductive and steric effects of alkyl groups (R) in understanding the nature of the $\text{Co}^{\text{III}}\text{-C}$ bond in alkyl cobaloximes, and comments on related alkyl cobalamins, 115–8

COBALT

Tetra-aza macrocyclic complexes containing a pendant amino group. Copper(II), nickel(II), and cobalt(III) complexes of 5-aminomethyl-2,5,10,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane (L) and X -ray structural characterisation of the cobalt(III) complex $[\text{Co}(\text{L})\text{Cl}](\text{ClO}_4)\text{Cl}\cdot\text{EtOH}$, 85–92

Relative importance of the inductive and steric effects of alkyl groups (R) in understanding the nature of the $\text{Co}^{\text{III}}\text{-C}$ bond in alkyl cobaloximes, and comments on related alkyl cobalamins, 115–8

Metal complexes ($\text{Co}^{3+}, \text{Fe}^{3+}, \text{Mn}^{3+}$) of the quinquedentate salen analogue, 2-methyl-4-benzylthio-*N,N'*-butane-1,2-diylibis(salicylideneimine): crystal structures and sulphur coordination effects, 179–84

Possible biological activity in proton, metal, and aminohydroxamic acid equilibria. Protonation and complex formation reactions between *N*-hydroxy-D-asparagine and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution (1988, 953), 383 Co-ordination chemistry of higher oxidation states. Part 29.

Cobalt(III) complexes of bi- and tetra-thioethers: synthesis, properties, and cobalt-59 nuclear magnetic resonance data, 453–6

Base-catalysed aquation of $\alpha\beta\text{-syn}$ and $\alpha\beta\text{-anti}$ -chloro- and bromo-[1,9-bis(2'-pyridyl)-2,5,8-triazaundecane]- and -[1,11-bis(2'-pyridyl)-2,6,10-triazaundecane]-cobalt(III) cations. An unusually base-sensitive halogenopentamine cobalt(III) system, 645–54

Kinetics of aquation and base hydrolysis of some *trans*- $[\text{Co}(\text{tn})_2(\text{B})\text{Cl}]^{n+}$ ($\text{tn} = 1,3\text{-diaminopropane}; \text{B} = \text{OH}, \text{NCS}, \text{N}_3, \text{NO}_2, \text{NH}_3, \text{or CN}$) complexes. The effect of ring size on reactivity, 853–8

New carbide clusters in the cobalt subgroup. Part 17. Preparation and structural characterization of the mixed-metal octahedral dianion $[\text{Co}_2\text{Rh}_4\text{C}(\text{CO})_{13}]^{2-}$ as its $[\text{PPh}_4]^+$ salt, 879–84

Synthesis and spectroscopic studies of cobalt(III) complexes of the chiral pendant-arm macrocycle *N,N,N'*-tris[(2*S*)-2-hydroxypropyl]-1,4,7-triazacyclononane (H_3L); crystal structure of the hydrogen-bridged dimer $[\text{CoLH}_3\text{LCo}](\text{PF}_6)_3$, 931–6

Lewis base adducts of Group 11 metal(I) complexes. Part 48.

Heterobimetallic tris(dithiocarbamato)cobalt(III)-copper(I) halide adducts as a vehicle for novel copper(I) halide clusters (structurally characterized), 971–6

6,13-Diamino-6,13-dimethyl-1,4,8,11-tetra-azacyclotetradecane, L^7 , a new, potentially sexidentate polyamine ligand. Variable co-ordination to cobalt(III) and crystal structure of the complex $[\text{Co}(\text{L}^7)]\text{Cl}_2[\text{ClO}_4]$, 1059–66

Isotopic multiplets in the carbon-13 nuclear magnetic resonance spectra due to partial deuteration of co-ordinated NH_3 and NH groups of cobalt(III) complexes containing an N-glycoside. Importance of sugar ring oxygen atoms in the long-range isotope shifts, 1241–8

Thermodynamic and spectrophotometric investigation of complex formation between hydrogen, cobalt(II), nickel(II), and copper(II) ions and 2-amino-N-hydroxypropanamide in aqueous solution, 1299–310

Heteronuclear transition metal–alkyne clusters. Part 1. Reactions of tris(alkyne)monocarbonyltungsten complexes with octacarbonyldicobalt. Synthesis, X -ray crystal structure, and two-dimensional nuclear magnetic resonance studies of $[\text{WC}_{20}(\mu\text{-C}_5\text{Et}_2)_2(\mu\text{-C}_4\text{Et}_4)(\text{CO})_8]$, 1375–80

Synthesis and characterization of cobalt(III) and cobalt(IV) complexes of *N*-(2-hydroxyphenyl)salicylamide and homologues, 1613–6

Neutron diffraction study of the cation distributions in the systems $\text{Fe}_{7-x}\text{M}_x(\text{PO}_4)_6$ ($\text{M} = \text{Mn}$ or Co), 1765–70

Steric effect of the adamantyl ligand on crystal structure and dissociation rates in organocobalt B_{12} models, 1941–6

Rearrangements of $>\text{C}=\text{N}$ -functionalities promoted by cobalt(I): reactivity of cyclopentadienylbis(ethylene)cobalt(I) with dibenzophenone azine, benzophenone oxime, benzylidene(phenyl)amine and crystal structure determinations of the products, 1957–62

Spectral intensity and charge distributions in the tetrahedral chromophores, bis(3-amino-2-methylpropane-2-thiolato)- and diacetatobis(ethylenethiourea)-cobalt(II), 2109–14

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 92. Synthesis of alkynylmethylidyne-molybdenum and -tungsten complexes and their reactions with $[\text{Co}_2(\text{CO})_8]$ and $[\text{Mo}_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2]$, 2261–8

Functionalized amino acid complexation to a cobalt(III) complex of $(2R,5R,8R,11R)-2,5,8,11$ -tetraethyl-1,4,7,10-tetra-azacyclododecane. Crystal structures of complexes containing (*R*)- or (*S*)-serine and (*R*)- α -methylserine, and release of amino acid from the complexes, 2359–66

On the existence of the anion trichloro(η -cyclopentadienyl)titana(1–). Crystal structure of the salt $[\text{Co}(\text{C}_5\text{H}_5)_2]\text{Ti}(\text{C}_5\text{H}_5)\text{Cl}_3$, 2453–6

COBALT-59

Co-ordination chemistry of higher oxidation states. Part 29.

Cobalt(III) complexes of bi- and tetra-thioethers: synthesis, properties, and cobalt-59 nuclear magnetic resonance data, 453–6

CONDUCTIVITY

Crystal structure of the new low-dimensional metal $\text{Cs}_{0.82}[\text{Pd}(\text{S}_2\text{C}_2(\text{CN})_2)_2]\cdot 0.5\text{H}_2\text{O}$, 1101–4

COPPER

The interaction of 1-methylimidazoline-2(3*H*)-thione with copper(ii) salts, 39–42

Crystal structure, spectroscopic and magnetic properties of the complex $[\text{Cu}(\text{paphy})(\text{NCS})(\text{SCN})]$ (paphy = pyridine-2-carbaldehyde 2'-pyridylhydrazone). An unusual di- μ -thiocyanato-*N* bridged copper(ii) dimer, 53–6

Tetra-aza macrocyclic complexes containing a pendant amino group. Copper(ii), nickel(ii), and cobalt(iii) complexes of 5-aminomethyl-2,5,10,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane (L) and X-ray structural characterisation of the cobalt(iii) complex $[\text{Co}(\text{L})\text{Cl}]_2\text{ClO}_4\cdot \text{EtOH}$, 85–92

The unusual co-ordination ability of vasopressin-like peptides; potentiometric and spectroscopic studies of some copper(ii) and nickel(ii) complexes, 173–8

The copper(ii) promoted hydrolysis of benzylpenicillin. Evidence for the participation of a Cu-OH species in the hydrolysis of the β -lactam ring, 197–202

Crystal structure and magnetic properties of 1-aqua- μ -hydroxo-1,2,2-tris(perchlorato)-1,2-bis(2,2',6',2"-terpyridine)dicopper(ii), 237–42

The co-ordination chemistry of divalent cobalt, nickel, and copper. Part 13. Characterization and stability constants of copper(ii) species isolated from mixed-ligand solutions; crystal structure of the complex (pyridine-2,6-dicarboxylato)(*N,N,N',N'*-tetramethyl-1,3-diaminopropane)copper(ii), 253–8

Thiosemicarbazones as co-ordinating agents. Part 3. Synthesis, spectroscopic characterization, and X-ray structure of methyl pyruvate thiosemicarbazone hemihydrate, chloro(ethyl pyruvate thiosemicarbazone)copper(ii) (green form), and chloro(pyruvic acid thiosemicarbazone)copper(ii) dihydrate (blue form), 361–6

Possible biological activity in proton, metal, and aminohydroxamic acid equilibria. Protonation and complex formation reactions between *N*-hydroxy-D-asparagine and cobalt(ii), nickel(ii), copper(ii), and hydrogen ions in aqueous solution (1988, 953, 383)

Polyhedral rearrangements in undeca- and dodeca-nuclear Group 1B metal osmium clusters, 505–10

Interligand interaction in mixed amino acid complexes, 567–8

Spectrophotometric study of complex formation in copper(ii) mono-, di-, and tri-ethanolamine systems, 569–74

Tetrathiafulvalenium salts of planar Pt^{II}, Pd^{II}, and Cu^{III} 1,2-dithio-oxalato-S,S' anions. Synthesis, chemistry and molecular structures of bis(tetrathiafulvalenium) bis(1,2-dithio-oxalato-S,S')palladate(ii), $[\text{ttf}]_2[\text{Pd}(\text{S}_2\text{C}_2\text{O}_2)_2]$, and of bis(tetrathiafulvalenium)tetrathiafulvalene bis(1,2-dithio-oxalato-S,S')platinate(ii), $[\text{ttf}]_3[\text{Pt}(\text{S}_2\text{C}_2\text{O}_2)_2]$, 719–28

Synthesis and X-ray crystal structure of chloro[2(1*H*)-pyridinethione-S]-bis(triphenylphosphine)copper(i), 749–52

Crystal structure of a novel tetrานuclear dithiocarboxylato-copper(i) complex, $[\{\text{Cu}(\text{S}_2\text{CC}_6\text{H}_4\text{Me}-p)\}_4\text{PPh}_3]_2$, 753–6

Reactivity of metallic copper with acidic organic compounds in the presence of molecular oxygen. Crystal structure of $[\{\text{Cu}(\text{NCO})(\text{py})_2\}_2]\cdot \text{H}_2\text{O}$, 921–6

Synthesis and characterization of imidazolate-bridged iron(iii)-copper(ii) and manganese(iii)-copper(ii) binuclear complexes, 943–6

Lewis base adducts of Group 11 metal(i) complexes. Part 48.

Heterobimetallic tris(dithiocarbamato)cobalt(iii)-copper(i) halide adducts as a vehicle for novel copper(i) halide clusters (structurally characterized), 971–6

Magnetic properties of mono-, bi-, and tri-nuclear copper(ii) complexes of novel oxamato and oxamido ligands. Crystal structure of a mononuclear precursor, 1017–26

Complexes of aminophosphonates. Part 3. Copper(ii) complexes of several monophosphono and diphosphono dipeptides, 1053–8

Iridium(iii) hydrido complexes with 2-pyridinaldazines. Crystal structure of $[\text{IrH}_2(\text{mpaa})(\text{PPh}_3)_2]\text{SbF}_6$. Synthesis and characterization of homo- and hetero-nuclear complexes, 1075–82

Low-lying electronic states in μ -phenolato copper(ii) binuclear compounds with hydroxo, ethanolato, azido, and cyanato exogenous bridges: crystal structures, magnetic properties, and interpretations, 1117–26

Blue copper models. Spectroscopic and electrochemical studies of copper(ii) complexes with new ligand systems containing sulphur

and nitrogen donor atoms, 1161–70

Metal complexes as ligands. Part 12. Synthesis and X-ray crystal structure of a hexanuclear nickel(ii) complex based on two isosceles trinuclear components with imidazolate bridges, 1187–92

Reaction of $[\text{Rh}_2\{\mu\text{-1,8-(NH)}_2\text{C}_10\text{H}_6\}(\text{CO})_2(\text{PPh}_3)_2]$ with $[\text{M}(\text{PPh}_3)]^+$ ($\text{M} = \text{Au, Ag, or Cu}$): formation of unusual 44-electron Rh_2M clusters. Crystal structure of $[\text{Rh}_2\{\mu\text{-Au}(\text{PPh}_3)\}\{\mu\text{-1,8-(NH)}_2\text{C}_10\text{H}_6\}(\text{CO})_2(\text{PPh}_3)_2]\text{ClO}_4\cdot 2\text{CH}_2\text{Cl}_2$, 1249–52

Composition, stability, and lability of copper(ii) dipeptide complexes, 1293–8

Thermodynamic and spectrophotometric investigation of complex formation between hydrogen, cobalt(ii), nickel(ii), and copper(ii) ions and 2-amino-*N*-hydroxypropanamide in aqueous solution, 1299–310

Synthesis, spectroscopic, and magnetic studies of copper(ii) with *N*-protected amino acids: crystal and molecular structure of diaquatetrakis(*μ*-benzoyl-*α*-alaninato)-dicopper(ii), 1341–4

Autoxidation of a dicopper(i) macrocyclic complex: evidence for a mixed-valence intermediate, 1351–60

Synthesis, structure, and reactivity of the heteronuclear clusters $[\text{Os}_5\text{H}(\text{CO})_{15}|\text{M}(\text{PR}_3)|]$ and $[\text{Os}_5(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}_2]$ ($\text{M} = \text{Cu, Ag, or Au}; \text{R} = \text{Ph or Me}$): the X-ray crystal structure of $[\text{Os}_5\text{H}(\text{CO})_{15}|\text{Au}(\text{PPh}_3)|]$, 1421–6

Dimeric mixed-ligand copper(ii) complexes with di-imines and salicylaldehyde, 1435–8

Synthesis of the cluster dianion $[\text{Os}_{11}\text{C}(\text{CO})_{27}]^{2-}$ by pyrolysis and its reactions with electrophiles; X-ray structure analysis of the mixed-metal derivative

$[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\{\text{Cu}(\text{NCMe})\}]\cdot \text{CH}_2\text{Cl}_2$ and the hydrido derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455–64

Synthesis and crystal structure of $[\text{NiOs}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-Cu}(\text{PPh}_3))(\text{CO})_9]$. Hydrogenation–dehydrogenation reactions of dienes and alcohols with heterogeneous catalysts derived from $[\text{NiM}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-X})(\text{CO})_9]$ [$\text{M} = \text{Ru, X} = \text{H}; \text{M} = \text{Os, X} = \text{Cu}(\text{PPh}_3)$], 1477–82

Binuclear copper(i) complexes containing two or three 2-(diphenylphosphine)pyridine (dppy) bridging ligands: X-ray crystal structure of $[\text{Cu}_2(\mu\text{-dppy})_3(\text{MeCN})]\text{BF}_4$ and variable-temperature ^1H and ^{31}P - $\{^1\text{H}\}$ nuclear magnetic resonance studies, 1499–506

X-Ray structure of a ternary complex, copper(ii)-inosine 5'-monophosphate–benzimidazole. The first example for the absence of direct metal–nucleotide interaction, 1507–12

Functional metallocacycles derivatives and their polymers. Part 23. Biaxially orientated Langmuir–Blodgett films of substituted metallophthalocyanines, 1543–6

Lewis-base adducts of Group 11 metal(i) compounds. Part 58.

Syntheses, X-ray structures, and far-infrared spectra of the novel $[\text{CuX}_2(\text{bidentate ligand})]^-$ ($\text{X} = \text{Cl or Br}$) anion salts of the $[\text{Cu}(\text{dpa})_2]^+$ cation and the dimeric iodide $[(\text{dpa})\text{CuI}_2\text{Cu}(\text{dpa})]$ [$\text{dpa} = \text{di}(2\text{-pyridyl)amine}$], 1639–44

Copper(ii)-assisted hydrolysis of 2,4,6-tris(2-pyridyl)-1,3,5-triazine. Part 2. Crystal structures of $[\text{bis}(2\text{-pyridylcarbonyl})\text{amido}](\text{pyridine-2-carboxamide})\text{copper(ii)}$ trifluoromethanesulphonate and $[\text{bis}(2\text{-pyridylcarbonyl})\text{amido}][2,4,6\text{-tris}(2\text{-pyridyl)-1,3,5-triazine}]\text{copper(ii)}$ trifluoromethanesulphonate, 1681–8

Closely related macrocyclic and acyclic tridentate, pyridine derivatives, containing sulphur, and their complexes. Crystal structures of {dichloro-3,10-dithia-16-azabicyclo[10.3.1]hexadeca-1(16),12,14-triene}copper(ii) and $[\text{2,6-bis(ethylthiomethyl)pyridine}]$ dichlorocopper(ii), 1739–44

Mixed-metal complexes in solution. A potentiometric and spectrophotometric study of copper(ii)–nickel(ii)– δ -hydroxylysine complexes in aqueous solution, 1745–50

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 91. Steric implications in the reactions of the complexes $[\text{M}(\equiv\text{CR})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ ($\text{M} = \text{Mo or W, R} = \text{C}_6\text{H}_4\text{Me-4, C}_6\text{H}_4\text{OMe-2, or C}_6\text{H}_3\text{Me}_2\text{-2,6}$) with the compounds $[\text{Mo}(\text{CO})_3(\text{NCMe})_3]$ and $[\text{Cu}(\text{thf})(\eta\text{-C}_5\text{Me}_5)]$ ($\text{thf} = \text{tetrahydrofuran}$): crystal structure of $[\text{Mo}_3(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)]$, 1871–8

Electron spin resonance study of copper(ii) complexes of X-glycine and glycyl-X type dipeptides, and related tripeptides. Variation of co-ordination modes with ligand excess and pH in fluid and frozen aqueous solutions, 1925–32

Spectroscopic and electrochemical study of copper and zinc complexes with a N_2S_2 ligand donor set. Crystal structure of the

COPPER (contd)

- copper(II) complex derived from 1,3-propylenediamine and 3-formyl-1-phenyl-2(1*H*)-pyridinethione, 1979–86
- A barium dithiocyanate complex with an unusual macrocycle containing pendant arms, the amine precursor, and related copper(II) diperchlorate complexes: synthesis and structure, 1993–2002
- The heteronuclear cluster chemistry of the Group 1B metals. Part 12. Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds $[M_2Ru_4(\mu_3-H)_2(\mu-L)(CO)_{12}]$ [$M = Cu$ or Ag , $L = Ph_2P(CH_2)_nPPh_2$ ($n = 1–6$) or $Rs-Ph_2PCH=CHPPh_2$; $M = Cu$, $L = Ph_2As(CH_2)_nPPh_2$ ($n = 1$ or 2)] in solution, 2039–48
- Binuclear copper(II) complexes of new dinucleating ligands with a pyrazolate group as an endogenous bridge. Effects of exogenous azide and acetate bridges on magnetic properties, 2077–82
- Crystal structures and magnetic properties of bi- and tetra-nuclear copper(II) complexes of 2,6-disformyl-4-methylphenol di(benzoylhydrazone), 2083–8
- Synthesis, structure, and redox properties of dicopper(II,II) complexes of *N*-(hydroxyphenyl)salicylamide and homologues. Facile oxidation to dicopper(II,III) and -(III,III) species; crystal structure of bis(tetrabutylphosphonium) bis[{(2-oxidobenzoyl)(2'-oxidophenyl)amido(3-)NO²⁻,μ-O²⁻}cuprate(II)}, 2089–94
- Crystal structure and magnetic properties of the heterobinuclear complex $[(CH_3CN)LNi(mnt)Cu(mnt)]CH_3CN$ ($L =$ tetrabeno[b,f,j,n][1,5,9,13]tetra-azacyclohexadecine and mnt = maleonitriledithiolate), 2095–8
- Interaction of benzo-1,3-thiazoline-2-thione and related ligands with copper(II) salts and the single-crystal X-ray structure of μ -[1,2-bis(diphenylphosphino)ethane]-bis[(benzo-1,3-thiazole-2-thiolato-S)[1,2-bis(diphenylphosphino)ethane]copper(I)}, 2167–72
- Oxidative addition of dibenzoyl peroxide on metallic copper. The crystal structure of [{Cu(C₆H₅CO₂)₂(py)}₂], 2331–4
- Copper anti-inflammatory drugs in rheumatoid arthritis. Part 2. A potentiometric and spectroscopic study of copper(II) polyaminodicarboxylate complexes, 2429–34

CROWN ETHER

- The nature of a sodium–crown interaction studied by sodium-23 nuclear magnetic resonance and by crystal structure determination of a complex with co-ordinated solvent, 203–10
- Optical absorption and magnetic circular dichroism studies of hydrogen, copper(II), zinc(II), nickel(II), and cobalt(II) crown ether-substituted monomeric and dimeric phthalocyanines, 2397–406

CRYPTATE

- Thermodynamic and electrochemical behaviour of lanthanide cryptates in *N,N*-dimethylformamide, 367–70

CRYSTAL STRUCTURE

- Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene ($C=CHOEt$) and ethoxyethyne ($CH\equiv COEt$) ligands respectively, 5–12
- Lead(II) and barium(II) complexes of a potentially octadentate macrocyclic ligand (H_4L), capable of providing endogenous alkoxy and phenoxy bridges. The X-ray crystal structures of $[Pb(H_4L)][ClO_4]_2$, $[Pb_2(H_2L)][ClO_4]_2$, and $[H_6L](H_2O)_2][ClO_4]_2$, 19–24
- Rhodium complexes of the binucleating ligands pyridine-2-thiolate and benzothiazole-2-thiolate. Crystal structures of [{Rh(μ -SC₅H₄N)(CO)}₂] and [{Rh(μ -SC₅H₄N)(tfbb)}₂]·Me₂CO (tfbb = tetrafluorobenzobarrelene), 25–32
- Five-co-ordinated silicon compounds: geometry of formation by intramolecular co-ordination. Crystal structure of 2-(dimethylaminomethyl)phenyl-1-naphthylsilane, 43–52
- Crystal structure, spectroscopic and magnetic properties of the complex $[Cu(paphy)(NCS)(SCN)]$ (paphy = pyridine-2-carbaldehyde 2'-pyridylhydrazone). An unusual di- μ -thiocyanato-*N* bridged copper(II) dimer, 53–6
- An investigation of halide abstraction by arsenic(III): reactions of AsCl₃ with MCl₂ ($M = Mg$ or Zn) and TiCl₄ in donor solvents ($MeCN$ or $POCl_3$). Formation and crystal structure of the dimeric titanium(IV) compound $[Ti_2Cl_6(POCl_3)_2(PO_2Cl_2)_2]$, 57–60
- Comparison of the structures and properties of $[N(C_2H_5)_4][Pt\{S_2C_2(CN)_2\}_2]$ and $[H_3O]_x[NH_4]_{1-x}[Pt\{S_2C_2(CN)_2\}_2] \cdot 2 - xH_2O$, 61–6
- The crystal structures of $K[Pd\{S_2C_2(CN)_2\}_2]H_2O$ and $[NH_4][Pd\{S_2C_2(CN)_2\}_2]H_2O$, 67–72

- (Diphenylphosphinomethyl)diphenylphosphine sulphide ($Ph_2PCH_2PPh_2S$) and its methanide anion $Ph_2PCHPPh_2S^-$ as ligands in organogold chemistry. X-Ray crystal structure of $Rs-[Au(C_6F_5)_2(Ph_2CHPPh_2S)]$, 73–8
- Halogeno, hydride, and alkyl carbonyl-monocyclopentadienyl niobium derivatives. Crystal and molecular structure of $[Nb(\eta-C_5H_5)Cl(H)(CO)(PMo_3)_2]$, 79–84
- Tetra-aza macrocyclic complexes containing a pendant amino group. Copper(II), nickel(II), and cobalt(III) complexes of 5-aminomethyl-2,5,10,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane (L) and X-ray structural characterisation of the cobalt(III) complex $[Co(L)Cl][ClO_4]Cl\cdot EtOH$, 85–92
- Structural, spectroscopic, and electrochemical characterization of tetrakis- μ -(2-pyrrolidinonato)-dirhodium(II) and tetrakis- μ -(δ -valerolactamato)-dirhodium(II), 93–100
- Synthesis of α -, γ -phosphorus functionalized alkyl lithium species; X-ray structures of $[\{Li(L)(CH_2PMeR)\}_2]$ [$L = N,N,N',N'$ -tetramethylethylenediamine (tmen), $R = Me$ or Ph ; $L = (-)$ -sparteine, $R = Ph$] and $[\bar{Li}(tmen)\{CH(SiMe_3)C_6H_4\bar{P}Ph_2-o\}]$, 105–14
- Studies in Werner clathrates. Part 9. Structures with bis(isothiocyanato)tetrakis(4-phenylpyridine)nickel(II) as host, 119–26
- Synthesis, electrochemistry, and electron spin resonance spectra of some seventeen-electron complexes containing the $\{Mo(NO)\}^{2+}$ core and the molecular structure of $[Mo(NO)\{HB(Me_2pz)_3\}(pyr)_2]$, 127–32
- Chelate-stabilised primary, secondary, and tertiary alcohol and alkoxo platinum(II) complexes. X-Ray crystal structure of $[Pt(Ph_2PCH_2CH_2O)]H_2O$, 139–44
- Oxovanadium(IV) and dioxomolybdenum(VI) complexes: synthesis from the corresponding acetylacetone complexes and X-Ray structures, 145–50
- Reactivity of cationic molybdenum(II) complexes. Part 3. Synthesis of methoxide and carbomethoxide derivatives of η -pentamethylcyclopentadienyl-molybdenum(II) and the crystal structure determination of the oxidation product *trans*- $[\{Mo(\eta-C_5Me_5)(O)_2\}_2(\mu-O)]$, 155–60
- Metal complexes (Co^{3+} , Fe^{3+} , Mn^{3+}) of the quinquedentate salen analogue, 2-methyl-4-benzylthio-*N,N'*-butane-1,2-diylibis(salicylideneimine): crystal structures and sulphur co-ordination effects, 179–84
- Highly polydentate ligands. Part 4. Crystal structures of neodymium(III) and erbium(III) complexes of 3,12-bis(carboxymethyl)-6,9-dioxa-3,12-diazatetradecanedioate(4-), 185–92
- The nature of a sodium–crown interaction studied by sodium-23 nuclear magnetic resonance and by crystal structure determination of a complex with co-ordinated solvent, 203–10
- Palladium complexes with the tripodal phosphine tris(2-diphenylphosphinoethyl)amine. Synthesis and structure of trigonal, tetrahedral, trigonal bipyramidal, and square planar complexes, 211–6
- The structure of $PO(OPF_2)_3$ in the gaseous and crystalline phases, 223–8
- Correlations between nuclear magnetic resonance spectra and crystal structures. A carbon-13 nuclear magnetic resonance study in the solid state of the Group 15 xanthates $[A(S_2COR)_3]$ ($A = As^{III}$, Sb^{III} , or Bi^{III} ; $R = Me$, Et , or Pr^i), 233–6
- Crystal structure and magnetic properties of 1-aqua- μ -hydroxo-1,2,2-tris(perchlorato)-1,2-bis(2,2'-6,2''-terpyridine)dicopper(II), 237–42
- Synthesis and structural studies on silver–tin complex salts with bis(diphenylarsino)methane and bis(diphenylphosphino)methane, 247–52
- The co-ordination chemistry of divalent cobalt, nickel, and copper. Part 13. Characterization and stability constants of copper(II) species isolated from mixed-ligand solutions; crystal structure of the complex (pyridine-2,6-dicarboxylato)(*N,N,N',N'*-tetramethyl-1,3-diaminopropane)copper(II), 253–8
- Metallacyclic complexes. Part 2. Synthesis of metallathietane-3,3-dioxide complexes of platinum(II) and palladium(II): the crystal structures of $[Pt\{CH(COPh)S(O)_2CH(COPh)\}(PPh_3)_2]\cdot 2CH_2Cl_2$ and $[Pd\{CH(COPh)S(O)_2CH(COPh)\}(PMoPh_2)_2]\cdot CH_2Cl_2\cdot H_2O$, 259–66
- A reinvestigation of the structures of organotin sulphates and

CRYSTAL STRUCTURE (contd)

- chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74
- Synthesis, infrared, and tin-119 Mössbauer spectroscopic characterization of seven-co-ordinate diorganotin(IV) adducts with 2,6-diacetylpyridine acylhydrazones, including the X-ray crystal structure of $\text{SnEt}_2(\text{dapt})[\text{H}_2\text{dapt} = 2,6\text{-diacetylpyridine bis}(2\text{-thenoylhydrazone})]$, 289–94
- Reactions of the bis(alkyne) complexes $[\text{WI}_2(\text{CO})(\text{NCMe})(\eta^2-\text{RC}_2\text{R}_2)_2]$ ($\text{R} = \text{Me}$ or Ph) with mono- and bi-dentate phosphine-donor ligands and the X-ray crystal structure of $[\text{WI}_2(\text{CO})(\text{Ph}_2\text{P}(\text{CH}_3)\text{PPh}_2)(\eta^2\text{-MeC}_2\text{Me})]$, 295–302
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 84. Carbaborane tungsten–platinum complexes having a $\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6}$ ligand; crystal structures of $[\text{WPt}(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_n(\text{PEt}_3)(\mu\text{-}\sigma\text{-}\eta^5\text{-C}_2\text{B}_9\text{H}_8\text{Me}_2)]$ ($n = 2$ or 3), 303–12
- Novel five-co-ordinate diorganorhodium compounds. X-Ray structure of bis-*ortho*-chelated $[\text{Rh}^{III}(\text{Me})\{\text{C}_6\text{H}_3(\text{CH}_2\text{NMe}_2)_2\text{-2,6}\}]$, 317–24
- A comparison of the rates of intramolecular hydrogen migration in the molecules $[\text{Mo}(\eta\text{-C}_7\text{H}_8)_2]^{n+}$ to give $[\text{Mo}(\eta\text{-C}_7\text{H}_7)(\eta\text{-C}_7\text{H}_9)]^{n+}$, $n = 0$ or 1; crystal structures of $[\text{Mo}(\eta\text{-C}_7\text{H}_8)_2]$, $[\text{Mo}(\eta\text{-C}_7\text{H}_7)(\eta\text{-C}_7\text{H}_9)]$, and $[\text{Mo}(\eta\text{-C}_7\text{H}_8)_2][\text{BF}_4^-]$, 331–44
- Chemistry of metallacyclobutanones (η^3 -oxodimethylenemethane complexes). Part 5. Ligand substitution reactions of metallacyclobutan-3-ones of platinum(II) and palladium(II) with tertiary phosphines and t-butyl isocyanide; crystal structures of $[\text{Pt}^{\{\text{CH}(\text{CO}_2\text{Me})\text{COCH}(\text{CO}_2\text{Me})\}}(\text{CNBu}^{\text{i}})(\text{AsPh}_3)]$ and $[\text{Pt}^{\{\text{CH}(\text{CO}_2\text{Me})\text{CO}\bar{\text{C}}(\text{CO}_2\text{Me})\text{C}(\text{NHBu}^{\text{i}})\}}(\text{CNBu}^{\text{i}})_2]$, 345–56
- Thiosemicarbazones as co-ordinating agents. Part 3. Synthesis, spectroscopic characterization, and X-ray structure of methyl pyruvate thiosemicarbazone hemihydrate, chloro(ethyl pyruvate thiosemicarbazone)copper(II) (green form), and chloro(pyruvic acid thiosemicarbazone)copper(II) dihydrate (blue form), 361–6
- Metallaheteroborane chemistry. Part 3. Synthesis of $[2,2\text{-}(\text{PR}_3)_2\text{-}1,2\text{-TePtB}_{10}\text{H}_{10}]$ ($\text{R}_3 = \text{Et}_3$, Bu^{n}_3 , or Me_2Ph), their characterisation by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of $[2,2\text{-}(\text{PEt}_3)_2\text{-}1,2\text{-TePtB}_{10}\text{H}_{10}]$ (1988, 2555), 383
- Thiolate complexes of osmium(III) with tertiary phosphines and their dicarbonyl derivatives of osmium(II); X-ray crystal structure of *trans,trans,trans*- $[\text{Os}(\text{SC}_6\text{F}_5)_2(\text{CO})_2(\text{PEt}_2\text{Ph})_2]$, 419–24
- One-electron transfer processes in the reaction of tin(II) halides with substituted *o*-quinones; crystal structure of bis(3,5-di-t-butylcatecholato)tin-1,10-phenanthroline-dimethylformamide (1/2), 439–46
- Tris(trimethylsilyl)methyl and tris(dimethylphenylsilyl)methyl derivatives of boron. Crystal structures of dihydroxy[tris(trimethylsilyl)methyl]borane and of the lithium–boron complex $[(\text{MeOH})_2\text{Li}(\mu\text{-OMe})_2\text{B}(\text{OMe})_2]$, 447–52
- Actinide structural studies. Part 17. The crystal and molecular structures of four uranyl(vi) complexes with *N,N*-dialkylthiocarbamate ligands, 471–80
- Synthesis of sulphimide complexes of palladium and platinum. Crystal and molecular structure of *cis*-dichloro[S,S -dimethyl-*N*-(2-pyrimidinyl)sulphimide](triphenylphosphine)palladium(II), 497–504
- Reactions of co-ordinated ligands. Part 44. The reactivity of the carbene complex $[\text{Mo}(\equiv\text{CCH}_2\text{Bu}^{\text{i}})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$ towards protonic acids, diphenylphosphine, and the π -acid tetracyanoethylene; formation and crystal structures of $[\text{MoH}(\equiv\text{CCH}_2\text{Bu}^{\text{i}})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)][\text{BF}_4^-]$, $[\text{Mo}(\sigma\text{-OC(O)CF}_3)\{\eta^2\text{-HOC(O)CF}_3\}\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, and $[\text{Mo}(\equiv\text{CCH}_2\text{Bu}^{\text{i}})\{\eta^2\text{-C}_2(\text{CN})_4\}\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, 511–8
- Reaction of elemental sulphur with $(5\text{H},14\text{H}-6,8,15,17\text{-tetramethylbenzo}[b,i][1,4,8,11]\text{tetra-azacyclotetradecinato})\text{iron(II)}$; synthesis and X-ray structure of a μ -sulphido iron(III) complex, 551–4
- Oxidation of a $\text{Pt}_2(\mu\text{-SO}_2)$ moiety to $\mu\text{-SO}_3$ and the crystal structure of $[\text{Pt}_2(\mu\text{-SO}_3)(\mu\text{-SO}_2)(\text{C}_8\text{H}_{12})_2]$, 557–60
- Syntheses, electrochemistry, and spectroscopy of dirhodium(II) tetra-acetamide and tetrakis(trifluoroacetamide) complexes with axial Group 15 substituents. The X-ray crystal structures of $[\text{Rh}_2(\text{CH}_3\text{CONH})_4(\text{AsPh}_3)_2]$ and $[\text{Rh}_2(\text{CH}_3\text{CONH})_{4-n}(\text{CH}_3\text{CO}_2)_n(\text{MPh}_3)_2]$, $\text{M} = \text{As}$ or Sb , $n \approx 1$, 581–8

- Lewis-base adducts of lead(II) compounds. Part 4. Novel isomeric dimers of lead(II) thiocyanate-1,10-phenanthroline(1/2). Crystal structure determinations, 595–600
- Chemistry of multi-site bound unsaturated ligands on metal carbonyl clusters. Intramolecular reductive coupling in the conversion of $[\text{Ru}_3(\mu\text{-H})(\text{CO})_9(\mu_3\text{-}\eta^2\text{-C}=\text{CBu}^{\text{i}})]$ to the butadienyl cluster $[\text{Ru}_3(\text{CO})_8\{\mu_3\text{-}\eta^4\text{-C}(\text{Bu}^{\text{i}})\text{C}(\text{H})(\text{C}(\text{Ph})(\text{PPh}_2))\}]$, and its crystal structure. An example of ligand isomerism on metal clusters, 601–6
- Carbonyl, hydridoacarbonyl, and hydridoalkoxycarbonyl complexes of iridium with phosphorus–sulphur hybrid ligands, and the crystal and molecular structure of $[\text{IrH}(\text{CO}_2\text{Me})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{SEt})_2]\text{BPh}_4$, 607–14
- Crystallographic and nuclear magnetic resonance studies of carbonyl(η^3 -methylallyl)(trichlorostannyl) complexes of platinum(II) and palladium(II), 615–22
- Preparation of the bis(trimethylsilyl)amido lanthanide chlorides $[\{\text{Ln}[\text{N}(\text{SiMe}_3)_2\}_2(\mu\text{-Cl})(\text{thf})\}_2]$ ($\text{thf} = \text{tetrahydrofuran}$), and the crystal and molecular structures of the gadolinium and ytterbium complexes, 623–6
- Chemistry of subvalent main group metals with dithiolates. Part 2. X-Ray structural, thermal, and theoretical investigations on the novel one-dimensional stacking structure $\text{K}_2[\text{Pb}(\text{S}_2\text{C}=\text{C}(\text{CN})_2)_2]\text{H}_2\text{O}$, 627–30
- Studies on olefin-co-ordinating transition-metal carbene complexes. Part 14. An unusual reaction of tricarbonyl(η^4 -cyclopenta-1,3-diene)iron: a new approach to acyldicarbonyl(η -cyclopentadienyl)iron complexes, and the X-ray crystal structure of $[\text{Fe}(\text{COC}_6\text{H}_4\text{CF}_3\text{-}p)(\eta\text{-C}_5\text{H}_5)(\text{CO})_2]$, 635–8
- Synthesis of diphenylphosphido-bridged manganese-gold complexes. Crystal structures of $[\text{N}(\text{PPh}_3)_2][\text{Au}_2\text{Mn}(\mu\text{-PPh}_2)_2(\text{CO})_4(\text{C}_6\text{F}_5)_2]$ and $[\text{Au}_2\text{Mn}_2(\mu\text{-PPh}_2)_4(\text{CO})_8]$, 639–44
- Synthesis, characterization, and crystal structures of bismuth complexes with 2,6-diacetylpyridine bis(2-thienylhydrazone) (H_2dapt): $[\text{Bi}(\text{H}_2\text{dapt})\text{Cl}_2]\text{-dmso}\cdot\text{H}_2\text{O}$ and $[\text{Bi}(\text{H}_2\text{dapt})\text{Cl}]\text{-dmso}$, containing a quinquedentate ligand, 671–6
- Tri(η -cyclopentadienyl)uranium(IV) silyl and siloxide compounds. Crystal structure of $[\text{U}(\eta^5\text{-C}_5\text{H}_5)_3(\text{OSiPh}_3)]$. Insertion of isocyanide into a uranium–silicon bond, 677–82
- Synthesis, structure, and chemical reactivity of a stable C_5Me_5 -substituted phosphanylium ion: (pentamethylcyclopentadienyl)-(t-butylamino)phosphanylium tetrachloroaluminat e, 693–700
- A dynamic nuclear magnetic resonance study of trimethylplatinum(IV) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part I. Low-temperature solution properties, and X-ray crystal structure of $[\text{PtIMe}_3\{\text{MeS}_2\text{CHCH}(\text{SMe})_2\}]$, 701–10
- Tetrathiafulvalenium salts of planar Pt^{II} , Pd^{II} , and Cu^{III} 1,2-dithio-oxalato- S,S' anions. Synthesis, chemistry and molecular structures of bis(tetrathiafulvalenium) bis(1,2-dithio-oxalato- S,S')palladate(II), $[\text{ttf}]_2[\text{Pd}(\text{S}_2\text{C}_2\text{O}_2)_2]$, and of bis(tetrathiafulvalenium)tetrathiafulvalene bis(1,2-dithio-oxalato- S,S')platinate(II), $[\text{ttf}]_3[\text{Pt}(\text{S}_2\text{C}_2\text{O}_2)_2]$, 719–28
- Formation in solution, synthesis, and electrochemical study of oxalato complexes of N,N' -ethylenebis(salicylideneiminato)-chromium(III) and -iron(III): crystal structures of piperidinium [N,N' -ethylenebis(salicylideneiminato)](oxalato- O^2- -chromate(III) and -ferrate(III)), 729–38
- Synthesis and X-ray crystal structure of $[\text{Rh}(\text{NO})(\text{PPh}_3)_2(p\text{-MeC}_6\text{H}_4\text{NNNC}_6\text{H}_4\text{Me}-p)][\text{BF}_4^-]\text{-CH}_2\text{Cl}_2$, 745–8
- Synthesis and X-ray crystal structure of chloro[2(1*H*)-pyridine-*S*]-bis(triphenylphosphine)copper(I), 749–52
- Crystal structure of a novel tetrานuclear dithiocarboxylato-copper(I) complex, $[\text{Cu}(\text{S}_2\text{CC}_6\text{H}_4\text{Me}-p)_4(\text{PPh}_3)]$, 753–6
- Solid-state structure and solution conformation of 2,2,4,4,6,6-hexa(*p*-chlorophenoxy)cyclotri- λ^5 -phosphazatriene. X-Ray and dipole moment studies, 757–60
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 86. Alkylidyne(carborane)-molybdenum–gold, –rhodium and –iron complexes; crystal structure of $[\text{NEt}_4][\text{MoFe}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)(\mu\text{-}\sigma\text{-}\eta^5\text{-C}_2\text{B}_9\text{H}_8\text{Me}_2)(\text{CO})_8]$, 797–808
- Binuclear platinum(II) complexes bridged by pyridine-2-thiolate and related ligands. Synthesis and crystal structure, 815–20
- Reactions of amine hydrochlorides with phosphorus trichloride oxide; preparation and X-ray crystal structure of 2,4,6-trichloro-1,3,5-triphenyl-1,3,5,2,4,4,6,6,6-triazatrichrophosphorinane 2,4,6-trioxide (PhNPOCl_3), 821–8
- Tri-aza analogue of an acetylacetone-complex formed by

CRYSTAL STRUCTURE (contd)

- nucleophilic attack of a co-ordinated nitrile by amidines. *X*-Ray crystal structure of $[\text{Pt}\{\text{HNC}(\text{Ph})\text{NC}(\text{Ph})\text{NH}\}_2]_2\text{PhMe}$, 837–42
- Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NN'N'*-tetra-acetate (cdta). Part 3. Structural and magnetic characterization of the dinuclear $[\text{M}(\text{OH}_2)_2][\text{M}'(\text{cdta})]\cdot\text{H}_2\text{O}$ ($\text{M}, \text{M}' = \text{Ni}, \text{Ni}; \text{Mn}, \text{Ni}; \text{Mn}, \text{Cu}; \text{Co}, \text{Ni}; \text{or Co}, \text{Cu}$) and the tetrานuclear $[\text{M}(\text{OH}_2)_2][\text{M}'(\text{cdta})(\text{OH}_2)]\cdot 4\text{H}_2\text{O}$ ($\text{M}, \text{M}' = \text{Zn}, \text{Zn}; \text{Zn}-\text{Co}; \text{Co}, \text{Co}; \text{or Mn}, \text{Co}$) complexes, 863–72
- New carbide clusters in the cobalt subgroup. Part 17. Preparation and structural characterization of the mixed-metal octahedral dianion $[\text{Co}_2\text{Rh}_4\text{C}(\text{CO})_3]^{2-}$ as its $[\text{PPH}_4]^+$ salt, 879–84
- Gold(i) complexes derived from secondary phosphines: $[\{\text{Au}(\mu-\text{PR}_2)\}_n], [\{\text{Au}_2(\mu-\text{PPH}_2)\}]^+, [\{\text{AuX}(\text{PR}_2)\}]$, and $[\{\text{Au}(\text{PHR}_2)\}_n]^+$. Crystal structure of $[\text{AuBr}(\text{PHP}_2)]$, 907–14
- Crystal structures and solution dynamics of mono- and di-nuclear rhodium complexes of heterodinucleating macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 915–20
- Reactivity of metallic copper with acidic organic compounds in the presence of molecular oxygen. Crystal structure of $[\{\text{Cu}(\text{NCO})(\text{py})_2\}_2]\cdot\text{H}_2\text{O}$, 921–6
- Crystal structure and mass spectrometry of dibromodimethyl[1-methyl-2(3*H*)-imidazolinethione-*S*]tin(iv), 927–30
- Synthesis and spectroscopic studies of cobalt(iii) complexes of the chiral pendant-arm macrocycle *N,N',N''*-tris[(2*S*)-2-hydroxypropyl]-1,4,7-triazacyclononane (H_3L); crystal structure of the hydrogen-bridged dimer $[\text{CoL}_3\text{LCo}][\text{PF}_6]_3$, 931–6
- Synthesis and characterization of imidazolate-bridged iron(ii)-copper(ii) and manganese(iii)-copper(ii) binuclear complexes, 943–6
- N,N'*-Ethylenebis(acetylacetoneiminato)titanium complexes: synthesis, crystal structures, and properties of titanium(iii) and titanium(iv) derivatives, 953–8
- Reactivity of cationic molybdenum(ii) complexes. Part 4. Isolation and crystal structure determination of *cis*- $[\text{Mo}(\text{CO})_2(\eta-\text{C}_5\text{Me}_5)(\text{PPh}_3)(\text{N}_3)]$ and *cis*- $[\text{Mo}(\text{CO})_2(\eta-\text{C}_5\text{Me}_5)(\text{PPh}_3)(\text{NCO})]$, 959–64
- Precursor catenand complexes: synthesis, structure, and electrochemistry of bis(2,6-di-iminopyridyl) complexes of nickel(ii). The single-crystal *X*-ray structure of $[\text{NiL}^4]_2[\text{BF}_4]_2$, 965–70
- Lewis base adducts of Group 11 metal(i) complexes. Part 48. Heterobimetallic tris(dithiocarbamato)cobalt(iii)-copper(i) halide adducts as a vehicle for novel copper(i) halide clusters (structurally characterized), 971–6
- Platinum metal complexes of potentially chelating alkene-thioether and alkene selenoether ligands: synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{MeE}(\text{CH}_2)_n\text{CH}=\text{CH}_2\}]$ ($\text{M} = \text{Pt}$ or Pd ; $\text{X} = \text{Cl}, \text{Br}$, or I ; $\text{E} = \text{S}$ or Se ; $n = 2$ or 3) and the *X*-ray structure of *cis*-dibromo(2-thia-6-heptene)platinum(ii), $[\text{PtBr}_2\{\text{MeS}(\text{CH}_2)_3\text{CH}=\text{CH}_2\}]$, 985–90
- Monocyclopentadienyl pentafluorothiophenolate complexes of molybdenum and their interactions with thallium(i) ions; dynamic nuclear magnetic resonance studies and crystal and molecular structures of $[\text{TiMo}(\text{SC}_6\text{F}_5)_2\text{L}_2(\text{cp})]$ ($\text{L} = \text{CO}$ or SC_6F_5 ; $\text{cp} = \eta^5\text{-C}_5\text{H}_5$), 991–1002
- Synthesis and reactivity of trinuclear complexes of platinum containing the single bridging ligand SC_4H_8 . Molecular structure of $[\text{NBu}_4]_2[\text{trans-PtCl}_2\{(\mu-\text{SC}_4\text{H}_8)\text{Pt}(\text{C}_6\text{F}_5)_3\}_2]$, 1011–6
- Magnetic properties of mono-, bi-, and tri-nuclear copper(ii) complexes of novel oxamato and oxamido ligands. Crystal structure of a mononuclear precursor, 1017–26
- Six- and five-co-ordinate ruthenium(ii) complexes with 1,2-bis(dicyclohexylphosphino)ethane. *X*-Ray crystal structure of the d^6 five-co-ordinate $[\text{RuCl}_2\{(\text{C}_6\text{H}_{11})_2\text{PCH}_2\text{CH}_2\text{P}(\text{C}_6\text{H}_{11})_2\}_2\text{PF}_6]$, 1045–52
- 6,13-Diamino-6,13-dimethyl-1,4,8,11-tetra-azacyclotetradecane, L^7 , a new, potentially sexidentate polyamine ligand. Variable coordination to cobalt(ii) and crystal structure of the complex $[\text{Co}(\text{L}^7)]\text{Cl}_2[\text{ClO}_4]$, 1059–66
- Synthesis of a new P_2N_2 ligand *N,N'*-bis[2-(diphenylphosphino)phenyl]propane-1,3-diamine, H_2L^2 , and some of its complexes with elements of the nickel triad and rhodium: *X*-ray structure analyses of the neutral complex $[\text{NiL}^2]$ and the *trans*-spanned $[\text{Rh}(\text{CO})\text{Cl}(\text{H}_2\text{L}^2)]$, 1067–74
- Iridium(iii) hydrido complexes with 2-pyridinaldazines. Crystal structure of $[\text{IrH}_2(\text{mpaa})(\text{PPh}_3)_2]\text{SbF}_6$. Synthesis and characterization of homo- and hetero-nuclear complexes, 1075–82

- Crystal structures of di-iron μ -η²-(C,C)-ketene complexes, $[\{\text{Fe}(\eta^5-\text{C}_5\text{H}_5)(\text{CO})_2\}_2(\mu-\text{CH}_2\text{CO})]$ and $[\{\text{Fe}(\eta^5-\text{C}_5\text{H}_5)(\text{CO})_2\}_2(\mu-\text{CH}_2\text{CO})]$. Synergistic metal-to-ligand back donation as revealed by comparison with $[\{\text{Fe}(\eta^5-\text{C}_5\text{H}_5)(\text{CO})_2\}_2(\mu-\text{CH}_2\text{CH}_2\text{CO})]$, 1083–8
- Monohydrido-bridged platinum(ii)-gold(i) complexes. *X*-Ray crystal structure of $[(\text{Ph}_3\text{P})_2(\text{C}_6\text{Cl}_5)\text{Pt}(\mu-\text{H})\text{Au}(\text{PPh}_3)]\cdot\text{ClO}_4\cdot 2\text{Et}_2\text{O}$, 1089–92
- Synthesis and reactivity of binuclear pyrazolate-bridged ruthenium(i) complexes. Crystal structures of bis[μ -(3,5-dimethylpyrazolato-*NN'*)tricarbonylruthenium(i)] ($\text{Ru}-\text{Ru}$) and bis[μ -(3,5-dimethylpyrazolato-*NN'*)-μ-iodo-tricarbonylruthenium(ii)] tri-iodide, 1093–100
- Crystal structure of the new low-dimensional metal $\text{Cs}_{0.82}[\text{Pd}(\text{S}_2\text{C}_2(\text{CN})_2)]\cdot 0.5\text{H}_2\text{O}$, 1101–4
- Synthesis and characterization of pentanuclear Fe–Ir carbonyl clusters. Solid-state structures of $[\text{N}(\text{PPh}_3)_2]_2[\text{FeIr}_4(\text{CO})_{15}]$ and $[\text{PPh}_4][\text{Fe}_2\text{Ir}_3(\text{CO})_{14}]$, 1109–16
- Low-lying electronic states in μ-phenolato copper(ii) binuclear compounds with hydroxo, ethanolato, azido, and cyanato exogenous bridges: crystal structures, magnetic properties, and interpretations, 1117–26
- Preparation and characterisation of linear mixed-metal trinuclear carboxylate complexes of general formula $[\text{M}_2\text{M}'(\text{O}_2\text{CR})_6(\text{C}_9\text{H}_7\text{N})_2]$; crystal structure of $[\text{Zn}_2\text{Ba}(\text{O}_2\text{CCMe}_3)_6(\text{C}_9\text{H}_7\text{N})_2]$, 1127–32
- Trinuclear pyrazolate complexes. Crystal structure of $[(\text{OC})_2\text{Rh}(\mu-\text{pz})_2\text{Pd}(\mu-\text{pz})_2\text{Rh}(\text{CO})_2]$, 1133–8
- Synthesis and characterization of a family of nitrile-substituted triangular hydrocobaltnum clusters, $[\text{Re}_3(\mu-\text{H})(\text{CO})_{12-n}(\text{NCMe})_n]$ ($n = 1$ –3). *X*-Ray crystal structures of $[\text{Re}_3(\mu-\text{H})_3(\text{CO})_9(\text{NCMe})]$ and $[\text{Re}_3(\mu-\text{H})_3(\text{CO})_9(\text{NCMe})_3]$, 1143–8
- New metal-sulphur–nitrogen compounds from reactions in liquid ammonia. The *X*-ray structures of *trans*-bis(acetophenone dimethylhydrazone-*N*^a)dichloropalladium(ii) and [di(azathien)-1-yl-S¹N⁴]²⁻(2-hydrazonoethyl)phenyl]palladium(ii), 1179–86
- Metal complexes as ligands. Part 12. Synthesis and *X*-ray crystal structure of a hexanuclear nickel(ii) complex based on two isosceles trinuclear components with imidazolate bridges, 1187–92
- Reactivity of the iridium nitrosyl complex $[\text{Ir}(\text{NO})(\text{paa})(\text{PPh}_3)_2][\text{PF}_6]_2$ ($\text{paa} = 2$ -pyridinecarbaldehyde azine) with nucleophiles: insertion of NO into a methinic C–H bond; crystal structure of $[\text{Ir}((2-\text{C}_5\text{H}_4\text{N})\text{CH}=\text{N}-\text{N}=\text{C}(\text{NO})(2-\text{C}_5\text{H}_4\text{N})\{\text{Me}(\text{EtO})\text{C}=\text{NH}\}(\text{PPh}_3)_2][\text{PF}_6]_2$, 1217–26
- The heteronuclear cluster chemistry of the Group 1B metals. Part 11. Effect of the nature of the bidentate diphosphine ligand on the metal framework structures of the gold heteronuclear cluster compounds $[\text{Au}_2\text{Ru}_4(\mu_3-\text{H})(\mu-\text{H})\{\mu-\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2\}(\text{CO})_{12}]$ ($n = 1$ –6). *X*-Ray crystal structures of $[\text{Au}_2\text{Ru}_4(\mu_3-\text{H})(\mu-\text{H})\{\mu-\text{Ph}_2\text{P}(\text{CH}_2)_2\text{PPh}_2\}(\text{CO})_{12}]$ ($n = 1$ or 2), 1227–36
- Reaction of $[\text{Rh}_2(\mu-1,8-(\text{NH})_2\text{C}_{10}\text{H}_6)(\text{CO})_2(\text{PPh}_3)_2]$ with $[\text{L}^1, \text{PPh}_3]^+$ ($\text{M} = \text{Au}, \text{Ag}$, or Cu): formation of unusual 44-electron Rh_2M clusters. Crystal structure of $[\text{Rh}_2(\mu-\text{Au}(\text{PPh}_3))\{\mu-1,8-(\text{NH})_2\text{C}_{10}\text{H}_6\}(\text{CO})_2(\text{PPh}_3)_2]\text{ClO}_4\cdot 2\text{CH}_2\text{Cl}_2$, 1249–52
- Mode of binding of dimethyl sulphoxide in platinum(ii) and palladium(ii) cationic complexes containing a chelated diamine. *X*-Ray crystal structures of (2,2'-bipyridyl)chloro(dimethyl sulphoxide-*O*)palladium(ii) tetrafluoroborate and chloro(dimethyl sulphoxide-*S*)(ethylenediamine)palladium(ii) perchlorate, 1265–72
- Structure of μ-fluoro-bis[2,2'-bipyridyl]fluoronicnickel(ii)] fluoride–ethanol (1/2) trihydrate, revealing several unique features including strong hydrogen bonds between fluoride ions and ethanol molecules, 1273–6
- Contributions to the chemistry of boron. Part 194. Synthesis and characterization of mono-, bi-, and tri-cyclic boron–nitrogen–sulphur ring systems. Crystal structure of 2,7,9,14-tetrakis(diethylamino)-3,5,6,10,12,13-hexathia-1,8-diazab-2,4,7,9,11,14-hexaboratricyclo[9.3.0.0^{4,8}]tetradecane, 1311–20
- Reactions of co-ordinated ligands. Part 45. Ligand displacement and oxidative reactions of the carbyne complex $[\text{Mo}(\equiv\text{CCH}_2\text{Bu}')\{\text{P}(\text{OMe})_3\}_2(\eta^5-\text{C}_5\text{H}_5)]$; crystal and molecular structures of $[\text{Mo}(\eta^3-\text{RN}=\text{C}(\text{CH}_2\text{Bu}')\cdots\text{C}(\text{CH}_2\text{Bu}')\cdots\text{C}=\text{NR}]\{\text{CNR}\}_2(\eta^5-\text{C}_5\text{H}_5)]$ ($\text{R} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), $[\text{Mo}(\text{C}=\text{CHBu}')\text{I}\{\text{P}(\text{OMe})_3\}_2(\eta^5-\text{C}_5\text{H}_5)]$, and $[\text{Mo}\{\text{C}=\text{CH}_2\text{Bu}'\}\text{P}(\text{O})(\text{OMe})_2]\{\text{P}(\text{OMe})_3\}(\eta^5-\text{C}_5\text{H}_5)]$, 1321–32

CRYSTAL STRUCTURE (contd)

- Reactions of co-ordinated ligands. Part 46. Formation of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}=\text{CMe}_2)(\text{CO})_{10}]$ by carbon–carbon σ -bond and carbon–hydrogen activation of 3,3-dimethylcyclopropene; crystal and molecular structure of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and evidence for its facile rearrangement to $[\text{Os}_3(\mu\text{-H})(\mu\text{-CH}=\text{CHCHMe}_2)(\text{CO})_{10}]$, 1333–40
- Synthesis, spectroscopic, and magnetic studies of copper(II) with *N*-protected amino acids: crystal and molecular structure of diaquatetrakis(μ -*N*-benzoyl- α -alaninato)-dicopper(II), 1341–4
- Sulphonamide nitrogen-containing, nitrogen protected amino acids interacting with the Cd^{2+} ion. A cadmium-113 nuclear magnetic resonance and potentiometric study. *X*-Ray diffraction of $[\text{Cd}(\text{ts}\text{-}\beta\text{-alaO})_2(\text{H}_2\text{O})_4]$, 1345–50
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 87. Docosahedral carbaborane(alkylidyne)tungsten complexes as reagents for the synthesis of compounds with heteronuclear metal–metal bonds: crystal structures of $[\text{NEt}_4][\text{W}(\equiv\text{CC}_6\text{H}_3\text{Me}_2\text{-}2,6)(\text{CO})_2(\eta^6\text{-C}_{2\text{B}_{10}\text{H}_{10}\text{Me}_2})]$ and $[\text{NEt}_4][\text{WFe}(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-}2,6)(\text{CO})_4(\eta^6\text{-C}_{2\text{B}_{10}\text{H}_{10}\text{Me}_2})]$, 1363–74
- Heteronuclear transition metal–alkyne clusters. Part 1. Reactions of tris(alkyne)monocarbonyltungsten complexes with octacarbonyldicobalt. Synthesis, *X*-ray crystal structure, and two-dimensional nuclear magnetic resonance studies of $[\text{WC}_{20}(\mu\text{-C}_2\text{Et}_2)(\mu\text{-C}_4\text{Et}_4)(\text{CO})_8]$, 1375–80
- Complexes of the pyridine-based, tridentate, sulphur-containing ligands 2,6-bis(ethylthiomethyl)pyridine, 2,6-bis(methoxycarbonylethylthiomethyl)pyridine, and 2,6-bis(benzylthiomethyl)pyridine. Crystal structure of [2,6-bis(ethylthiomethyl)pyridine]dichlorocadmium(II) hydrate, 1381–4
- Heteroleptic poly(pyrazolyl)borate derivatives of the lanthanide ions. The synthesis of acetylacetone complexes and the molecular structures of $[\text{Ln}\{\text{HB}(\text{C}_3\text{N}_2\text{H}_3)_3\}_2\{\text{MeC}(\text{O})\text{CHC}(\text{O})\text{Me}\}]$ ($\text{Ln} = \text{Ce}$ or Yb), 1393–400
- A planar, binucleating, pyrazole derivative ligand. Crystal structure of bis[μ -3,5-bis(2'-pyridyl)pyrazolato- $N^1\text{N}^2\text{N}^2'$]-bis[dimethanolnickel(II)] dichloride dihydrate, 1401–4
- Synthesis and *X*-ray crystal structure of $[\text{ReO}(\text{Cl})_2(2\text{-OC}_6\text{H}_4\text{CH}=\text{NCH}_2\text{CH}_2\text{OCH}_2)]$, an example of four-coordination by a potentially sexidentate ligand, 1405–8
- Preparation and crystal structure of a binuclear compound of methylmercury(II) containing bismuthiol I as a bridging ligand, 1409–12
- Hydrazido-complexes of titanium(IV) with tris(pyrazolyl)borates. Crystal structures of $[\text{TiLCl}_2(\text{NMeNMe}_2)]$ and $[\text{TiLCl}_2(\text{NPhNH}_2)]$, L = tris(pyrazolyl)borate, 1413–6
- Crystal and molecular structure of [1,2-bis(diethylphosphino)ethane mono-oxide-*P*]trichloro-oxomolybdenum(V), 1417–20
- Synthesis, structure, and reactivity of the heteronuclear clusters $[\text{Os}_5\text{H}(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}]$ and $[\text{Os}_5(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}_2]$ ($\text{M} = \text{Cu, Ag, or Au; R} = \text{Ph or Me}$): the *X*-ray crystal structure of $[\text{Os}_5\text{H}(\text{CO})_{15}\{\text{Au}(\text{PPh}_3)\}]$, 1421–6
- Reactivity of binuclear pyrazolate-bridged rhodium complexes with isocyanides. Molecular structure of $[\text{Rh}(\mu\text{-pz})\text{I}_2(\text{CNBu}')_2(\mu\text{-Ph}_2\text{PCH}_2\text{PPh}_2)]\text{BF}_4$ (pz = pyrazolate), 1427–34
- Synthesis, *X*-ray characterization, and Q-band electron spin resonance studies of heterometallic, polymeric compounds formed by yttrium and lanthanide(II) nitrates with HgL_2 ($\text{L} = \text{C}_4\text{H}_6\text{NO}$, the anion of 2-pyrrolidone); crystal structures of $[(\text{Hg}_3\text{EuL}_4(\text{NO}_3)_3)_n]$ and $[(\text{Hg}_3\text{Tb}_2\text{L}_6(\text{NO}_3)_6)_n]$, 1439–48
- Crystal and molecular structures of μ -hydrido-bis[*fac*-(2,2'-bipyridine)tricarbonylrhenium(II)] chloride, *fac*-(2,2'-bipyridine)tricarbonylformatorhenium(II), and *fac*-(2,2'-bipyridine)tricarbonyl(cyanotrihydroborato)rhenium(II), 1449–54
- Synthesis of the cluster dianion $[\text{Os}_{11}\text{C}(\text{CO})_{27}]^{2-}$ by pyrolysis and its reactions with electrophiles; *X*-ray structure analysis of the mixed-metal derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\{\text{Cu}(\text{NCMe})\}]\text{CH}_2\text{Cl}_2$ and the hydrido derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455–64
- Polyhedral organoiridaborane chemistry. The reaction of MeNC with $[6-(\eta^5\text{-C}_5\text{Me}_5)\text{-}nido\text{-}6\text{-IrB}_9\text{H}_{13}]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[4,4-(\eta^5\text{-C}_5\text{Me}_5)(\text{MeNC})\text{-}arachno\text{-}4\text{-IrB}_8\text{H}_{12}]$, 1465–70
- Synthesis and crystal structure of $[\text{NiOs}_3(\eta^5\text{-C}_5\text{H}_5)(\mu\text{-H})_2\{\mu\text{-Cu}(\text{PPh}_3)\}(\text{CO})_6]$. Hydrogenation–dehydrogenation reactions of

- dienes and alcohols with heterogeneous catalysts derived from $[\text{NiM}(\eta^5\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-X})(\text{CO})_9]$ [$\text{M} = \text{Ru, X} = \text{H; M} = \text{Os, X} = \text{Cu}(\text{PPh}_3)$], 1477–82
- The structure of 18-molybdochisulphate(vi)(4–) ion in $(\text{NEt}_4)_4\text{S}_2\text{Mo}_{18}\text{O}_{62}\text{CH}_3\text{CN}$, 1491–8
- Binuclear copper(I) complexes containing two or three 2-(diphenylphosphine)pyridine (dppy) bridging ligands: *X*-ray crystal structure of $[\text{Cu}_2(\mu\text{-dppy})_3(\text{MeCN})][\text{BF}_4]_2$ and variable-temperature ^1H and ^{31}P – ^1H nuclear magnetic resonance studies, 1499–506
- X*-Ray structure of a ternary complex, copper(II)-inosine 5'-monophosphate–benzimidazole. The first example for the absence of direct metal–nucleotide interaction, 1507–12
- Crystal and molecular structures of diaqua(nitrilotriacetato)aluminium(III) and di- μ -hydroxo-bis(nitrilotriacetato)-dialuminato(III) dianion, 1513–8
- Synthesis, characterization, and thermolytic behaviour of η^3 -allyl(methyl)(tertiary phosphine)palladium(II), and *X*-ray structure of the cluster $[\text{Pd}_4(\mu\text{-C}_3\text{H}_5)(\mu\text{-Cl})_2(\mu\text{-dppm})_2]\text{-}4\text{C}_4\text{H}_8\text{O}$ (dppm = $\text{Ph}_2\text{PCH}_2\text{PPh}_3$), 1519–26
- Tertiary phosphine complexes of triply bonded ditungsten bis(1-butylamide) tetrachloride. The *X*-ray crystal structures of $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMe}_3)_2]$, $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PEt}_3)_2]$, and *cis*- $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMe}_2\text{Ph})_2]$, 1537–42
- Chemistry of phosphido-bridged dimolybdenum complexes. Part 3. Reinvestigation of the reaction between $[\text{Mo}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_6]$ and P_2Ph_4 ; *X*-ray structures of $[\text{Mo}_2(\eta^5\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\mu\text{-CO})]$ and *trans*- $[\text{Mo}_2(\eta^5\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2\text{O}(\text{CO})]$, 1555–64
- Insertion of phenylethyne-thiolate or –selenolate into metal–carbene bonds: synths for co-ordinated thioacyl or selenoacyl anions and for anionic thiocarbene or selenocarbene complexes, 1565–78
- Synthesis and reactivity of binuclear perhalogenophenyl derivatives of platinum(II). Molecular structure of $[\text{Pt}_2(\text{C}_6\text{F}_5)_2(\text{CO})_2\text{-}0.4\text{CHCl}_3$, 1583–8
- Phenyltin trichloroacetate complexes: preparations and structures, 1589–98
- Molecular structure of heterobimetallic $\mu\text{-}\eta^2\text{-}(\text{C},\text{C})\text{-ketene}$ complexes, 1627–30
- Stacked planar molecules in antiferromagnetic di-isothiocyanatobis(thiourea)chromium(II), a new structural type for chromium(II), 1631–4
- Lewis-base adducts of Group 11 metal(II) compounds. Part 58. Syntheses, *X*-ray structures, and far-infrared spectra of the novel $[\text{CuX}_2(\text{bidentate ligand})]^-$ ($\text{X} = \text{Cl}$ or Br) anion salts of the $[\text{Cu}(\text{dpa})_2]^+$ cation and the dimeric iodide $[(\text{dpa})\text{CuI}_2\text{Cu}(\text{dpa})]$ [$\text{dpa} = \text{di}(2\text{-pyridyl)}\text{amine}$], 1639–44
- Metalla-heteroborane chemistry. Part 5. Synthesis of *cis*- $[(\text{PR}_3)_2\text{-}1,2\text{-SePtB}_{10}\text{H}_{10}]$ ($\text{R}_3 = \text{Et}_3$, $\text{Bu}^n{}_3$, or Me_2Ph) and *arachno*- $[(9,9\text{-}(\text{PMe}_2\text{Ph})_2\text{-}6,9\text{-SePtB}_8\text{H}_{10})]$ and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of *cis*- $[(2,2\text{-}(\text{PMe}_2\text{Ph})_2\text{-}1,2\text{-SePtB}_{10}\text{H}_{10})]$ and *arachno*- $[(9,9\text{-}(\text{PMe}_2\text{Ph})_2\text{-}6,9\text{-SePtB}_8\text{H}_{10})]$, 1657–68
- Platinum metal thioether macrocyclic complexes: synthesis, electrochemistry, and single-crystal *X*-ray structures of *cis*- $[\text{RhCl}_2\text{L}^2]\text{PF}_6$ and *trans*- $[\text{RhCl}_2\text{L}^3]\text{PF}_6$ ($\text{L}^2 = 1,4,8,11$ -tetraathiacyclotetradecane, $\text{L}^3 = 1,5,9,13$ -tetraathiacyclohexadecane), 1675–80
- Copper(II)-assisted hydrolysis of 2,4,6-tris(2-pyridyl)-1,3,5-triazine. Part 2. Crystal structures of [bis(2-pyridylcarbonyl)amido]pyridine-2-carboxamide)copper(II) trifluoromethanesulphonate and [bis(2-pyridylcarbonyl)amido] J_2 2,4,6-tris(2-pyridyl)-1,3,5-triazine]copper(II) trifluoromethanesulphonate, 1681–8
- Syntheses, crystal structures, and Raman and resonance Raman spectroscopy of platinum(II), platinum(IV), and halogen-bridged chain platinum(II,IV) complexes involving the terdentate ligand 3-methyl-3-azapentane-1,5-diamine (4Me-dien), 1689–96
- Synthesis and reactions of 1,6-bis(2-butyltellurophenyl)-2,5-diazahexa-1,5-diene and related compounds. The crystal and molecular structures of 2-(butyldichlorotelluro)benzaldehyde and bis[2-(hydroxyiminomethyl)phenyl] ditelluride, 1697–704
- Direct insertion of a nitrogen atom into the S–S bond of a 1,2,3,5-dithiadiazole ring in a direct current nitrogen glow discharge, and *X*-ray crystal structure of 4-methyl-1,2,3,5-dithiadiazole, 1705–14
- Complexes of ligands providing endogenous bridges. Part 7. The concurrent formation of 1 + 1 and 2 + 2 oxa-azamacrocycles;

CRYSTAL STRUCTURE (contd)

- the X-ray crystal structure of an exocyclic dinuclear copper(II) complex, 1727–38
- Closely related macrocyclic and acyclic tridentate, pyridine derivatives, containing sulphur, and their complexes. Crystal structures of {dichloro-3,10-dithia-16-azabicyclo[10.3.1]hexadeca-1(16),12,14-triene}copper(II) and [2,6-bis(ethylthiomethyl)pyridine]dichlorocopper(II), 1739–44
- Structural effects of one-electron oxidation on a cyclobutadieneiron derivative: X-ray crystal structures of [Fe(CO){P(OMe)₃}₂(η-C₄Ph₄)] and [Fe(CO){P(OMe)₃}₂(η-C₄Ph₄)BF₄], 1751–8
- Synthetic, spectroscopic, and X-ray studies on methylmercury(II) and dimethylthallium(III) complexes with cyclopentanone thiosemicarbazone, 1787–92
- Phenoxo and diphenoxo complexes of titanium(III) and vanadium(IV): synthesis and X-ray structure, 1793–8
- Organic chemistry of dinuclear metal centres. Part 12. Synthesis, X-ray crystal structure, and reactivity of the di-μ-alkylidene complex [Ru₂(CO)₂(μ-CHMe)(μ-CMe₂)(η-C₅H₅)₂]: alkylidene linking, 1799–806
- Organic chemistry of dinuclear metal centres. Part 13. Synthesis, structure, and reactivity of [Ru₂(CO)₄(η⁵:η⁵-C₅H₄CH₂C₅H₄)], 1807–14
- Preparation of thionylimide complexes of titanium, zirconium, and hafnium. Crystal structure of [Zr(cp)(η-C₅Me₅)(NSO)₂], 1815–8
- The mechanisms of fluxionality of [Fe₃(CO)_{12-n}{P(OR)₃}_n] (R = Me, n = 0–3; R = Pr^t, n = 2 or 3) and the X-ray structure of [Fe₃(CO)₁₀{P(OMe)₃}₂], 1831–44
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 88. Carbaboranetungsteniridium compounds; crystal structure of the complex [WIr(μ-CC₆H₄Me₄)(CO)₂(PEt₃)₂(η⁵-C₂B₉H₉Me₂)], 1845–54
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 89. Tetra- and penta-nuclear tungsten-rhodium complexes: crystal structures of [W₃Rh₂(μ-CO)₂(μ-CMe){μ-C(Me)C(O)}(μ-PPPh₂)₂(μ₃-CMe)(CO)₂(η-C₅H₅)₃] and [W₃Rh₂(μ-CO)₃(μ-CMe){μ-C(Me)PPPh₂}(<μ₃-CMe)(CO)₂(η-C₅H₅)₃], 1855–64
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 91. Steric implications in the reactions of the complexes [M₂(C≡R)(CO)₂(η-C₅H₅)] (M = Mo or W, R = C₆H₄Me-4, C₆H₄OMe-2, or C₆H₃Me₂-2,6) with the compounds [Mo(CO)₃(NCMe)₃] and [Cu(thf)(η-C₅Me₅)] (thf = tetrahydrofuran); crystal structure of [Mo₃(μ-CC₆H₃Me₂-2,6)₂(CO)₆(η-C₅H₅)₂], 1871–8
- Synthetic and structural studies on organomolybdenum and organotungsten complexes containing bismuth. X-Ray crystal structures of [BiCl{Mo(CO)₃(η-C₅H₅)₂}]₂ and [BiCl{Mo(CO)₂(CNBu')₂}(η-C₅H₅)₂]₂ (1988, 2941), 1887
- Preparation, characterisation, and reactions with phosphite donor ligands of the new cationic monoalkyne complexes [WI(CO)(NCMe){Ph₂P(CH₂)_nPPH₂}](η²-RC₂R)][BF₄]_n (n = 1–6; R = Me or Ph). X-Ray crystal structure of [WI(CO)-{P(OPr)₃}₂{Ph₂P(CH₂)₂PPH₂}](η²-MeC₂Me)][BF₄]_{0.5}[OH]_{0.5}·0.5H₂O, 1903–10
- Metal complexes of α -hydroxyiminophosphonic acid derivatives. Preparation and crystal structures of calcium and cadmium complexes of methyl (E)- α -hydroxyiminobenzylphosphonate, 1911–4
- Steric effect of the adamantyl ligand on crystal structure and dissociation rates in organocobalt B₁₂ models, 1941–6
- Chemistry of [{Cr(cp)(CO)₃}₂] (cp = η⁵-C₅H₅). Reaction with elemental P₄: A high yield synthesis and crystal structures of [{Cr(cp)(CO)₂}₂(μ-η²-P₂)] and [{Cr(cp)(CO)₂}₂(η³-P₃)], 1951–6
- Rearrangements of >C=N– functionalities promoted by cobalt(I): reactivity of cyclopentadienylbis(ethylene)cobalt(I) with dibenzophenone azine, benzophenone oxime, benzylidene(phenyl)amine and crystal structure determinations of the products, 1957–62
- Spectroscopic and electrochemical study of copper and zinc complexes with a N₂S₂ ligand donor set. Crystal structure of the copper(II) complex derived from 1,3-propylenediamine and 3-formyl-1-phenyl-2(1H)-pyridinethione, 1979–86
- Preparation and structural studies of diphenylthiophosphinito rhodium(III) complexes. X-Ray structure analysis of [(C₅Me₅)Rh₂(μ-Cl)₂(μ-Ph₂PS)][BPh₄]₂, 1987–92
- A barium dithiocyanate complex with an unusual macrocycle containing pendant arms, the amine precursor, and related copper(II) diperchlorate complexes: synthesis and structure, 1993–2002
- Synthesis and characterization of bis(xanthate) complexes of Mo^{II} and W^{II}. Crystal structure of [W(S₂COPr^t-S)(S₂COPr^t-SS')(CO)₂(PM₃e)₂], 2003–10
- Ruthenium oxo complexes of macrocyclic tertiary amines. Synthesis, electrochemistry, and reactivities of ruthenium oxo complexes of meso-2,3,7,11,12-pentamethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L¹), and X-ray crystal structure of trans-[Ru^{IV}L¹(O)(NCO)]ClO₄, 2011–6
- Cluster activation via cyclometallation. Formation of the cyclometallated triosmium carbonyl cluster [Os₃H₂(CO)₉{P(C₆H₄)Ph}] and its reactions with phosphorus donor ligands. The X-ray crystal structures of [Os₃H₂(CO)₉{P(C₆H₄)Ph}] and [Os₃H₂(CO)₉{P(OMe)₃}(PPh₃)], 2023–32
- Reactions of the cyclometallated triosmium cluster [Os₃H₂(CO)₉{P(C₆H₄)Ph}] with Brønsted acids and dihydrogen: synthesis and X-ray crystal structure of [Os₃H₂(CF₃CO₂)(CO)₉(μ-PPh₃)], 2033–8
- Reduction–oxidation properties of organotransition-metal complexes. Part 30. Triazenido-bridged carbonylrhodium bipyridyl complexes with [Rh₂]²⁺, [Rh₂]³⁺, and [Rh₂]⁴⁺ cores, and the X-ray structures of [Rh₂(CO)₂(bipy)(μ-RNNNR)₂][BF₄]₂·CH₂Cl₂ and [Rh₂I(CO)(bipy)(μ-RNNNR)₂]₂[PF₆]₂·2.5CH₂Cl₂ (R = p-tolyl), 2049–60
- Synthesis and properties of the bis(μ-thiosulphato-S)-bis(dinitrosylferrate)(2–) anion. Crystal and molecular structure of bis[bis(triphenylphosphoranylidene)ammonium] bis(μ-thiosulphato-S)-bis(dinitrosylferrate), 2061–4
- Chelate-stabilised tetrahydrofuran complexes of platinum(II). Crystal structure of [PtCl(Ph₂PCH₂C₄H₇O)₂(Ph₂C₄H₇O)][PF₆]₂, 2069–72
- Activation of a vinyl carbon–hydrogen bond in a tris(pyrazolyl)boratoiridium complex. The X-ray crystal structure of [IrH{HB(pz)₃}({σ}-C₈H₁₃)(η²-C₈H₁₄)], 2073–6
- Crystal structures and magnetic properties of bi- and tetra-nuclear copper(II) complexes of 2,6-diformyl-4-methylphenol di(benzoylhydrazone), 2083–8
- Synthesis, structure, and redox properties of dicopper(II,II) complexes of N-(hydroxyphenyl)salicylamide and homologues. Facile oxidation to dicopper-(II,II) and -(II,III) species; crystal structure of bis(tetrabutylphosphonium) bis{[(2-oxidobenzoyl)(2'-oxidophenyl)amido(3-)-NO²,μ-O²⁻]cuprate(II)}, 2089–94
- Crystal structure and magnetic properties of the heterobinuclear complex [(CH₃CN)LNi(mnt)Cu(mnt)]·CH₃CN (L = tetrabenzo[b,f,j,n][1,5,9,13]tetra-azacyclohexadecine and mnt = maleonitriledithiolate), 2095–8
- Mesitylgold(I) complexes. Crystal structures of NEt₄[Au(mes)Cl] and [N(PPh₃)₂]₂[mes]Au(CN)Au(mes)], 2127–32
- Stereochemical role of lone pairs in main-group elements. Part 4. The crystal structure at 120 K of bis(1,4,7,10,13-pentaoxacyclopentadecane)tin(II) bis[trichlorostannate(II)], confirming the sterically inactive tin lone pair, 2155–60
- Low-valent isocyanide complexes and clusters of palladium and platinum. Crystal structure of [Pt{C(=NR)N(R)C(=NR)}(RNC)₂] (R = 2,6-Me₂C₆H₃), 2161–6
- Interaction of benzo-1,3-thiazoline-2-thione and related ligands with copper(II) salts and the single-crystal X-ray structure of μ-[1,2-bis(diphenylphosphino)ethane]-bis[benzo-1,3-thiazole-2-thiolato-S][1,2-bis(diphenylphosphino)ethane]copper(I), 2167–72
- Synthesis and crystal structures of three four-membered ring compounds containing PN₂Ti skeletons, 2173–8
- Characterisation of tridentate macrocyclic complexes of molybdenum(0) with tetra-aza macrocycles. Crystal structure of [Mo(CO)₃L¹] (L¹ = C-meso-5,5,7,12,12,14-hexamethyl-1,4,8,11-tetra-azacyclooctadecane), 2183–6
- Iron carbene complexes. Part 3. Aminolysis of di-iron alkoxy carbene complexes with ammonia and dimethylamine. X-Ray crystal structure of [Fe₂(μ-SPh)₂(CO)₅{C(NMe₂)Ph}], 2199–204
- Electrochemistry of the quasi-reversible bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate-(V) and -(IV) and bis[2-hydroxy-2-methylbutanoato(2–)]oxochromate-(V) and -(IV) redox couples and the crystal and molecular structure of sodium bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate(V) sesquihydrate, 2205–10

CRYSTAL STRUCTURE (contd)

- Stepwise conversion of $[\text{Os}(\text{CF}_3\text{CO}_2)_2(\text{CO})_4]$ into $[\text{Os}(\text{pyS})_2(\text{CO})_2]$: X-ray crystal structures of the complexes $[\text{Os}(\text{pyS})_2(\text{CO})_x]$ where $x = 2$ or 3 and pyS is the pyridine-2-thionato ligand, 2211–6
- Lanthanide ion complexes of the calixarenes. Part 4. Double inclusion by *p*-t-butylcalix[4]arene (H_4L). Crystal structures of $[\text{Eu}_2(\text{HL})_2(\text{dmf})_4] \cdot 7\text{dmf}$ (dmf = dimethylformamide) and $\text{H}_4\text{L}\cdot\text{dmso}$ (dmso = dimethyl sulphoxide), 2217–22
- The 4-phenyl-1,2,3,5-dithiadiazole complex $[\text{Fe}_2(\text{CO})_6(\text{PhCN}_2\text{S}_2)]$: its preparation, crystal structure, and an extended-Hückel molecular-orbital study of the bonding, 2229–34
- Synthesis of arylthallium complexes using organomercury compounds. Crystal and molecular structures of $[\text{NMe}_4][\text{Ti}(\text{C}_6\text{H}_4\text{N}=\text{NPh}-2)\text{Cl}_3]$ and $[\text{Ti}(\text{C}_6\text{H}_4\text{CH}_2\text{NMe}_2-2)\text{Cl}_2]$, 2241–6
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 93. Synthesis of heteropolynuclear metal compounds with chains of seven to eleven metal atoms; crystal structure of $[\text{Mo}_2\text{W}_3\text{Pt}_6(\mu_3\text{-CMe})_3(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\text{CO})_{10}(\text{PMe}_2\text{Ph})_4(\eta\text{-C}_5\text{H}_5)_5]$, 2269–80
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 95. Synthesis and crystal structures of $[\text{W}_2\text{RuPt}(\mu\text{-CC}_6\text{H}_4\text{Me}-4)(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)(\text{CO})_6(\text{PMe}_2\text{Ph})(\eta\text{-C}_5\text{H}_5)_2]$ and $[\text{W}_2\text{Ru}_2\text{Pt}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\mu\text{-CO})_2(\text{CO})_9(\eta\text{-C}_5\text{H}_5)_2]$, 2289–96
- Insertion of CS_2 into Ag-S bonds. Preparations and crystal structures of $[\{\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}^i_3-2,4,6)\}_4\cdot\text{CHCl}_3]_n$ and its insertion product with CS_2 , $[\{\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}^i_3-2,4,6)\}_2\{\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}^i_3-2,4,6)\}_6]\cdot8\text{CHCl}_3$, 2297–302
- A convenient route to binuclear osmium(i) complexes containing diphosphine and arsinophosphine ligands. X-Ray crystal structure of $[\text{Os}_2(\text{MeCO}_2)_2(\text{CO})_4(\text{dppm})_2]$, 2303–8
- Unusual chelating properties of the ligand 1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-dithiaoctane (bddo). Crystal structures of $\text{Ni}(\text{bddo})(\text{NCS})_2$, $\text{Zn}(\text{bddo})(\text{NCS})_2$ and $\text{Cd}_2(\text{bddo})(\text{NCS})_4$, 2309–14
- Platinum metal complexes of potentially chelating alkene thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{E}[(\text{CH}_2)_n\text{CR=CR}_2]\}_2]$ ($\text{M} = \text{Pt}$ or Pd ; $\text{X} = \text{Cl}$, Br , or I ; $\text{E} = \text{S}$ or Se ; $n = 2$ or 3; $\text{R} = \text{H}$ or Me) and the X-ray crystal structure of *cis*-di-iodo(5-thianona-1,8-diene)platinum(ii), $[\text{PtI}_2\{\text{S}[(\text{CH}_2)_2\text{CH=CH}_2]\}_2]$, 2315–22
- Reactions of chloro(triphenylphosphine)gold(i) and $[\mu\text{-1,2-bis(diphenylphosphine)-ethane}]$ -bis[bromogold(i)] with oxopurine bases. Molecular and crystal structure and bonding of (theophyllinato)(triphenylphosphine)gold(i), 2323–30
- Oxidative addition of dibenzoyl peroxide on metallic copper. The crystal structure of $[\text{Cu}(\text{C}_6\text{H}_5\text{CO}_2)_2(\text{py})_2]$, 2331–4
- Crystal and molecular structure of a lattice adduct, $\text{HgPh}_2\cdot2[\text{Hg}(\text{SCN})_2(\text{PPh}_3)_2]$: an unusual case of two- and four-co-ordinate mercury in the same unit cell, 2339–42
- Mixed-metal carbido carbonyl clusters. Part 4. Synthesis and structural characterization of $[\text{Rh}_6\text{C}(\text{CO})_{13}(\text{Au}_2(\text{PPh}_3)_2)]$, 2343–6
- Cyclic metallasiloxane compounds of Ti^{IV} and Zr^{IV} . X-Ray crystal structures of a *cis*-bis(pyridine) bis(cyclosiloxy)titanate and a pyridinelithiated tris(cyclosiloxy)zirconate, 2347–52
- Functionalized amino acid complexation to a cobalt(iii) complex of $(2R,5R,8R,11R)-2,5,8,11$ -tetraethyl-1,4,7,10-tetraazacyclododecane. Crystal structures of complexes containing (*R*)- or (*S*)-serine and (*R*)- α -methylserine, and release of amino acid from the complexes, 2359–66
- Reactions of 1-alkynes with trans-[$\text{ReCl}(\text{N}_2)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2$]: preparation of the vinylidene compounds *trans*-[$\text{ReCl}(-\text{C=CHR})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2$] ($\text{R} = \text{alkyl or aryl}$) and X-ray structure of *trans*-[$\text{ReCl}(-\text{C=CHPh})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2$], 2381–8
- The chemistry of dinitrogen residues. Part 7. Hydrazido(1–)- and *N,N*-dimethylhydrazinato(1–)-complexes of titanium(iv), 2389–96
- Substituted cyclopentadienyl complexes. Part 4. The catalysed synthesis and proton nuclear magnetic resonance spectra of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_4\text{Me})(\text{CO})(\text{L})\text{I}]$ and $[\text{Ru}(\eta^5\text{-C}_5\text{H}_7)(\text{CO})(\text{L})\text{I}]$ ($\text{L} = \text{phosphine or isocyanide}$) and the crystal structure determination of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_7)(\text{CO})(\text{P}(\text{CH}_2\text{Ph})_3)\text{I}]$, 2407–14
- Cyclometallation of unsaturated carboxylic acids by pentamethylcyclopentadienyl-rhodium and -iridium complexes. Crystal structures of $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh=CHCO-O})(\text{Me}_2\text{SO})]$ and $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh=CHCO-OMe})\text{I}]^+$, 2415–22

Complexes of lead(ii) bis(O,O' -dialkyl dithiophosphates) with nitrogen donor ligands. The crystal structures of $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\cdot\text{en}$ ($\text{en} = \text{ethylenediamine}$), $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\cdot\text{bipy}$ ($\text{bipy} = 2,2'\text{-bipyridine}$), and $\{\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\}_2\cdot\text{en}$, 2443–8

Preparation and X-ray crystal structure of $[\text{Ru}_3\text{RhH}_3(\text{CO})_{12}]$, 2449–52

On the existence of the anion trichloro(η -cyclopentadienyl)titanate(1–). Crystal structure of the salt $[\text{Co}(\text{C}_5\text{H}_5)_2\text{Ti}(\text{C}_5\text{H}_5)\text{Cl}_3]$, 2453–6

Tables of bond lengths determined by X-ray and neutron diffraction. Part 2. Organometallic compounds and co-ordination complexes of the *d*- and *f*-block metals, S1–S84

CYANATE

Reactivity of cationic molybdenum(ii) complexes. Part 4. Isolation and crystal structure determination of *cis*- $[\text{Mo}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)(\text{PPh}_3)(\text{N}_3)]$ and *cis*- $[\text{Mo}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)(\text{PPh}_3)(\text{NCO})]$, 959–64

CYANIDE

Reductive nitrosylation of tetraoxometalates. Part 15. Generation of $\text{Mn}(\text{NO})^{3+}$, $\text{Mn}(\text{NO})^{2+}$, and $\text{Mn}(\text{NO})^{2+}$ moieties directly from MnO_4^- . Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese, 1963–72

Kinetics of formation, dissociation, and isomerization reactions in complexes of cyanopyridines and benzonitrile with pentacyanoruthenate(ii), 1973–8

Mesitylgold(i) complexes. Crystal structures of $\text{NEt}_4[\text{Au}(\text{mes})\text{Cl}]$ and $[\text{N}(\text{PPh}_3)_2][(\text{mes})\text{Au}(\text{CN})\text{Au}(\text{mes})]$, 2127–32

Reductive nitrosylation of tetraoxometalates. Part 15. Generation of $\text{Mn}(\text{NO})^{3+}$, $\text{Mn}(\text{NO})^{2+}$, and $\text{Mn}(\text{NO})^{2+}$ moieties directly from MnO_4^- . Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese (1989, 1963), 2461

CYANOPYRIDINE

Kinetics of formation, dissociation, and isomerization reactions in complexes of cyanopyridines and benzonitrile with pentacyanoruthenate(ii), 1973–8

CYANOTRIHYDROBORATE

Structural and molecular structures of $\mu\text{-hydrido-bis}[fac\text{-}(2,2'\text{-bipyridine})tricarbonylrhenium(i)]$ chloride, *fac*-(2,2'-bipyridine)tricarbonylformatorhenium(i), and *fac*-(2,2'-bipyridine)tricarbonyl(cyanotrihydroborato)rhenium(i), 1449–54

CYCLOBUTADIENE

Structural effects of one-electron oxidation on a cyclobutadieneiron derivative: X-ray crystal structures of $[\text{Fe}(\text{CO})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_4\text{Ph}_4)]$ and $[\text{Fe}(\text{CO})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_4\text{Ph}_4)]\text{BF}_4^-$, 1751–8

CYCLOHEPTADIENYL

Addition of anilines to the $[\text{Fe}(\text{CO})_3(1\text{--}\eta\text{-C}_7\text{H}_9)]\text{BF}_4^-$ complex, and the 'Ordered Transition State Mechanism,' 561–6

CYCLOHEPTATRIENE

A comparison of the rates of intramolecular hydrogen migration in the molecules $[\text{Mo}(\eta\text{-C}_7\text{H}_8)_2]^{n+}$ to give $[\text{Mo}(\eta\text{-C}_7\text{H}_7)(\eta\text{-C}_7\text{H}_9)]^{n+}$, $n = 0$ or 1; crystal structures of $[\text{Mo}(\eta\text{-C}_7\text{H}_8)_2]$, $[\text{Mo}(\eta\text{-C}_7\text{H}_7)(\eta\text{-C}_7\text{H}_9)]$, and $[\text{Mo}(\eta\text{-C}_7\text{H}_8)_2]\text{BF}_4^-$, 331–44

CYCLOHEXANEDIAMINETETRA-ACETATE

Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNN'N'*-tetra-acetate (cdta). Part 3. Structural and magnetic characterization of the dinuclear $[\text{M}(\text{OH}_2)_5][\text{M}'(\text{cdta})\cdot\text{H}_2\text{O}]$ ($\text{M}, \text{M}' = \text{Ni}, \text{Ni}; \text{Mn}, \text{Ni}; \text{Mn}, \text{Cu}; \text{Co}, \text{Ni}; \text{or Co}, \text{Cu}$) and the tetrานuclear $[\text{M}(\text{OH}_2)_4][\text{M}'(\text{cdta})(\text{OH}_2)\cdot4\text{H}_2\text{O}] (\text{M}, \text{M}' = \text{Zn}, \text{Zn}; \text{Zn-Co}; \text{Co}, \text{Co}; \text{or Mn}, \text{Co})$ complexes, 863–72

CYCLOMETALLATION

Cyclometallation of unsaturated carboxylic acids by pentamethylcyclopentadienyl-rhodium and -iridium complexes. Crystal structures of $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh=CHCO-O})(\text{Me}_2\text{SO})]$ and $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh=CHCO-OMe})\text{I}]^+$, 2415–22

CYCLO-OCTENE

Activation of a vinyl carbon–hydrogen bond in a tris(pyrazolyl)boratoiridium complex. The X-ray crystal structure of $[[\text{IrH}\{\text{HB}(\text{pz})_3\}(\sigma\text{-C}_8\text{H}_{14})]\cdot2\text{C}_2\text{H}_2\text{O}]$, 2073–6

CYCЛОПЕНТАДИЕНИДИЙ

Organic chemistry of dinuclear metal centres. Part 13. Synthesis, structure, and reactivity of $[\text{Ru}_2(\text{CO})_4(\eta^5\text{:}\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{C}_5\text{H}_4)]$, 1807–14

CYCЛОПЕНТАДИЕНИЛ

Halogeno, hydride, and alkyl carbonyl-monocyclopentadienyl niobium derivatives. Crystal and molecular structure of $[\text{Nb}(\eta\text{-C}_5\text{H}_5)\text{Cl}(\text{H})(\text{CO})(\text{PMe}_3)_2]$, 79–84

Studies on olefin-co-ordinating transition-metal carbene complexes.

CYCLOPENTADIENYL (contd)

Part 14. An unusual reaction of tricarbonyl(η^4 -cyclopenta-1,3-diene)iron: a new approach to acyldicarbonyl(η -cyclopentadienyl)iron complexes, and the X-ray crystal structure of $[\text{Fe}(\text{COC}_6\text{H}_4\text{CF}_3-p)(\eta\text{-C}_5\text{H}_5)(\text{CO})_2]$, 635–8
 On the relationship between crystallographic and spectroscopic evidence of dynamic processes in the solid state: the cases of *cis*- and *trans*- $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_4]$, 1721–6
 Organic chemistry of dinuclear metal centres. Part 12. Synthesis, X-ray crystal structure, and reactivity of the di- μ -alkylidene complex $[\text{Ru}_2(\text{CO})_2(\mu\text{-CHMe})(\mu\text{-CMe}_2)(\eta\text{-C}_5\text{H}_5)_2]$: alkylidene linking, 1799–806
 Preparation of thionylimide complexes of titanium, zirconium, and hafnium. Crystal structure of $[\text{Zr}(\text{cp})(\eta\text{-C}_5\text{Me}_5)(\text{NSO})_2]$, 1815–8
 Chemistry of $[\{\text{Cr}(\text{cp})(\text{CO})_3\}_2]$ ($\text{cp} = \eta^5\text{-C}_5\text{H}_5$). Reaction with elemental P₄. A high yield synthesis and crystal structures of $[\{\text{Cr}(\text{cp})(\text{CO})_2\}_2(\mu\text{-}\eta^2\text{-P}_2)]$ and $[\text{Cr}(\text{cp})(\text{CO})_2(\eta^3\text{-P}_3)]$, 1951–6
 Rearrangements of $\geq \text{C}-\text{N}$ -functionalities promoted by cobalt(I): reactivity of cyclopentadienylbis(ethylene)cobalt(I) with dibenzophenone azine, benzophenone oxime, benzylidene(phenyl)amine and crystal structure determinations of the products, 1957–62

Hydrogen scrambling processes in $[\text{Ta}(\eta\text{-C}_5\text{Me}_5)(\eta\text{-C}_5\text{H}_5)\{(\mu\text{-H}_2\text{BH}_2)\}]$: a surprising observation, 2133–8

Substituted cyclopentadienyl complexes. Part 4. The catalysed synthesis and proton nuclear magnetic resonance spectra of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_4\text{Me})(\text{CO})(\text{L})\text{I}]$ and $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)(\text{CO})(\text{L})\text{I}]$ ($\text{L} =$ phosphine or isocyanide) and the crystal structure determination of $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)(\text{CO})\{\text{P}(\text{CH}_2\text{Ph})_3\}\text{I}]$, 2407–14

On the existence of the anion trichloro(η -cyclopentadienyl)titanate(1–). Crystal structure of the salt $[\text{Co}(\text{C}_5\text{H}_5)_2]\text{Ti}(\text{C}_5\text{H}_5)\text{Cl}_3$, 2453–6

CYCLOPENTANONE

Synthetic, spectroscopic, and X-ray studies on methylmercury(II) and dimethylthallium(III) complexes with cyclopentanone thiosemicarbazone, 1787–92

CYCLOPHOSPHAZANE

Reactions of amine hydrochlorides with phosphorus trichloride oxide; preparation and X-ray crystal structure of 2,4,6-trichloro-1,3,5-triphenyl-1,3,5,2 λ^5 ,4 λ^5 ,6 λ^5 -triazatrichosphorinane 2,4,6-trioxide (PhNPOCl_3), 821–8

CYCLOPHOSPHAZENE

Solid-state structure and solution conformation of 2,2,4,4,6,6-hexa(*p*-chlorophenoxy)cyclotri- λ^5 -phosphazatriene. X-Ray and dipole moment studies, 757–60

CYCLOPROPENE

Reactions of co-ordinated ligands. Part 46. Formation of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH=CMe}_2)(\text{CO})_{10}]$ by carbon–carbon σ -bond and carbon–hydrogen activation of 3,3-dimethylcyclopropene; crystal and molecular structure of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and evidence for its facile rearrangement to $[\text{Os}_3(\mu\text{-H})(\mu\text{-CH=CHCHMe}_2)(\text{CO})_{10}]$, 1333–40

DECOMPOSITION

A kinetic study of the gas-phase thermolysis of pentaborane(11), 385–90

Influence of added hydrogen on the kinetics and mechanism of thermal decomposition of tetraborane(10) and of pentaborane(11) in the gas phase, 391–8

DETERGENT

Electronic spectral study of the aqua \rightleftharpoons hydroxo equilibrium of model iron(III) haems encapsulated in aqueous detergent micelles, 1003–6

DIAMINE

The co-ordination chemistry of divalent cobalt, nickel, and copper. Part 13. Characterization and stability constants of copper(II) species isolated from mixed-ligand solutions; crystal structure of the complex (pyridine-2,6-dicarboxylato)(*N,N,N',N'*-tetramethyl-1,3-diaminopropane)copper(II), 253–8

DIARSINE

Synthesis and structural studies on silver–tin complex salts with bis(diphenylarsino)methane and bis(diphenylphosphino)methane, 247–52

Reactivity of binuclear pyrazolate-bridged rhodium complexes with isocyanides. Molecular structure of $[\text{Rh}_2(\mu\text{-pz})\text{I}_2(\text{CNBu}')_2(\mu\text{-Ph}_2\text{PCH}_2\text{PPh}_2)_2]\text{BF}_4$ (pz = pyrazolate), 1427–34

DIAZAPHOSPHETIDINE

Synthesis and crystal structures of three four-membered ring compounds containing PN₂Ti skeletons, 2173–8

DIAZENE

Reactivity of new osmium dihydrides with arenediazonium cations: preparation and properties of bis(aryldiazene) and mono(aryldiazenido) complexes. A comparison with analogous iron and ruthenium derivatives, 2353–8

DIAZOMETHANE

Preparation of (chloromethyl)palladium(II) derivatives from complexes of palladium dichloride by reaction with diazomethane or bis(chloromethyl)mercury, 761–6

DICARBAUNDECABORATE

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 84. Carbaborane tungsten–platinum complexes having a $\mu\text{-CC}_6\text{H}_3\text{Me}_2$ -2,6 ligand: crystal structures of $[\text{WP}(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_n(\text{PEt}_3)(\mu\text{-}\sigma\text{:}\eta^5\text{-C}_2\text{B}_9\text{H}_8\text{Me}_2)]$ ($n = 2$ or 3), 303–12

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 86. Alkylidyne(carbaborane)-molybdenum–gold, –rhodium and –iron complexes; crystal structure of $[\text{NEt}_4]_2[\text{MoFe}_2(\mu\text{-C}_6\text{H}_4\text{Me-4})(\mu\text{-}\sigma\text{:}\sigma'\text{:}\eta^5\text{-C}_2\text{B}_9\text{H}_8\text{Me}_2)]$, 797–808

DICHLOROETHANE

Formation of mixed-ligand complexes of cadmium(II) halide with tri-n-octylphosphine oxide and pyridine or substituted pyridines in 1,2-dichloroethane, 1259–64

DIETHYLCARBAMATE

Formation of alkylurethanes from carbon dioxide by regioselective O-alkylation of alkali-metal *N,N*-diethylcarbamates in the presence of complexing agents, 1007–10

DIHYDROGEN

Influence of added hydrogen on the kinetics and mechanism of thermal decomposition of tetraborane(10) and of pentaborane(11) in the gas phase, 391–8

Hydrogen production from ethanol catalysed by Group 8 metal complexes, 489–96

Kinetics of reactions of hydrogen with $[\text{Os}_3(\text{CO})_{11}(\text{NCMe})]$ and $[\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2]$, 2235–40

Photochemical reactivity of $2[\text{Ir}^{\text{III}}\text{HCl}_2(\text{PPr}^{\text{I}}_3)_2]\text{-H}_2$. Formation of monohydrido isomers of Ir^{III} and evidence of photoinduced oxidative addition of H₂ to Ir^{III}, 2367–70

DI-IMINE

Dimeric mixed-ligand copper(II) complexes with di-imines and salicylaldehyde, 1435–8

DIKETONE

Ligand additivity in the oxidation potentials of bidentate mixed-ligand ruthenium(II) complexes, 371–4

DIMETHYLFORMAMIDE

Spectrophotometric and calorimetric studies on the formation of binary (2,2'-bipyridine)nickel(II) and ternary (2,2'-bipyridine)chloronickel(II) complexes in *N,N*-dimethylformamide, 655–60

DIMETHYLGLYOXIMATE

Relative importance of the inductive and steric effects of alkyl groups (R) in understanding the nature of the Co^{III}–C bond in alkyl cobaloximes, and comments on related alkyl cobalamins, 115–8

Steric effect of the adamantyl ligand on crystal structure and dissociation rates in organocobalt B₁₂ models, 1941–6

DIMETHYL SULPHOXIDE

Studies in Werner clathrates. Part 9. Structures with bis(isothiocyanato)tetrakis(4-phenylpyridine)nickel(II) as host, 119–26

The nickel(II)–bromide system in dimethyl sulphoxide: a detailed study of the influences of temperature and mole ratio, 325–30

Mode of binding of dimethyl sulphoxide in platinum(II) and palladium(II) cationic complexes containing a chelated diamine. X-Ray crystal structures of (2,2'-bipyridyl)chloro(dimethyl sulphoxide-*O*)palladium(II) tetrafluoroborate and chloro(dimethyl sulphoxide-*S*)(ethylenediamine)palladium(II) perchlorate, 1265–72

Lanthanide ion complexes of the calixarenes. Part 4. Double inclusion by *p-t*-butylcalix[4]arene (H₄L). Crystal structures of $[\text{Eu}_2(\text{H}_4\text{L})_2(\text{dmf})_4]\cdot 7\text{dmf}$ (dmf = dimethylformamide) and $\text{H}_4\text{L}\cdot \text{dmso}$ (dmso = dimethyl sulphoxide), 2217–22

DINITROGEN

A theoretical study on the co-ordination of dinitrogen and related molecules to nickel(0): ‘*ab initio*’ calculations on the model compounds $[\text{Ni}(\text{PH}_3)_2(\text{N}_2)]$, $[\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{CH}_2)]$, and $[\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{H}_2)]$, 33–8

DIOXANE

Reactivity studies on the hydrolysis of iron(II) complexes of substituted phenanthrolines in dioxane–water solvents, 1105–8

DIOXYGEN

Reactivity of metallic copper with acidic organic compounds in the presence of molecular oxygen. Crystal structure of $[\text{Cu}(\text{NCO})(\text{py})_2] \cdot \text{H}_2\text{O}$, 921–6

Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(III)–ethanol–carbon tetrachloride system, 1197–202

Autoxidation of a dicopper(I) macrocyclic complex: evidence for a mixed-valence intermediate, 1351–60

Reactions of heteropolyanions in non-polar solvents. Part 3.

Activation of dioxygen by manganese(II) centres in polytungstates. Oxidation of hindered phenols, 1483–90

DIPHENYLTIOPHOSPHINITE

Preparation and structural studies of diphenylthiophosphinito rhodium(II) complexes. X-Ray structure analysis of $[(\text{C}_5\text{Me}_5)\text{Rh}]_2(\mu\text{-Cl})_2(\mu\text{-Ph}_2\text{PS})[\text{BPh}_4]$, 1987–92

DIPHOSPHINE

Synthesis and structural studies on silver–tin complex salts with bis(diphenylarsino)methane and bis(diphenylphosphino)methane, 247–52

Reactions of the bis(alkyne) complexes $[\text{WI}_2(\text{CO})(\text{NCMe})(\eta^2\text{-RC}_2\text{R})_2]$ ($\text{R} = \text{Me}$ or Ph) with mono- and bi-dentate phosphine-donor ligands and the X-ray crystal structure of $[\text{WI}_2(\text{CO})(\text{Ph}_2\text{P}(\text{CH}_2)\text{PPPh}_2)(\eta^2\text{-MeC}_2\text{Me})]$, 295–302

Mechanistic studies on hydrazido(2–)complexes. Cleavage of the nitrogen–nitrogen bond in the reaction of

$[\text{Mo}\{\text{NN}(\text{CH}_2)_4\text{CH}_2\}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPPh}_2)_2]$ with acid, 425–30
Six- and five-co-ordinate ruthenium(II) complexes with 1,2-bis(dicyclohexylphosphino)ethane. X-Ray crystal structure of the d^6 five-co-ordinate $[\text{RuCl}\{(\text{C}_6\text{H}_{11})_2\text{PCH}_2\text{CH}_2\text{P}(\text{C}_6\text{H}_{11})_2\}_2]\text{PF}_6$, 1045–52

Synthesis of hydridotrifluoromethyl complexes of platinum(II). A spectroscopic investigation of the *trans* and *cis* influence of ligands in hydridoplatinum(II) compounds, 1149–60

Syntheses and redox properties of mixed isocyanide, carbonyl, or nitrile complexes of rhodium(I) *trans*- $[\text{Re}(\text{CNMe})\text{L}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPPh}_2)_2\text{JX}$ [$\text{L} = \text{CNR}$ ($\text{R} = \text{alkyl}$ or aryl), CO , or NCMe ; $\text{X} = \text{Cl}$, BF_4^- , or PF_6^-], 1209–16

The heteronuclear cluster chemistry of the Group 1B metals. Part 11. Effect of the nature of the bidentate diphosphine ligand on the metal framework structures of the gold heteronuclear cluster compounds $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})(\mu\text{-H})\{\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPPh}_2\}(\text{CO})_{12}]$ ($n = 1\text{--}6$). X-Ray crystal structures of $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})(\mu\text{-H})\{\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPPh}_2\}(\text{CO})_{12}]$ ($n = 1$ or 2), 1227–36

Substitution reactions of the zerovalent iron complexes $[\text{Fe}(\eta\text{-C}_6\text{H}_6)(\text{PMMe}_3)_2]$ and $[\text{Fe}(\eta\text{-C}_6\text{H}_6)(\text{dmpe})]$ ($\text{dmpe} = \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_2$), 1385–92

Crystal and molecular structure of [1,2-bis(diethylphosphino)ethane mono-oxide-*P,O*]trichloro-oxomolybdenum(V), 1417–20

Reactivity of binuclear pyrazolate-bridged rhodium complexes with isocyanides. Molecular structure of $[\text{Rh}_2(\mu\text{-pz})\text{I}_2(\text{CNBu}')_2(\mu\text{-Ph}_2\text{PCH}_2\text{PPPh}_2)]\text{BF}_4^-$ ($\text{pz} = \text{pyrazolate}$), 1427–34

Synthesis, characterization, and thermolytic behaviour of η^3 -allyl(methyl)(tertiary phosphine)palladium(II), and X-ray structure of the cluster $[\text{Pd}_4(\mu\text{-C}_3\text{H}_5)_2(\mu\text{-Cl})_2(\mu\text{-dppm})_2]\text{4C}_4\text{H}_4\text{O}$ ($\text{dppm} = \text{Ph}_2\text{PCH}_2\text{PPPh}_2$), 1519–26

Nucleophilic (Michaeli) additions to $(\text{Ph}_2\text{P})_2\text{C}=\text{CH}_2$ when chelated to platinum(II) dichloride or di-iodide, 1527–30

Base-promoted cleavage of a P–C bond in $[\text{PtCl}_2(\text{Ph}_2\text{PCH}_2\text{PPPh}_2)\text{PP}']$ under mild conditions, 2017–22

The heteronuclear cluster chemistry of the Group 1B metals. Part 12. Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds $[\text{M}_2\text{Ru}_4(\mu_3\text{-H})_2(\mu\text{-L})(\text{CO})_{12}]$ [$\text{M} = \text{Cu}$ or Ag ; $\text{L} = \text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPPh}_2$ ($n = 1\text{--}6$) or *cis*- $\text{Ph}_2\text{PCH}=\text{CHPPPh}_2$; $\text{M} = \text{Cu}$, $\text{L} = \text{Ph}_2\text{As}(\text{CH}_2)_n\text{PPPh}_2$ ($n = 1$ or 2)] in solution, 2039–48

Interaction of benzo-1,3-thiazoline-2-thione and related ligands with copper(II) salts and the single-crystal X-ray structure of μ -[1,2-bis(diphenylphosphino)ethane]-bis{(benzo-1,3-thiazole-2-thiolato-*S*}[1,2-bis(diphenylphosphino)ethane]copper(I)}, 2167–72

A convenient route to binuclear osmium(I) complexes containing diphosphine and arsinophosphine ligands. X-Ray crystal structure of $[\text{Os}_2(\text{MeCO}_2)_2(\text{CO})_4(\text{dppm})_2]$, 2303–8

Reactions of 1-alkynes with *trans*-

$[\text{ReCl}(\text{N}_2)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPPh}_2)_2]$: preparation of the vinylidene compounds *trans*- $[\text{ReCl}(\text{C}=\text{CHR})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPPh}_2)_2]$ ($\text{R} = \text{alkyl}$ or aryl) and X-ray structure of *trans*- $[\text{ReCl}(\text{C}=\text{CHPh})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPPh}_2)_2]$, 2381–8

DIPOLÉ MOMENT

Solid-state structure and solution conformation of 2,2,4,6,6-hexa(*p*-chlorophenoxy)cyclotri- λ^5 -phosphazatriene. X-Ray and dipole moment studies, 757–60

DIPYRIDYLAMINE

Lewis-base adducts of Group 11 metal(I) compounds. Part 58.

Syntheses, X-ray structures, and far-infrared spectra of the novel $[\text{CuX}_2(\text{bidentate ligand})]^-$ ($\text{X} = \text{Cl}$ or Br) anion salts of the $[\text{Cu}(\text{dpa})_2]^+$ cation and the dimeric iodide $[(\text{dpa})\text{CuI}_2\text{Cu}(\text{dpa})]$ ($\text{dpa} = \text{di}(2\text{-pyridyl})\text{amine}$), 1639–44

DITHIADIAZOLE

Direct insertion of a nitrogen atom into the S–S bond of a 1,2,3,5-dithiadiazole ring in a direct current nitrogen glow discharge, and X-ray crystal structure of 4-methyl-1,2,3,5-dithiadiazole, 1705–14

DITHIADIAZOLYL

The 4-phenyl-1,2,3,5-dithiadiazole complex $[\text{Fe}_2(\text{CO})_6(\text{PhCN}_2\text{S}_2)]$: its preparation, crystal structure, and an extended-Hückel molecular-orbital study of the bonding, 2229–34

DITHIAOCTANE

Unusual chelating properties of the ligand 1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-dithiaoctane (bddo). Crystal structures of $\text{Ni}(\text{bddo})(\text{NCS})_2$, $\text{Zn}(\text{bddo})(\text{NCS})_2$, and $\text{Cd}_2(\text{bddo})(\text{NCS})_4$, 2309–14

DITHIARSOLANE

Some transition-metal complexes of 1,3,2-dithiarsolane, 885–8

DITHIOACETATE

Solid-state high-resolution carbon-13 magnetic resonance spectra of dithioacetate complexes of platinum, palladium, and nickel, 2457–60

DITHIOCARBAMATE

Actinide structural studies. Part 17. The crystal and molecular structures of four uranyl(vi) complexes with *N,N*-dialkylthiocarbamate ligands, 471–80

Lewis base adducts of Group 11 metal(I) complexes. Part 48.

Heterobimetallic tris(dithiocarbamato)cobalt(III)-copper(I) halide adducts as a vehicle for novel copper(I) halide clusters (structurally characterized), 971–6

DITHIOCARBONATE

Correlations between nuclear magnetic resonance spectra and crystal structures. A carbon-13 nuclear magnetic resonance study in the solid state of the Group 15 xanthates $[\text{A}(\text{S}_2\text{COR})_3]$ ($\text{A} = \text{As}^{III}$, Sb^{III} , or Bi^{III} ; $\text{R} = \text{Me}$, Et , or Pr^i), 233–6

Synthesis and characterization of bis(xanthate) complexes of Mo^{IV} and W^{VI} . Crystal structure of $[\text{W}(\text{S}_2\text{COPr}^i\text{-S})(\text{S}_2\text{COPr}^i\text{-S})\text{CO}_2(\text{PMe}_3)_2]$, 2003–10

DITHIOCARBOXYLATE

Crystal structure of a novel tetrานuclear dithiocarboxylato copper(I) complex, $[\text{Cu}(\text{S}_2\text{CC}_6\text{H}_4\text{Me}-p)_4(\text{PPh}_3)_2]$, 753–6

DITHIOLATE

Crystal structure of the new low-dimensional metal $\text{Cs}_{0.82}[\text{Pd}(\text{S}_2\text{C}_2\text{CN})_2] \cdot 0.5\text{H}_2\text{O}$, 1101–4

DITHIO-OXALATE

Tetrahydrofulvalenium salts of planar Pt^{II} , Pd^{II} , and Cu^{III} 1,2-dithio-oxalato-*S,S'* anions. Synthesis, chemistry and molecular structures of bis(tetrahydrofulvalenium) bis(1,2-dithio-oxalato-*S,S'*)palladate(II), $[\text{ttf}]_2[\text{Pd}(\text{S}_2\text{C}_2\text{O}_2)_2]$, and of bis(tetrahydrofulvalenium)tetrahydrofulvalene bis(1,2-dithio-oxalato-*S,S'*)platinate(II), $[\text{ttf}]_3[\text{Pt}(\text{S}_2\text{C}_2\text{O}_2)_2]$, 719–28

DITHIOPHOSPHATE

Complexes of lead(II) bis(*O,O'*-dialkyl dithiophosphates) with nitrogen donor ligands. The crystal structures of $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\text{en}$ ($\text{en} = \text{ethylenediamine}$), $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\text{bipy}$ ($\text{bipy} = 2,2'\text{-bipyridine}$), and $\{\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\text{en}\}_2$, 2443–8

DYNAMICS

Solution structure and dynamic behaviour of two isomers of $[\text{Fe}_3(\text{CO})_9\{\text{P}(\text{OR}_3)_3\}]$ ($\text{R} = \text{Me}$, Et , or Ph) derivatives, 1277–82

On the relationship between crystallographic and spectroscopic evidence of dynamic processes in the solid state: the cases of *cis*- and *trans*- $[\text{Fe}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_4]$, 1721–6

ELECTROCHEMISTRY

Structural, spectroscopic, and electrochemical characterization of tetrakis- μ -(2-pyrrolidinone)-dirhodium(II) and tetrakis- μ -(δ -valerolactamato)-dirhodium(II), 93–100

ELECTROCHEMISTRY (contd)

- Synthesis, electrochemistry, and electron spin resonance spectra of some seventeen-electron complexes containing the $\{\text{Mo}(\text{NO})\}^{2+}$ core and the molecular structure of $[\text{Mo}(\text{NO})\{\text{HB}(\text{Me}_2\text{pz})_3\}(\text{pyr})_2]$, 127–32
- Thermodynamic and electrochemical behaviour of lanthanide cryptates in *N,N*-dimethylformamide, 367–70
- Ligand additivity in the oxidation potentials of bidentate mixed-ligand ruthenium(II) complexes, 371–4
- Syntheses, electrochemistry, and spectroscopy of dirhodium(II) tetra-acetamide and tetrakis(trifluoroacetamide) complexes with axial Group 15 substituents. The X-ray crystal structures of $[\text{Rh}_2(\text{CH}_3\text{CONH})_4(\text{AsPh}_3)_2]$ and $[\text{Rh}_2(\text{CH}_3\text{CONH})_{4-n}(\text{CH}_3\text{CO}_2)_n(\text{MPh}_3)_2]$, M = As or Sb, $n \approx 1$, 581–8
- The chemical and electrochemical one-electron oxidation of diamidonaphthalene-bridged complexes: paramagnetic $[\text{Rh}_2]^{3+}$ containing A-frames, 683–8
- Trinuclear mixed-valence complexes with the linear $[\text{Rh}_3]^{4+}$ core, 689–92
- Formation in solution, synthesis, and electrochemical study of oxalato complexes of *N,N'*-ethylenebis(salicylideneiminato)-chromium(III) and -iron(III): crystal structures of piperidinium [*N,N'*-ethylenebis(salicylideneiminato)](oxalato- $O^1\text{O}^2$)-chromate(III) and -ferrate(III), 729–38
- Precursor catenand complexes: synthesis, structure, and electrochemistry of bis(2,6-di-iminopyridyl) complexes of nickel(II). The single-crystal X-ray structure of $[\text{NiL}_2][\text{BF}_4]_2$, 965–70
- Blue copper models. Spectroscopic and electrochemical studies of copper(II) complexes with new ligand systems containing sulphur and nitrogen donor atoms, 1161–70
- Autoxidation of a dicopper(I) macrocyclic complex: evidence for a mixed-valence intermediate, 1351–60
- Effect of axially bound anions on the electroreduction of tin(IV) porphyrins in tetrahydrofuran, 1531–6
- Synthesis, structure, and electrochemical reduction of 2-(arylazo)pyridine complexes of rhodium(III), 1635–8
- Platinum metal thioether macrocyclic complexes: synthesis, electrochemistry, and single-crystal X-ray structures of *cis*- $[\text{RhCl}_2\text{L}^2]\text{PF}_6$ and *trans*- $[\text{RhCl}_2\text{L}^3]\text{PF}_6$ ($\text{L}^2 = 1,4,8,11$ -tetrahydroacyclohexadecane, $\text{L}^3 = 1,5,9,13$ -tetrahydroacyclohexadecane), 1675–80
- Reductive nitrosylation of tetraoxometalates. Part 15. Generation of $\text{Mn}(\text{NO})^{3+}$, $\text{Mn}(\text{NO})^{2+}$ and $\text{Mn}(\text{NO})^{2+}$ moieties directly from MnO_4^- . Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese, 1963–72
- Spectroscopic and electrochemical study of copper and zinc complexes with a N_3S_2 ligand donor set. Crystal structure of the copper(II) complex derived from 1,3-propylenediamine and 3-formyl-1-phenyl-2(1*H*)-pyridinethione, 1979–86
- Ruthenium oxo complexes of macrocyclic tertiary amines. Synthesis, electrochemistry, and reactivities of ruthenium oxo complexes of *meso*-2,3,7,11,12-pentamethyl-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), and X-ray crystal structure of *trans*- $[\text{Ru}^{\text{IV}}\text{L}^1(\text{O})(\text{NCO})]\text{ClO}_4$, 2011–6
- Reduction–oxidation properties of organotransition-metal complexes. Part 30. Triazenediyl-bridged carbonylrhodium bipyridyl complexes with $[\text{Rh}_2]^{4+}$, $[\text{Rh}_2]^{3+}$, and $[\text{Rh}_2]^{4+}$ cores, and the X-ray structures of $[\text{Rh}_2(\text{C}\ddot{\text{O}}_2)(\text{bipy})(\mu-\text{RNNNR}_2)_2][\text{BF}_4] \cdot \text{CH}_2\text{Cl}_2$ and $[\{\text{Rh}_2\text{I}(\text{CO})(\text{bipy})(\mu-\text{RNNNR}_2)_2\}_2][\text{PF}_6]_2 \cdot 2.5\text{CH}_2\text{Cl}_2$ (R = *p*-tolyl), 2049–60
- Synthesis, structure, and redox properties of dicopper(II,III) complexes of *N*-(hydroxyphenyl)salicylamide and homologues. Facile oxidation to dicopper(II,II) and -(II,III) species; crystal structure of bis(tetrabutylphosphonium) bis{[(2-oxidobenzoyl)(2'-oxidophenyl)amido(3-)– NO_2^2 , $\mu\text{-O}^2$]cuprate(II)}, 2089–94
- Complexes of technetium, rhenium, and rhodium with sexidentate Schiff-base ligands, 2115–20
- Electrochemistry of the quasi-reversible bis[2-ethyl-2-hydroxybutanoato(2-)Joxochromate(–V) and (–IV) and bis[2-hydroxy-2-methylbutanoato(2-)Joxochromate(–V) and (–IV) redox couples and the crystal and molecular structure of sodium bis[2-ethyl-2-hydroxybutanoato(2-)Joxochromate(V) sesquihydrate, 2205–10
- Synthesis, spectroscopic, electrochemical, and magnetic properties of dimolybdenum(II,II), diruthenium(II,III) and -(II,II) complexes containing bridging aspirinate (2-acetoxybenzoate) ligands, 2435–42

Reductive nitrosylation of tetraoxometalates. Part 15. Generation of $\text{Mn}(\text{NO})^{3+}$, $\text{Mn}(\text{NO})^{2+}$, and $\text{Mn}(\text{NO})^{2+}$ moieties directly from MnO_4^- . Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese (1989, 1963), 2461

ELECTRODE

Comparative study on the multielectron reduction of $\text{HOC}_2\text{H}_4\text{N}_3$ catalyzed by $[\text{NBu}_4^n]_4[\text{MoFe}_3\text{S}_4(\text{SPh})_3(o-\text{O}_2\text{C}_6\text{Cl}_4)]_2^-$ and $[\text{NBu}_4^n]_2[\text{Fe}_4\text{S}_4(\text{SPh})_4]$ -modified glassy carbon electrodes, 1547–54

ELECTRODE POTENTIAL

Ligand additivity in the oxidation potentials of bidentate mixed-ligand ruthenium(II) complexes, 371–4

Syntheses and redox properties of mixed isocyanide, carbonyl, or nitrile complexes of rhenium(I) *trans*- $[\text{Re}(\text{CNMe})\text{L}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPH}_2)_2]\text{X}$ [L = CNR (R = alkyl or aryl), CO, or NCMe; X = Cl, BF_4^- , or PF_6^-], 1209–16

ELECTRON DIFFRACTION

The structure of $\text{PO}(\text{OPF}_2)_3$ in the gaseous and crystalline phases, 223–8

The molecular structures of gaseous tetrafluorohydridophosphorane, HPF_4 , and trifluorodihydridophosphorane, H_2PF_3 , as determined by electron diffraction, 545–50

Isopropylsilylamine and disilyl-*t*-butylamine: preparation, spectroscopic properties, and molecular structure in the gas phase, determined by electron diffraction, 779–84

Gas-phase molecular structure of *NN'*-bis(trimethylsilyl)sulphur diimide, determined by electron diffraction, 859–62

ELECTRONIC

A further investigation of the polarized single-crystal electronic absorption spectrum of $[\text{AsPh}_4]_4[\text{CrOCl}_4]$, 1–4

Spectroscopic studies in matrix isolated tungsten chlorides and bromides, 313–6

The nickel(II)-bromide system in dimethyl sulphoxide: a detailed study of the influences of temperature and mole ratio, 325–30

Spectrophotometric study of complex formation in copper(II) mono-, di-, and tri-ethanolamine systems, 569–74

Syntheses, electrochemistry, and spectroscopy of dirhodium(II) tetra-acetamide and tetrakis(trifluoroacetamide) complexes with axial Group 15 substituents. The X-ray crystal structures of $[\text{Rh}_2(\text{CH}_3\text{CONH})_4(\text{AsPh}_3)_2]$ and $[\text{Rh}_2(\text{CH}_3\text{CONH})_{4-n}(\text{CH}_3\text{CO}_2)_n(\text{MPh}_3)_2]$, M = As or Sb, $n \approx 1$, 581–8

Spectrophotometric and calorimetric studies on the formation of binary (2,2'-bipyridine)nickel(II) and ternary (2,2'-bipyridine)chloronickel(II) complexes in *N,N*-dimethylformamide, 655–60

Electronic spectral study of the aqua \rightleftharpoons hydroxo equilibrium of model iron(III) haems encapsulated in aqueous detergent micelles, 1003–6

Thermodynamic and spectrophotometric investigation of complex formation between hydrogen, cobalt(II), nickel(II), and copper(II) ions and 2-amino-*N*-hydroxypropanamide in aqueous solution, 1299–310

Mixed-metal complexes in solution. A potentiometric and spectrophotometric study of copper(II)-nickel(II)- δ -hydroxylysine complexes in aqueous solution, 1745–50

Electron spin resonance study of copper(II) complexes of X-glycine and glycyl-X type dipeptides, and related tripeptides. Variation of co-ordination modes with ligand excess and pH in fluid and frozen aqueous solutions, 1925–32

Spectral intensity and charge distributions in the tetrahedral chromophores, bis(3-amino-2-methylpropane-2-thiolato)- and diacetatobis(ethylenethiourea)-cobalt(II), 2109–14

Optical absorption and magnetic circular dichroism studies of hydrogen, copper(II), zinc(II), nickel(II), and cobalt(II) crown ether-substituted monomeric and dimeric phthalocyanines, 2397–406

ELECTRON TRANSFER

Kinetics of the oxidation of chloride ions by tris(2,2'-bipyridine)nickel(III) ions in aqueous perchlorate media, 151–4

Kinetics and mechanism of the reaction between 4-*t*-butylphenolate anion and tetrahydroxoargentate(III) in aqueous alkaline media, 217–22

Stabilisation of trivalent nickel through 1:2 co-ordination by cyclic terdentate ligands $\text{CH}_2\text{CH}_2\text{NH}(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{X}$ (X = NH, O, or S), 229–32

Reduction–oxidation properties of organotransition-metal

ELECTRON TRANSFER (contd)

- complexes. Part 29. Pentaphenylcyclopentadienyl complexes of ruthenium, 283–8
 Kinetic studies of the oxidation reaction of arsenic(III) to arsenic(V) by peroxodisulphate ion in aqueous alkaline media, 357–60
 Ligand additivity in the oxidation potentials of bidentate mixed-ligand ruthenium(II) complexes, 371–4
 One-electron transfer processes in the reaction of tin(II) halides with substituted *o*-quinones; crystal structure of bis(3,5-di-*t*-butylcatecholato)tin-1,10-phenanthroline-dimethylformamide (1/1/2), 439–46
 Oxidation of silver metal and silver(I) cations by molybdenum, tungsten, and uranium hexafluorides and by the nitrosonium cation in acetonitrile. Preparation and properties of silver(III) hexafluorometalates(V), 461–4
 Oxidation of a Pt₂(μ-SO₂) moiety to μ-SO₃ and the crystal structure of [Pt₂(μ-SO₃)(μ-SO₂)(C₈H₁₂)₂], 557–60
 The chemical and electrochemical one-electron oxidation of diamidonaphthalene-bridged complexes: paramagnetic [Rh₂]³⁺ containing A-frames, 683–8
 Trinuclear mixed-valence complexes with the linear [Rh₃]⁴⁺ core, 689–92
 Catalytic oxidation of water mediated by rhenium oxides, 849–52
 Blue copper models. Spectroscopic and electrochemical studies of copper(II) complexes with new ligand systems containing sulphur and nitrogen donor atoms, 1161–70
 Syntheses and redox properties of mixed isocyanide, carbonyl, or nitrile complexes of rhenium(I) *trans*-[Re(CNMe)L(Ph₂PCH₂CH₂PPPh₂)₂]X [L = CNR (R = alkyl or aryl), CO, or NCMe; X = Cl, BF₄, or PF₆], 1209–16
 Photo-oxidation of alcohols by hexachlorometallate(IV) ions (M = Pt, Pd, or Ir): spin trapping and matrix isolation electron spin resonances studies, 1237–40
 Autoxidation of a dicopper(I) macrocyclic complex: evidence for a mixed-valence intermediate, 1351–60
 Reactions of heteropolyanions in non-polar solvents. Part 3. Activation of dioxygen by manganese(II) centres in polytungstates. Oxidation of hindered phenols, 1483–90
 Effect of axially bound anions on the electroreduction of tin(IV) porphyrins in tetrahydrofuran, 1531–6
 Comparative study on the multielectron reduction of HOC₂H₄N₃ catalyzed by [NBuⁿ]₄[MoFe₃S₄(SPh)₃(*o*-O₂C₆Cl₄)₂] and [NBuⁿ]₄[Fe₄S₄(SPh)₄]-modified glassy carbon electrodes, 1547–54
 Synthesis, structure, and electrochemical reduction of 2-(arylaquo)pyridine complexes of rhodium(III), 1635–8
 Kinetics of the uncatalysed and silver(I)-catalysed oxidation of hydrogen peroxide by [ethylenebis(biguaine)]silver(III) ion in acid perchlorate media, 1645–50
 Structural effects of one-electron oxidation on a cyclobutadieneiron derivative: X-ray crystal structures of [Fe(CO){P(OMe)₃}₂(η-C₄Ph₄)] and [Fe(CO){P(OMe)₃}₂(η-C₄Ph₄)]BF₄, 1751–8
 Co-ordination compounds of indium. Part 47. Indium(II) and thallium(II) derivatives of alkanedithiols, 1759–64
 Ruthenium oxo complexes of macrocyclic tertiary amines. Synthesis, electrochemistry, and reactivities of ruthenium oxo complexes of *meso*-2,3,7,11,12-pentamethyl-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L¹), and X-ray crystal structure of *trans*-[Ru^{IV}L¹(O)(NCO)]ClO₄, 2011–6
 Reduction–oxidation properties of organotransition-metal complexes. Part 30. Triazenido-bridged carbonylrhodium bipyridyl complexes with [Rh₂]²⁺, [Rh₂]³⁺, and [Rh₂]⁴⁺ cores, and the X-ray structures of [Rh₂(CO)₂(bipy)(μ-RNNNR)₂][BF₄]₂CH₂Cl₂ and [{Rh₂I(CO)(bipy)(μ-RNNNR)₂}][PF₆]₂·2.5CH₂Cl₂ (R = *p*-tolyl), 2049–60
 Oxo transfer reactions. Mechanistic studies on the oxidation of triphenylphosphine by *trans*-[Ru^{VI}L(O₂)₂]²⁺ (L = 1,4,8,11-tetramethyl-1,4,8,11-tetra-azacyclotetradecane), 2065–8
 Synthesis, structure, and redox properties of dicopper(II,II) complexes of *N*-(hydroxyphenyl)salicylamide and homologues. Facile oxidation to dicopper(II,II) and -(III,III) species; crystal structure of bis(tetrabutylphosphonium) bis{[(2'-oxidoibenzoyl)-(2'-oxidophenyl)amido(3-)-NO₂,μ-O²⁻]cuprate(II)}, 2089–94
 Synthesis and X-ray crystal structure of tetra(2-methylphenyl)molybdenum(IV), Mo(2-MeC₆H₄)₄. Redox chemistry of M(2-MeC₆H₄)₄ compounds of molybdenum, ruthenium, and osmium, 2149–54
 Electrochemistry of the quasi-reversible bis[2-ethyl-2-hydroxybutanoato(2-)]oxochromate-(V) and -(IV) and bis[2-

hydroxy-2-methylbutanoato(2-)]oxochromate-(V) and -(IV) redox couples and the crystal and molecular structure of sodium bis[2-ethyl-2-hydroxybutanoato(2-)]oxochromate(V) sesquihydrate, 2205–10

ELECTROPHILICITY

Metal cluster-induced electrophilicity of arenes, 481–8

EQUILIBRIUM

The equilibrium of strontium ion exchange of sodium titanate, Na₄Ti₉O₂₀·xH₂O, 101–4

The nickel(II)-bromide system in dimethyl sulphoxide: a detailed study of the influences of temperature and mole ratio, 325–30

Electronic spectral study of the aqua ⇌ hydroxo equilibrium of model iron(III) haems encapsulated in aqueous detergent micelles, 1003–6

ES.R.

Synthesis, electrochemistry, and electron spin resonance spectra of some seventeen-electron complexes containing the {Mo(NO)}²⁺ core and the molecular structure of [Mo(NO){HB(Me₂Pz)₃}(pyr)₂], 127–32

One-electron transfer processes in the reaction of tin(II) halides with substituted *o*-quinones; crystal structure of bis(3,5-di-*t*-butylcatecholato)tin-1,10-phenanthroline-dimethylformamide (1/1/2), 439–46

Photochemical studies of alkylammonium molybdates. Part 8. Location of protons interacting with the paramagnetic electron in a single crystal of photoirradiated [NH₃Pr⁺]₆[Mo₈O₂₆(OH)₂]·2H₂O, 661–70

The chemical and electrochemical one-electron oxidation of diamidonaphthalene-bridged complexes: paramagnetic [Rh₂]³⁺ containing A-frames, 683–8

Photo-oxidation of alcohols by hexachlorometallate(IV) ions (M = Pt, Pd, or Ir): spin trapping and matrix isolation electron spin resonances studies, 1237–40

Proton electron nuclear double resonance study of oxovanadium(IV) complexes of D-galacturonic and polygalacturonic acids, 1283–8

Proton electron nuclear double resonance study of oxovanadium(IV) complexes of *o*-diphenolic ligands, 1289–92

Synthesis, X-ray characterization, and Q-band electron spin resonance studies of heterometallic, polymeric compounds formed by yttrium and lanthanide(III) nitrates with HgL₂ (L = C₄H₆NO, the anion of 2-pyrrolidone); crystal structures of [Hg₂EuL₄(NO₃)₃]_n and [Hg₃Tb₂L₆(NO₃)₆]_n, 1439–48

Electron spin resonance study of copper(II) complexes of X-glycine and glycyl-X type dipeptides, and related tripeptides. Variation of co-ordination modes with ligand excess and pH in fluid and frozen aqueous solutions, 1925–32

Binuclear copper(II) complexes of new dinucleating ligands with a pyrazolate group as an endogenous bridge. Effects of exogenous azide and acetate bridges on magnetic properties, 2077–82

ETHANOL

Hydrogen production from ethanol catalysed by Group 8 metal complexes, 489–96

Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(II)-ethanol–carbon tetrachloride system, 1197–202

Structure of μ-fluoro-bis[bis(2,2'-bipyridyl)fluoroniobel(IV)] fluoride–ethanol (1/2) trihydrate, revealing several unique features including strong hydrogen bonds between fluoride ions and ethanol molecules, 1273–6

ETHANOLAMINE

Spectrophotometric study of complex formation in copper(II) mono-, di-, and tri-ethanolamine systems, 569–74

ETHYLENE

Rearrangements of >C=N– functionalities promoted by cobalt(II): reactivity of cyclopentadienylbis(ethylene)cobalt(I) with dibenzophenone azine, benzophenone oxime, benzylidene(phenyl)amine and crystal structure determinations of the products, 1957–62

ETHYLENEDIAMINE

Mode of binding of dimethyl sulphoxide in platinum(II) and palladium(II) cationic complexes containing a chelated diamine.

X-Ray crystal structures of (2,2'-bipyridyl)chloro(dimethyl sulphoxide-*O*)palladium(II) tetrafluoroborate and chloro(dimethyl sulphoxide-*S*)(ethylenediamine)palladium(II) perchlorate, 1265–72

Complexes of lead(II) bis(*O,O'*-dialkyl dithiophosphates) with nitrogen donor ligands. The crystal structures of

Pb[S₂P(OEt)₂]₂·en (en = ethylenediamine), Pb[S₂P(OEt)₂]₂·bipy (bipy = 2,2'-bipyridine), and {Pb[S₂P(OEt)₂]₂·en}, 2443–8

ETHYNE

Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene ($C=CHOEt$) and ethoxyethyne ($CH\equiv COEt$) ligands respectively, 5–12

EUROPIUM

Lanthanide ion complexes of the calixarenes. Part 4. Double inclusion by *p*-t-butylcalix[4]arene (H_4L). Crystal structures of $[Eu_2(HL)_2(dmf)_4] \cdot 7dmf$ ($dmf =$ dimethylformamide) and $H_4L \cdot dmsO$ ($dmsO =$ dimethyl sulphoxide), 2217–22

EXCHANGE

Zirconium phosphate ester interchange reactions, 1617–24

FILM

Functional metallocacyclic derivatives and their polymers. Part 23. Biaxially orientated Langmuir–Blodgett films of substituted metallophthalocyanines, 1543–6

FLUORIDE

Oxidation of silver metal and silver(i) cations by molybdenum, tungsten, and uranium hexafluorides and by the nitrosonium cation in acetonitrile. Preparation and properties of silver(III) hexafluorometalates(v), 461–4

Radiotracers in fluorine chemistry. Part 12. Reactions between caesium fluoride and arsenic pentafluoride, boron trifluoride, sulphur tetrafluoride, carbonyl fluoride, and carbon dioxide under heterogeneous conditions at room temperature.

Radiotracer studies using fluorine-18, sulphur-35, and carbon-14, 937–42

Structure of μ -fluoro-bis[bis(2,2'-bipyridyl)fluoronickel(II)] fluoride–ethanol (1/2) trihydrate, revealing several unique features including strong hydrogen bonds between fluoride ions and ethanol molecules, 1273–6

FLUXIONALITY

Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene ($C=CHOEt$) and ethoxyethyne ($CH\equiv COEt$) ligands respectively, 5–12

A dynamic nuclear magnetic resonance study of trimethylplatinum(IV) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part 2. High-temperature fluxionality, 711–8

The mechanisms of fluxionality of $[Ir_4(CO)_{11}(PEt_3)]$; a reappraisal of the mechanism of carbonyl fluxionality on clusters, 889–94

Platinum metal complexes of potentially chelating alkene–thioether and alkene selenoether ligands: synthesis and dynamic nuclear magnetic resonance study of $[MX_2(MeE(CH_2)_nCH=CH_2)]$ ($M = Pt$ or Pd ; $X = Cl$, Br , or I ; $E = S$ or Se ; $n = 2$ or 3) and the X-ray structure of *cis*-dibromo(2-thia-6-heptene)platinum(II), $[PtBr_2(MeS(CH_2)_3CH=CH_2)]$, 985–90

Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten $[M\{\eta^3-SC(CF_3)=C(CF_3)H\}(CF_3C\equiv CCF_3)(\eta^5-C_5H_5)]$ ($M = Mo$ or W). Their relevance to the mechanism of metal promoted isomerisation of carbon–carbon double bonds, 1027–36

Preparation and characterisation of some nickel(II) tetra-aza macrocyclic complexes bearing pendant polymerisable groups as part of the ligand superstructure, 1599–608

The mechanisms of fluxionality of $[Fe_3(CO)_{12-n}\{P(OR)_3\}_n]$ ($R = Me$, $n = 0$ –3; $R = Pr$, $n = 2$ or 3) and the X-ray structure of $[Fe_3(CO)_{10}\{P(OMe)_3\}_2]$, 1831–44

The heteronuclear cluster chemistry of the Group 1B metals. Part 12.

Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds $[M_2Ru_4(\mu_3-H)_2(\mu-L)(CO)_{12}]$ ($M = Cu$ or Ag , $L = Ph_2P(CH_2)_nPPh_2$, $n = 1$ –6) or *cis*- $Ph_2PCH=CHPPh_2$; $M = Cu$, $L = Ph_2As(CH_2)_nPPh_2$ ($n = 1$ or 2) in solution, 2039–48

Hydrogen scrambling processes in $[Ta(\eta-C_5Me_5)(\eta-C_5H_5)\{\mu-H_2BH_2\}]$: surprising observation, 2133–8

Platinum metal complexes of potentially chelating alkene thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of $[MX_2(Et(CH_2)_nCR=CR_2)_2]$ ($M = Pt$ or Pd ; $X = Cl$, Br , or I ; $E = S$ or Se ; $n = 2$ or 3; $R = H$ or Me) and the X-ray crystal structure of *cis*-di-iodo(5-thianona-1,8-diene)platinum(II), $[PtI_2\{S(CH_2)_2CH=CH_2\}_2]$, 2315–22

FORMATE

Crystal and molecular structures of μ -hydrido-bis[*fac*-(2,2'-bipyridine)tricarbonylrhenium(I)] chloride, *fac*-(2,2'-bipyridine)tricarbonylformatorhenium(I), and *fac*-(2,2'-bipyridine)tricarbonyl(cyanotrihydroborato)rhenium(I), 1449–54

GALACTURONIC ACID

Proton electron nuclear double resonance study of oxovanadium(IV) complexes of D-galacturonic and polygalacturonic acids, 1283–8

GALLIUM

The hydrolysis of metal ions. Part 11. The ionic strength dependence of gallium(III), 399–402

GEL

Aqueous chemistry and thermodynamic modelling of $CaO-SiO_2-H_2O$ gels, 2371–80

GLYCOSIDE

Isotopic multiplets in the carbon-13 nuclear magnetic resonance spectra due to partial deuteration of co-ordinated NH_2 and NH groups of cobalt(III) complexes containing an N-glycoside. Importance of sugar ring oxygen atoms in the long-range isotope shifts, 1241–8

GOLD

(Diphenylphosphinomethyl)diphenylphosphine sulphide ($(Ph_2PCH_2)PPh_2S$) and its methanide anion $Ph_2PCHPPh_2S^-$ as ligands in organogold chemistry. X-Ray crystal structure of *cis*- $[\Au(C_6F_5)_2(PPh_2CHPPh_2S)]$, 73–8

Polyhedral rearrangements in undeca- and dodeca-nuclear Group 1B metal osmium clusters, 505–10

Gold-197 Mössbauer spectra and the bonding of some gold–gold and gold–platinum clusters, 539–44

Studies on gold(II) complexes with hard and soft donor ligands.

Part 1. Complexes with *o*-aminobenzenethiol (1988, 2607), 555

Synthesis of diphenylphosphido-bridged manganese–gold complexes. Crystal structures of $[N(PPh_3)_2]Au_2Mn(\mu-PPh_2)_2(CO)_4(C_6F_5)_2$ and $[Au_2Mn_2(\mu-PPh_2)_4(\tilde{CO})_8]$, 639–44

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 86.

Alkyldyne(carborane)molybdenum–gold–rhodium and –iron complexes; crystal structure of $[NEt_4]MoFe_2(\mu_3-CC_6H_4Me-4(\mu-\sigma:\sigma:\eta^5-C_2B_9H_7Me_2)CO_8]$, 797–808

Gold(I) complexes derived from secondary phosphines: $[\{Au(\mu-PR_2)_n\}, [AuBr_2(\mu-PPh_2)]^-, [AuX(PHR_2)]$, and $[\{Au(PHR_2)_n\}]^+$. Crystal structure of $[AuBr_2(PPh_2)]$, 907–14

Monohydrido-bridged platinum(II)–gold(I) complexes. X-Ray crystal structure of $[(Ph_3P)_2(C_6Cl_5)Pt(\mu-H)Au(PPh_3)-ClO_4 \cdot 2Et_2O]$, 1089–92

The heteronuclear cluster chemistry of the Group 1B metals. Part 11. Effect of the nature of the bidentate diphosphine ligand on the metal framework structures of the gold heteronuclear cluster compounds $[Au_2Ru_4(\mu_3-H)(\mu-H)\{\mu-Ph_2P(CH_2)_nPPh_2\}(CO)_{12}]$ ($n = 1$ –6). X-Ray crystal structures of $[Au_2Ru_4(\mu_3-H)(\mu-H)\{\mu-Ph_2P(CH_2)_nPPh_2\}(CO)_{12}]$ ($n = 1$ or 2), 1227–36

Reaction of $[Rh_2(\mu-1,8-(NH)_2C_{10}H_6)_2(CO)_2(PPh_3)_2]$ with $[\{M(PPh_3)_4\}^+]$ ($M = Au$, Ag , or Cu): formation of unusual 44-electron Rh_2M clusters. Crystal structure of $[Rh_2(\mu-Au(PPh_3))_2\{\mu-1,8-(NH)_2C_{10}H_6\}(CO)_2(PPh_3)_2]ClO_4 \cdot 2CH_2Cl_2$, 1249–52

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 87. Dicosahedral carbaborane(alkylidyne)–tungsten complexes as reagents for the synthesis of compounds with heteronuclear metal–metal bonds: crystal structures of $[NEt_4][W(\equiv CC_6H_3Me_2-2,6)(CO)_2(\eta^6-C_2B_{10}H_{10}Me_2)]$ and $[NEt_4][WFe(\mu-CC_6H_3Me_2-2,6)(CO)_4(\eta^6-C_2B_{10}H_{10}Me_2)]$, 1363–74

Synthesis, structure, and reactivity of the heteronuclear clusters $[Os_4H(CO)_{15}\{M(PR_3)\}]$ and $[Os_5(CO)_{15}\{M(PR_3)\}_2]$ ($M = Cu$, Ag , or Au ; $R = Ph$ or Me); the X-ray crystal structure of $[Os_4H(CO)_{15}\{Au(PPh_3)\}]$, 1421–6

Bi-, tetra-, and poly-nuclear complexes of gold(I) and silver(I) with bis(diphenylphosphino)-methanide or -amide acting as six-electron donors, 1883–6

Mesitylgold(I) complexes. Crystal structures of $NEt_4[Au(mes)Cl]$ and $[N(PPh_3)_2](mes)Au(CN)Au(mes)]$, 2127–32

Reactions of chloro(triphenylphosphine)gold(I) and $[\mu-1,2-bis(diphenylphosphino)ethane]-bis[bromogold(I)]$ with oxopurine bases. Molecular and crystal structure and bonding of (theophyllinato)(triphenylphosphine)gold(I), 2323–30

Mixed-metal carbido carbonyl clusters. Part 4. Synthesis and structural characterization of $[Rh_6C(CO)_{13}\{Au_2(PPh_3)_2\}]$, 2343–6

HAEMOGLOBIN

The binding of trialkyltin(IV) moieties to rat haemoglobin, and the structure of model systems, studied by tin-119 Mössbauer spectroscopy, 519–26

HAFNIUM

Preparation of thionylimide complexes of titanium, zirconium, and hafnium. Crystal structure of $[Zr(cp)(\eta-C_5Me_5)(NSO)_2]$, 1815–8

HALIDE

- Halogeno, hydride, and alkyl carbonyl-monocyclopentadienyl niobium derivatives. Crystal and molecular structure of $[\text{Nb}(\eta\text{-C}_5\text{H}_5)\text{Cl}(\text{H})(\text{CO})(\text{PMe}_3)_2]$, 79–84
 Spectroscopic studies in matrix isolated tungsten chlorides and bromides, 313–16
 A kinetic and mechanistic study of the solvolysis of aryltellurium trihalides in organic solvents, 457–60
 Structural and motional study of pyridinium tetrahalogenoantimonates(IV) by means of nuclear quadrupole resonance spectroscopy, 631–4
 Lewis base adducts of Group 11 metal(I) complexes. Part 48. Heterobimetallic tris(dithiocarbamato)cobalt(II)–copper(I) halide adducts as a vehicle for novel copper(I) halide clusters (structurally characterized), 971–6
 Formation of mixed-ligand complexes of cadmium(II) halide with tri-n-octylphosphine oxide and pyridine or substituted pyridines in 1,2-dichloroethane, 1259–64
 Investigations of tin(II) halide aqueous solutions and solvent extracts by ^{119}Sn nuclear magnetic resonance spectroscopy: characterisation of the trihalogenostannate(II) anions $[\text{SnX}_3]^-$, $[\text{SnX}_2\text{Y}]^-$, and $[\text{SnX(Y)}Z]^-$ (X, Y , or $\text{Z} = \text{Cl}, \text{Br}$, or I), 2223–8
 Platinum metal complexes of potentially chelating alkene thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{E}[(\text{CH}_2)_n\text{CR=CR}_2]\}_2]$ ($\text{M} = \text{Pt}$ or Pd ; $\text{X} = \text{Cl}, \text{Br}$, or I; $\text{E} = \text{S}$ or Se ; $n = 2$ or 3; $\text{R} = \text{H}$ or Me) and the X-ray crystal structure of *cis*-di-iodo(5-thianona-1,8-diene)platinum(II), $[\text{PtI}_2\{\text{S}[(\text{CH}_2)_2\text{CH=CH}_2]\}_2]$, 2315–22

HETEROCYCLE

- Contributions to the chemistry of boron. Part 194. Synthesis and characterization of mono-, bi-, and tri-cyclic boron–nitrogen–sulphur ring systems. Crystal structure of 2,7,9,14-tetrakis(diethylamino)-3'5,6,10,12,13-hexathia-1,8-diazab-2,4,7,9,11,14-hexaboratricyclo[9.3.0.0^{4,8}]tetradecane, 1311–20

HETEROPOLYANION

- Reactions of heteropolyanions in non-polar solvents. Part 3. Activation of dioxygen by manganese(II) centres in polytungstates. Oxidation of hindered phenols, 1483–90
 The structure of 18-molybdodisulphate(VI) ($4-$) ion in $(\text{NEt}_4)_2\text{S}_2\text{Mo}_{18}\text{O}_{62}\cdot\text{CH}_3\text{CN}$, 1491–8
 Monomolybdonavanadate and *cis*- and *trans*-dimolybdo-octavanadate, 1915–24

HEXA DIENEDIYL

- Chemistry of multi-site bound unsaturated ligands on metal carbonyl clusters. Intramolecular reductive coupling in the conversion of $[\text{Ru}_3(\mu\text{-H})(\text{CO})_9(\mu_3,\eta^2\text{-C}\equiv\text{CBu}')]_2$ to the butadienyl cluster $[\text{Ru}_3(\text{CO})_8\{\mu_3,\eta^2\text{-C}(\text{Bu}')\text{CC}(\text{H})\text{C}(\text{Ph})(\text{PPh}_3)\}]$, and its crystal structure. An example of ligand isomerism on metal clusters, 601–6

HYDRAZIDE

- Mechanistic studies on hydrazido($2-$)-complexes. Cleavage of the nitrogen–nitrogen bond in the reaction of $[\text{Mo}\{\text{NN}(\text{CH}_2)_4\text{CH}_2\}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$ with acid, 425–30
 Hydrazido-complexes of titanium(IV) with tris(pyrazolyl)borates. Crystal structures of $[\text{TiLCl}_2(\text{NMeNM}_2)]$ and $[\text{TiLCl}_2(\text{NPhNH}_2)]$, L = tris(pyrazolyl)borate, 1413–16

HYDRAZINE

- Nucleophilic (Michaeli) additions to $(\text{Ph}_2\text{P})_2\text{C=CH}_2$ when chelated to platinum(II) dichloride or di-iodide, 1527–30

HYDRAZONATE

- Synthesis, infrared, and tin-119 Mössbauer spectroscopic characterization of seven-co-ordinate diorganotin(IV) adducts with 2,6-diacylpyridine acetylhydrazone, including the X-ray crystal structure of $\text{SnEt}_2\text{(dapt)}$ [$\text{H}_2\text{dapt} = 2,6\text{-diacetylpyridine bis}(2\text{-thenoylhydrazone})$], 289–94

HYDRAZONE

- Crystal structure, spectroscopic and magnetic properties of the complex $[\text{Cu}(\text{paphy})(\text{NCS})(\text{SCN})]$ (paphy = pyridine-2-carbaldehyde 2-pyridylhydrazone). An unusual di- μ -thiocyanato-*N* bridged copper(II) dimer, 53–6
 Synthesis, characterization, and crystal structures of bismuth complexes with 2,6-diacylpyridine bis(2-thenoylhydrazone) (H_2dapt): $[\text{Bi}(\text{Hdapt})\text{Cl}_2]\text{-dmsO-H}_2\text{O}$ and $[\text{Bi}(\text{dapt})\text{Cl}]\text{-dmsO}$, containing a quinquedentate ligand, 671–6
 New metal-sulphur–nitrogen compounds from reactions in liquid ammonia. The X-ray structures of *trans*-bis(acetophenone dimethylhydrazone- N^2)dichloropalladium(II) and [di(azathien)-1-yl- $S^1\text{N}^4$][2-(hydrazoneoethyl)phenyl]palladium(II), 1179–86

- Crystal structures and magnetic properties of bi- and tetra-nuclear copper(II) complexes of 2,6-diformyl-4-methylphenol di(benzoylhydrazone), 2083–8

HYDRIDE

- Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene (C=CHOEt) and ethoxyethyne (CH=COEt) ligands respectively, 5–12
 Halogeno, hydride, and alkyl carbonyl-monocyclopentadienyl niobium derivatives. Crystal and molecular structure of $[\text{Nb}(\eta\text{-C}_5\text{H}_5)\text{Cl}(\text{H})(\text{CO})(\text{PMe}_3)_2]$, 79–84
 Reactions of hydride complexes of ruthenium(II) with alkynes, 589–94
 Chemistry of multi-site bound unsaturated ligands on metal carbonyl clusters. Intramolecular reductive coupling in the conversion of $[\text{Ru}_3(\mu\text{-H})(\text{CO})_9(\mu_3,\eta^2\text{-C}\equiv\text{CBu}')]_2$ to the butadienyl cluster $[\text{Ru}_3(\text{CO})_8\{\mu_3,\eta^2\text{-C}(\text{Bu}')\text{CC}(\text{H})\text{C}(\text{Ph})(\text{PPh}_3)\}]$, and its crystal structure. An example of ligand isomerism on metal clusters, 601–6

- Iridium(III) hydrido complexes with 2-pyridinaldazines. Crystal structure of $[\text{IrH}_2(\text{mpaa})(\text{PPh}_3)_2]\text{SbF}_6$. Synthesis and characterization of homo- and hetero-nuclear complexes, 1075–82

- Monohydrido-bridged platinum(II)–gold(I) complexes. X-Ray crystal structure of $[(\text{Ph}_3\text{P})_2(\text{C}_6\text{Cl}_5\text{Pt}(\mu\text{-H})\text{Au}(\text{PPh}_3))\text{ClO}_4\cdot 2\text{Et}_2\text{O}$, 1089–92

- Synthesis and characterization of a family of nitrile-substituted triangular hydridocarbonyl rhodium clusters, $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{12-n}(\text{NCMe})_n]$ ($n = 1$ –3). X-Ray crystal structures of $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_{11}(\text{NCMe})]$ and $[\text{Re}_3(\mu\text{-H})_3(\text{CO})_9(\text{NCMe})_3]$, 1143–8

- Synthesis of hydridotrifluoromethyl complexes of platinum(II). A spectroscopic investigation of the *trans* and *cis* influence of ligands in hydridoplatinum(II) compounds, 1149–60

- The heteronuclear cluster chemistry of the Group 1B metals. Part 11. Effect of the nature of the bidentate diphosphine ligand on the metal framework structures of the gold heteronuclear cluster compounds $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})(\mu\text{-H})(\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2)(\text{CO})_{12}]$ ($n = 1$ –6). X-Ray crystal structures of $[\text{Au}_2\text{Ru}_4(\mu_3\text{-H})(\mu\text{-H})(\mu\text{-Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2)(\text{CO})_{12}]$ ($n = 1$ or 2), 1227–36

- Reactions of co-ordinated ligands. Part 46. Formation of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH=CMe}_2)(\text{CO})_{10}]$ by carbon–carbon σ -bond and carbon–hydrogen activation of 3,3-dimethylcyclopropene; crystal and molecular structure of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and evidence for its facile rearrangement to $[\text{Os}_3(\mu\text{-H})(\mu\text{-CH=CHCHMe}_2)(\text{CO})_{10}]$, 1333–40

- Synthesis, structure, and reactivity of the heteronuclear clusters $[\text{Os}_5\text{H}(\text{CO})_{15}\text{M}(\text{PR}_3)]$ and $[\text{Os}_5(\text{CO})_{15}\text{M}(\text{PR}_3)_2]$ ($\text{M} = \text{Cu}, \text{Ag}$, or Au ; R = Ph or Me); the X-ray crystal structure of $[\text{Os}_5\text{H}(\text{CO})_{15}\{\text{Au}(\text{PPh}_3)\}]$, 1421–6

- Crystal and molecular structures of μ -hydrido-bis[*fac*-(2,2'-bipyridine)tricarbonylrhodium(I)] chloride, *fac*-(2,2'-bipyridine)tricarbonylformatorhenium(I), and *fac*-(2,2'-bipyridine)tricarbonyl(cyanotrihydroborato)rhodium(I), 1449–54

- Synthesis of the cluster dianion $[\text{Os}_{11}\text{C}(\text{CO})_{27}]^{2-}$ by pyrolysis and its reactions with electrophiles; X-ray structure analysis of the mixed-metal derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}(\text{Cu}(\text{NCMe}))]\text{CH}_2\text{Cl}_2$ and the hydrido derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455–64

- Synthesis and crystal structure of $[\text{NiOs}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\text{Cu}(\text{PPh}_3))\text{CO}_6]$. Hydrogenation–dehydrogenation reactions of dienes and alcohols with heterogeneous catalysts derived from $[\text{NiM}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-X})(\text{CO})_6]$ [$\text{M} = \text{Ru}, \text{X} = \text{H}; \text{M} = \text{Os}, \text{X} = \text{Cu}(\text{PPh}_3)$], 1477–82

- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 88. Carbaboranetungsteniridium compounds; crystal structure of the complex $[\text{WIr}(\mu\text{-CC}_6\text{H}_4\text{Me}_4)(\text{CO})_2(\text{PEt}_3)_2(\eta^5\text{-C}_2\text{B}_9\text{H}_9\text{Me}_2)]$, 1845–54

- Cluster activation *via* cyclometallation. Formation of the cyclometallated triosmium carbonyl cluster $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ and its reactions with phosphorus donor ligands. The X-ray crystal structures of $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ and $[\text{Os}_3\text{H}(\text{CO})_9\{\text{P}(\text{OMe})_3\}\text{PPh}_2]$, 2023–32

- Reactions of the cyclometallated triosmium cluster $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ with Brønsted acids and dihydrogen: synthesis and X-ray crystal structure of $[\text{Os}_3\text{H}_2(\text{CF}_3\text{CO}_2)(\text{CO})_9(\mu\text{-PPh}_3)]$, 2033–8

- The heteronuclear cluster chemistry of the Group 1B metals. Part

HYDRIDE (contd)

12. Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds $[M_2Ru_4(\mu_3\text{-H})_2(\mu\text{-L})(\text{CO})_{12}]$ [$M = \text{Cu}$ or Ag , $\text{L} = \text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2$ ($n = 1\text{--}6$) or *cis*- $\text{Ph}_2\text{PCH}=\text{CHPPh}_2$; $M = \text{Cu}$, $\text{L} = \text{Ph}_2\text{As}(\text{CH}_2)_n\text{PPh}_2$ ($n = 1$ or 2)] in solution, 2039–48
 Activation of a vinyl carbon–hydrogen bond in a tris(pyrazolyl)boratoiridium complex. The *X*-ray crystal structure of $[\text{IrH}(\text{HB}(\text{pz})_3)_3(\sigma^2\text{-C}_8\text{H}_{13})(\eta^2\text{-C}_8\text{H}_{14})]$, 2073–6
 Reactivity of new osmium dihydrides with arenediazonium cations: preparation and properties of bis(aryldiazene) and mono(aryldiazenido) complexes. A comparison with analogous iron and ruthenium derivatives, 2353–8
 Photochemical reactivity of $2[\text{Ir}^{III}\text{HCl}_2(\text{PPr}')_2]\text{H}_2$. Formation of monohydrido isomers of Ir^{III} and evidence of photoinduced oxidative addition of H_2 to Ir^{III} , 2367–70
 Preparation and *X*-ray crystal structure of $[\text{Ru}_3\text{RhH}_3(\text{CO})_{12}]$, 2449–52

HYDRIDO

- Carbonyl, hydridocarbonyl, and hydridoalkoxycarbonyl complexes of iridium with phosphorus–sulphur hybrid ligands, and the crystal and molecular structure of $[\text{IrH}(\text{CO}_2\text{Me})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{SEt}_2)\text{BPh}_4]$, 607–14

HYDROFORMYLATION

- Cationic dinuclear rhodium complexes as catalyst precursors for the hydroformylation of alkenes, 1579–82

HYDROGEN-2

- Relation between the torsion angles of acetate chelate rings and the deuteron nuclear magnetic resonance chemical shifts for polyamine-N-carboxylatochromate(III) complexes, 1947–50

HYDROGENATION

- Cationic dinuclear rhodium complexes as catalyst precursors for the hydroformylation of alkenes, 1579–82

HYDROGEN BONDING

- Structure of μ -fluoro-bis[bis(2,2'-bipyridyl)fluoroniobium(IV)] fluoride–ethanol (1/2) trihydrate, revealing several unique features including strong hydrogen bonds between fluoride ions and ethanol molecules, 1273–6

HYDROGEN PEROXIDE

- Kinetics of the uncatalysed and silver(I)-catalysed oxidation of hydrogen peroxide by [ethylenebis(biguanide)]silver(III) ion in acid perchlorate media, 1645–50

HYDROLYSIS

- The copper(II) promoted hydrolysis of benzylpenicillin. Evidence for the participation of a Cu–OH species in the hydrolysis of the β -lactam ring, 197–202

- Thermodynamics of hydroxy complex formation of dialkyltin(IV) ions in aqueous solution, 773–8

- Kinetics of aquation and base hydrolysis of some *trans*- $[\text{Co}(\text{tn})_2(\text{B})\text{Cl}]^{n+}$ ($\text{tn} = 1,3\text{-diaminopropane}$; $\text{B} = \text{OH}$, NCS , N_3 , NO_2 , NH_3 , or CN) complexes. The effect of ring size on reactivity, 853–8

- Reactivity studies on the hydrolysis of iron(II) complexes of substituted phenanthrolines in dioxane–water solvents, 1105–8

- Multinuclear magnetic resonance studies of the hydrolysis of aluminium(III). Part 9. Prolonged hydrolysis with aluminium metal monitored at very high magnetic field, 1889–96

- Multinuclear magnetic resonance studies of the hydrolysis of aluminium(III). Part 10. Proton, carbon-13, and aluminium-27 spectra of aluminium acetate at very high magnetic field, 1897–902

- Base-promoted cleavage of a P–C bond in $[\text{PtCl}_2(\text{Ph}_2\text{PCH}_2\text{PPh}_2)_2\text{PF}_6]$ under mild conditions, 2017–22

HYDROXAMIC ACID

- Complex-forming properties of L- α -alaninehydroxamic acid (2-amino-N-hydroxypropanamide), 2247–52

HYDROXIDE

- Crystal structure and magnetic properties of 1-aqua- μ -hydroxo-1,2,2-tris(perchlorato)-1,2-bis(2,2';6',2"-terpyridine)dicopper(II), 237–42

- The reaction of hydroxo-species of palladium(II) with bidentate nucleophiles, 375–8

- Preparation and interconversion of dimeric di- μ -hydroxo and tri- μ -hydroxo complexes of platinum(II) and palladium(II) with 2,2'-bipyridine and 1,10-phenanthroline, 403–12

- Thermodynamics of hydroxy complex formation of dialkyltin(IV) ions in aqueous solution, 773–8

- Crystal and molecular structures of

- diaqua(nitrilotriacetato)aluminium(III) and di- μ -hydroxo-bis(nitrilotriacetato)dialuminate(III) dianion, 1513–8

- Phenyltin trichloroacetate complexes: preparations and structures, 1589–98

- Preparation and structural studies of diphenylthiophosphinito rhodium(III) complexes. X-Ray structure analysis of $[(\text{C}_5\text{Me}_5)_2\text{Rh}](\mu\text{-Cl})_2(\mu\text{-Ph}_2\text{PS})][\text{BPh}_4]$, 1987–92

- The chemistry of dinitrogen residues. Part 7. Hydrazido(1–)- and *N,N*-dimethylhydroxylaminato(1–)-complexes of titanium(IV), 2389–96

HYDROXYLAMINATE

- The chemistry of dinitrogen residues. Part 7. Hydrazido(1–)- and *N,N*-dimethylhydroxylaminato(1–)-complexes of titanium(IV), 2389–96

HYPONITROUS ACID

- Stoichiometric and nitrogen-15 labelling studies on the hyponitrous acid–nitrous acid reaction, 527–32

- The kinetics and mechanism of the hyponitrous acid–nitrous acid reaction at 0 °C, 533–8

HYROXIDE

- The copper(II) promoted hydrolysis of benzylpenicillin. Evidence for the participation of a Cu–OH species in the hydrolysis of the β -lactam ring, 197–202

IMIDAZOLE

- Synthesis and characterization of imidazole-bridged iron(III)–copper(II) and manganese(II)–copper(II) binuclear complexes, 943–6

- Metal complexes as ligands. Part 12. Synthesis and *X*-ray crystal structure of a hexanuclear nickel(II) complex based on two isosceles trinuclear components with imidazole bridges, 1187–92

IMIDAZOLINE

- The interaction of 1-methylimidazoline-2(3*H*)-thione with copper(II) salts, 39–42

- Crystal structure and mass spectrometry of dibromodimethyl[1-methyl-2(3*H*)-imidazolinethione-*S*]tin(IV), 927–30

IMIDE

- Gas-phase molecular structure of *NN'*-bis(trimethylsilyl)sulphur di-imide, determined by electron diffraction, 859–62

IMINE

- Reactivity of the iridium nitrosyl complex

- $[\text{Ir}(\text{NO})(\text{paa})(\text{PPh}_3)_2][\text{PF}_6]_2$ ($\text{paa} = 2\text{-pyridinecarbaldehyde azine}$) with nucleophiles: insertion of NO into a methinic C–H bond; crystal structure of $[\text{Ir}((2\text{-C}_5\text{H}_4\text{N})\text{CH}=\text{N}=\text{N}=\text{C}(\text{NOH})(2\text{-C}_5\text{H}_4\text{N})\{\text{Me}(\text{EtO})\text{C}=\text{NH}\}(\text{PPh}_3)_2][\text{PF}_6]_2$, 1217–26

- Spectroscopic and electrochemical study of copper and zinc complexes with a N_2S_2 ligand donor set. Crystal structure of the copper(II) complex derived from 1,3-propylenediamine and 3-formyl-1-phenyl-2(1*H*)-pyridinethione, 1979–86

INDENYL

- Electronic structure of transition-metal indenyl compounds: a He I and He II photoelectron spectroscopic study of $[\text{Mn}(\eta^5\text{-C}_9\text{H}_7)(\text{CO})_3]$, $[\text{Fe}(\eta^5\text{-C}_9\text{H}_7)_2]$, $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)_2]$, and $[\text{Ru}(\eta^5\text{-C}_9\text{H}_7)(\eta\text{-C}_5\text{Me}_5)_2]$, 2139–48

- Substituted cyclopentadienyl complexes. Part 4. The catalysed synthesis and proton nuclear magnetic resonance spectra of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_4\text{Me})(\text{CO})(\text{L})]$ and $[\text{Ru}(\eta^5\text{-C}_5\text{H}_7)(\text{CO})(\text{L})]$ ($\text{L} = \text{phosphine or isocyanide}$) and the crystal structure determination of $[\text{Ru}(\eta^5\text{-C}_5\text{H}_7)(\text{CO})\{\text{P}(\text{CH}_2\text{Ph})_3\}\text{I}]$, 2407–14

INDIUM

- Synthesis and characterization of benzylindium compounds, 1625–6
 Co-ordination compounds of indium. Part 47. Indium(I) and thallium(I) derivatives of alkanedithiols, 1759–64

INOSINE

- X-Ray structure of a ternary complex, copper(II)–inosine 5'-monophosphate–benzimidazole. The first example for the absence of direct metal–nucleotide interaction, 1507–12

ION EXCHANGE

- The equilibrium of strontium ion exchange of sodium titanate, $\text{Na}_4\text{Ti}_9\text{O}_{20}\cdot x\text{H}_2\text{O}$, 101–4

- The nature of a sodium–crown interaction studied by sodium-23 nuclear magnetic resonance and by crystal structure determination of a complex with co-ordinated solvent, 203–10

- Lamellar inorganic ion exchangers. Hydrogen–lithium ion exchange in γ -titanium phosphate, 829–36

- Formation of chromium interlayers in α -zirconium phosphate, 895–900

- Lamellar inorganic ion exchangers. H^+/Cs^+ ion exchange in γ -titanium phosphate, 1825–30

IONIC STRENGTH

The hydrolysis of metal ions. Part 11. The ionic strength dependence of gallium(III), 399–402

ION PAIRING

Vibrational study of ionic association in aprotic solvents. Part 12. Isothiocyanate M(NCS) complexes of non-transition-metal ions, 2193–8

I.R.

Synthesis, infrared, and tin-119 Mössbauer spectroscopic characterization of seven-co-ordinate diorganotin(IV) adducts with 2,6-diacylpyridine acylhydrazone, including the X-ray crystal structure of $\text{SnEt}_2(\text{dapt})$ [$\text{H}_2\text{dapt} = 2,6\text{-diacylpyridine bis}(2\text{-thenoylhydrazone})$], 289–94

Spectroscopic studies in matrix isolated tungsten chlorides and bromides, 313–6

Lewis-base adducts of Group 11 metal(I) compounds. Part 58.

Syntheses, X-ray structures, and far-infrared spectra of the novel $[\text{CuX}_2(\text{bidentate ligand})]^{+}$ ($\text{X} = \text{Cl}$ or Br) anion salts of the $[\text{Cu}(\text{dpa})_2]^{+}$ cation and the dimeric iodide $[(\text{dpa})\text{CuI}_2\text{Cu}(\text{dpa})]$ [$\text{dpa} = \text{di}(2\text{-pyridyl)amine}$], 1639–44

Vibrational study of ionic association in aprotic solvents. Part 12. Isothiocyanate M(NCS) complexes of non-transition-metal ions, 2193–8

Bonding in clusters. Part 10. The nature of the BH groups in *closoborane* anions. A study by vibrational spectroscopy, electronic spectroscopy for chemical analysis, and *ab initio* computation, 2253–60

IRIDIUM

Carbonyl, hydrido carbonyl, and hydridoalkoxycarbonyl complexes of iridium with phosphorus-sulphur hybrid ligands, and the crystal and molecular structure of

$[\text{IrH}(\text{CO}_2\text{Me})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{SEt})_2]\text{BPh}_4$, 607–14

The mechanisms of fluxionality of $[\text{Ir}_4(\text{CO})_{11}(\text{PEt}_3)]$: a reappraisal of the mechanism of carbonyl fluxionality on clusters, 889–94

Iridium(III) hydrido complexes with 2-pyridinaldazines. Crystal structure of $[\text{IrH}_2(\text{mpaa})(\text{PPH}_3)_2]\text{SbF}_6$. Synthesis and characterization of homo- and hetero-nuclear complexes, 1075–82

Synthesis and characterization of pentanuclear Fe–Ir carbonyl clusters. Solid-state structures of $[\text{N}(\text{PPh}_3)_2]_2[\text{FeIr}_4(\text{CO})_{15}]$ and $[\text{PPh}_4][\text{Fe}_2\text{Ir}_3(\text{CO})_{14}]$, 1109–16

Trinuclear pyrazolate complexes. Crystal structure of $[(\text{OC})_2\text{Rh}(\mu\text{-pz})_2\text{Pd}(\mu\text{-pz})_2\text{Rh}(\text{CO})_2]$, 1133–8

Synthesis of ammonium ions and nitrosylation reactions using nitrosyl chloride and alkyl nitrites, 1139–42

Reactivity of the iridium nitrosyl complex

$[\text{Ir}(\text{NO})(\text{paa})(\text{PPh}_3)_2]\text{[PF}_6\text{]}_2$ (paa = 2-pyridinecarbaldehyde azine) with nucleophiles: insertion of NO into a methinic C–H bond; crystal structure of $[\text{Ir}\{(\text{2-C}_5\text{H}_4\text{N})\text{CH}=\text{N}-\text{N}=\text{C}(\text{NOH})-(2\text{-C}_5\text{H}_4\text{N})\}\{\text{Me}(\text{EtO})\text{C}=\text{NH}\}(\text{PPh}_3)_2]\text{[PF}_6\text{]}_2$, 1217–26

Photo-oxidation of alcohols by hexachlorometalate(IV) ions ($M = \text{Pt, Pd, or Ir}$): spin trapping and matrix isolation electron spin resonances studies, 1237–40

Polyhedral organoiridaborane chemistry. The reaction of MeNC with $[6\text{-}(\eta^5\text{-C}_5\text{Me}_5)\text{-}nido\text{-}6\text{-IrB}_9\text{H}_{13}]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[4,4\text{-}(\eta^5\text{-C}_5\text{Me}_5)(\text{MeNC})\text{-arachno-4-IrB}_9\text{H}_{12}]$, 1465–70

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 88. Carbaboranetungsteniridium compounds; crystal structure of the complex $[\text{WIr}(\mu\text{-CC}_6\text{H}_4\text{Me}_4)(\text{CO})_2(\text{PEt}_3)_2(\eta^5\text{-C}_5\text{B}_9\text{H}_9\text{Me}_2)]$, 1845–54

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 90. Synthesis of the complexes $[\text{W}_2\text{M}_2(\mu\text{-CO})(\mu\text{-CMe})(\mu_3\text{-CMe})(\mu\text{-PPH}_3)_2(\text{CO})_3(\mu\text{-C}_5\text{H}_5)_2]$ ($M = \text{Rh or Ir}$) and related tetra- and penta-nuclear metal compounds; interconversion of ketenyl and λ^5 -phospho-alkyne ligands bridging tungsten–rhodium bonds, 1865–70

Activation of a vinyl carbon–hydrogen bond in a tris(pyrazolyl)boratoiridium complex. The X-ray crystal structure of $[\text{IrH}(\text{HB(pz)}_3)(\sigma\text{-C}_8\text{H}_{13})(\eta^2\text{-C}_8\text{H}_{14})]$, 2073–6

Photochemical reactivity of $2[\text{Ir}^{\text{III}}\text{HCl}_3(\text{PPR}'_3)_2]\text{H}_2$. Formation of monohydrido isomers of Ir^{III} and evidence of photoinduced oxidative addition of H_2 to Ir^{III} , 2367–70

Cyclometallation of unsaturated carboxylic acids by pentamethylcyclopentadienyl-rhodium and -iridium complexes.

Crystal structures of $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh}=\text{CHCO}-\text{O})(\text{Me}_2\text{SO})]$ and $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh}=\text{CHCO}-\text{OMe})\text{I}]^+$, 2415–22

IRON

Metal complexes (Co^{3+} , Fe^{3+} , Mn^{3+}) of the quinquedentate salen analogue, 2-methyl-4-benzylthio-N,N'-butane-1,2-diybis(salicylideneimine): crystal structures and sulphur coordination effects, 179–84

The binding of trialkyltin(IV) moieties to rat haemoglobin, and the structure of model systems, studied by tin-119 Mössbauer spectroscopy, 519–26

Reaction of elemental sulphur with $(5H,14H\text{-}6,8,15,17\text{-tetramethylidibenz}[b,i][1,4,8,11]\text{tetra-azacyclotetradecinato})\text{iron(II)}$: synthesis and X-ray structure of a μ -sulphido iron(II) complex, 551–4

Addition of anilines to the $[\text{Fe}(\text{CO})_3(1\text{-}5\text{-}\eta\text{-C}_7\text{H}_9)]\text{BF}_4^-$ complex, and the 'Ordered Transition State Mechanism', 561–6

Studies on olefin-co-ordinating transition-metal carbene complexes.

Part 14. An unusual reaction of tricarbonyl($\eta^4\text{-cyclopenta-1,3-diene})\text{iron}$: a new approach to acyldicarbonyl($\eta^4\text{-cyclopentadienyl})\text{iron}$ complexes, and the X-ray crystal structure of $[\text{Fe}(\text{COC}_6\text{H}_4\text{CF}_3\text{-}p)(\eta^4\text{-C}_5\text{H}_5)(\text{CO})_2]$, 635–8

Formation in solution, synthesis, and electrochemical study of oxalato complexes of N,N' -ethylenabis(salicylideneiminato)-chromium(III) and -iron(III); crystal structures of piperidinium $[N,N'$ -ethylenabis(salicylideneiminato)](oxalato-O¹O²)-chromate(III) and -ferrate(III), 729–38

Solvent influence on trifluoromethylation reactions with $\text{Cd}(\text{CF}_3)_2$: one-pot synthesis of trifluoromethyl iron complexes, 739–44

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 86. Alkylidyne(carbaborane)-molybdenum–gold, –rhodium and –iron complexes; crystal structure of $[\text{NEt}_4][\text{MoFe}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me-4})(\mu\text{-}\sigma\text{:}\sigma'\text{-}\eta^5\text{-C}_2\text{B}_9\text{H}_7\text{Me}_2)\text{(CO)}_8]$, 797–808

Synthesis and characterization of imidazolate-bridged iron(III)–copper(II) and manganese(III)–copper(II) binuclear complexes, 943–6

Electronic spectral study of the aqua --- hydroxo equilibrium of model iron(III) haems encapsulated in aqueous detergent micelles, 1003–6

Crystal structures of di-iron $\mu\text{-}\eta^2\text{-}(\text{C,C})\text{-ketene}$ complexes, $[\{\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\}_2(\mu\text{-CH}_2\text{CO})]$ and $[\{\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\}_2(\mu\text{-CH}_2\text{CO})]$. Synergistic metal-to-ligand back donation as revealed by comparison with $[\{\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\}_2(\mu\text{-CH}_2\text{CH}_2\text{CO})]$, 1083–8

Reactivity studies on the hydrolysis of iron(II) complexes of substituted phenanthrolines in dioxane–water solvents, 1105–8

Synthesis and characterization of pentanuclear Fe–Ir carbonyl clusters. Solid-state structures of $[\text{N}(\text{PPh}_3)_2]_2[\text{FeIr}_4(\text{CO})_{15}]$ and $[\text{PPh}_4][\text{Fe}_2\text{Ir}_3(\text{CO})_{14}]$, 1109–16

Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(III)–ethanol–carbon tetrachloride system, 1197–202

Solution structure and dynamic behaviour of two isomers of $[\text{Fe}_3(\text{CO})_9\text{P}(\text{OR}_3)_3]$ ($R = \text{Me, Et, or Ph}$) derivatives, 1277–82

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 87. Dodecahedral carbaborane(alkylidyne)tungsten complexes as reagents for the synthesis of compounds with heteronuclear metal–metal bonds: crystal structures of $[\text{NEt}_4][\text{W}(\equiv\text{CC}_6\text{H}_4\text{Me}_2\text{-2,6})(\text{CO})_2(\eta^6\text{-C}_2\text{B}_{10}\text{H}_{10}\text{Me}_2)]$ and $[\text{NEt}_4][\text{WF}_6(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_4(\eta^6\text{-C}_2\text{B}_{10}\text{H}_{10}\text{Me}_2)]$, 1363–74

Substitution reactions of the zerovalent iron complexes $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{PMc}_3)_2]$ and $[\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{dmpe})]$ ($\text{dmpe} = \text{Me}_2\text{PCH}_2\text{CH}_2\text{PMe}_3$), 1385–92

Comparative study on the multielectron reduction of $\text{HOC}_2\text{H}_4\text{N}_3$ catalyzed by $[\text{NBu}_4]^+[\text{MoFe}_3\text{S}_4(\text{SPh})_3(o\text{-O}_2\text{C}_6\text{Cl}_4)]_2$ and $[\text{NBu}_4]^+[\text{Fe}_4\text{S}_4(\text{SPh})_4]$ -modified glassy carbon electrodes, 1547–54

Molecular structure of heterobimetallic $\mu\text{-}\eta^2\text{-}(\text{C,C})\text{-ketene}$ complexes, 1627–30

On the relationship between crystallographic and spectroscopic evidence of dynamic processes in the solid state: the cases of *cis*- and *trans*- $[\text{Fe}_2(\eta^5\text{-C}_5\text{H}_5)_2(\text{CO})_4]$, 1721–6

Structural effects of one-electron oxidation on a cyclobutadieneiron derivative: X-ray crystal structures of $[\text{Fe}(\text{CO})_4\text{P}(\text{OMe})_3]_2(\eta^4\text{-C}_4\text{Ph}_4)$ and $[\text{Fe}(\text{CO})_4\text{P}(\text{OMe})_3]_2(\eta^4\text{-C}_4\text{Ph}_4)$, 1751–8

Neutron diffraction study of the cation distributions in the systems $\text{Fe}_7\text{xM}_x(\text{PO}_4)_6$ ($M = \text{Mn or Co}$), 1765–70

The mechanisms of fluxionality of $[\text{Fe}_3(\text{CO})_{12-n}\text{P}(\text{OR})_3]_n$ ($R = \text{Me, } n = 0\text{--}3; R = \text{Pr}^i, n = 2\text{ or } 3$) and the X-ray structure of $[\text{Fe}_3(\text{CO})_{10}\text{P}(\text{OMe})_3]_2$, 1831–44

IRON (contd)

Synthesis and properties of the bis(μ -thiosulphato-*S*)-bis(dinitrosylferrate)(2-) anion. Crystal and molecular structure of bis[bis(triphenylphosphoranylidene)ammonium] bis(μ -thiosulphato-*S*)-bis(dinitrosylferrate), 2061-4

Transition metal-cyanocarbon chemistry. Part 8. Cyano-substituted buta-1,3-dienylidene bridged di-iron complexes from unprecedented cyanoalkyne insertion into the carbon-hydrogen bond of a bridging alkenylidene ligand, 2187-92

Iron carbene complexes. Part 3. Aminolysis of di-iron alkoxycarbene complexes with ammonia and dimethylamine. X-Ray crystal structure of $[\text{Fe}_2(\mu\text{-SP}_2)(\text{CO})_5(\text{C}(\text{NM}_2)\text{Ph})]$, 2199-2204

The 4-phenyl-1,2,3,5-dithiadiazole complex $[\text{Fe}_2(\text{CO})_6(\text{PhCN}_2\text{S}_2)]$: its preparation, crystal structure, and an extended-Hückel molecular-orbital study of the bonding, 2229-34

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 94. Synthesis of di- and tri-metal compounds via the reagents $[\text{M}(\equiv\text{C}-\text{C}\equiv\text{CBu}')(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ ($\text{M} = \text{Mo}$ or W), 2281-8

ISOCYANATE

Reactivity of metallic copper with acidic organic compounds in the presence of molecular oxygen. Crystal structure of $[\text{Cu}(\text{NCO})(\text{py})_2]_2\cdot\text{H}_2\text{O}$, 921-6

Ruthenium oxo complexes of macrocyclic tertiary amines.

Synthesis, electrochemistry, and reactivities of ruthenium oxo complexes of *meso*-2,3,7,11,12-pentamethyl-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), and X-ray crystal structure of *trans*- $[\text{Ru}^{\text{IV}}\text{L}^1(\text{O})(\text{NCO})]\text{ClO}_4$, 2011-16

ISOCYANIDE

Chemistry of metallacyclobutanones (η^3 -oxodimethylenemethane complexes). Part 5. Ligand substitution reactions of metallacyclobutan-3-ones of platinum(II) and palladium(II) with tertiary phosphines and t-butyl isocyanide; crystal structures of $[\text{Pt}(\text{CH}(\text{CO}_2\text{Me})\text{COCH}(\text{CO}_2\text{Me})\text{C}(\text{CNBu}')\text{(AsPh}_3)]$ and $[\text{Pt}(\text{CH}(\text{CO}_2\text{Me})\text{CO}\bar{\text{C}}(\text{CO}_2\text{Me})\text{C}(\text{NHBu}')\text{(CNBu}'_2)]$, 345-56

Syntheses and redox properties of mixed isocyanide, carbonyl, or nitrile complexes of rhodium(I) *trans*- $[\text{Re}(\text{CNMe})\text{L}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPPh}_2)_2]\text{X}$ [$\text{L} = \text{CNR}$ ($\text{R} = \text{alkyl}$ or aryl), CO , or NCMe ; $\text{X} = \text{Cl}$, BF_4^- , or PF_6^-], 1209-16

Reactions of co-ordinated ligands. Part 45. Ligand displacement and oxidative reactions of the carbyne complex $[\text{Mo}(\equiv\text{CCH}_2\text{Bu}')\text{P}(\text{OMe})_3](\eta\text{-C}_5\text{H}_5)$; crystal and molecular structures of $[\text{Mo}(\eta^3\text{-RN-C}\cdots\text{C}(\text{CH}_2\text{Bu}')\cdots\text{C=NR})(\text{CNR})_2(\eta\text{-C}_5\text{H}_5)]$ ($\text{R} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), $[\text{Mo}(\text{C}=\text{CHBu}')\text{I}\text{P}(\text{OMe})_3](\eta\text{-C}_5\text{H}_5)$, and $[\text{Mo}=\text{C}(\text{CH}_2\text{Bu}')\text{P}(\text{O})(\text{OMe})_2]\text{P}(\text{OMe})_3(\eta\text{-C}_5\text{H}_5)$, 1321-32

Reactivity of binuclear pyrazolate-bridged rhodium complexes with isocyanides. Molecular structure of $[\text{Rh}_2(\mu\text{-pz})\text{I}_2(\text{CNBu}')_2(\mu\text{-Ph}_2\text{PCH}_2\text{PPPh}_2)_2]\text{BF}_4^-$ (pz = pyrazolate), 1427-34

Synthesis and reactivity of binuclear perhalogenophenyl derivatives of platinum(II). Molecular structure of $[\text{Pt}_2(\text{C}_6\text{F}_5)_2(\text{CO})_2(\text{PPPh}_3)_2]\text{0.4CHCl}_3$, 1583-8

Mechanism and energetics of reductive elimination of ethane from some tetramethylplatinum(IV) complexes, 2121-6

Low-valent isocyanide complexes and clusters of palladium and platinum. Crystal structure of $[\text{Pt}(\text{C}=\text{NR})\text{N}(\text{R})\text{C}(\equiv\text{NR})(\text{RNC})_2]$ ($\text{R} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), 2161-6

ISOMERIZATION

Nucleophilic attack of methanol on bis(benzonitrile)dichloroplatinum: formation of mono- and bis-imido ester derivatives, 947-52

Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten $[\text{M}(\eta^3\text{-SC}(\text{CF}_3)=\text{C}(\text{CF}_3)\text{H})(\text{CF}_3\text{C}\equiv\text{CCF}_3)(\eta^3\text{-C}_5\text{H}_5)]$ ($\text{M} = \text{Mo}$ or W). Their relevance to the mechanism of metal-promoted isomerisation of carbon-carbon double bonds, 1027-36

Kinetics of formation, dissociation, and isomerization reactions in complexes of cyanopyridines and benzonitrile with pentacyanoruthenate(II), 1973-8

ISOTHIOCYANATE

Studies in Werner clathrates. Part 9. Structures with bis(isothiocyanato)tetrakis(4-phenylpyridine)nickel(II) as host, 119-26

Stacked planar molecules in antiferromagnetic di-isothiocyanatobis(thiourea)chromium(II), a new structural type for chromium(II), 1631-4

KETENE

Crystal structures of di-iron $\mu\text{-}\eta^2\text{-}(\text{C},\text{C})$ -ketene complexes, $[\{\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\}_2(\mu\text{-CH}_2\text{CO})]$ and $[\{\text{Fe}(\eta^5\text{-C}_5\text{Me}_5)(\text{CO})_2\}_2(\mu\text{-CH}_2\text{CO})]$. Synergistic metal-to-ligand back donation as revealed by comparison with $[\{\text{Fe}(\eta^5\text{-C}_5\text{H}_5)(\text{CO})_2\}_2(\mu\text{-CH}_2\text{CH}_2\text{CO})]$, 1083-8

Molecular structure of heterobimetallic $\mu\text{-}\eta^2\text{-}(\text{C},\text{C})$ -ketene complexes, 1627-30

KETONATE

Reactions of chloro(triphenylphosphine)gold(I) and $[\mu\text{-1,2-bis(diphenylphosphine)ethane}]$ -bis[bromogold(I)] with oxopurine bases. Molecular and crystal structure and bonding of (theophyllinato)(triphenylphosphine)gold(I), 2323-30

KINETICS

Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene ($\text{C}=\text{CHOEt}$) and ethoxyethyne ($\text{CH}=\text{COEt}$) ligands respectively, 5-12

Kinetics of the oxidation of chloride ions by tris(2,2'-bipyridine)nickel(III) ions in aqueous perchlorate media, 151-4

The copper(II) promoted hydrolysis of benzylpenicillin. Evidence for the participation of a Cu-OH species in the hydrolysis of the β -lactam ring, 197-202

Kinetics and mechanism of the reaction between 4-t-butylphenolate anion and tetrahydroxoargentate(III) in aqueous alkaline media, 217-22

Kinetic studies of the oxidation reaction of arsenic(III) to arsenic(V) by peroxodisulphate ion in aqueous alkaline media, 357-60

The reaction of hydroxo-species of palladium(II) with bidentate nucleophiles, 375-8

A kinetic study of the gas-phase thermolysis of pentaborane(11), 385-90

Influence of added hydrogen on the kinetics and mechanism of thermal decomposition of tetraborane(10) and of pentaborane(11) in the gas phase, 391-8

Mechanistic studies on hydrazido(2-) complexes. Cleavage of the nitrogen-nitrogen bond in the reaction of $[\text{Mo}(\text{NN}(\text{CH}_2)_4\text{CH}_2)_2(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$ with acid, 425-30

A kinetic and mechanistic study of the solvolysis of aryltellurium trihalides in organic solvents, 457-60

The kinetics and mechanism of the hyponitrous acid-nitrous acid reaction at 0 °C, 533-8

Addition of anilines to the $[\text{Fe}(\text{CO})_3(1-\text{5-}\eta\text{-C}_7\text{H}_9)]\text{BF}_4^-$ complex, and the 'Ordered Transition State Mechanism', 561-6

Base-catalysed aquation of $\alpha\beta\text{-syn}$ - and $\alpha\beta\text{-anti}$ -chloro- and bromo-[1,9-bis(2'-pyridyl)-2,5,8-triazaanonane]- and -[1,11-bis(2'-pyridyl)-2,6,10-triazaundecane]-cobalt(III) cations. An unusually base-sensitive halogenopentamine cobalt(III) system, 645-54

A kinetic study of the transmission of ligand effects across the quadruple bond in $[\text{Re}_2\text{Cl}_8]^{2-}$, 767-72

Kinetics of aquation and base hydrolysis of some *trans*- $[\text{Co}(\text{tn})_2(\text{B})\text{Cl}]^{n+}$ ($\text{tn} = 1,3\text{-diaminopropane}$; $\text{B} = \text{OH}$, NCS , N_3^- , NO_2 , NH_3 , or CN) complexes. The effect of ring size on reactivity, 853-8

Variable-temperature and -pressure proton nuclear magnetic resonance studies of the rates and mechanisms of *N,N*-dimethylformamide and acetonitrile exchange with the square-pyramidal complexes mono(solvento){3,7,11-tribenzyl-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene}nickel(II), 873-8

Reactivity studies on the hydrolysis of iron(II) complexes of substituted phenanthrolines in dioxane-water solvents, 1105-8

Composition, stability, and lability of copper(II) dipeptide complexes, 1293-8

Kinetics of the uncatalysed and silver(I)-catalysed oxidation of hydrogen peroxide by [ethylenebis(biguaniide)]silver(III) ion in acid perchlorate media, 1645-50

Substitution reactions on sterically hindered square-planar *trans*- $[\text{NiBr}(\text{R})(\text{PR}'_3)_2]$ ($\text{R} = \text{aryl}$) complexes. Effects of the substituents of the aryl ligand, 1669-74

Photofragmentation kinetics of some triruthenium carbonyl clusters, 1933-40

Kinetics of formation, dissociation, and isomerization reactions in complexes of cyanopyridines and benzonitrile with pentacyanoruthenate(II), 1973-8

Oxo transfer reactions. Mechanistic studies on the oxidation of triphenylphosphine by *trans*- $[\text{Ru}^{\text{VI}}\text{L}(\text{O})_2]^{2+}$ ($\text{L} = 1,4,8,11\text{-tetramethyl-1,4,8,11-tetra-azacyclotetradecane}$), 2065-8

KINETICS (contd)

Mechanism and energetics of reductive elimination of ethane from some tetramethylplatinum(IV) complexes, 2121–6
 Kinetics of reactions of hydrogen with $[Os_3(CO)_{11}(NCMe)]$ and $[Os_3(CO)_{10}(NCMe)_2]$, 2235–40

LABELLING

Stoichiometric and nitrogen-15 labelling studies on the hyponitrous acid–nitrous acid reaction, 527–32
 Radiotracers in fluorine chemistry. Part 12. Reactions between caesium fluoride and arsenic pentafluoride, boron trifluoride, sulphur tetrafluoride, carbonyl fluoride, and carbon dioxide under heterogeneous conditions at room temperature.
 Radiotracer studies using fluorine-18, sulphur-35, and carbon-14, 937–42

LANTHANIDE

Highly polydentate ligands. Part 4. Crystal structures of neodymium(III) and erbium(III) complexes of 3,12-bis(carboxymethyl)-6,9-dioxa-3,12-diazatetradecanedioate(4-), 185–92

Thermodynamic and electrochemical behaviour of lanthanide cryptates in *N,N*-dimethylformamide, 367–70

Preparation of the bis(trimethylsilyl)amido lanthanide chlorides $[\text{Ln}(\text{N}(\text{SiMe}_3)_2)_2(\mu\text{-Cl})(\text{thf})_2]$ (thf = tetrahydrofuran), and the crystal and molecular structures of the gadolinium and ytterbium complexes, 623–6

Heteroleptic poly(pyrazolyl)borate derivatives of the lanthanide ions. The synthesis of acetylacetone complexes and the molecular structures of $[\text{Ln}\{\text{HB}(\text{C}_3\text{N}_2\text{H}_3)_3\}_2\{\text{MeC(O)CHC(O)Me}\}]$ (Ln = Ce or Yb), 1393–400

Synthesis, X-ray characterization, and *Q*-band electron spin resonance studies of heterometallic, polymeric compounds formed by yttrium and lanthanide(III) nitrates with HgL_2 (L = $\text{C}_4\text{H}_6\text{NO}$, the anion of 2-pyrrolidone); crystal structures of $[(\text{Hg}_2\text{EuL}_4(\text{NO}_3)_3)_n]$ and $[(\text{Hg}_3\text{Tb}_2\text{L}_6(\text{NO}_3)_6)_n]$, 1439–48

Lanthanide ion complexes of the calixarenes. Part 4. Double inclusion by *p*-t-butylcalix[4]arene (H_4L). Crystal structures of $[\text{Eu}_2(\text{HL})_2(\text{dmf})_7\text{dmf}]$ (dmf = dimethylformamide) and $\text{H}_4\text{L}\text{·dmsO}$ (dmsO = dimethyl sulfoxide), 2217–22

LEAD

Lead(II) and barium(II) complexes of a potentially octadentate macrocyclic ligand (H_4L), capable of providing endogenous alkoxy and phenoxy bridges. The X-ray crystal structures of $[\text{Pb}(\text{H}_4\text{L})][\text{ClO}_4]_2$, $[\text{Pb}_2(\text{H}_2\text{L})][\text{ClO}_4]_2$, and $[(\text{H}_6\text{L})(\text{H}_2\text{O})_2][\text{ClO}_4]_2$, 19–24

Lewis-base adducts of lead(II) compounds. Part 4. Novel isomeric dimers of lead(II) thiocyanate-1,10-phenanthroline(1/2). Crystal structure determinations, 595–600

Chemistry of subvalent main group metals with dithiolates. Part 2. X-Ray structural, thermal, and theoretical investigations on the novel one-dimensional stacking structure $\text{K}_2[\text{Pb}\{\text{S},\text{C}=\text{C}(\text{CN})_2\}_2]\text{·}2\text{H}_2\text{O}$, 627–30

Complexes of lead(II) bis(O,O' -dialkyl dithiophosphates) with nitrogen donor ligands. The crystal structures of $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\text{·en}$ (en = ethylenediamine), $\text{Pb}[\text{S}_2\text{P}(\text{OEt})_2]_2\text{·bipy}$ (bipy = 2,2'-bipyridine), and $[\text{Pb}(\text{S}_2\text{P}(\text{OEt})_2)_2\text{·en}]$, 2443–8

LIGAND FIELD

Spectral intensity and charge distributions in the tetrahedral chromophores, bis(3-amino-2-methylpropane-2-thiolato)- and diacetatobis(ethylenethiourea)-cobalt(II), 2109–14

LITHIUM

Synthesis of α , γ -phosphorus functionalized alkyl lithium species; X-ray structures of $[(\text{Li}(\text{L})(\text{CH}_2\text{PMeR})_2)]$ [L = *N,N,N',N'*-tetramethylethylenediamine (tmen), R = Me or Ph; L = (-)-sparteine, R = Ph] and $[\text{Li}(\text{tmn})\{\text{CH}(\text{SiMe}_3)\text{C}_6\text{H}_4\text{PPh}_2\text{-}o\}]$, 105–14

Tris(trimethylsilyl)methyl and tris(dimethylphenylsilyl)methyl derivatives of boron. Crystal structures of dihydroxy[tris(trimethylsilyl)methyl]borane and of the lithium-boron complex $[(\text{MeOH})_2\text{Li}(\mu\text{-OMe})_2\text{B}(\text{OMe})_2]$, 447–52

MACROCYCLE

Lead(II) and barium(II) complexes of a potentially octadentate macrocyclic ligand (H_4L), capable of providing endogenous alkoxy and phenoxy bridges. The X-ray crystal structures of $[\text{Pb}(\text{H}_4\text{L})][\text{ClO}_4]_2$, $[\text{Pb}_2(\text{H}_2\text{L})][\text{ClO}_4]_2$, and $[(\text{H}_6\text{L})(\text{H}_2\text{O})_2][\text{ClO}_4]_2$, 19–24

Tetra-aza macrocyclic complexes containing a pendant amino

group. Copper(II), nickel(II), and cobalt(III) complexes of 5-aminomethyl-2,5,10,12-tetramethyl-1,4,8,11-tetraazacyclotetradecane (L) and X-ray structural characterisation of the cobalt(III) complex $[\text{Co}(\text{L})\text{Cl}][\text{ClO}_4]\text{Cl}\text{-EtOH}$, 85–92

Thermodynamics of the formation of metal complexes of 1,4,10-trioxa-7,13-diazacyclpentadecane-*N,N*'-diacetic acid and of 1,4,10,13-tetraoxa-7,16-diazacyclo-octadecane-*N,N*'-diacetic acid, 133–8

Oxovanadium(IV) and dioxomolybdenum(VI) complexes: synthesis from the corresponding acetylacetone complexes and X-Ray structures, 145–50

Ruthenium phthalocyanine chemistry: synthesis and properties of a mixed-valence nitrido-bridged ruthenium phthalocyanine dimer, 193–6

Stabilisation of trivalent nickel through 1:2 co-ordination by cyclic terdentate ligands $\text{CH}_2\text{CH}_2\text{NH}(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{X}$ (X = NH, O, or S), 229–32

Thermodynamic and electrochemical behaviour of lanthanide cryptates in *N,N*-dimethylformamide, 367–70

The binding of trialkyltin(IV) moieties to rat haemoglobin, and the structure of model systems, studied by tin-119 Mössbauer spectroscopy, 519–26

Reaction of elemental sulphur with $(5\text{H},14\text{H}\text{-}6,8,15,17\text{-tetramethylidbenzo}[b,i][1,4,8,11]\text{tetraazacyclotetradecinato})\text{iron(II)}$: synthesis and X-ray structure of a μ -sulphido iron(III) complex, 551–4

Variable-temperature and -pressure proton nuclear magnetic resonance studies of the rates and mechanisms of *N,N*-dimethylformamide and acetonitrile exchange with the square-pyramidal complexes mono(solvento){3,7,11-tribenzyl-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene}nickel(II), 873–8

Crystal structures and solution dynamics of mono- and di-nuclear rhodium complexes of heterodinucleating macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 915–20

Synthesis and spectroscopic studies of cobalt(III) complexes of the chiral pendant-arm macrocycle *N,N,N',N'*-tris[(2*S*)-2-hydroxypropyl]-1,4,7-triazacyclononane (H_3L): crystal structure of the hydrogen-bridged dimer $[\text{CoL}_3][\text{PF}_6]_3$, 931–6

Electronic spectral study of the aqua H_2O hydroxo equilibrium of model iron(III) haems encapsulated in aqueous detergent micelles, 1003–6

6,13-Diamino-6,13-dimethyl-1,4,8,11-tetraazacyclotetradecane, L^7 , a new, potentially sexidentate polyamine ligand. Variable co-ordination to cobalt(III) and crystal structure of the complex $[\text{Co}(\text{L}^7)]\text{Cl}_2[\text{ClO}_4]_2$, 1059–66

Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(III)–ethanol–carbon tetrachloride system, 1197–202

Autoxidation of a dicopper(I) macrocyclic complex: evidence for a mixed-valence intermediate, 1351–60

Effect of axially bound anions on the electroreduction of tin(IV) porphyrins in tetrahydrofuran, 1531–6

Functional metallomacrocycles and their polymers. Part 23. Biaxially orientated Langmuir–Blodgett films of substituted metallophthalocyanines, 1543–6

Preparation and characterisation of some nickel(II) tetra-aza macrocyclic complexes bearing pendant polymerisable groups as part of the ligand superstructure, 1599–608

Platinum metal thioether macrocyclic complexes: synthesis, electrochemistry, and single-crystal X-ray structures of *cis*- $[\text{RhCl}_2\text{L}^2]\text{PF}_6$ and *trans*- $[\text{RhCl}_2\text{L}^3]\text{PF}_6$ (L^2 = 1,4,8,11-tetraazacyclotetradecane, L^3 = 1,5,9,13-tetraazacyclohexadecane), 1675–80

Complexes of ligands providing endogenous bridges. Part 7. The concurrent formation of 1 + 1 and 2 + 2 oxa-azamacrocycles; the X-ray crystal structure of an exocyclic dinuclear copper(II) complex, 1727–38

Closely related macrocyclic and acyclic tridentate, pyridine derivatives, containing sulphur, and their complexes. Crystal structures of {dichloro-3,10-dithia-16-azabicyclo[10.3.1]hexadeca-1(16),12,14-triene}copper(II) and [2,6-bis(ethylthiomethyl)pyridine]dichlorocuppper(II), 1739–44

A barium dithiocyanate complex with an unusual macrocycle containing pendant arms, the amine precursor, and related copper(II) diperchlorate complexes: synthesis and structure, 1993–2002

Ruthenium oxo complexes of macrocyclic tertiary amines.

MACROCYCLE (contd)

- Synthesis, electrochemistry, and reactivities of ruthenium oxo complexes of *meso*-2,3,7,11,12-pentamethyl-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), and *X*-ray crystal structure of *trans*-[Ru^{IV} L^1 (O)(NCO)]ClO₄, 2011–16
- Oxo transfer reactions. Mechanistic studies on the oxidation of triphenylphosphine by *trans*-[Ru^{VI} L (O)₂]²⁺ ($L = 1,4,8,11$ -tetramethyl-1,4,8,11-tetra-azacyclotetradecane), 2065–8
- Crystal structure and magnetic properties of the heterobinuclear complex [(CH₃CN)LNi(mnt)Cu(mnt)]·CH₃CN ($L =$ tetrabenzo[*b,f,j,n*][1,5,9,13]tetra-azacyclohexadecine and mnt = maleonitriledithiolate), 2095–8
- Stereochemical role of lone pairs in main-group elements. Part 4. The crystal structure at 120 K of bis(1,4,7,10,13-pentaoxacyclopentadecane)tin(II) bis[trichlorostannate(II)], confirming the sterically inactive tin lone pair, 2155–60
- Characterisation of tridentate macrocyclic complexes of molybdenum(0) with tetra-aza macrocycles. Crystal structure of [Mo(CO)₃ L^1] ($L^1 = C$ -*meso*-5,5,7,12,12,14-hexamethyl-1,4,8,11-tetra- α -azacyclotetradecane), 2183–6
- Functionalized amino acid complexation to a cobalt(III) complex of (2R,5R,8R,11R)-2,5,8,11-tetraethyl-1,4,7,10-tetra-azacyclododecane. Crystal structures of complexes containing (*R*)- or (*S*)-serine and (*R*)- α -methylserine, and release of amino acid from the complexes, 2359–66
- Optical absorption and magnetic circular dichroism studies of hydrogen, copper(II), zinc(II), nickel(II), and cobalt(II) crown ether-substituted monomeric and dimeric phthalocyanines, 2397–406

MAGNESIUM

- An investigation of halide abstraction by arsenic(III); reactions of AsCl₃ with MCl₂ (M = Mg or Zn) and TiCl₄ in donor solvents (MeCN or POCl₃). Formation and crystal structure of the dimeric titanium(IV) compound [Ti₂Cl₆-(POCl₃)₂(PO₂Cl₂)₂], 57–60
- Preparation of the ordered perovskite (SrLa)(MgMo)O₆ and its physicochemical characterization, 2335–8

MAGNETISM

- Crystal structure, spectroscopic and magnetic properties of the complex [Cu(paphy)(NCS)(SCN)] (paphy = pyridine-2-carbaldehyde 2'-pyridylhydrazone). An unusual di- μ -thiocyanato-*N* bridged copper(II) dimer, 53–6
- Crystal structure and magnetic properties of 1-aqua- μ -hydroxo-1,2,2-tris(perchlorato)-1,2-bis(2,2';6,2"-terpyridine)dicopper(II), 237–42
- Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NNNN'*-tetra-acetate (cdta). Part 3. Structural and magnetic characterization of the dinuclear [M(OH₂)₅][M'(cdta)]·H₂O (M,M' = Ni,Ni; Mn,Ni; Mn,Cu; Co,Ni; or Co,Cu) and the tetranuclear [M(OH₂)₅][M'(cdta)(OH₂)₄H₂O (M,M' = Zn,Zn; Zn,Co; Co,Co; or Mn,Co) complexes, 863–72
- Synthesis and characterization of imidazolate-bridged iron(III)–copper(II) and manganese(III)–copper(II) binuclear complexes, 943–6
- Magnetic properties of mono-, bi-, and tri-nuclear copper(II) complexes of novel oxamato and oxamido ligands. Crystal structure of a mononuclear precursor, 1017–26
- Low-lying electronic states in μ -phenolato copper(II) binuclear compounds with hydroxo, ethanolato, azido, and cyanato exogenous bridges: crystal structures, magnetic properties, and interpretations, 1117–26
- Preparation and characterisation of linear mixed-metal trinuclear carboxylate complexes of general formula [M₂M'(O₂Cr)₆(C₉H₇N)₂]_n; crystal structure of [Zn₂Ba(O₂CCMe₃)₆(C₉H₇N)₂]_n, 1127–32
- Synthesis, spectroscopic, and magnetic studies of copper(II) with *N*-protected amino acids: crystal and molecular structure of diaquatetrakis(μ -*N*-benzoyl-*z*-alaninato)-dicopper(II), 1341–4
- Dimeric mixed-ligand copper(II) complexes with di-imines and salicylaldehydes, 1435–8
- Stacked planar molecules in antiferromagnetic di-isothiocyanatobis(thiourea)chromium(II), a new structural type for chromium(II), 1631–4
- Binuclear copper(II) complexes of new dinucleating ligands with a pyrazolate group as an endogenous bridge. Effects of exogenous azido and acetate bridges on magnetic properties, 2077–82
- Crystal structures and magnetic properties of bi- and tetra-nuclear copper(II) complexes of 2,6-diformyl-4-methylphenol di(benzoylhydrazone), 2083–8

Crystal structure and magnetic properties of the heterobinuclear complex [(CH₃CN)LNi(mnt)Cu(mnt)]·CH₃CN ($L =$ tetrabenzo[*b,f,j,n*][1,5,9,13]tetra-azacyclohexadecine and mnt = maleonitriledithiolate), 2095–8

Synthesis, spectroscopic, electrochemical, and magnetic properties of dimolybdenum(II,II), diruthenium-(II,III) and -(II,II) complexes containing bridging aspirinate (2-acetoxybenzoate) ligands, 2435–42

MANGANESE

- Metal complexes (Co³⁺, Fe³⁺, Mn³⁺) of the quinquedentate salen analogue, 2-methyl-4-benzylthio-*N,N'*-butane-1,2-diybis(salicylideneimine): crystal structures and sulphur coordination effects, 179–84
- Synthesis of diphenylphosphido-bridged manganese-gold complexes. Crystal structures of [N(PPh₃)₂]₂[Au₂Mn(μ -PPh₂)₂(CO)₂]₂ and [Au₂Mn₂(μ -PPh₂)₄(CO)₈]₂, 639–44
- Synthesis and characterization of imidazolate-bridged iron(III)–copper(II) and manganese(III)–copper(II) binuclear complexes, 943–6
- Reactions of heteropolyanions in non-polar solvents. Part 3. Activation of dioxygen by manganese(II) centres in polytungstates. Oxidation of hindered phenols, 1483–90
- Molecular structure of heterobimetallic μ - η^2 -(C,C)-ketene complexes, 1627–30
- Neutron diffraction study of the cation distributions in the systems Fe_{7-x}M_x(PO₄)₆ (M = Mn or Co), 1765–70
- Reductive nitrosylation of tetraoxometalates. Part 15. Generation of Mn(NO)³⁺, Mn(NO)²⁺ and Mn(NO)₂²⁺ moieties directly from MnO₄⁻. Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese, 1963–72
- Reductive nitrosylation of tetraoxometalates. Part 15. Generation of Mn(NO)³⁺, Mn(NO)²⁺, and Mn(NO)₂²⁺ moieties directly from MnO₄⁻. Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese (1989, 1963), 2461

MASS SPECTROMETRY

- A kinetic study of the gas-phase thermolysis of pentaborane(11), 385–90
- Influence of added hydrogen on the kinetics and mechanism of thermal decomposition of tetraborane(10) and of pentaborane(11) in the gas phase, 391–8
- Crystal structure and mass spectrometry of dibromodimethyl[1-methyl-2(3*H*)-imidazolinethione-*S*]tin(IV), 927–30

MATRIX ISOLATION

- Spectroscopic studies in matrix isolated tungsten chlorides and bromides, 313–16
- Pt-oxidation of alcohols by hexachlorometalate(IV) ions (M = Pt, Pd, or Ir): spin trapping and matrix isolation electron spin resonance studies, 1237–40

MECHANISM

- Kinetics and mechanism of the reaction between 4-*t*-butylphenolate anion and tetrahydroxoargentate(III) in aqueous alkaline media, 217–22
- Mechanistic studies on hydrazido(2-)–complexes. Cleavage of the nitrogen–nitrogen bond in the reaction of [Mo{NN(CH₂)₄CH₂}₂(Ph₂PCH₂CH₂PPh₂)₂] with acid, 425–30
- The kinetics and mechanism of the hyponitrous acid–nitrous acid reaction at 0 °C, 533–8
- The mechanisms of fluxionality of [Ir₄(CO)₁₁(PEt₃)]; a reappraisal of the mechanism of carbonyl fluxionality on clusters, 889–94
- Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(III)–ethanol–carbon tetrachloride system, 1197–1202

MERCURY

- Preparation of (chloromethyl)palladium(II) derivatives from complexes of palladium dichloride by reaction with diazomethane or bis(chloromethyl)mercury, 761–6
- Preparation and crystal structure of a binuclear compound of methylmercury(II) containing bismuthiol I as a bridging ligand, 1409–12
- Synthesis, X-ray characterization, and Q-band electron spin resonance studies of heterometallic, polymeric compounds formed by yttrium and lanthanide(III) nitrates with HgL₂ ($L = C_4H_6NO$, the anion of 2-pyrrolidone); crystal structures of [{Hg₂EuL₄(NO₃)₃}] and [{Hg₂Tb₂L₆(NO₃)₆}], 1439–48
- Synthesis of the cluster dianion [Os₁₁C(CO)₂₇]²⁻ by pyrolysis and its reactions with electrophiles; X-ray structure analysis of the mixed-metal derivative [PMePh₃][Os₁₁C(CO)₂₇](Cu(NCMe))₂],

MERCURY (contd)

- CH_2Cl_2 and the hydrido derivative $[\text{PMePh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455-64
 Synthetic, spectroscopic, and X-ray studies on methylmercury(II) and dimethylthallium(III) complexes with cyclopentanone thiosemicarbazone, 1787-92
 Synthesis of arylthallium complexes using organomercury compounds. Crystal and molecular structures of $[\text{NMe}_4][\text{Tl}(\text{C}_6\text{H}_4\text{N}=\text{NPh}-2)\text{Cl}_3]$ and $[\text{Tl}(\text{C}_6\text{H}_4\text{CH}_2\text{NMe}_2-2)\text{Cl}_2]$, 2241-6
 Crystal and molecular structure of a lattice adduct, $\text{HgPh}_3 \cdot 2[\text{Hg}(\text{SCN})_2(\text{PPh}_3)_2]$: an unusual case of two- and four-co-ordinate mercury in the same unit cell, 2339-42

MESITYL

- Mesitylgold(I) complexes. Crystal structures of $\text{NEt}_4[\text{Au}(\text{mes})\text{Cl}]$ and $[\text{N}(\text{PPh}_3)_2][(\text{mes})\text{Au}(\text{CN})\text{Au}(\text{mes})]$, 2127-32

METALLABORANE

- Metallaheteroborane chemistry. Part 3. Synthesis of $[2.2-(\text{PR}_3)_2-1,2-\text{TePtB}_{10}\text{H}_{10}]$ ($\text{R}_3 = \text{Et}_3$, $\text{Bu}^n{}_3$, or Me_2Ph), their characterisation by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of $[2.2-(\text{PEt}_3)_2-1,2-\text{TePtB}_{10}\text{H}_{10}]$ (1988, 2555), 383
 Polyhedral organoiridaborane chemistry. The reaction of MeNC with $[6-(\eta^5-\text{C}_5\text{Me}_5)-nido-6-\text{IrB}_9\text{H}_{13}]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[4,4-(\eta^5-\text{C}_5\text{Me}_5)(\text{MeNC})-arachno-4-\text{IrB}_9\text{H}_{12}]$, 1465-70
 Metallaheteroborane chemistry. Part 5. Synthesis of *cis*- $[2.2-(\text{PR}_3)_2-1,2-\text{SePtB}_{10}\text{H}_{10}]$ ($\text{R}_3 = \text{Et}_3$, $\text{Bu}^n{}_3$, or Me_2Ph) and *arachno*- $[9,9-(\text{PMe}_2\text{Ph})_2-6,9-\text{SePtB}_8\text{H}_{10}]$ and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of *cis*- $[2.2-(\text{PEt}_3)_2-1,2-\text{SePtB}_{10}\text{H}_{10}]$ and *arachno*- $[9,9-(\text{PMe}_2\text{Ph})_2-6,9-\text{SePtB}_8\text{H}_{10}]$, 1657-68

METALLACYCLE

- Metallacyclic complexes. Part 2. Synthesis of metallathietane-3,3-dioxide complexes of platinum(II) and palladium(II); the crystal structures of $[\text{Pt}\{\text{CH}(\text{COPh})\text{S}(\text{O})_2\}\overset{\text{C}}{\text{CH}}(\text{COPh})-\text{(PPh}_3)_2\cdot 2\text{CH}_2\text{Cl}_2$ and $[\text{Pd}\{\text{CH}(\text{COPh})\text{S}(\text{O})_2\}\overset{\text{C}}{\text{CH}}(\text{COPh})-\text{(PMePh}_2)_2\cdot \text{CH}_2\text{Cl}_2\cdot \text{H}_2\text{O}$, 259-66
 Chemistry of metallacyclobutanones (η^3 -oxodimethylenemethane complexes). Part 5. Ligand substitution reactions of metallacyclobutan-3-ones of platinum(II) and palladium(II) with tertiary phosphines and t-butyl isocyanide; crystal structures of $[\text{Pt}\{\text{CH}(\text{CO}_2\text{Me})\text{CO}\overset{\text{C}}{\text{CH}}(\text{CO}_2\text{Me})\}\{\text{CNBu}'\}(\text{AsPh}_3)]$ and $[\text{Pt}\{\text{CH}(\text{CO}_2\text{Me})\text{CO}\overset{\text{C}}{\text{CH}}(\text{CO}_2\text{Me})\}\overset{\text{C}}{\text{C}}(\text{NHBu}')\{\text{CNBu}'\}]$, 345-56

METHANIDE

- (Diphenylphosphinomethyl)diphenylphosphine sulphide ($\text{Ph}_2\text{PCH}_2\text{PPh}_2\text{S}$) and its methanide anion $\text{Ph}_2\text{PCHPPh}_2\text{S}^-$ as ligands in organogold chemistry. X-Ray crystal structure of *cis*- $[\text{Au}(\text{C}_6\text{F}_5)_2(\text{PPh}_2\text{CHPPh}_2\text{S})]$, 73-8
 Bi-, tetra-, and poly-nuclear complexes of gold(I) and silver(I) with bis(diphenylphosphino)-methanide or -amide acting as six-electron donors, 1883-6

METHANOL

- Nucleophilic attack of methanol on bis(benzonitrile)dichloroplatinum: formation of mono- and bis-imido ester derivatives, 947-52

METHOXIDE

- Reactivity of cationic molybdenum(II) complexes. Part 3. Synthesis of methoxide and carbomethoxide derivatives of η^5 -pentamethylcyclopentadienyl-molybdenum(II) and the crystal structure determination of the oxidation product *trans*- $[\{\text{Mo}(\eta^5-\text{C}_5\text{Me}_5)(\text{O})_2\}_2(\mu-\text{O})]$, 155-60

- Tris(trimethylsilyl)methyl and tris(dimethylphenylsilyl)methyl derivatives of boron. Crystal structures of dihydroxy[tris(trimethylsilyl)methyl]borane and of the lithium boron complex $[(\text{MeOH})_2\text{Li}(\mu-\text{OMe})_2\text{B}(\text{OMe})_2]$, 447-52

METHYL

- Halogeno, hydride, and alkyl carbonyl-monocyclopentadienyl niobium derivatives. Crystal and molecular structure of $[\text{Nb}(\eta^5-\text{C}_5\text{H}_5)\text{Cl}(\text{H})(\text{CO})(\text{PMe}_3)_2]$, 79-84
 Syntheses and dynamic stereochemistry of methyl and phenyl carbonyl heterobinuclear metal-metal bonded complexes, 379-82
 A dynamic nuclear magnetic resonance study of trimethylplatinum(IV) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part 1. Low-temperature solution

properties, and X-ray crystal structure of $[\text{Pt}(\text{Me}_3)_2(\text{Me})_2\text{CHCH}(\text{SMe})_2]$, 701-10

A dynamic nuclear magnetic resonance study of trimethylplatinum(IV) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part 2. High-temperature fluxionality, 711-18

Crystal structure and mass spectrometry of dibromodimethyl[1-methyl-2(3*H*)-imidazolinethione-*S*]tin(IV), 927-30

Preparation and crystal structure of a binuclear compound of methylmercury(II) containing bismuthiol I as a bridging ligand, 1409-12

Synthesis, characterization, and thermolytic behaviour of η^3 -allyl(methyl)(tertiary phosphine)palladium(II), and X-ray structure of the cluster $[\text{Pd}_4(\mu-\text{C}_3\text{H}_5)_2(\mu-\text{Cl})_2(\mu-\text{dppm})_2] \cdot 4\text{C}_4\text{H}_8\text{O}$ (dppm = $\text{Ph}_2\text{PCH}_2\text{PPh}_2$), 1519-26

Synthetic, spectroscopic, and X-ray studies on methylmercury(II) and dimethylthallium(III) complexes with cyclopentanone thiosemicarbazone, 1787-92

Synthesis and properties of trimethyl(trifluorosilyl)stannane, 1879-82

Mechanism and energetics of reductive elimination of ethane from some tetramethylplatinum(IV) complexes, 2121-6

METHYLLALLYL
\eta^3-甲基(三甲基乙酰基)(三氯烷基)配合物的铂(II)和钯(II)的晶状学和核磁共振研究, 615-22
MICELLE

电子光谱研究模型铁(III)血红蛋白在洗涤剂微乳液中的水合平衡, 1003-6

MIXED VALENCE

钌(II) phthalocyanine化学:合成和性质, 193-6

三核混合价态配合物的线性 $[\text{Rh}_3]^{4+}$ 核心, 689-92

二铜(II)大环配合物的自动氧化:证据为一个混合价态中间体, 1351-60

合成, 晶状学, Raman 和共振拉曼光谱研究铂(II), 铂(IV)和卤代链铂(II, IV)配合物涉及三齿配体 3-甲基-3-亚丙基-1,5-二胺 (4Me-dien), 1689-96

MOLECULAR ORBITAL

理论研究二氮化物和相关分子与镍(0)的共配位:从模型化合物 $[\text{Ni}(\text{PH}_3)_2(\text{N}_2)]$, $[\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{CH}_2)]$, 和 $[\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{H}_2)]$ 的 *ab initio* 计算, 33-8

光化学研究季铵盐钼盐。第8部分:光活性作用于单晶中, 661-70

对 $[\text{NH}_3\text{Pr}^+]_6[\text{Mo}_8\text{O}_{26}(\text{OH})_2] \cdot 2\text{H}_2\text{O}$ 的研究, 1037-44

对 $[\text{Ru}_2(\text{O}_2\text{CCF}_3)_4]$ 和 $[\text{Ru}_2(\text{O}_2\text{CCF}_3)_4(\text{NO})_2]$ 的 *ab initio* 自洽场分子轨道研究:对新型电荷转移效应的研究, 线性 $\text{R}_2\text{S}_3\text{N}_2$ 和环形 XS_3N_2 系统, 1609-12

电子自旋共振研究铜(II)配合物 X-甘氨酸和甘氨-X型二肽, 和相关的三肽。变化的共配位模式与 pH 在液体和冷冻水溶液中的影响, 1925-32

一些含碳异构键的氧化加成到钨和铂的某些配合物, 2099-108

4-苯基-1,2,3,5-四硫杂环丁烷配合物 $[\text{Fe}_2(\text{CO})_6(\text{PhCN}_2\text{S}_2)]$:其制备, 晶状学, 和一个扩展的 Hückel 分子轨道研究, 2229-34

MOLECULAR SIEVE

热和水热稳定性 SAPO-5 和 SAPO-37 分子筛, 1361-2

MOLYBDENUM

合成, 电化学, 和电子自旋共振光谱研究十七个电子配合物的某些含 Mo(NO)²⁺ 核心和分子结构, 127-32

从相应的乙酰乙酸盐配合物合成的二茂钼(VI)配合物:合成和 X-射线结构, 145-50

阳离子钼(II)配合物的反应性。第3部分:含甲氧基和乙酰乙酸盐的 η^1 -

MOLYBDENUM (contd)

- pentamethylcyclopentadienyl-molybdenum(II) and the crystal structure determination of the oxidation product *trans*-[$\{\text{Mo}(\eta\text{-C}_5\text{Me}_5)_2\}_2(\mu\text{-O})$], 155–60
- A comparison of the rates of intramolecular hydrogen migration in the molecules [$\text{Mo}(\eta\text{-C}_7\text{H}_8)_2\}_2^+$ to give [$\text{Mo}(\eta\text{-C}_7\text{H}_7)(\eta\text{-C}_7\text{H}_9)\}_n^+$, $n = 0$ or 1; crystal structures of [$\text{Mo}(\eta\text{-C}_7\text{H}_8)_2$], [$\text{Mo}(\eta\text{-C}_7\text{H}_7)(\eta\text{-C}_7\text{H}_9)$], and [$\text{Mo}(\eta\text{-C}_7\text{H}_8)_2\}$] [BF_4^-], 331–44
- Synthesis and spectroscopic characterisation of [$\text{MoO}(\text{SR})_4^-$] ($\text{R} = \text{C}_6\text{H}_5$, C_{10}H_7 , 4-CH₃C₆H₄, 4-CIC₆H₄, 3-CIC₆H₄, 4-FC₆H₄, 2,6-Cl₂C₆H₃, or C₆F₅) complexes and [$\text{MoO}(\text{SC}_6\text{F}_5)_4^-$], 413–8
- Mechanistic studies on hydrazido(2–)-complexes. Cleavage of the nitrogen–nitrogen bond in the reaction of [$\text{Mo}(\text{NN}(\text{CH}_2)_4\text{CH}_2\}$] ($\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPH}_2$) with acid, 425–30
- Reactions of co-ordinated ligands. Part 44. The reactivity of the carbyne complex [$\text{Mo}(\equiv\text{CCH}_2\text{Bu}')\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)$] towards protonic acids, diphenylphosphine, and the π-acid tetracyanoethylene; formation and crystal structures of [$\text{MoH}(\equiv\text{CCH}_2\text{Bu}')\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)\}$] [BF_4^-], [$\text{Mo}\{\sigma\text{-OC(O)CF}_3\}_2(\eta^2\text{-HOC(O)CF}_3)\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)$], and [$\text{Mo}(\equiv\text{CCH}_2\text{Bu}')\{\eta^2\text{-C}_2(\text{CN})_4\}\{\text{P}(\text{OMe})_3\}(\eta\text{-C}_5\text{H}_5)$], 511–8
- Photochemical studies of alkylammonium molybdates. Part 8. Location of protons interacting with the paramagnetic electron in a single crystal of photoirradiated [$\text{NH}_3\text{Pr}'_6[\text{Mo}_8\text{O}_{26}(\text{OH})_2]\cdot 2\text{H}_2\text{O}$], 661–70
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 85. Synthesis of chain and ring compounds containing molybdenum, 785–96
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 86.
- Alkylidyne(carborane)molybdenum–gold, –rhodium and –iron complexes; crystal structure of [$\text{NEt}_4\text{][MoFe}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)\text{-(}\mu\text{-}\sigma\text{:}\sigma\text{-}\eta^5\text{-C}_2\text{B}_9\text{H}_7\text{Me}_2\text{)}(\text{CO})_8$], 797–808
- Reactivity of cationic molybdenum(II) complexes. Part 4. Isolation and crystal structure determination of *cis*-[$\text{Mo}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)(\text{PPh}_3)(\text{N}_3)$] and *cis*-[$\text{Mo}(\text{CO})_2(\eta\text{-C}_5\text{Me}_5)(\text{PPh}_3)(\text{NCO})$], 959–64
- Monocyclopentadienyl pentafluorothiophenate complexes of molybdenum and their interactions with thallium(I) ions; dynamic nuclear magnetic resonance studies and crystal and molecular structures of [$\text{TiMo}(\text{SC}_6\text{F}_5)_2\text{L}_2(\text{cp})$] ($\text{L} = \text{CO}$ or SC_6F_5 ; $\text{cp} = \eta^5\text{-C}_5\text{H}_5$), 991–1002
- Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten [$\text{M}(\eta^3\text{-SC(CF}_3)=\text{C}(\text{CF}_3)\text{H})(\text{CF}_3\text{C}\equiv\text{CCF}_3)(\eta^5\text{-C}_5\text{H}_5)$] ($\text{M} = \text{Mo}$ or W). Their relevance to the mechanism of metal-promoted isomerisation of carbon–carbon double bonds, 1027–36
- Studies on transition metal peroxy complexes. Part 8. The nature of peroxomolybdates and peroxotungstates in aqueous solution, 1203–8
- Reactions of co-ordinated ligands. Part 45. Ligand displacement and oxidative reactions of the carbyne complex [$\text{Mo}(\equiv\text{CCH}_2\text{Bu}')\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)$]; crystal and molecular structures of [$\text{Mo}(\eta^3\text{-RN=C-C(CH}_2\text{Bu)}\text{=C(RNR)}(\text{CNR})_2(\eta\text{-C}_5\text{H}_5)$] ($\text{R} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), [$\text{Mo}(\text{C=CHBu}')\{\text{P}(\text{OMe})_3\}(\eta\text{-C}_5\text{H}_5)$], and [$\text{Mo}=\text{C}(\text{CH}_2\text{Bu}')\text{P}(\text{O})(\text{OMe})_2\}$ $\{\text{P}(\text{OMe})_3\}(\eta\text{-C}_5\text{H}_5)$], 1321–32
- Crystal and molecular structure of [1,2-bis(diethylphosphino)ethane mono-oxide-*P*O]trichloro-oxomolybdenum(V), 1417–20
- The structure of 18-molybdodisulphate(vi)(4–) ion in ($\text{NEt}_4\text{)_4S}_2\text{Mo}_6\text{O}_{62}\cdot\text{CH}_3\text{CN}$), 1491–8
- Comparative study on the multielectron reduction of $\text{HOC}_2\text{H}_4\text{N}_3$ catalyzed by [$\text{NBu}^n\text{}_4\text{][MoFe}_2\text{S}_4(\text{SPh})_3(o\text{-O}_2\text{C}_6\text{Cl}_4)_2$] and [$\text{NBu}^n\text{}_4\text{][Fe}_4\text{S}_4(\text{SPh})_4$]-modified glassy carbon electrodes, 1547–54
- Chemistry of phosphido-bridged dimolybdenum complexes. Part 3. Reinvestigation of the reaction between [$\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_6$] and P_2Ph_4 ; X-ray structures of [$\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\text{CO})_2$], [$\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\mu\text{-CO})$] and *trans*-[$\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2\text{O}(\text{CO})$], 1555–64
- Charge density in the [$\text{Mo}(\text{NCS})_6$]^{3–} ion in $\text{Cs}_3\text{Mo}(\text{NCS})_6$, 1779–86
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 91. Steric implications in the reactions of the complexes [$\text{M}(\equiv\text{CR})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)$] ($\text{M} = \text{Mo}$ or W , $\text{R} = \text{C}_6\text{H}_4\text{Me}-4$, $\text{C}_6\text{H}_4\text{OMe}-2$, or $\text{C}_6\text{H}_3\text{Me}_2-2,6$) with the

compounds [$\text{Mo}(\text{CO})_3(\text{NCMe})_3$] and [$\text{Cu}(\text{thf})(\eta\text{-C}_5\text{Me}_5)$] ($\text{thf} = \text{tetrahydrofuran}$); crystal structure of [$\text{Mo}_3(\mu\text{-CC}_6\text{H}_3\text{Me}_2-2,6)_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2$], 1871–8

Synthetic and structural studies on organomolybdenum and organotungsten complexes containing bismuth. X-Ray crystal structures of [$\text{BiCl}_4\text{Mo}(\text{CO})_3(\eta\text{-C}_5\text{H}_5)_2$] and [$\text{BiCl}_4\text{Mo}(\text{CO})_2(\text{CNBu}^i)(\eta\text{-C}_5\text{H}_5)_2$] (1988, 2941). 1887

Monomolybdonovanadate and *cis*- and *trans*-dimolybdo-octavanadate, 1915–24

Synthesis and characterization of bis(xanthate) complexes of Mo^{II} and W^{II} . Crystal structure of [$\text{W}(\text{S}_2\text{COPr}^i\text{-S})(\text{S}_2\text{COPr}^i\text{-S})\text{CO}_2(\text{PMMe}_3)_2$], 2003–10

Synthesis and X-ray crystal structure of tetra(2-methylphenyl)molybdenum(IV), $\text{Mo}(2\text{-MeC}_6\text{H}_4)_4$. Redox chemistry of $\text{M}(2\text{-MeC}_6\text{H}_4)_4$ compounds of molybdenum, rhenium, ruthenium, and osmium, 2149–54

Characterisation of tridentate macrocyclic complexes of molybdenum(0) with tetra-aza macrocycles. Crystal structure of [$\text{Mo}(\text{CO})_3\text{L}^1$] ($\text{L}^1 = \text{C-meso-5,5,7,12,12-hexamethyl-1,4,8,11-tetra-azacyclotetradecane}$), 2183–6

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 92. Synthesis of alkynylmethylidyne-molybdenum and -tungsten complexes and their reactions with [$\text{Co}_2(\text{CO})_8$] and [$\text{Mo}_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2$], 2261–8

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 93. Synthesis of heteropolynuclear metal compounds with chains of seven to eleven metal atoms; crystal structure of [$\text{Mo}_3\text{W}_3\text{Pt}_6(\mu_3\text{-CMe})_3(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\text{CO})_{10}\text{-}(\text{PM}_2\text{Ph})_4(\eta\text{-C}_5\text{H}_5)_5$], 2269–80

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 94. Synthesis of di- and tri-metal compounds *i/via* the reagents [$\text{M}(\equiv\text{C-C}\equiv\text{CBu}^i)(\text{CO})_2(\eta\text{-C}_5\text{H}_5)$] ($\text{M} = \text{Mo}$ or W), 2281–8

Preparation of the ordered perovskite ($\text{SrLa}(\text{MgMo})\text{O}_6$ and its physicochemical characterization, 2335–8

Synthesis, spectroscopic, electrochemical, and magnetic properties of dimolybdenum(II,III), diruthenium-(II,II) and -(II,II) complexes containing bridging aspirinate (2-acetoxybenzoate) ligands, 2435–42

MOLYBDENUM-95

Monomolybdonovanadate and *cis*- and *trans*-dimolybdo-octavanadate, 1915–24

MÖSSBAUER

A reinvestigation of the structures of organotin sulphates and chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74

Synthesis, infrared, and tin-119 Mössbauer spectroscopic characterization of seven-co-ordinate diorganotin(IV) adducts with 2,6-diacetylpyridine acylhydrazones, including the X-ray crystal structure of $\text{SnEt}_2(\text{dapt})$ [$\text{H}_2\text{dapt} = 2,6\text{-diacetylpyridine bis(2-thienylhydrazone)}$], 289–94

The binding of trialkyltin(IV) moieties to rat haemoglobin, and the structure of model systems, studied by tin-119 Mössbauer spectroscopy, 519–26

Gold-197 Mössbauer spectra and the bonding of some gold–gold and gold–platinum clusters, 539–44

NAPHTHALENE

The chemical and electrochemical one-electron oxidation of diamidonaphthalene-bridged complexes: paramagnetic [Rh_2]³⁺ containing A-frames, 683–8

Reaction of [$\text{Rh}_2\{\mu\text{-1,8-(NH)}_2\text{C}_10\text{H}_6\}_2(\text{CO})_2(\text{PPh}_3)_2$] with [$\text{M}(\text{PPh}_3)_2$] ($\text{M} = \text{Au}$, Ag , or Cu): formation of unusual 44-electron Rh_2M clusters. Crystal structure of [$\text{Rh}_2\{\mu\text{-Au}(\text{PPh}_3)_2\}\{\mu\text{-1,8-(NH)}_2\text{C}_10\text{H}_6\}_2(\text{CO})_2(\text{PPh}_3)_2\text{ClO}_4\cdot 2\text{CH}_2\text{Cl}_2$], 1249–52

NAPHTHYRIDONATE

Trinuclear mixed-valence complexes with the linear [Rh_3]⁴⁺ core, 689–92

NEUTRON DIFFRACTION

Neutron diffraction study of the cation distributions in the systems $\text{Fe}_{7-x}\text{M}_x(\text{PO}_4)_6$ ($\text{M} = \text{Mn}$ or Co), 1765–70

Tables of bond lengths determined by X-ray and neutron diffraction. Part 2. Organometallic compounds and co-ordination complexes of the *d*- and *f*-block metals, S1–S84

NICKEL

A theoretical study on the co-ordination of dinitrogen and related molecules to nickel(0): ‘*ab initio*’ calculations on the model compounds [$\text{Ni}(\text{PH}_3)_2(\text{N}_2)$], [$\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{CH}_2)$], and [$\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{H}_2)$], 33–8

NICKEL (contd)

- Tetra-aza macrocyclic complexes containing a pendant amino group. Copper(II), nickel(II), and cobalt(III) complexes of 5-aminomethyl-2,5,10,12-tetramethyl-1,4,8,11-tetra-azacyclotetradecane (L) and X-ray structural characterisation of the cobalt(III) complex $[\text{Co}(\text{L})\text{Cl}] (\text{ClO}_4)^-$ -EtOH, 85-92
- Studies in Werner clathrates. Part 9. Structures with bis(isothiocyanato)tetrakis(4-phenylpyridine)nickel(II) as host, 119-26
- Kinetics of the oxidation of chloride ions by tris(2,2'-bipyridine)nickel(III) ions in aqueous perchlorate media, 151-4
- The unusual co-ordination ability of vasopressin-like peptides; potentiometric and spectroscopic studies of some copper(II) and nickel(II) complexes, 173-8
- Stabilisation of trivalent nickel through 1:2 co-ordination by cyclic terdentate ligands $\text{CH}_2\text{CH}_2\text{NH}(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{X}$ ($\text{X} = \text{NH}, \text{O}$, or S), 229-32
- The nickel(II)-bromide system in dimethyl sulphoxide: a detailed study of the influences of temperature and mole ratio, 325-30
- Possible biological activity in proton, metal, and aminohydroxamic acid equilibria. Protonation and complex formation reactions between N-hydroxy-D-asparagine and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution (1988, 953, 383)
- Spectrophotometric and calorimetric studies on the formation of binary (2,2'-bipyridine)nickel(II) and ternary (2,2'-bipyridine)chloronickel(II) complexes in *N,N*-dimethylformamide, 655-60
- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 85. Synthesis of chain and ring compounds containing molybdenum, 785-96
- Variable-temperature and -pressure proton nuclear magnetic resonance studies of the rates and mechanisms of *N,N*-dimethylformamide and acetonitrile exchange with the square-pyramidal complexes mono(solvato){3,7,11-tribenzo-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene}nickel(II), 873-8
- Precursor catenand complexes: synthesis, structure, and electrochemistry of bis(2,6-di-iminopyridyl) complexes of nickel(II). The single-crystal X-ray structure of $[\text{NiL}^4_2][\text{BF}_4]^2$, 965-70
- Synthesis of a new P_2N_2 ligand *N,N'*-bis[2-(diphenylphosphino)phenyl]propane-1,3-diamine, H_2L^2 , and some of its complexes with elements of the nickel triad and rhodium: X-ray structure analyses of the neutral complex $[\text{NiL}^2]$ and the *trans*-spanned $[\text{Rh}(\text{CO})\text{Cl}(\text{H}_2\text{L}^2)]$, 1067-74
- Structure of μ -fluoro-bis[bis(2,2'-bipyridyl)fluoronickel(II)] fluoride-ethanol (1/2) trihydrate, revealing several unique features including strong hydrogen bonds between fluoride ions and ethanol molecules, 1273-6
- Thermodynamic and spectrophotometric investigation of complex formation between hydrogen, cobalt(II), nickel(II), and copper(II) ions and 2-amino-N-hydroxypropanamide in aqueous solution, 1299-310
- Co-ordination of nickel(II) ions by angiotensin II and its peptide fragments. A potentiometric, proton nuclear magnetic resonance and circular dichroism spectroscopic study, 1471-6
- Synthesis and crystal structure of $[\text{NiOs}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2\{\mu\text{-Cu}(\text{PPPh}_3)\}(\text{CO})_9]$. Hydrogenation-dehydrogenation reactions of dienes and alcohols with heterogeneous catalysts derived from $[\text{NiM}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-X})(\text{CO})_9]$ [$\text{M} = \text{Ru}, \text{X} = \text{H}; \text{M} = \text{Os}, \text{X} = \text{Cu}(\text{PPPh}_3)$], 1477-82
- Functional metallocacycle derivatives and their polymers. Part 23. Biaxially orientated Langmuir-Blodgett films of substituted metallophthalocyanines, 1543-6
- Preparation and characterisation of some nickel(II) tetra-aza macrocyclic complexes bearing pendant polymerisable groups as part of the ligand superstructure, 1599-608
- Molecular structure of heterobimetallic $\mu\text{-}\eta^2\text{-(C,C)-ketene}$ complexes, 1627-30
- Substitution reactions on sterically hindered square-planar *trans*- $[\text{NiBr}(\text{R})(\text{PR}')_2]$ ($\text{R} = \text{aryl}$) complexes. Effects of the substituents of the aryl ligand, 1669-74
- Mixed-metal complexes in solution. A potentiometric and spectrophotometric study of copper(II)-nickel(II)- δ -hydroxylysine complexes in aqueous solution, 1745-50
- Solid-state high-resolution carbon-13 magnetic resonance spectra of dithioacetate complexes of platinum, palladium, and nickel, 2457-60

NIOBIUM

Halogeno, hydride, and alkyl carbonyl-monocyclopentadienyl niobium derivatives. Crystal and molecular structure of $[\text{Nb}(\eta\text{-C}_5\text{H}_5)\text{Cl}(\text{H})(\text{CO})(\text{PMe}_3)_2]$, 79-84

NITRATE

Preparation and interconversion of dimeric di- μ -hydroxo and tri- μ -hydroxo complexes of platinum(II) and palladium(II) with 2,2'-bipyridine and 1,10-phenanthroline, 403-12

Synthesis, X-ray characterization, and Q-band electron spin resonance studies of heterometallic, polymeric compounds formed by yttrium and lanthanide(III) nitrates with HgI_2 ($\text{L} = \text{C}_6\text{H}_5\text{NO}$, the anion of 2-pyrrolidone); crystal structures of $[(\text{Hg}_2\text{Eu}_4(\text{NO}_3)_3)_n]$ and $[(\text{Hg}_3\text{Tb}_2\text{L}_6(\text{NO}_3)_6)_n]$, 1439-48

NITRIDE

Ruthenium phthalocyanine chemistry: synthesis and properties of a mixed-valence nitrido-bridged ruthenium phthalocyanine dimer, 193-6

NITRILE

Tri-aza analogue of an acetylacetone-complex formed by nucleophilic attack of a co-ordinated nitrile by amidines. X-Ray crystal structure of $[\text{Pt}\{(\text{HNC}(\text{Ph})\text{NC}(\text{Ph})\text{NH}\}_2]^2\text{PhMe}$, 837-42

NITRILOTRIACETATE

Crystal and molecular structures of diaqua(nitrilotriacetato)aluminium(III) and di- μ -hydroxo-bis(nitrilotriacetato)-dialuminato(III) dianion, 1513-8

NITRITE

Synthesis of ammonium ions and nitrosylation reactions using nitrosyl chloride and alkyl nitrates, 1139-42

NITROGEN

A theoretical study on the co-ordination of dinitrogen and related molecules to nickel(0): 'ab initio' calculations on the model compounds $[\text{Ni}(\text{PH}_3)_2(\text{N}_2)]$, $[\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{CH}_2)]$, and $[\text{Ni}(\text{PH}_3)_2(\text{N}_2\text{H}_2)]$, 33-8

Stoichiometric and nitrogen-15 labelling studies on the hyponitrous acid-nitrous acid reaction, 527-32

The kinetics and mechanism of the hyponitrous acid-nitrous acid reaction at 0 °C, 533-8

Solid-state structure and solution conformation of 2,2,4,4,6,6-hex(*p*-chlorophenoxy)cyclotri- λ^5 -phosphazatriene. X-Ray and dipole moment studies, 757-60

Isopropylidilsilylamine and disilyl-*t*-butylamine: preparation, spectroscopic properties, and molecular structure in the gas phase, determined by electron diffraction, 779-84

Reactions of amine hydrochlorides with phosphorus trichloride oxide; preparation and X-ray crystal structure of 2,4,6-trichloro-1,3,5-triphenyl-1,3,5,2 λ^5 ,4 λ^5 ,6 λ^5 -triazatrichrophosphorinan-2,4,6-trioxide (PhNPOCl_3), 821-8

Gas-phase molecular structure of *NN'*-bis(trimethylsilyl)sulphur di-imide, determined by electron diffraction, 859-62

New metal-sulphur nitrogen compounds from reactions in liquid ammonia. The X-ray structures of *trans*-bis(acetophenone dimethylhydrazone- N^+)dichloropalladium(II) and [di(azathien)-1-yl-S $^1\text{N}^4$][2-(hydrazonoethyl)phenyl]palladium(II), 1179-86

Contributions to the chemistry of boron. Part 194. Synthesis and characterization of mono-, bi-, and tri-cyclic boron-nitrogen-sulphur ring systems. Crystal structure of 2,7,9,14-tetrakis(diethylamino)-3,5,6,10,12,13-hexathia-1,8-diazab-2,4,7,9,11,14-hexaboratricyclo[9.3.0.0^{4,8}]tetradecane, 1311-20

An *ab initio* self-consistent field molecular-orbital study of novel stereoelectronic effects in linear $\text{R}_2\text{S}_3\text{N}_2$ and cyclic XS_3N_2 systems, 1609-12

Direct insertion of a nitrogen atom into the S-S bond of a 1,2,3,5-dithiadiazole ring in a direct current nitrogen glow discharge, and X-ray crystal structure of 4-methyl-1,2,3,5-dithiadiazole, 1705-14

Preparation and properties of $[\text{M}(\text{S}_2\text{N}_2\text{H})\text{Cl}(\text{L})]$ ($\text{M} = \text{Pt}, \text{L} = \text{PMe}_2\text{Ph}$ or C_2H_4 ; $\text{M} = \text{Pd}, \text{L} = \text{PMe}_2\text{Ph}$, $[\text{Pt}(\text{S}_3\text{N})\text{Cl}(\text{PMe}_2\text{Ph})]$, and corresponding ^{15}N -enriched complexes, 2179-82

NITROGEN-15

Stoichiometric and nitrogen-15 labelling studies on the hyponitrous acid-nitrous acid reaction, 527-32

Preparation and properties of $[\text{M}(\text{S}_2\text{N}_2\text{H})\text{Cl}(\text{L})]$ ($\text{M} = \text{Pt}, \text{L} = \text{PMe}_2\text{Ph}$ or C_2H_4 ; $\text{M} = \text{Pd}, \text{L} = \text{PMe}_2\text{Ph}$, $[\text{Pt}(\text{S}_3\text{N})\text{Cl}(\text{PMe}_2\text{Ph})]$, and corresponding ^{15}N -enriched complexes, 2179-82

NITROSYL

Synthesis, electrochemistry, and electron spin resonance spectra of some seventeen-electron complexes containing the $[\text{Mo}(\text{NO})]^{2+}$

NITROSYL (contd)

- core and the molecular structure of $[\text{Mo}(\text{NO})\{\text{HB}(\text{Me}_2\text{pz})_3\}(\text{pyr})_2]$, 127–32
- Synthesis and X-ray crystal structure of $[\text{Rh}(\text{NO})(\text{PPh}_3)_2 \cdot (p\text{-MeC}_6\text{H}_4\text{NNNC}_6\text{H}_4\text{Me}-p)]\text{[BF}_4^- \cdot \text{CH}_2\text{Cl}_2$, 745–8
- An investigation of the bonding in $[\text{Ru}_2(\text{O}_2\text{CCF}_3)_4]$ and $[\text{Ru}_2(\text{O}_2\text{CCF}_3)_4(\text{NO})_2]$, 1037–44
- Synthesis of ammonium ions and nitrosylation reactions using nitrosyl chloride and alkyl nitrites, 1139–42
- Reactivity of the iridium nitrosyl complex $[\text{Ir}(\text{NO})(\text{paa})(\text{PPh}_3)_2]\text{[PF}_6^-]_2$ ($\text{paa} = 2\text{-pyridinecarbaldehyde azine}$) with nucleophiles: insertion of NO into a methinic C–H bond; crystal structure of $[\text{Ir}\{(\text{2-C}_5\text{H}_4\text{N})\text{CH}=\text{N}-\text{N}=\text{C}(\text{NOH})-(\text{2-C}_5\text{H}_4\text{N})\}\{\text{Me}(\text{EtO})\text{C}=\text{NH}\}(\text{PPh}_3)_2]\text{[PF}_6^-]_2$, 1217–26
- Reductive nitrosylation of tetraoxometalates. Part 15. Generation of $\text{Mn}(\text{NO})^{3+}$, $\text{Mn}(\text{NO})^{2+}$, and $\text{Mn}(\text{NO})_2^{2+}$ moieties directly from MnO_4^- . Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese, 1963–72
- Synthesis and properties of the bis(μ -thiosulphato-*S*)-bis(dinitrosylferrate)(2–) anion. Crystal and molecular structure of bis[bis(triphenylphosphoranylidene)ammonium] bis(μ -thiosulphato-*S*)-bis(dinitrosylferrate), 2061–4
- Synthesis, spectroscopic, electrochemical, and magnetic properties of dimolybdenum(i,i), diruthenium-(ii,iii) and -(i,ii) complexes containing bridging aspirinate (2-acetoxybenzoate) ligands, 2435–42
- Reductive nitrosylation of tetraoxometalates. Part 15. Generation of $\text{Mn}(\text{NO})^{3+}$, $\text{Mn}(\text{NO})^{2+}$, and $\text{Mn}(\text{NO})_2^{2+}$ moieties directly from MnO_4^- . Synthesis, characterization, and electrochemistry of cyanonitrosyl derivatives of manganese (1989, 1963), 2461

NITROUS ACID

- Stoichiometric and nitrogen-15 labelling studies on the hyponitrous acid–nitrous acid reaction, 527–32
- The kinetics and mechanism of the hyponitrous acid–nitrous acid reaction at 0 °C, 533–8
- N.M.R.**
- Synthesis and stereochemical studies of η^3 -allyl palladium(ii) complexes containing a chiral chelating ligand, 161–8
- The nature of a sodium–crown interaction studied by sodium-23 nuclear magnetic resonance and by crystal structure determination of a complex with co-ordinated solvent, 203–10
- Correlations between nuclear magnetic resonance spectra and crystal structures. A carbon-13 nuclear magnetic resonance study in the solid state of the Group 15 xanthates $[\text{A}(\text{S}_2\text{COR})_3]$ ($\text{A} = \text{As}^{III}, \text{Sb}^{III}, \text{or Bi}^{III}; \text{R} = \text{Me, Et, or Pr}^I$), 233–6
- A reinvestigation of the structures of organotin sulphates and chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74
- Oxygen-17 nuclear magnetic resonance spectroscopic studies of aqueous alkaline silicate solutions, 275–82
- Metalla heteroborane chemistry. Part 3. Synthesis of $[2,2\text{-(PR}_3)_2\text{,1-TePtB}_{10}\text{H}_{10}]$ ($\text{R}_3 = \text{Et}_3, \text{Bu}^n{}_3, \text{or Me}_2\text{Ph}$), their characterisation by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of $[2,2\text{-(PEt}_3)_2\text{,1-2-TePtB}_{10}\text{H}_{10}]$ (1988, 2555), 383
- Co-ordination chemistry of higher oxidation states. Part 29. Cobalt(iii) complexes of bi- and tetra-thioethers: synthesis, properties, and cobalt-59 nuclear magnetic resonance data, 453–6
- Crystallographic and nuclear magnetic resonance studies of carbonyl(η^3 -methylallyl)(trichlorostannyl) complexes of platinum(ii) and palladium(ii), 615–22
- Base-catalysed aquation of $\alpha\beta$ -syn- and $\alpha\beta$ -anti-chloro- and bromo-[1,9-bis(2'-pyridyl)-2,5,8-triazaanone]- and -[1,11-bis(2'-pyridyl)-2,6,10-triazaundecane]-cobalt(iii) cations. An unusually base-sensitive halogenopentamine cobalt(iii) system, 645–54
- A dynamic nuclear magnetic resonance study of trimethylplatinum(iv) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part 1. Low-temperature solution properties, and X-ray crystal structure of $[\text{Pt}(\text{IMe}_3\{\text{MeS}\}_2)\text{CHCH(SMe)}_2]$, 701–10
- A dynamic nuclear magnetic resonance study of trimethylplatinum(iv) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part 2. High-temperature fluxionality, 711–8
- Solid-state phosphorus-31 nuclear magnetic resonance spectroscopy of phosphorus sulphides, 809–14
- High-resolution solid-state tin-119 nuclear magnetic resonance spectroscopy of ternary tin oxides, 843–8
- Variable-temperature and -pressure proton nuclear magnetic resonance studies of the rates and mechanisms of *N,N*-dimethyl-

- formamide and acetonitrile exchange with the square-pyramidal complexes mono(solvento){3,7,11-tribenzylo-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene}nickel(ii), 873–8
- Phosphorus-31 and selenium-77 nuclear magnetic resonance spectra of sulphur- or arsenic-substituted derivatives of tetraphosphorus triselenide; fitting of chemical shifts and coupling constants to molecular parameters for cage compounds containing a phosphorus–selenium–phosphorus linkage, 977–84
- Platinum metal complexes of potentially chelating alkene-thioether and alkene-selenoether ligands: synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{MeE}(\text{CH}_2)_n\text{CH}=\text{CH}_2\}]$ ($\text{M} = \text{Pt or Pd}; \text{X} = \text{Cl, Br, or I; E} = \text{S or Se}; n = 2 \text{ or } 3$) and the X-ray structure of *cis*-dibromo(2-thia-6-heptene)platinum(ii), $[\text{PtBr}_2\{\text{MeS}(\text{CH}_2)_3\text{CH}=\text{CH}_2\}]$, 985–90
- Monocyclopentadienyl pentafluorophenolate complexes of molybdenum and their interactions with thallium(i) ions; dynamic nuclear magnetic resonance studies and crystal and molecular structures of $[\text{TIMo}(\text{SC}_6\text{F}_5)_2\text{L}_2(\text{cp})]$ ($\text{L} = \text{CO or SC}_6\text{F}_5; \text{cp} = \eta^5\text{-C}_5\text{H}_5$), 991–1002
- Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten $[\text{M}\{\eta^3\text{-SC}(\text{CF}_3)=\text{C}(\text{CF}_3)\text{H}](\text{CF}_3\text{C}\equiv\text{C}\text{F}_3)\{\eta^5\text{-C}_5\text{H}_5\}]$ ($\text{M} = \text{Mo or W}$). Their relevance to the mechanism of metal-promoted isomerisation of carbon–carbon double bonds, 1027–36
- Zinc(ii) complexes of 4,7-diazadecane-1,10-diamine, 4,8-diazaundecane-1,11-diamine, and tetraethylenepentamine: a carbon-13 nuclear magnetic resonance study, 1171–8
- Isotopic multiplets in the carbon-13 nuclear magnetic resonance spectra due to partial deuteration of co-ordinated NH_2 and NH groups of cobalt(iii) complexes containing an N-glycoside. Importance of sugar ring oxygen atoms in the long-range isotope shifts, 1241–8
- Structure and dynamic behaviour of two isomers of $[\text{Fe}_3(\text{CO})_9\{\text{P}(\text{OR}_3)\}_3]$ ($\text{R} = \text{Me, Et, or Ph}$) derivatives, 1277–82
- Sulphonamide nitrogen-containing, nitrogen protected amino acids interacting with the Cd^{2+} ion. A cadmium-113 nuclear magnetic resonance and potentiometric study. X-Ray diffraction of $[\text{Cd}(\text{ts-}\beta\text{-alaO})_2(\text{H}_2\text{O})_4]$, 1345–50
- Polyhedral organoiridaborane chemistry. The reaction of MeNC with $[6-(\eta^5\text{-C}_5\text{Me}_5)\text{-nido-6-IrB}_9\text{H}_{12}]$. Nuclear magnetic resonance studies and the crystal and molecular structure of $[4,4-(\eta^5\text{-C}_5\text{Me}_5)(\text{MeNC})\text{-arachno-4-IrB}_9\text{H}_{12}]$, 1465–70
- Co-ordination of nickel(ii) ions by angiotensin II and its peptide fragments. A potentiometric, proton nuclear magnetic resonance and circular dichroism spectroscopic study, 1471–6
- Binuclear copper(i) complexes containing two or three 2-(diphenylphosphine)pyridine (dpyp) bridging ligands: X-ray crystal structure of $[\text{Cu}_2(\mu\text{-dpyp})_3(\text{MeCN})][\text{BF}_4^-]_2$ and variable-temperature ^1H and $^{31}\text{P}\{-^1\text{H}\}$ nuclear magnetic resonance studies, 1499–506
- Preparation and characterisation of some nickel(ii) tetra-aza macrocyclic complexes bearing pendant polymerisable groups as part of the ligand superstructure, 1599–608
- Nature of peroxyborate species in aqueous solution: a study by boron-11 nuclear magnetic resonance and Raman spectroscopy, 1651–6
- Metalla heteroborane chemistry. Part 5. Synthesis of *closo*-[2,2-(PR_3)₂1,2-SePtB₁₀H₁₀] ($\text{R}_3 = \text{Et}_3, \text{Bu}^n{}_3, \text{or Me}_2\text{Ph}$) and *arachno*-[9,9-(PMo_2Ph)₂6,9-SePtB₈H₁₀] and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of *closo*-[2,2-(PEt_3)₂1,2-SePtB₁₀H₁₀] and *arachno*-[9,9-(PMo_2Ph)₂6,9-SePtB₈H₁₀], 1657–68
- The mechanisms of fluxionality of $[\text{Fe}_3(\text{CO})_{12-n}\{\text{P}(\text{OR})_3\}_n]$ ($\text{R} = \text{Me, n = 0–3; R = Pr}^I, n = 2 \text{ or } 3$) and the X-ray structure of $[\text{Fe}_3(\text{CO})_{10}\{\text{P}(\text{OMe})_3\}_2]$, 1831–44
- Multinuclear magnetic resonance studies of the hydrolysis of aluminium(iii). Part 9. Prolonged hydrolysis with aluminium metal monitored at very high magnetic field, 1889–96
- Multinuclear magnetic resonance studies of the hydrolysis of aluminium(iii). Part 10. Proton, carbon-13, and aluminium-27 spectra of aluminium acetate at very high magnetic field, 1897–902
- Monomolybdonavanadate and *cis*- and *trans*-dimolybdo-octavanadate, 1915–24
- Relation between the torsion angles of acetate chelate rings and the deuteron nuclear magnetic resonance chemical shifts for polyamine-N-carboxylatochromate(iii) complexes, 1947–50
- Hydrogen scrambling processes in $[\text{Ta}(\eta\text{-C}_5\text{Me}_5)(\eta\text{-C}_5\text{H}_5)\{(\mu\text{-H})_2\text{BH}_2\}]$: a surprising observation, 2133–8

N.M.R. (contd)

- Investigations of tin(II) halide aqueous solutions and solvent extracts by ^{119}Sn nuclear magnetic resonance spectroscopy: characterisation of the trihalogenostannate(II) anions $[\text{SnX}_3]^-$, $[\text{SnX}_2\text{Y}]^-$, and $[\text{SnX}(\text{Y})\text{Z}]^-$ ($\text{X}, \text{Y}, \text{or Z} = \text{Cl}, \text{Br}, \text{or I}$), 2223-8
 Platinum metal complexes of potentially chelating alkene thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{E}[(\text{CH}_2)_2\text{CR=CR}_2]\}_2]$ ($\text{M} = \text{Pt or Pd}; \text{X} = \text{Cl, Br, or I}; \text{E} = \text{S or Se}; n = 2 \text{ or } 3; \text{R} = \text{H or Me}$) and the X -ray crystal structure of *cis*-di-iodo(5-thianona-1,8-diene)platinum(II), $[\text{PtI}_2\{\text{S}[(\text{CH}_2)_2\text{CH=CH}_2]\}_2]$, 2315-22
 Substituted cyclopentadienyl complexes. Part 4. The catalysed synthesis and proton nuclear magnetic resonance spectra of $[\text{Ru}(\eta^5-\text{C}_5\text{H}_4\text{Me})(\text{CO})(\text{L})\text{I}]$ and $[\text{Ru}(\eta^5-\text{C}_5\text{H}_4)(\text{CO})(\text{L})\text{I}]$ ($\text{L} = \text{phosphine or isocyanide}$) and the crystal structure determination of $[\text{Ru}(\eta^5-\text{C}_5\text{H}_4)(\text{CO})\{\text{P}(\text{CH}_2\text{Ph})_3\}\text{I}]$, 2407-14
 Solid-state high-resolution carbon-13 magnetic resonance spectra of dithioacetate complexes of platinum, palladium, and nickel, 2457-60

NON-TRANSITION METAL

- Vibrational study of ionic association in aprotic solvents. Part 12. Isothiocyanate M(NCS) complexes of non-transition-metal ions, 2193-8

N.Q.R.

- Structural and motional study of pyridinium tetrahalogenoantimonates(III) by means of nuclear quadrupole resonance spectroscopy, 631-4

NUCLEOTIDE

- Potentiometric and electron spin resonance investigation on the interactions between adenosine 5'-triphosphate (ATP) and triphosphate ligands with vanadyl ion. Synthesis and characterization of solid vanadyl-ATP compounds from acidic aqueous solution, 575-80
 X -Ray structure of a ternary complex, copper(II)-inosine 5'-monophosphate-benzimidazole. The first example for the absence of direct metal-nucleotide interaction, 1507-12

OSMIUM

- Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene ($\text{C}=\text{CHOEt}$) and ethoxyethyne ($\text{CH}\equiv\text{COEt}$) ligands respectively, 5-12
 Thiolate complexes of osmium(II) with tertiary phosphines and their dicarbonyl derivatives of osmium(II): X -ray crystal structure of *trans,trans,trans*- $[\text{Os}(\text{SC}_6\text{F}_5)_2(\text{CO})_2(\text{PEt}_2\text{Ph})_2]$, 419-24
 Metal cluster-induced electrophilicity of arenes, 481-8
 Polyhedral rearrangements in undeca- and dodeca-nuclear Group 1B metal osmium clusters, 505-10
 Studies on transition-metal oxo and nitrido complexes. Part 10. New oxo-ruthenium and oxo-osmium pyridine complexes, and use of the former as catalysts for oxidation of alcohols, 901-6
 Reactions of co-ordinated ligands. Part 46. Formation of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH=CMe}_2)(\text{CO})_{10}]$ by carbon-carbon σ -bond and carbon-hydrogen activation of 3,3-dimethylcyclopropene; crystal and molecular structure of $[\text{Os}_3(\mu\text{-H})(\mu\text{-CCH}_2\text{CHMe}_2)(\text{CO})_{10}]$ and evidence for its facile rearrangement to $[\text{Os}_3(\mu\text{-H})(\mu\text{-CH=CHCHMe}_2)(\text{CO})_{10}]$, 1333-40

- Synthesis, structure, and reactivity of the heteronuclear clusters $[\text{Os}_3\text{H}(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}]$ and $[\text{Os}_3(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}_2]$ ($\text{M} = \text{Cu, Ag, or Au}; \text{R} = \text{Ph or Me}$); the X -ray crystal structure of $[\text{Os}_3\text{H}(\text{CO})_{15}\{\text{Au}(\text{PPh}_3)\}]$, 1421-6

- Synthesis of the cluster dianion $[\text{Os}_{11}\text{C}(\text{CO})_{27}]^{2-}$ by pyrolysis and its reactions with electrophiles; X -ray structure analysis of the mixed-metal derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\{\text{Cu}(\text{NCMe})\}\cdot\text{CH}_2\text{Cl}_2$ and the hydrido derivative $[\text{PMcPh}_3][\text{Os}_{11}\text{C}(\text{CO})_{27}\text{H}]$, 1455-64

- Synthesis and crystal structure of $[\text{NiOs}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-Cu}(\text{PPh}_3))(\text{CO})_9]$. Hydrogenation-dehydrogenation reactions of dienes and alcohols with heterogeneous catalysts derived from $[\text{NiM}_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2(\mu\text{-X})(\text{CO})_9]$ ($\text{M} = \text{Ru, X} = \text{H}; \text{M} = \text{Os, X} = \text{Cu}(\text{PPh}_3)$), 1477-82

- Cluster activation *via* cyclometallation. Formation of the cyclometallated triosmium carbonyl cluster $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ and its reactions with phosphorus donor ligands. The X -ray crystal structures of $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ and $[\text{Os}_3\text{H}(\text{CO})_9\{\text{P}(\text{OMe})_3\}(\text{PPh}_3)]$, 2023-32

- Reactions of the cyclometallated triosmium cluster $[\text{Os}_3\text{H}_2(\text{CO})_9\{\text{P}(\text{C}_6\text{H}_4)\text{Ph}\}]$ with Brønsted acids and dihydrogen: synthesis and X -ray crystal structure of

- $[\text{Os}_3\text{H}_2(\text{CF}_3\text{CO}_2)(\text{CO})_9(\mu\text{-PPh}_2)]$, 2033-8

- Synthesis and X -ray crystal structure of tetra(2-methylphenyl)molybdenum(IV), $\text{Mo}(2\text{-MeC}_6\text{H}_4)_4$. Redox chemistry of $\text{M}(2\text{-MeC}_6\text{H}_4)_4$ compounds of molybdenum, rhenium, ruthenium, and osmium, 2149-54

- Stepwise conversion of $[\text{Os}(\text{CF}_3\text{CO}_2)_2(\text{CO})_4]$ into $[\text{Os}(\text{pyS})_2(\text{CO})_2]$: X -ray crystal structures of the complexes $[\text{Os}(\text{pyS})_x(\text{CO})_2]$ where $x = 2$ or 3 and pyS is the pyridine-2-thionato ligand, 2211-6

- Kinetics of reactions of hydrogen with $[\text{Os}_3(\text{CO})_{11}(\text{NCMe})]$ and $[\text{Os}_3(\text{CO})_{10}(\text{NCMe})_2]$, 2235-40

- A convenient route to binuclear osmium(I) complexes containing diphosphine and arsinophosphine ligands. X -Ray crystal structure of $[\text{Os}_2(\text{MeCO}_2)_2(\text{CO})_4(\text{dpmm})_2]$, 2303-8

- Reactivity of new osmium dihydrides with arenediazonium cations: preparation and properties of bis(aryldiazene) and mono(aryldiazido) complexes. A comparison with analogous iron and ruthenium derivatives, 2353-8

OXALATE

- Formation in solution, synthesis, and electrochemical study of oxalato complexes of *N,N'*-ethylenebis(salicylideneiminato)-chromium(III) and -iron(III): crystal structures of piperidinium [*N,N'*-ethylenebis(salicylideneiminato)](oxalato- O^2)-chromate(III) and -ferrate(III), 729-38

OXAMATE

- Magnetic properties of mono-, bi-, and tri-nuclear copper(II) complexes of novel oxamato and oxamido ligands. Crystal structure of a mononuclear precursor, 1017-26

OXAMIDE

- Magnetic properties of mono-, bi-, and tri-nuclear copper(II) complexes of novel oxamato and oxamido ligands. Crystal structure of a mononuclear precursor, 1017-26

OXIDATIVE ADDITION

- Redox activation of the hexanuclear ruthenium cluster $[\text{Ru}_6(\text{CO})_{16}]^{2-}$ using oxidative addition, 243-6

- Studies of the oxidative addition of some substrates containing carbon-heteroatom bonds to some tungsten and platinum complexes, 2099-108

- Oxidative addition of dibenzoyl peroxide on metallic copper. The crystal structure of $[[\text{Cu}(\text{C}_6\text{H}_5\text{CO}_2)_2(\text{py})_2]]$, 2331-4

OXIDE

- A further investigation of the polarized single-crystal electronic absorption spectrum of $[\text{AsPh}_4][\text{CrOCl}_4]$, 1-4

- Oxovanadium(IV) and dioxomolybdenum(VI) complexes: synthesis from the corresponding acetylacetone complexes and X -Ray structures, 145-50

- Reactivity of cationic molybdenum(II) complexes. Part 3. Synthesis of methoxide and carbomethoxide derivatives of η -pentamethylcyclopentadienyl-molybdenum(II) and the crystal structure determination of the oxidation product *trans*- $[\{\text{Mo}(\eta\text{-C}_5\text{Me}_5)(\text{O})_2\}_2(\mu\text{-O})]$, 155-60

- Synthesis and spectroscopic characterisation of $[\text{MoO}(\text{SR})_4]^-$ ($\text{R} = \text{C}_6\text{H}_5, \text{C}_{10}\text{H}_7, 4\text{-CH}_3\text{C}_6\text{H}_4, 4\text{-ClC}_6\text{H}_4, 3\text{-ClC}_6\text{H}_4, 4\text{-FC}_6\text{H}_4, 2,6\text{-Cl}_2\text{C}_6\text{H}_3$, or C_6F_5) complexes and $[\text{MoO}(\text{SC}_6\text{F}_5)_4]^{2-}$, 413-8

- Actinide structural studies. Part 17. The crystal and molecular structures of four uranyl(VI) complexes with *N,N'*-dialkylthiocarbamate ligands, 471-80

- High-resolution solid-state tin-119 nuclear magnetic resonance spectroscopy of ternary tin oxides, 843-8

- Catalytic oxidation of water mediated by rhenium oxides, 849-52

- Studies on transition-metal oxo and nitrido complexes. Part 10.

- New oxo-ruthenium and oxo-osmium pyridine complexes, and use of the former as catalysts for oxidation of alcohols, 901-6

- Synthesis and X -ray crystal structure of $[\text{ReO}(\text{Cl})\{2\text{-OC}_6\text{H}_4\text{CH=NCH}_2\text{CH}_2\text{OCH}_2\}_2]$, an example of four-coordination by a potentially sexidentate ligand, 1405-8

- Crystal and molecular structure of $[\text{L}_2\text{bis}(\text{diethylphosphino})\text{ethane}-\text{mono-oxide-P}_2\text{O}]$ trichloro-oxomolybdenum(V), 1417-20

- Chemistry of phosphido-bridged dimolybdenum complexes. Part 3.

- Reinvestigation of the reaction between $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_6]$ and P_2Ph_4 . X -ray structures of $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\text{CO})_2]$, $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\mu\text{-CO})]$ and *trans*- $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2\text{O}(\text{CO})]$, 1555-64

- Phenylin trichloroacetate complexes: preparations and structures, 1589-98

- Ruthenium oxo complexes of macrocyclic tertiary amines.

- Synthesis, electrochemistry, and reactivities of ruthenium oxo complexes of *meso*-2,3,7,11,12-pentamethyl-3,7,11,17-tetra-

OXIDE (contd)

- azabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), and X-ray crystal structure of $trans$ -[Ru^{IV} L^1 (O)(NCO)]ClO₄, 2011–6
Oxo transfer reactions. Mechanistic studies on the oxidation of triphenylphosphine by $trans$ -[Ru^{VI} L (O)₂]²⁺ (L = 1,4,8,11-tetramethyl-1,4,8,11-tetra-azacyclotetradecane), 2065–8
Electrochemistry of the quasi-reversible bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate(v) and -(iv) and bis[2-hydroxy-2-methylbutanoato(2–)]oxochromate(v) and -(iv) redox couples and the crystal and molecular structure of sodium bis[2-ethyl-2-hydroxybutanoato(2–)]oxochromate(v) sesquihydrate, 2205–10
Preparation of the ordered perovskite (SrLa)(MgMo)O₆ and its physicochemical characterization, 2335–8
Aqueous chemistry and thermodynamic modelling of CaO-SiO₂-H₂O gels, 2371–80

OXIME

- Rearrangements of C=N-functionalities promoted by cobalt(i): reactivity of cyclopentadienylbis(ethylene)cobalt(i) with dibenzophenone azine, benzophenone oxime, benzylidene(phenyl)amine and crystal structure determinations of the products, 1957–62

OXYGEN-17

- Oxygen-17 nuclear magnetic resonance spectroscopic studies of aqueous alkaline silicate solutions, 275–82
Monomolybdonavanadate and *cis*- and *trans*-dimolybdo-octavanadate, 1915–24

PALLADIATION

- Reactions of cyclopalladated compounds. Part 21. Various examples of sulphur-assisted intramolecular palladation of aryl and alkyl groups, 1715–20

PALLADIUM

- The crystal structures of K[Pd{S₂C₂(CN)₂}₂]·H₂O and [NH₄][Pd{S₂C₂(CN)₂}₂]·H₂O, 67–72
Synthesis and stereochemical studies of η^3 -allyl palladium(ii) complexes containing a chiral chelating ligand, 161–8
Synthesis and reactivity of binuclear homo- or hetero-metallic complexes [NBu₄]₂[MM'(μ-C₆F₅)₂(C₆F₅)₄] (M = M' = Pd or Pt; M = Pt, M' = Pd) with bridging pentafluorophenyl groups, 169–72
Palladium complexes with the tripodal phosphine tris(2-diphenylphosphinoethoxy)amine. Synthesis and structure of trigonal, tetrahedral, trigonal bipyramidal, and square planar complexes, 211–6

- Metallacyclic complexes. Part 2. Synthesis of metallathietane-3,3-dioxide complexes of platinum(ii) and palladium(ii); the crystal structures of [Pt{CH(COPh)S(O)₂CH(COPh)}₂(PPPh₃)₂·2CH₂Cl₂ and [Pd{CH(COPh)S(O)₂CH(COPh)}₂(PMcPh₃)₂·2CH₂Cl₂·H₂O, 259–66

- The reaction of hydroxo-species of palladium(ii) with bidentate nucleophiles, 375–8

- Preparation and interconversion of dimeric di-μ-hydroxo and tri-μ-hydroxo complexes of platinum(ii) and palladium(ii) with 2,2'-bipyridine and 1,10-phenanthroline, 403–12

- Synthesis of sulphimide complexes of palladium and platinum. Crystal and molecular structure of *cis*-dichloro[S,S-dimethyl-N-(2-pyrimidinyl)sulphimide](triphenylphosphine)palladium(ii), 497–504

- Crystallographic and nuclear magnetic resonance studies of carbonyl(η^3 -methylallyl)(trichlorostannyl) complexes of platinum(ii) and palladium(ii), 615–22

- Tetrathiafulvalenium salts of planar Pt^{II}, Pd^{II}, and Cu^{III} 1,2-dithio-oxalato-S,S anions. Synthesis, chemistry, and molecular structures of bis(tetrathiafulvalenium) bis(1,2-dithio-oxalato-S,S) palladate(ii), [ttf]₂[Pd(S₂C₂O₂)₂], and of bis(tetrathiafulvalenium)tetrathiafulvalene bis(1,2-dithio-oxalato-S,S) platinate(ii), [ttf]₂[Pt(S₂C₂O₂)₂], 719–28

- Preparation of (chloromethyl)palladium(ii) derivatives from complexes of palladium dichloride by reaction with diazomethane or bis(chloromethyl)mercury, 761–6

- Tri-aza analogue of an acetylacetato-complex formed by nucleophilic attack of a co-ordinated nitrile by amidines. X-Ray crystal structure of [Pt(HNC(Ph)NC(Ph)NH)₂]₂· $\frac{1}{2}$ PtMe, 837–42

- Platinum metal complexes of potentially chelating alkene-thioether and alkene-selenoether ligands: synthesis and dynamic nuclear magnetic resonance study of [MX₂{MeE(CH₂)_nCH=CH₂}₂] (M = Pt or Pd; X = Cl, Br, or I; E = S or Se; n = 2 or 3) and the X-ray structure of *cis*-dibromo(2-thia-6-heptene)platinum(ii),

[PtBr₂{MeS(CH₂)₃CH=CH₂}₂] 985–90

- Iridium(iii) hydrido complexes with 2-pyridinaldazines. Crystal structure of [IrH₂(mpaa)(PPh₃)₂]SbF₆. Synthesis and characterization of homo- and hetero-nuclear complexes, 1075–82

Crystal structure of the new low-dimensional metal Cs_{0.82}[Pd{S₂C₂(CN)₂}₂]·0.5H₂O 1101–4

- Trinuclear pyrazolate complexes. Crystal structure of [(OC)₂Rh(μ-pz)₂Pd(μ-pz)₂Rh(CO)₂], 1133–8

- New metal-sulphur-nitrogen compounds from reactions in liquid ammonia. The X-ray structures of *trans*-bis(acetophenone dimethylhydrazone-*N*³)dichloropalladium(ii) and [di(azathien)-1-yl-S¹N⁴]₂[2-(hydrazonoethyl)phenyl]palladium(ii), 1179–86

- Chelate-stabilised alcohol- and alkoxo-palladium(ii) complexes derived from the ligands Ph₂PCH₂CR₂OH (R = Me or H), 1193–6

- Photo-oxidation of alcohols by hexachlorometalate(iv) ions (M = Pt, Pd, or Ir): spin trapping and matrix isolation electron spin resonances studies, 1237–40

- Mode of binding of dimethyl sulphoxide in platinum(ii) and palladium(ii) cationic complexes containing a chelated diamine. X-Ray crystal structures of (2,2'-bipyridyl)chloro(dimethyl sulphoxide-O)palladium(ii) tetrafluoroborate and chloro(dimethyl sulphoxide-S)(ethylenediamine)palladium(ii) perchlorate, 1265–72

- Synthesis, characterization, and thermolytic behaviour of η^3 -allyl(methyl)(tertiary phosphine)palladium(ii), and X-ray structure of the cluster [Pd₄(μ-C₃H₅)₂(μ-Cl)₂(μ-dppm)₂]₂4C₄H₈O (dppm = Ph₂PCH₂PPh₃), 1519–26

- Reactions of cyclopalladated compounds. Part 21. Various examples of sulphur-assisted intramolecular palladation of aryl and alkyl groups, 1715–20

- Closely related macrocyclic and acyclic tridentate, pyridine derivatives, containing sulphur, and their complexes. Crystal structures of {dichloro-3,10-dithia-16-azabicyclo[10.3.1]hexadeca-1(16),12,14-triene}copper(ii) and [2,6-bis(ethylthiomethyl)pyridine]dichlorocopper(ii), 1739–44

- Low-valent isocyanide complexes and clusters of palladium and platinum crystal structure of [Pt{C=(NR)N(R)C(=NR)}(RNC)₂] (R = 2,6-Me₂C₆H₃), 2161–6

- Preparation and properties of [M(S₂N₂H)Cl(L)] (M = Pt, L = PMe₂Ph or C₆H₄; M = Pd, L = PMe₂Ph), [Pt(S₃N)Cl(PMe₂Ph)], and corresponding ¹⁵N-enriched complexes, 2179–82

- Platinum metal complexes of potentially chelating alkene thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of [MX₂E{(CH₂)_nCR=CR₂}₂] (M = Pt or Pd; X = Cl, Br, or I; E = S or Se; n = 2 or 3; R = H or Me) and the X-ray crystal structure of *cis*-di-iodo(5-thianona-1,8-diene)platinum(ii), [PII₂{S[(CH₂)₂CH=CH₂}₂]₂], 2315–22

- Solid-state high-resolution carbon-13 magnetic resonance spectra of dithioacetate complexes of platinum, palladium, and nickel, 2457–60

PENICILLIN

- The copper(ii) promoted hydrolysis of benzylpenicillin. Evidence for the participation of a Cu-OH species in the hydrolysis of the β-lactam ring, 197–202

PENTACHLOROPHENYL

- Synthesis and reactivity of binuclear perhalogenophenyl derivatives of platinum(i). Molecular structure of [Pt₂(C₆F₅)₂(CO)₂(PPh₃)₂]·0.4CHCl₃, 1583–8

PENTAFLUOROPHENYL

- (Diphenylphosphinomethyl)diphenylphosphine sulphide (Ph₂PCH₂PPh₂S) and its methanide anion Ph₂PCHPPh₂S⁻ as ligands in organogold chemistry. X-Ray crystal structure of *cis*-[Au(C₆F₅)₂(PPPh₂CHPPh₂S)], 73–8

- Synthesis and reactivity of binuclear homo- or hetero-metallic complexes [NBu₄]₂[MM'(μ-C₆F₅)₂(C₆F₅)₄] (M = M' = Pd or Pt; M = Pt, M' = Pd) with bridging pentafluorophenyl groups, 169–72

- Synthesis and reactivity of trinuclear complexes of platinum containing the single bridging ligand SC₄H₈. Molecular structure of [NBu₄]₂[*trans*-PtCl₂{(μ-SC₄H₈)Pt(C₆F₅)₃}₂], 1011–6

- Synthesis and reactivity of binuclear perhalogenophenyl derivatives of platinum(i). Molecular structure of [Pt₂(C₆F₅)₂(CO)₂(PPh₃)₂]·0.4CHCl₃, 1583–8

PENTAFLUOROTHIOPHENOLATE

- Monocyclopentadienyl pentafluorothiophenolate complexes of molybdenum and their interactions with thallium(i) ions;

PENTAFLUOROTHIOPHENOLATE (contd)

dynamic nuclear magnetic resonance studies and crystal and molecular structures of $[\text{TiMo}(\text{SC}_6\text{F}_5)_2\text{L}_2(\text{cp})]$ ($\text{L} = \text{CO}$ or SC_6F_5 ; $\text{cp} = \eta^5\text{-C}_5\text{H}_5$), 991–1002

PENTAMETHYLCYCLOCYCLOPENTADIENYL

Reactivity of cationic molybdenum(II) complexes. Part 3. Synthesis of methoxide and carbomethoxide derivatives of η -pentamethylcyclopentadienyl-molybdenum(II) and the crystal structure determination of the oxidation product *trans*- $[\{\text{Mo}(\eta\text{-C}_5\text{Me}_5)(\text{O})_2\}_2(\mu\text{-O})]$, 155–60

Preparation and structural studies of diphenylthiophosphinito rhodium(III) complexes. X-Ray structure analysis of $[(\text{C}_5\text{Me}_5)_2\text{Rh}]_2(\mu\text{-Cl})_2(\mu\text{-Ph}_2\text{PS})[\text{BPh}_4]$, 1987–92

PENTAOXACYCLOCYCLOPENTADECANE

Stereochemical role of lone pairs in main-group elements. Part 4. The crystal structure at 120 K of bis(1,4,7,10,13-pentaoxacyclopentadecane)tin(II) bis[trichlorostannate(IV)], confirming the sterically inactive tin lone pair, 2155–60

PENTAPHENYLCYCLOCYCLOPENTADIENYL

Reduction–oxidation properties of organotransition-metal complexes. Part 29. Pentaphenylcyclopentadienyl complexes of ruthenium, 283–8

PEPTIDE

The unusual co-ordination ability of vasopressin-like peptides: potentiometric and spectroscopic studies of some copper(II) and nickel(II) complexes, 173–8

Complexes of aminophosphonates. Part 3. Copper(II) complexes of several monophosphono and diphosphono dipeptides, 1053–8

Composition, stability, and lability of copper(II) dipeptide complexes, 1293–8

Electron spin resonance study of copper(II) complexes of X-glycine and glycyl-X type dipeptides, and related tripeptides. Variation of co-ordination modes with ligand excess and pH in fluid and frozen aqueous solutions, 1925–32

PEROVSKITE

Preparation of the ordered perovskite $(\text{SrLa})(\text{MgMo})\text{O}_6$ and its physicochemical characterization, 2335–8

PEROXIDE

Studies on transition metal peroxy complexes. Part 8. The nature of peroxyomolybdates and peroxytungstates in aqueous solution, 1203–8

Oxidative addition of dibenzoyl peroxide on metallic copper. The crystal structure of $[(\text{Cu}(\text{C}_6\text{H}_5\text{CO}_2)_2(\text{py}))_2]$, 2331–4

PEROXOBORATE

Nature of peroxyborate species in aqueous solution: a study by boron-11 nuclear magnetic resonance and Raman spectroscopy, 1651–6

PEROXODISULPHATE

Kinetic studies of the oxidation reaction of arsenic(III) to arsenic(V) by peroxodisulphate ion in aqueous alkaline media, 357–60

PHENANTHROLINE

Preparation and interconversion of dimeric di- μ -hydroxo and tri- μ -hydroxo complexes of platinum(II) and palladium(II) with 2,2'-bipyridine and 1,10-phenanthroline, 403–12

Reactivity studies on the hydrolysis of iron(II) complexes of substituted phenanthrolines in dioxane–water solvents, 1105–8

Mode of binding of dimethyl sulphoxide in platinum(II) and palladium(II) cationic complexes containing a chelated diamine. X-Ray crystal structures of (2,2'-bipyridyl)chloro(dimethyl sulphoxide-O)palladium(II) tetrafluoroborate and chloro(dimethyl sulphoxide-S)(ethylenediamine)palladium(II) perchlorate, 1265–72

PHENOL

Reactions of heteropolyanions in non-polar solvents. Part 3. Activation of dioxygen by manganese(II) centres in polytungstates. Oxidation of hindered phenols, 1483–90

Crystal structures and magnetic properties of bi- and tetra-nuclear copper(II) complexes of 2,6-disformyl-4-methylphenol di(benzoylhydrazone), 2083–8

PHENOLATE

Kinetics and mechanism of the reaction between 4-t-butylphenolate anion and tetrahydroxoargentate(III) in aqueous alkaline media, 217–22

Low-lying electronic states in μ -phenolato copper(II) binuclear compounds with hydroxo, ethanolato, azido, and cyanato exogenous bridges: crystal structures, magnetic properties, and interpretations, 1117–26

Proton electron nuclear double resonance study of oxovanadium(IV) complexes of *o*-diphenolic ligands, 1289–92

PHENOXIDE

Phenoxy and diphenoxo complexes of titanium(III) and vanadium(IV): synthesis and X-ray structure, 1793–8

PHENYL

Syntheses and dynamic stereochemistry of methyl- and phenyl-carbonyl heterobinuclear metal–metal bonded complexes, 379–82

Phenyltin trichloroacetate complexes: preparations and structures, 1589–98

Crystal and molecular structure of a lattice adduct, $\text{HgPh}_2\cdot 2[\text{Hg}(\text{SCN})_2(\text{PPh}_3)_2]$: an unusual case of two- and four-co-ordinate mercury in the same unit cell, 2339–42

PHOSPHANE

Chemistry of $[(\text{Cr}(\text{cp})(\text{CO})_3)_2]$ ($\text{cp} = \eta^5\text{-C}_5\text{H}_5$). Reaction with elemental P_4 . A high yield synthesis and crystal structures of $[(\text{Cr}(\text{cp})(\text{CO})_2)_2(\mu\text{-}\eta^2\text{-P}_2)]$ and $[\text{Cr}(\text{cp})(\text{CO})_2(\eta^3\text{-P}_3)]$, 1951–6

PHOSPHANYLIUM

Synthesis, structure, and chemical reactivity of a stable C_5Me_5 -substituted phosphanylium ion: (pentamethylcyclopentadienyl)(*t*-butylamino)phosphanylium tetrachloroaluminate, 693–700

PHOSPHATE

An investigation of halide abstraction by arsenic(III): reactions of AsCl_3 with MCl_2 ($\text{M} = \text{Mg}$ or Zn) and TiCl_4 in donor solvents (MeCN or POCl_3). Formation and crystal structure of the dimeric titanium(IV) compound $[\text{Ti}_2\text{Cl}_6(\text{POCl}_3)_2(\text{PO}_2\text{Cl}_2)_2]$, 57–60

Potentiometric and electron spin resonance investigation on the interactions between adenosine 5'-triphosphate (atp) and triphosphate ligands with vanadyl ion. Synthesis and characterization of solid vanadyl-atp compounds from acidic aqueous solution, 575–80

Lamellar inorganic ion exchangers. Hydrogen-lithium ion exchange in γ -titanium phosphate, 829–36

Formation of chromium interlayers in α -zirconium phosphate, 895–900

Zirconium phosphate ester interchange reactions, 1617–24

Neutron diffraction study of the cation distributions in the systems $\text{Fe}_{7-x}\text{M}_x(\text{PO}_4)_6$ ($\text{M} = \text{Mn}$ or Co), 1765–70

Lamellar inorganic ion exchangers. H^+/Cs^+ ion exchange in γ -titanium phosphate, 1825–30

PHOSPHIDE

Synthesis of diphenylphosphido-bridged manganese-gold complexes. Crystal structures of $[\text{N}(\text{PPh}_3)_2][\text{Au}_2\text{Mn}(\mu\text{-PPh}_2)_2(\text{CO}_4)_2]$ and $[\text{Au}_2\text{Mn}_2(\mu\text{-PPh}_2)_4(\text{CO})_8]$, 639–44

Gold(I) complexes derived from secondary phosphines: $[(\text{Au}(\mu\text{-PR}_2)_n], [(\text{AuBr})_2(\mu\text{-PPh}_2)]^+, [\text{AuX}(\text{PPh}_2)_n]$, and $[(\text{Au}(\text{PPh}_2)_n)]^+$. Crystal structure of $[\text{AuBr}(\text{PPh}_2)_2]$, 907–14

Chemistry of phosphido-bridged dimolybdenum complexes. Part 3.

Reinvestigation of the reaction between $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\text{CO})_6]$ and P_2Ph_4 : X-ray structures of $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\text{CO})_6]$, $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2(\mu\text{-CO})]$ and *trans*- $[\text{Mo}_2(\eta\text{-C}_5\text{H}_5)_2(\mu\text{-PPh}_2)_2\text{O}(\text{CO})]$, 1555–64

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 89. Tetra- and penta-nuclear tungsten-rhodium complexes: crystal structures of $[\text{W}_3\text{Rh}_2(\mu\text{-CO})_2(\mu\text{-CMe})_2\text{C}(\text{Me})\text{C}(\text{O})](\mu\text{-PPh}_2)_2(\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3$ and $[\text{W}_3\text{Rh}_2(\mu\text{-CO})_3(\mu\text{-CMe})_2\text{C}(\text{Me})\text{PPh}_2](\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3$, 1855–64

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 90. Synthesis of the complexes $[\text{W}_2\text{M}_2(\mu\text{-CMe})(\mu_3\text{-CMe})(\mu\text{-PPh}_2)_2(\text{CO})_3(\mu\text{-C}_5\text{H}_5)]$ ($\text{M} = \text{Rh}$ or Ir) and related tetra- and penta-nuclear metal compounds; interconversion of ketenyl and λ^5 -phospha-alkyne ligands bridging tungsten-rhodium bonds, 1865–70

Cluster activation *via* cyclometallation. Formation of the cyclometallated triosmium carbonyl cluster

$[\text{Os}_3\text{H}_2(\text{CO})_9\text{P}(\text{C}_6\text{H}_4)\text{Ph}]$ and its reactions with phosphorus donor ligands. The X-ray crystal structures of

$[\text{Os}_3\text{H}_2(\text{CO})_9\text{P}(\text{C}_6\text{H}_4)\text{Ph}]$ and $[\text{Os}_3\text{H}(\text{CO})_9\text{P}(\text{OMe})_3](\text{PPh}_2)]$, 2023–32

Reactions of the cyclometallated triosmium cluster

$[\text{Os}_3\text{H}_2(\text{CO})_9\text{P}(\text{C}_6\text{H}_4)\text{Ph}]$ with Brønsted acids and dihydrogen: synthesis and X-ray crystal structure of $[\text{Os}_3\text{H}_2(\text{CF}_3\text{CO}_2)(\text{CO})_9(\mu\text{-PPh}_2)]$, 2033–8

PHOSPHINE

(Diphenylphosphinomethyl)diphenylphosphine sulphide ($\text{Ph}_2\text{PCH}_2\text{PPh}_2\text{S}$) and its methanide anion $\text{Ph}_2\text{PCH}(\text{PPh}_2)\text{S}^-$ as

PHOSPHINE (contd)

ligands in organogold chemistry. X-Ray crystal structure of *cis*-
 $[\text{Au}(\text{C}_6\text{F}_5)_2(\text{PPh}_2\text{CHPPh}_2\text{S})]$, 73–8
 Palladium complexes with the tripodal phosphine tris(2-diphenylphosphinoethyl)amine. Synthesis and structure of trigonal tetrahedral, trigonal bipyramidal, and square planar complexes, 211–6

Metallacyclic complexes. Part 2. Synthesis of metallathietane-3,3-dioxide complexes of platinum(ii) and palladium(ii); the crystal structures of $[\text{Pt}(\text{CH}(\text{COPh})\text{S}(\text{O})_2\text{CH}(\text{COPh})}$
 $(\text{PPh}_3)_2\cdot 2\text{CH}_2\text{Cl}_2$ and $[\text{Pd}(\text{CH}(\text{COPh})\text{S}(\text{O})_2\text{CH}(\text{COPh})}$
 $(\text{PMcPh}_2)_2\cdot \text{CH}_2\text{Cl}_2\cdot \text{H}_2\text{O}$, 259–66

Reactions of the bis(alkyne) complexes $[\text{WI}_2(\text{CO})(\text{NCMe})(\eta^2-\text{RC}_2\text{R})_2]$ ($\text{R} = \text{Me}$ or Ph) with mono- and bi-dentate phosphine-donor ligands and the X-ray crystal structure of $[\text{WI}_2(\text{CO})(\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2)(\eta^2-\text{MeC}_2\text{Me})]$, 295–302

Chemistry of metallacyclobutanones (η^3 -oxodimethylenemethane complexes). Part 5. Ligand substitution reactions of metallacyclobutan-3-ones of platinum(ii) and palladium(ii) with tertiary phosphines and t-butyl isocyanide; crystal structures of $[\text{Pt}(\text{CH}(\text{CO}_2\text{Me})\text{COCH}(\text{CO}_2\text{Me})\{\text{CNBu}'\}(\text{AsPh}_3)]$ and $[\text{Pt}(\text{CH}(\text{CO}_2\text{Me})\text{CO}\bar{\text{C}}(\text{CO}_2\text{Me})\text{C}(\text{NHBU}')\{\text{CNBu}'\}_2]$, 345–56

Thiolate complexes of osmium(iii) with tertiary phosphines and their dicarbonyl derivatives of osmium(ii); X-ray crystal structure of $[\text{trans,trans,trans-}[\text{Os}(\text{SC}_6\text{F}_5)_2(\text{CO})_2(\text{PEt}_2\text{Ph})_2]]$, 419–24
 Gold-197 Mössbauer spectra and the bonding of some gold–gold and gold–platinum clusters, 539–44

Reactions of hydride complexes of ruthenium(ii) with alkynes, 589–94
 A kinetic study of the transmission of ligand effects across the quadruple bond in $[\text{Re}_2\text{Cl}_8]^{2-}$, 767–72

Gold(i) complexes derived from secondary phosphines: $[\{\text{Au}(\mu-\text{PR}_2)_n\}_2]$, $[(\text{AuBr})_2(\mu-\text{PPH}_2)]^-$, $[\text{AuX}(\text{PhR}_2)]$, and $[(\text{Au}(\text{PhR}_2)_n)]^\pm$. Crystal structure of $[\text{AuBr}_2(\text{PhPPh}_2)]$, 907–14

Synthesis of a new P_2N_2 ligand N,N' -bis[2-(diphenylphosphino)phenyl]propane-1,3-diamine, H_2L^2 , and some of its complexes with elements of the nickel triad and rhodium; X-ray structure analyses of the neutral complex $[\text{NiL}^2]$ and the *trans*-spanned $[\text{Rh}(\text{CO})\text{Cl}(\text{H}_2\text{L}^2)]$, 1067–74

Synthesis of hydridotrifluoromethyl complexes of platinum(ii). A spectroscopic investigation of the *trans* and *cis* influence of ligands in hydridoplatinum(ii) compounds, 1149–60

Binuclear copper(i) complexes containing two or three 2-(diphenylphosphino)pyridine (dppy) bridging ligands; X-ray crystal structure of $[\text{Cu}_2(\mu\text{-dppy})_3(\text{MeCN})][\text{BF}_4]_2$ and variable-temperature ^1H and $^{31}\text{P}-\{^1\text{H}\}$ nuclear magnetic resonance studies, 1499–506

Synthesis, characterization, and thermolytic behaviour of η^3 -allyl(methyl)(tertiary phosphine)palladium(ii), and X-ray structure of the cluster $[\text{Pd}_4(\mu\text{-C}_3\text{H}_5)_2(\mu\text{-Cl})_2(\mu\text{-dppm})_2]\text{4C}_4\text{H}_8\text{O}$ (dppm = $\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2$), 1519–26

Tertiary phosphine complexes of triply bonded ditungsten bis(t-butylamide) tetrachloride. The X-ray crystal structures of $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMe}_3)_2]$, $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PEt}_3)_2]$, and *cis*- $[\text{W}_2\text{Cl}_4(\text{NHBu}')_2(\text{PMe}_2\text{Ph})_2]$, 1537–42

Substitution reactions on sterically hindered square-planar *trans*- $[\text{NiBr}(\text{R})(\text{PR}')_2]$ ($\text{R} = \text{aryl}$) complexes. Effects of the substituents of the aryl ligand, 1669–74

Bi-, tetra-, and poly-nuclear complexes of gold(i) and silver(i) with bis(diphenylphosphino)-methanide or -amide acting as six-electron donors, 1883–6

Preparation, characterisation, and reactions with phosphite donor ligands of the new cationic monoalkyne complexes $[\text{WI}(\text{CO})(\text{NCMe})(\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2)(\eta^2-\text{RC}_2\text{R})][\text{BF}_4]$ ($n = 1$ –6; $\text{R} = \text{Me}$ or Ph). X-Ray crystal structure of $[\text{WI}(\text{CO})\text{-}(\text{POPr}')_3\text{-}(\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2)(\eta^2-\text{MeC}_2\text{Me})][\text{BF}_4]_{0.5}[\text{OH}]_{0.5}\cdot 0.5\text{H}_2\text{O}$, 1903–10

Oxo transfer reactions. Mechanistic studies on the oxidation of triphenylphosphine by *trans*- $[\text{Ru}^{\text{IV}}\text{L}(\text{O})_2]^{2+}$ ($\text{L} = 1,4,8,11$ -tetramethyl-1,4,8,11-tetra-azacyclotetradecane), 2065–8

Studies of the oxidative addition of some substrates containing carbon heteroatom bonds to some tungsten and platinum complexes, 2099–108

Mesitylgold(i) complexes. Crystal structures of $\text{NEt}_4[\text{Au}(\text{mes})\text{Cl}]$ and $[\text{N}(\text{PPh}_3)_2][(\text{mes})\text{Au}(\text{CN})\text{Au}(\text{mes})]$, 2127–32

Reactions of chloro(triphenylphosphine)gold(i) and $[\mu\text{-1,2-bis(diphenylphosphine)ethane}]$ -bis[bromogold(i)] with oxopurine

bases. Molecular and crystal structure and bonding of (theophyllinato)(triphenylphosphine)gold(i), 2323–30

PHOSPHINE OXIDE

The structure of $\text{PO}(\text{OPF}_2)_3$ in the gaseous and crystalline phases, 223–8

Formation of mixed-ligand complexes of cadmium(ii) halide with tri-n-octylphosphine oxide and pyridine or substituted pyridines in 1,2-dichloroethane, 1259–64

PHOSPHINOALCOHOL

Chelate-stabilised primary, secondary, and tertiary alcohol and alkoxo platinum(ii) complexes. X-Ray crystal structure of $[\text{Pt}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{O})_2]\text{H}_2\text{O}$, 139–44

PHOSPHINOALKOXIDE

Chelate-stabilised primary, secondary, and tertiary alcohol and alkoxo platinum(ii) complexes. X-Ray crystal structure of $[\text{Pt}(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{O})_2]\text{H}_2\text{O}$, 139–44

PHOSPHINOALKYL

Synthesis of α , γ -phosphorus functionalized alkyl lithium species; X-ray structures of $[\{\text{Li}(\text{L})(\text{CH}_2\text{PMeR})_3\}_2]$ [$\text{L} = \text{N}, \text{N}, \text{N}'$ -tetramethylethylenediamine (tmen), $\text{R} = \text{Me}$ or Ph ; $\text{L} = (-)\text{sparteine}$, $\text{R} = \text{Ph}$] and $[\text{Li}(\text{tmen})\{\text{CH}(\text{SiMe}_3)\text{C}_6\text{H}_4\text{PPh}_2\}_2]$, 105–14

PHOSPHINOMETHYL

Chelate-stabilised tetrahydrofuran complexes of platinum(ii).

Crystal structure of $[\text{PtCl}(\text{Ph}_2\text{PCH}_2\text{C}_4\text{H}_7\text{O})\text{-}(\text{Ph}_2\text{PCH}_2\text{C}_4\text{H}_7\text{O})][\text{PF}_6]$, 2069–72

PHOSPHITE

Reactions of co-ordinated ligands. Part 44. The reactivity of the carbene complex $[\text{Mo}(\equiv\text{CCH}_2\text{Bu})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$ towards protonic acids, diphenylphosphine, and the π -acid tetracyanoethylene; formation and crystal structures of $[\text{MoH}(\equiv\text{CCH}_2\text{Bu})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)][\text{BF}_4]$, $[\text{Mo}\{\sigma\text{-OC(O)CF}_3\}(\eta^2\text{-HOC(O)CF}_3)\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, and $[\text{Mo}(\equiv\text{CCH}_2\text{Bu})\{\eta^2\text{-C}_2(\text{CN})_4\}\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, 511–8

Solution structure and dynamic behaviour of two isomers of $[\text{Fe}_3(\text{CO})_9(\text{POR}_3)_3]$ ($\text{R} = \text{Me}$, Et, or Ph) derivatives, 1277–82

Reactions of co-ordinated ligands. Part 45. Ligand displacement and oxidative reactions of the carbene complex

$[\text{Mo}(\equiv\text{CCH}_2\text{Bu})\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$; crystal and molecular structures of $[\text{Mo}\{\eta^2\text{-RN=C}\cdots\text{C}(\text{CH}_2\text{Bu})\cdots\text{C}=\text{NR}\}(\text{CNR})_2(\eta\text{-C}_5\text{H}_5)]$ ($\text{R} = 2,6\text{-Me}_2\text{C}_6\text{H}_3$), $[\text{Mo}(\text{C}=\text{CHBu})\text{I}\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, and $[\text{Mo}\{\text{C}(\text{CH}_2\text{Bu})\text{P}(\text{O})(\text{OMe})_2\}\{\text{P}(\text{OMe})_3\}_2(\eta\text{-C}_5\text{H}_5)]$, 1321–32

The mechanisms of fluxionality of $[\text{Fe}_3(\text{CO})_{12-n}\{\text{P}(\text{OR})_3\}_n]$ ($\text{R} = \text{Me}$, $n = 0$ –3; $\text{R} = \text{Pr}$, $n = 2$ or 3) and the X-ray structure of $[\text{Fe}_3(\text{CO})_{10}\{\text{P}(\text{OMe})_3\}_2]$, 1831–44

Preparation, characterisation, and reactions with phosphite donor ligands of the new cationic monoalkyne complexes

$[\text{WI}(\text{CO})(\text{NCMe})(\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2)(\eta^2\text{-RC}_2\text{R})][\text{BF}_4]$ ($n = 1$ –6; $\text{R} = \text{Me}$ or Ph). X-Ray crystal structure of $[\text{WI}(\text{CO})\text{-}(\text{POPr}')_3\text{-}(\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2)(\eta^2\text{-MeC}_2\text{Me})][\text{BF}_4]_{0.5}[\text{OH}]_{0.5}\cdot 0.5\text{H}_2\text{O}$, 1903–10

PHOSPHONATE

Complexes of aminophosphonates. Part 3. Copper(ii) complexes of several monophosphono and diphosphono dipeptides, 1053–8

Metal complexes of α -hydroxyiminophosphonic acid derivatives.

Preparation and crystal structures of calcium and cadmium complexes of methyl (*E*)- α -hydroxyiminobenzylphosphonate, 1911–4

PHOSPHONIC ACID

Organosulphur phosphorus acid compounds. Part 1. *m*-Sulphophenylphosphonic acid, 1819–24

PHOSPHORANE

The molecular structures of gaseous tetrafluorohydridophosphorane, HPF_4 , and trifluorodihydridophosphorane, H_2PF_3 , as determined by electron diffraction, 545–50

PHOSPHORUS

The structure of $\text{PO}(\text{OPF}_2)_3$ in the gaseous and crystalline phases, 223–8

The molecular structures of gaseous

tetrafluorohydridophosphorane, HPF_4 , and trifluorodihydridophosphorane, H_2PF_3 , as determined by electron diffraction, 545–50

Synthesis, structure, and chemical reactivity of a stable C_5Me_5 -substituted phosphanylium ion: (pentamethylcyclo-pentadienyl)-(t-butylamino)phosphanylium tetrachloroaluminate, 693–700

PLATINUM (contd)

- Synthesis of hydridotrifluoromethyl complexes of platinum(II). A spectroscopic investigation of the *trans* and *cis* influence of ligands in hydridoplatinum(II) compounds, 1149–60
- New metal-sulphur-nitrogen compounds from reactions in liquid ammonia. The X-ray structures of *trans*-bis(acetophenone dimethylhydrazone-*N'*²)dichloropalladium(II) and [di(azathien)-1-yl-*S'*²]²[2-(hydrazonoethyl)phenyl]palladium(II), 1179–86
- Photo-oxidation of alcohols by hexachlorometalate(IV) ions (M = Pt, Pd, or Ir): spin trapping and matrix isolation electron spin resonances studies, 1237–40
- Mode of binding of dimethylsulphoxide in platinum(II) and palladium(II) cationic complexes containing a chelated diamine. X-Ray crystal structures of (2,2'-bipyridyl)chloro(dimethyl sulphoxide-*O*)palladium(II) tetrafluoroborate and chloro(dimethyl sulphoxide-*S*)(ethylenediamine)palladium(II) perchlorate, 1265–72
- Nucleophilic (Michael) additions to (Ph_2P)₂C=CH₂ when chelated to platinum(II) dichloride or di-iodide, 1527–30
- Synthesis and reactivity of binuclear prehalogenophenyl derivatives of platinum(II). Molecular structure of [$\text{Pt}_2(\text{C}_6\text{F}_5)_2(\text{CO})_2(\text{PPPh}_3)_2\cdot 0.4\text{CHCl}_3$, 1583–8]
- Metalaheteroborane chemistry. Part 5. Synthesis of *cis*-[2,2-(PR_3)₂-1,2-SePtB₁₀H₁₀] (R_3 = Et₃, Buⁿ₃, or Me₂Ph) and *arachno*-[9,9-(PM₂Ph)₂-6,9-SePtB₈H₁₀] and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of *cis*-[2,2-(PEt₃)₂-1,2-SePtB₁₀H₁₀] and *arachno*-[9,9-(PM₂Ph)₂-6,9-SePtB₈H₁₀], 1657–68
- Syntheses, crystal structures, and Raman and resonance Raman spectroscopy of platinum(II), platinum(IV), and halogen-bridged chain platinum(II,IV) complexes involving the terdentate ligand 3-methyl-3-azapentane-1,5-diamine (4Me-dien), 1689–96
- Base-promoted cleavage of a P-C bond in [$\text{PtCl}_2(\text{Ph}_2\text{PCh}_2\text{PPh}_2\text{-}P\text{P}')

Chelate-stabilised tetrahydrofuran complexes of platinum(II). Crystal structure of [$\text{PtCl}(\text{Ph}_2\text{PCH}_2\text{C}_4\text{H}_7\text{O})\cdot (\text{Ph}_2\text{PCH}_2\text{C}_4\text{H}_7\text{O})\text{[PF}_6\text{]}$, 2069–72

Studies of the oxidative addition of some substrates containing carbon-heteroatom bonds to some tungsten and platinum complexes, 2099–108

Mechanism and energetics of reductive elimination of ethane from some tetramethylplatinum(IV) complexes, 2121–6

Low-valent isocyanide complexes and clusters of palladium and platinum. Crystal structure of [$\text{Pt}\{\text{C}(=\text{NR})\text{N}(\text{R})\text{C}(=\text{NR})\}\{\text{RNC}\}_2$] (R = 2,6-Me₂C₆H₃), 2161–6

Preparation and properties of [$\text{M}(\text{S}_2\text{N}_2\text{H})\text{Cl}(\text{L})$] (M = Pt, L = PMe₂Ph or C₂H₄; M = Pd, L = PMe₂Ph), [$\text{Pt}(\text{S}_2\text{N})\text{Cl}(\text{PMe}_2\text{Ph})$], and corresponding ¹⁵N-enriched complexes, 2179–82

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 93. Synthesis of heteropolynuclear metal compounds with chains of seven to eleven metal atoms; crystal structure of [$\text{Mo}_2\text{W}_3\text{Pt}_6(\mu_3\text{-CMe})_3(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\text{CO})_{10}(\text{PMMe}_2\text{Ph})_4(\eta\text{-C}_5\text{H}_5)_5$], 2269–80

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 95. Synthesis and crystal structures of [$\text{W}_2\text{RuPt}(\mu\text{-CC}_6\text{H}_4\text{Me}-4)(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)(\text{CO})_6(\text{PMMe}_2\text{Ph})(\eta\text{-C}_5\text{H}_5)_2$] and [$\text{W}_2\text{Ru}_2\text{Pt}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\mu\text{-CO})_2(\text{CO})_9(\eta\text{-C}_5\text{H}_5)_2$], 2289–96

Platinum metal complexes of potentially chelating alkene thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of [$\text{MX}_2\{\text{E}[(\text{CH}_2)\text{CR=CR}_2]\}_2$] (M = Pt or Pd; X = Cl, Br, or I; E = S or Se; n = 2 or 3; R = H or Me) and the X-ray crystal structure of *cis*-di-iodo(5-thianona-1,8-diene)platinum(II), [$\text{PtI}_2\{\text{S}[(\text{CH}_2)_2\text{CH=CH}_2]\}_2$], 2315–22

Solid-state high-resolution carbon-13 magnetic resonance spectra of dithioacetate complexes of platinum, palladium, and nickel, 2457–60$

POLYMER

- Synthesis, X-ray characterization, and Q-band electron spin resonance studies of heterometallic, polymeric compounds formed by yttrium and lanthanide(III) nitrates with HgL₂ (L = C₄H₆NO, the anion of 2-pyrrolidone); crystal structures of [$\{\text{Hg}_2\text{EuL}_4(\text{NO}_3)_3\}_n$] and [$\{\text{Hg}_3\text{Tb}_2\text{L}_6(\text{NO}_3)_6\}_n$], 1439–48

POLYMERIZATION

- Preparation and characterisation of some nickel(II) tetra-aza macrocyclic complexes bearing pendant polymerisable groups as part of the ligand superstructure, 1599–608

PORPHYRIN

- Photocatalytic reactions in the 2,3,7,8,12,13,17,18-octaethylporphyrinatoiron(III)-ethanol-carbon tetrachloride system, 1197–202
- Effect of axially bound anions on the electroreduction of tin(IV) porphyrins in tetrahydrofuran, 1531–6

PORPHYRINATE

- Electronic spectral study of the aqua \rightleftharpoons hydroxo equilibrium of model iron(III) haems encapsulated in aqueous detergent micelles, 1003–6

POTENTIAL ENERGY

- On the relationship between crystallographic and spectroscopic evidence of dynamic processes in the solid state: the cases of *cis*- and *trans*-[Fe($\eta\text{-C}_5\text{H}_5$)₂(CO)₄], 1721–6

PURINE

- Reactions of chloro(triphenylphosphine)gold(I) and [μ -1,2-bis(diphenylphosphine)ethane]-bis[bromogold(I)] with oxopurine bases. Molecular and crystal structure and bonding of (theophyllinato)(triphenylphosphine)gold(I), 2323–30

PYRAZOLE

- Synthesis and reactivity of binuclear pyrazolate-bridged ruthenium(II) complexes. Crystal structures of bis[μ -(3,5-dimethylpyrazolato-*NN'*)tricarbonylruthenium(I)] ($Ru-Ru$) and bis[μ -(3,5-dimethylpyrazolato-*NN'*)- μ -iodo-bis[tricarbonylruthenium(II)] tri-iodide, 1093–100

- Trinuclear pyrazolate complexes. Crystal structure of [(OC)₂Rh($\mu\text{-pz}$)₂Pd($\mu\text{-pz}$)₂Rh(CO)₂], 1133–8

- A planar, binucleating, pyrazole derivative ligand. Crystal structure of bis[μ -(3,5-bis(2'-pyridyl)pyrazolato-*N'*¹*N'*²*N'*³)- μ -dimethanolinickel(II)] dichloride dihydrate, 1401–4

- Reactivity of binuclear pyrazolate-bridged rhodium complexes with isocyanides. Molecular structure of [Rh₂($\mu\text{-pz}$)₂(CNBu^t)₂($\mu\text{-Ph}_2\text{PCH}_2\text{PPh}_2$)₂]BF₄ (pz = pyrazolate), 1427–34

- Preparation and structural studies of diphenylthiophosphinito rhodium(III) complexes. X-ray structure analysis of [(C₅Me₃)Rh₂($\mu\text{-Cl}$)₂($\mu\text{-Ph}_2\text{PS}$)][BPh₄], 1987–92

- Binuclear copper(II) complexes of new dinucleating ligands with a pyrazolate group as an endogenous bridge. Effects of exogenous azide and acetate bridges on magnetic properties, 2077–82

PYRAZOLYL

- Heteroleptic poly(pyrazolyl)borate derivatives of the lanthanide ions. The synthesis of acetylacetone complexes and the molecular structures of [Ln{HB(C₃N₂H₃)₃}₂{MeC(O)CHC(O)Me}] (Ln = Ce or Yb), 1393–400

- Unusual chelating properties of the ligand 1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-dithiaoctane (bddd). Crystal structures of Ni(bddd)(NCS)₂, Zn(bddd)(NCS)₂ and Cd₂(bddd)(NCS)₄, 2309–14

PYRAZOLYLBORATE

- Activation of a vinyl carbon–hydrogen bond in a tris(pyrazolyl)boratoiridium complex. The X-ray crystal structure of [IrH(HB(pz)₃)₂($\sigma\text{-C}_8\text{H}_{13}$)₂($\eta^2\text{-C}_8\text{H}_{14}$)], 2073–6

PYRIDINALDAZINE

- Iridium(III) hydrido complexes with 2-pyridinaldazines. Crystal structure of [IrH₂(mpaa)(PPh₃)₂]SbF₆. Synthesis and characterization of homo- and hetero-nuclear complexes, 1075–82

- Reactivity of the iridium nitrosyl complex

- [Ir(NO)(paa)(PPh₃)₂][PF₆]₂ (paa = 2-pyridinecarbaldehyde azine) with nucleophiles: insertion of NO into a methinic C–H bond; crystal structure of [Ir(2-C₅H₄N)CH=N–N=C(NOH)(2-C₅H₄N){Me(EtO)C=NH}(PPh₃)₂][PF₆]₂, 1217–26

PYRIDINE

- Studies in Werner clathrates. Part 9. Structures with bis(isothiocyanato)tetrakis(4-phenylpyridine)nickel(II) as host, 119–26

- Synthesis, infrared, and tin-119 Mössbauer spectroscopic characterization of seven-co-ordinate diorganotin(IV) adducts with 2,6-diacylpyridine acylhydrazones, including the X-ray crystal structure of SnEt₂(dap) [H_2dap = 2,6-diacylpyridine bis(2-thienoylhydrazone)], 289–94

- Ligand additivity in the oxidation potentials of bidentate mixed-ligand ruthenium(II) complexes, 371–4

- Synthesis, characterization, and crystal structures of bismuth complexes with 2,6-diacylpyridine bis(2-thienoylhydrazone) (H₂dapt): [Bi(Hdapt)Cl₂] \cdot dms₂ \cdot H₂O and [Bi(dapt)Cl] \cdot dms₂, containing a quinquedentate ligand, 671–6

PYRIDINE (contd.)

- Studies on transition-metal oxo and nitrido complexes. Part 10. New oxo-ruthenium and oxo-osmium pyridine complexes, and use of the former as catalysts for oxidation of alcohols, 901-6
- Reactivity of metallic copper with acidic organic compounds in the presence of molecular oxygen. Crystal structure of $\{Cu(NCO)(py)_2\}_2J\text{-H}_2\text{O}$, 921-6
- Precursor catenand complexes: synthesis, structure, and electrochemistry of bis(2,6-di-iminopyridyl) complexes of nickel(II). The single-crystal X-ray structure of $[NiL^4]_2[BF_4]_2$, 965-70
- Blue copper models. Spectroscopic and electrochemical studies of copper(II) complexes with new ligand systems containing sulphur and nitrogen donor atoms, 1161-70
- Formation of mixed-ligand complexes of cadmium(II) halide with tri-n-octylphosphine oxide and pyridine or substituted pyridines in 1,2-dichloroethane, 1259-64
- Complexes of the pyridine-based, tridentate, sulphur-containing ligands 2,6-bis(ethylthiomethyl)pyridine, 2,6-bis(methoxycarbonylthiomethyl)pyridine, and 2,6-bis(benzylthiomethyl)pyridine. Crystal structure of [2,6-bis(ethylthiomethyl)pyridine]dichlorocadmium(II) hydrate, 1381-4
- Binuclear copper(I) complexes containing two or three 2-(diphenylphosphine)pyridine (dpyp) bridging ligands: X-ray crystal structure of $[Cu_2(\mu\text{-dpyp})_3(MeCN)][BF_4]_2$ and variable-temperature ^1H and ^{31}P - ^{1}H nuclear magnetic resonance studies, 1499-506
- Copper(II)-assisted hydrolysis of 2,4,6-tris(2-pyridyl)-1,3,5-triazine. Part 2. Crystal structures of [bis(2-pyridylcarbonyl)amido](pyridine-2-carboxamide)copper(II) trifluoromethanesulphonate and [bis(2-pyridylcarbonyl)amido][2,4,6-tris(2-pyridyl)-1,3,5-triazine]copper(II) trifluoromethanesulphonate, 1681-8
- Closely related macrocyclic and acyclic tridentate, pyridine derivatives, containing sulphur, and their complexes. Crystal structures of {dichloro-3,10-dithia-16-azabicyclo[10.3.1]hexadeca-1(16),12,14-triene}copper(II) and [2,6-bis(ethylthiomethyl)pyridine]dichlorocopper(II), 1739-44
- Oxidative addition of dibenzoyl peroxide on metallic copper. The crystal structure of $[\{Cu(C_6H_5CO_2)_2(py)\}_2]$, 2331-4
- PYRIDINECARBALDEHYDE**
- Crystal structure, spectroscopic and magnetic properties of the complex $[Cu(paphy)(NCS)(SCN)]$ (paphy = pyridine-2-carbaldehyde 2'-pyridylhydrazone). An unusual di- μ -thiocyanato- N bridged copper(II) dimer, 53-6
- PYRIDINEDICARBOXYLATE**
- The co-ordination chemistry of divalent cobalt, nickel, and copper. Part 13. Characterization and stability constants of copper(II) species isolated from mixed-ligand solutions; crystal structure of the complex (pyridine-2,6-dicarboxylato)(*N,N,N',N'*-tetramethyl-1,3-diaminopropane)copper(II), 253-8
- PYRIDINE THIOLATE**
- Binuclear platinum(II) complexes bridged by pyridine-2-thiolate and related ligands. Synthesis and crystal structure, 815-20
- PYRIDINETHIONATE**
- Stepwise conversion of $[Os(CF_3CO_2)_2(CO)_4]$ into $[Os(pyS)_2(CO)_2]$; X-ray crystal structures of the complexes $[Os(pyS)_2(CO)_x]$ where $x = 2$ or 3 and pyS is the pyridine-2-thionato ligand, 2211-6
- PYRIDINETHIONE**
- Synthesis and X-ray crystal structure of chloro[2(1*H*)-pyridinethione-*S*]bis(triphenylphosphine)copper(I), 749-52
- Spectroscopic and electrochemical study of copper and zinc complexes with a N_2S_2 ligand donor set. Crystal structure of the copper(II) complex derived from 1,3-propylenediamine and 3-formyl-1-phenyl-2(1*H*)-pyridinethione, 1979-86
- PYRIDYL**
- A planar, binucleating, pyrazole derivative ligand. Crystal structure of bis[μ -3,5-bis(2-pyridyl)pyrazolato-*N*¹*N'*¹*N''*²] \cdot bis(dimethanolnickel(II)) dichloride dihydrate, 1401-4
- PYRIMIDINE**
- Reactions of chloro(triphenylphosphine)gold(I) and [μ -1,2-bis(diphenylphosphine)ethane]-bis[bromogold(I)] with oxopurine bases. Molecular and crystal structure and bonding of (theophyllinato)(triphenylphosphine)gold(I), 2323-30
- PYRROLIDINE**
- Synthesis and stereochemical studies of η^3 -allyl palladium(II) complexes containing a chiral chelating ligand, 161-8

PYRROLIDINONATE

Structural, spectroscopic, and electrochemical characterization of tetrakis- μ -(2-pyrrolidinonato)-dirhodium(II) and tetrakis- μ -(δ -valerolactamato)-dirhodium(II), 93-100

PYRROLIDONATE

Synthesis, X-ray characterization, and Q-band electron spin resonance studies of heterometallic, polymeric compounds formed by yttrium and lanthanide(III) nitrates with HgL_2 ($L = C_6H_6NO$, the anion of 2-pyrrolidone); crystal structures of $[\{Hg_2EuL_4(NO_3)_3\}_n]$ and $[\{Hg_3Tb_2L_6(NO_3)_6\}_n]$, 1439-48

PYRROLYL

Synthesis, electrochemistry, and electron spin resonance spectra of some seventeen-electron complexes containing the $\{Mo(NO)_3\}^{2+}$ core and the molecular structure of $[Mo(NO)_3(HB(Me_2pz)_3)](pyr)_2$, 127-32

PYRUVIC ACID

Thiosemicarbazones as co-ordinating agents. Part 3. Synthesis, spectroscopic characterization, and X-ray structure of methyl pyruvate thiosemicarbazone hemihydrate, chloro(ethyl pyruvate thiosemicarbazone)copper(II) (green form), and chloro(pyruvic acid thiosemicarbazone)copper(II) dihydrate (blue form), 361-6

QUINOLINE

Preparation and characterisation of linear mixed-metal trinuclear carboxylate complexes of general formula $[M_2M'(O_2CR)_6(C_9H_7N)_2]$; crystal structure of $[Zn_2Ba(O_2CCMe_3)_6(C_9H_7N)_2]$, 1127-32

QUINONE

One-electron transfer processes in the reaction of tin(II) halides with substituted *o*-quinones; crystal structure of bis(3,5-di-t-butylcatecholate)tin 1,10-phenanthroline-dimethylformamide (1/1/2), 439-46

RADIOTRACER

Radiotracers in fluorine chemistry. Part 12. Reactions between caesium fluoride and arsenic pentafluoride, boron trifluoride, sulphur tetrafluoride, carbonyl fluoride, and carbon dioxide under heterogeneous conditions at room temperature. Radiotracer studies using fluorine-18, sulphur-35, and carbon-14, 937-42

RAMAN

Syntheses, electrochemistry, and spectroscopy of dirhodium(II) tetra-acetamidate and tetrakis(trifluoroacetamide) complexes with axial Group 15 substituents. The X-ray crystal structures of $[Rh_2(CH_3CONH)_4(AsPh_3)_2]$ and $[Rh_2(CH_3CONH)_4(CH_3CO_2)_2(MPh_3)_2]$, M = As or Sb, $n \approx 1$, 581-8

Nature of peroxyborate species in aqueous solution: a study by boron-11 nuclear magnetic resonance and Raman spectroscopy, 1651-6

Syntheses, crystal structures, and Raman and resonance Raman spectroscopy of platinum(II), platinum(IV), and halogen-bridged chain platinum(II,IV) complexes involving the terdentate ligand 3-methyl-3-apzapentane-1,5-diamine (4Me-dien), 1689-96

Vibrational spectroscopic studies of tervalent hexa-aqua cations: oriented single-crystal Raman spectra between 275 and 1 200 cm⁻¹ of the caesium rhodium alums, $CsRh(SO_4)_2 \cdot 12H_2O$, $CsRh(SeO_4)_2 \cdot 12H_2O$, and $CsRh(SO_4)_2 \cdot 12D_2O$, 1771-8

Vibrational study of ionic association in aprotic solvents. Part 12. Isothiocyanate M(NCS) complexes of non-transition-metal ions, 2193-8

Bonding in clusters. Part 10. The nature of the BH groups in *closoborane* anions. A study by vibrational spectroscopy, electronic spectroscopy for chemical analysis, and *ab initio* computation, 2253-60

RHENIUM

A kinetic study of the transmission of ligand effects across the quadruple bond in $[Re_2Cl_8]^{2-}$, 767-72

Catalytic oxidation of water mediated by rhenium oxides, 849-52

Synthesis and characterization of a family of nitrile-substituted triangular hydridocarbonyl rhenium clusters, $[Re_3(\mu-H)_n(CO)_1{}_2 \cdot {}_n(NCMe)]$ ($n = 1-3$). X-Ray crystal structures of $[Re_3(\mu-H)_3(CO)_1{}_1(NCMe)]$ and $[Re_3(\mu-H)_3(CO)_1{}_2(NCMe)_3]$, 1143-8

Syntheses and redox properties of mixed isocyanide, carbonyl, or nitrile complexes of rhenium(I) *trans*- $[Re(CNMe)L(Ph_2PCH_2CH_2PPH_2)]X$ [L = CNR (R = alkyl or aryl), CO, or NCMe; X = Cl, BF_4^- , or PF_6^-], 1209-16

Synthesis and X-ray crystal structure of $[ReO(Cl)](OC_6H_4CH=NCH_2CH_2OCH_2)_3$, an example of four-co-ordination by a potentially sexidentate ligand, 1405-8

RHENIUM (contd)

- Crystal and molecular structures of μ -hydrido-bis[*fac*-(2,2'-bipyridine)tricarbonylrhenium(I)] chloride, *fac*-(2,2'-bipyridine)tricarbonylformatorhenium(I), and *fac*-(2,2'-bipyridine)tricarbonyl(cyanotrihydroborato)rhenium(I), 1449-54
 Complexes of technetium, rhenium, and rhodium with sexidentate Schiff-base ligands, 2115-20
 Synthesis and *X*-ray crystal structure of tetra(2-methylphenyl)molybdenum(IV), $\text{Mo}(2\text{-MeC}_6\text{H}_4)_4$. Redox chemistry of $\text{M}(2\text{-MeC}_6\text{H}_4)_4$ compounds of molybdenum, rhenium, ruthenium, and osmium, 2149-54
 Reactions of 1-alkynes with *trans*-[$\text{ReCl}(\text{N}_2)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2$]: preparation of the vinylidene compounds *trans*-[$\text{ReCl}(\text{C}=\text{C}=\text{CHR})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2$] (R = alkyl or aryl) and *X*-ray structure of *trans*-[$\text{ReCl}(\text{C}=\text{C}=\text{CHPh})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2$], 2381-8

RHODIUM

- Rhodium complexes of the binucleating ligands pyridine-2-thiolate and benzothiazole-2-thiolate. Crystal structures of [$\text{[Rh}(\mu-\text{SC}_5\text{H}_4\text{N})(\text{CO})_2]_2$] and [$\text{[Rh}(\mu-\text{SC}_5\text{H}_4\text{N})(\text{tfbb})]_2\text{-Me}_2\text{CO}$ (tfbb = tetrafluorobenzobarrelene)], 25-32
 Structural, spectroscopic, and electrochemical characterization of tetrakis- μ -(2-pyrrolidinone)-dirhodium(II) and tetrakis- μ -(δ -valerolactamato)-dirhodium(II), 93-100
 Novel five-co-ordinate diorganorhodium compounds. *X*-Ray structure of bis-*ortho*-chelated [$\text{Rh}^{III}(\text{Me})\{\text{C}_6\text{H}_3(\text{CH}_2\text{NMe}_2)_2\}_2\text{-2,6'_1}$], 317-24
 Syntheses, electrochemistry, and spectroscopy of dirhodium(II) tetra-acetamide and tetrakis(trifluoroacetamide) complexes with axial Group 15 substituents. The *X*-ray crystal structures of [$\text{Rh}_2(\text{CH}_3\text{CONH})_4(\text{AsPh}_3)_2$] and [$\text{Rh}_2(\text{CH}_3\text{CONH})_{4-n}(\text{CH}_3\text{CO}_2)_n(\text{MPh}_3)_2$], M = As or Sb, $n \approx 1.581-8$
 The chemical and electrochemical one-electron oxidation of diamidonaphthalene-bridged complexes: paramagnetic $[\text{Rh}_2]^{3+}$ containing A-frames, 683-8
 Trinuclear mixed-valence complexes with the linear $[\text{Rh}_3]^{4+}$ core, 689-92
 Synthesis and *X*-ray crystal structure of $[\text{Rh}(\text{NO})(\text{PPh}_3)_2(p\text{-MeC}_6\text{H}_4\text{NNNC}_6\text{H}_4\text{Me}-p)][\text{BF}_4]\text{-CH}_2\text{Cl}_2$, 745-8
 Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 86. Alkyldyne(carbaborane)molybdenum-gold, -rhodium and -iron complexes; crystal structure of $[\text{NEt}_4][\text{MoFe}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)\mu\text{-}\sigma:\sigma:\eta^5\text{-C}_2\text{B}_9\text{H}_7\text{Me}_2](\text{CO})_8$, 797-808
 New carbide clusters in the cobalt subgroup. Part 17. Preparation and structural characterization of the mixed-metal octahedral dianion $[\text{Co}_2\text{Rh}_4\text{C}(\text{CO})_{13}]^{2-}$ as its $[\text{PPh}_4]^+$ salt, 879-84
 Crystal structures and solution dynamics of mono- and di-nuclear rhodium complexes of heterodinucleating macrocyclic ligands containing the 2,6-di(thiomethyl)pyridine sub-unit, 915-20
 Synthesis of a new P_2N_2 ligand *N,N'*-bis[2-(diphenylphosphino)phenyl]propane-1,3-diamine, H_2L^2 , and some of its complexes with elements of the nickel triad and rhodium: *X*-ray structure analyses of the neutral complex $[\text{NiL}^2]$ and the *trans*-spanned $[\text{Rh}(\text{CO})\text{Cl}(\text{H}_2\text{L}^2)]$, 1067-74
 Trinuclear pyrazolate complexes. Crystal structure of $[(\text{OC})_2\text{Rh}(\mu\text{-pz})_2\text{Pd}(\mu\text{-pz})_2\text{Rh}(\text{CO})_2]$, 1133-8
 Synthesis of ammonium ions and nitrosylation reactions using nitrosyl chloride and aryl nitrites, 1139-42
 Reaction of $[\text{Rh}_2(\mu\text{-1,8-(NH)}_2\text{C}_{10}\text{H}_6)(\text{CO})_2(\text{PPh}_3)_2]$ with $[\text{M}(\text{PPh}_3)]^+$ (M = Au, Ag, or Cu): formation of unusual 44-electron Rh_2M clusters. Crystal structure of $[\text{Rh}_2(\mu\text{-Au}(\text{PPh}_3))(\mu\text{-1,8-(NH)}_2\text{C}_{10}\text{H}_6)(\text{CO})_2(\text{PPh}_3)_2]\text{ClO}_4\text{-2CH}_2\text{Cl}_2$, 1249-52
 Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 87. Docosahedral carbaborane(alkyldyne)tungsten complexes as reagents for the synthesis of compounds with heteronuclear metal-metal bonds: crystal structures of $[\text{NEt}_4][\text{W}(\equiv\text{CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_2(\eta^6\text{-C}_2\text{B}_{10}\text{H}_{10}\text{Me}_2)]$ and $[\text{NEt}_4][\text{WFe}(\mu\text{-CC}_6\text{H}_3\text{Me}_2\text{-2,6})(\text{CO})_4(\eta^6\text{-C}_2\text{B}_{10}\text{H}_{10}\text{Me}_2)]$, 1363-74
 Reactivity of binuclear pyrazolate-bridged rhodium complexes with isocyanides. Molecular structure of $[\text{Rh}_2(\mu\text{-pz})\text{I}_2(\text{CNBu}')_2(\mu\text{-Ph}_2\text{PCH}_2\text{PPh}_2)_2]\text{BF}_4$ (pz = pyrazolate), 1427-34
 Cationic dinuclear rhodium complexes as catalyst precursors for the hydroformylation of alkenes, 1579-82
 Synthesis, structure, and electrochemical reduction of 2-(arylaazo)pyridine complexes of rhodium(III), 1635-8

Platinum metal thioether macrocyclic complexes: synthesis, electrochemistry, and single-crystal *X*-ray structures of *cis*-[$\text{[RhCl}_2\text{L}^2]_2\text{PF}_6$ and *trans*-[$\text{[RhCl}_2\text{L}^3]_2\text{PF}_6$] ($\text{L}^2 = 1,4,8,11$ -tetraathiacyclohexadecane, $\text{L}^3 = 1,5,9,13$ -tetraathiacyclohexadecane), 1675-80

Vibrational spectroscopic studies of tervalent hexa-aqua cations: oriented single-crystal Raman spectra between 275 and 1 200 cm^{-1} of the caesium rhodium alums, $\text{CsRh}(\text{SO}_4)_2\text{-12H}_2\text{O}$, $\text{CsRh}(\text{SeO}_4)_2\text{-12H}_2\text{O}$, and $\text{CsRh}(\text{SO}_4)_2\text{-12D}_2\text{O}$, 1771-8

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 89. Tetra- and penta-nuclear tungsten-rhodium complexes: crystal structures of $[\text{W}_3\text{Rh}_2(\mu\text{-CO})(\mu\text{-CMe})(\mu\text{-C}(\text{Me})\text{C}(\text{O}))(\mu\text{-PPh}_2)_2(\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3]$ and $[\text{W}_3\text{Rh}_2(\mu\text{-CO})_3(\mu\text{-CMe})(\mu\text{-C}(\text{Me})\text{PPh}_2)(\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3]$, 1855-64

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 90. Synthesis of the complexes $[\text{W}_2\text{M}_2(\mu\text{-CO})(\mu\text{-CMe})(\mu_3\text{-CMe})(\mu\text{-PPh}_2)_2(\text{CO})_3(\mu\text{-C}_5\text{H}_5)_2]$ (M = Rh or Ir) and related tetra- and penta-nuclear metal compounds; interconversion of ketenyl and λ^5 -phospho-alkyne ligands bridging tungsten-rhodium bonds, 1865-70

Preparation and structural studies of diphenylthiophosphinito rhodium(III) complexes. *X*-Ray structure analysis of $[(\text{C}_5\text{Me}_5)\text{Rh}_2(\mu\text{-Cl})_2(\mu\text{-P}_2\text{PS})][\text{BPh}_4]$, 1987-92

Reduction-oxidation properties of organotransition-metal complexes. Part 30. Triazenido-bridged carbonylrhodium bipyridyl complexes with $[\text{Rh}_2]^{2+}$, $[\text{Rh}_2]^{3+}$, and $[\text{Rh}_2]^{4+}$ cores, and the *X*-ray structures of $[\text{Rh}_2(\text{CO})_2(\text{bipy})(\mu\text{-RNNNR})_2][\text{BF}_4]\text{-CH}_2\text{Cl}_2$ and $[\text{Rh}_2\text{I}(\text{CO})(\text{bipy})(\mu\text{-RNNNR})_2]_2[\text{PF}_6]\text{-2.5CH}_2\text{Cl}_2$ (R = *p*-tolyl), 2049-60

Complexes of technetium, rhenium, and rhodium with sexidentate Schiff-base ligands, 2115-20

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 94. Synthesis of di- and tri-metal compounds *via* the reagents $[\text{M}(\equiv\text{C-C}\equiv\text{CBu}^1)(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ (M = Mo or W), 2281-8

Mixed-metal carbido carbonyl clusters. Part 4. Synthesis and structural characterization of $[\text{Rh}_6\text{C}(\text{CO})_{13}\{\text{Au}_2(\text{PPh}_3)_2\}]$, 2343-6

Cyclometallation of unsaturated carboxylic acids by pentamethylcyclopentadienyl-rhodium and -iridium complexes.

Crystal structures of $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh}=\text{CHCO-O})(\text{Me}_2\text{SO})]$ and $[(\text{C}_5\text{Me}_5)\text{Ir}(\text{CPh}=\text{CHCO-OMe})\text{I}]^+$, 2415-22

Preparation and *X*-ray crystal structure of $[\text{Ru}_3\text{RhH}_3(\text{CO})_{12}]$, 2449-52

RUTHENIUM

Ruthenium phthalocyanine chemistry: synthesis and properties of a mixed-valence nitrido-bridged ruthenium phthalocyanine dimer, 193-6

Redox activation of the hexanuclear ruthenium cluster $[\text{Ru}_6\text{C}(\text{CO})_{16}]^{2-}$ using oxidative addition, 243-6

Reduction-oxidation properties of organotransition-metal complexes. Part 29. Pentaphenylcyclopentadienyl complexes of ruthenium, 283-8

Ligand additivity in the oxidation potentials of bidentate mixed-ligand ruthenium(II) complexes, 371-4

Alkyne insertion into ruthenium-carbon bonds: formation and ring-closing reactions of vinyl complexes, 465-70

Reactions of hydride complexes of ruthenium(II) with alkynes, 589-94

Chemistry of multi-site bound unsaturated ligands on metal carbonyl clusters. Intramolecular reductive coupling in the conversion of $[\text{Ru}_3(\mu\text{-H})(\text{CO})_9(\mu_3\text{-}\eta^2\text{-C}\equiv\text{CBu}^1)]$ to the butadienyl cluster $[\text{Ru}_3(\text{CO})_8(\mu_3\text{-}\eta^4\text{-C}(\text{Bu}^1)\text{CC}(\text{H})\text{-C}(\text{Ph})(\text{PPh}_2))]$, and its crystal structure. An example of ligand isomerism on metal clusters, 601-6

Studies on transition-metal oxo and nitrido complexes. Part 10. New oxo-ruthenium and oxo-osmium pyridine complexes, and use of the former as catalysts for oxidation of alcohols, 901-6

An investigation of the bonding in $[\text{Ru}_2(\text{O}_2\text{CCF}_3)_4]$ and $[\text{Ru}_2(\text{O}_2\text{CCF}_3)_4(\text{NO})_2]$, 1037-44

Six- and five-co-ordinate ruthenium(II) complexes with 1,2-bis(dicyclohexylphosphino)ethane. *X*-Ray crystal structure of the d^6 five-co-ordinate $[\text{RuCl}\{(\text{C}_6\text{H}_{11})_2\text{PCH}_2\text{CH}_2\text{P}(\text{C}_6\text{H}_{11})_2\}_2]\text{PF}_6$, 1045-52

Synthesis and reactivity of binuclear pyrazolate-bridged ruthenium(I) complexes. Crystal structures of bis[μ -(3,5-dimethylpyrazolato-*N,N'*)tricarbonylruthenium(I)] ($Ru-Ru$) and

RUTHENIUM (contd)

bis[μ -(3,5-dimethylpyrazolato-*NN'*)]- μ -iodo-
bis[tricarbonylruthenium(II) tri-iodide, 1093–100

Synthesis of ammonium ions and nitrosylation reactions using nitrosyl chloride and alkyl nitrites, 1139–42

The heteronuclear cluster chemistry of the Group 1B metals. Part 11. Effect of the nature of the bidentate diphosphine ligand on the metal framework structures of the gold heteronuclear cluster compounds $[Au_2Ru_4(\mu_3\text{-H})(\mu\text{-H})\{\mu\text{-Ph}_2P(\text{CH}_2)_nPPh_2\}\{\text{CO}\}_{12}]$ ($n = 1\text{--}6$). X -Ray crystal structures of $[Au_2Ru_4(\mu_3\text{-H})(\mu\text{-H})\{\mu\text{-Ph}_2P(\text{CH}_2)_nPPh_2\}\{\text{CO}\}_{12}]$ ($n = 1\text{ or }2$), 1227–36

Synthesis and crystal structure of $[NiOs_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2\{\mu\text{-Cu}(PPh_3)\}\{\text{CO}\}_9]$. Hydrogenation–dehydrogenation reactions of dienes and alcohols with heterogeneous catalysts derived from $[NiM_3(\eta\text{-C}_5\text{H}_5)(\mu\text{-H})_2\{\mu\text{-X}\}\{\text{CO}\}_9]$ [$M = Ru, X = H; M = Os, X = Cu(PPh_3)$], 1477–82

Organic chemistry of dinuclear metal centres. Part 12. Synthesis, X -ray crystal structure, and reactivity of the di- μ -alkylidene complex $[Ru_2(\text{CO})_2(\mu\text{-CHMe})(\mu\text{-CMe}_2)(\eta\text{-C}_5\text{H}_5)_2]$: alkylidene linking, 1799–806

Organic chemistry of dinuclear metal centres. Part 13. Synthesis, structure, and reactivity of $[Ru_2(\text{CO})_4(\eta^5\text{:}\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{C}_5\text{H}_4)]$, 1807–14

Photofragmentation kinetics of some triruthenium carbonyl clusters, 1933–40

Kinetics of formation, dissociation, and isomerization reactions in complexes of cyanopyridines and benzonitrile with pentacyanoruthenate(II), 1973–8

Ruthenium oxo complexes of macrocyclic tertiary amines.

Synthesis, electrochemistry, and reactivities of ruthenium oxo complexes of *meso*-2,3,7,11,12-pentamethyl-3,7,11,17-tetraazabicyclo[11.3.1]heptadeca-1(17),13,15-triene (L^1), and X -ray crystal structure of *trans*- $[Ru^{IV}L^1(\text{O})(\text{NCO})]\text{ClO}_4$, 2011–6

The heteronuclear cluster chemistry of the Group 1B metals. Part 12. Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds $[M_2Ru_4(\mu_3\text{-H})_2(\mu\text{-L})(\text{CO})_{12}]$ [$M = Cu$ or Ag , $L = Ph_2P(\text{CH}_2)_nPPh_2$ ($n = 1\text{--}6$) or *cis*- $Ph_2\text{PCH}=\text{CHPPh}_2$; $M = Cu$, $L = Ph_2\text{As}(\text{CH}_2)_nPPh_2$ ($n = 1$ or 2)] in solution, 2039–48

Oxo transfer reactions. Mechanistic studies on the oxidation of triphenylphosphine by *trans*- $[Ru^{VI}L(\text{O})_2]^{2+}$ ($L = 1,4,8,11$ -tetramethyl-1,4,8,11-tetra-azacyclotetradecane), 2065–8

Synthesis and X -ray crystal structure of tetra(2-methylphenyl)molybdenum(IV), $Mo(2\text{-MeC}_6\text{H}_4)_4$. Redox chemistry of $M(2\text{-MeC}_6\text{H}_4)_4$ compounds of molybdenum, rhenium, ruthenium, and osmium, 2149–54

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 95. Synthesis and crystal structures of $[W_2RuPt(\mu\text{-CC}_6\text{H}_4\text{Me}-4)(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)(\text{CO})_6(\text{PM}_2\text{Ph})(\eta\text{-C}_5\text{H}_5)_2]$ and $[W_2Ru_2Pt_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\mu\text{-CO})_2(\text{CO})_9(\eta\text{-C}_5\text{H}_5)_2]$, 2289–96

Substituted cyclopentadienyl complexes. Part 4. The catalysed synthesis and proton nuclear magnetic resonance spectra of $[Ru(\eta^5\text{-C}_5\text{H}_4\text{Me})(\text{CO})(L)]$ and $[Ru(\eta^5\text{-C}_5\text{H}_7)(\text{CO})(L)]$ ($L =$ phosphine or isocyanide) and the crystal structure determination of $[Ru(\eta^5\text{-C}_5\text{H}_7)(\text{CO})\{P(\text{CH}_2\text{Ph})_3\}]$, 2407–14

Synthesis, spectroscopic, electrochemical, and magnetic properties of dimolybdenum(II,II), diruthenium-(II,II) and -(II,II) complexes containing bridging aspirinate (2-acetoxybenzoate) ligands, 2435–42

Preparation and X -ray crystal structure of $[Ru_3RhH_3(\text{CO})_{12}]$, 2449–52

SALICYLALDEHYDATE

Dimeric mixed-ligand copper(II) complexes with di-imines and salicylaldehyde, 1435–8

SALICYLAMIDE

Synthesis, structure, and redox properties of dicopper(II,II) complexes of *N*-(hydroxyphenyl)salicylamide and homologues. Facile oxidation to dicopper-(II,III) and -(III,III) species; crystal structure of bis(tetrabutylphosphonium) bis{[(2-oxidobenzoyl)(2'-oxidophenyl)amido(3-) NO_2^+ , $\mu\text{-O}^2^-$]cuprate(II)}, 2089–94

SCHIFF BASE

Metal complexes (Co^{3+} , Fe^{3+} , Mn^{3+}) of the quinquedentate salen analogue, 2-methyl-4-benzylthio-*N,N'*-butane-1,2-diylbis(salicylideneimine): crystal structures and sulphur co-ordination effects, 179–84

Formation in solution, synthesis, and electrochemical study of

oxalato complexes of *N,N'*-ethylenebis(salicylideneiminato)-chromium(III) and -iron(III): crystal structures of piperidinium [*N,N'*-ethylenebis(salicylideneiminato)](oxalato- $O^1O^2^-$)-chromate(III) and ferrate(III), 729–38

N,N'-Ethylenebis(acetylacetoneiminato)titanium complexes: synthesis, crystal structures, and properties of titanium(III) and titanium(IV) derivatives, 953–8

Synthesis and X -ray crystal structure of $[\text{ReO}(\text{Cl})\{(2\text{-OC}_6\text{H}_4\text{CH}=\text{NCH}_2\text{CH}_2\text{OCH}_2)_2\}]$, an example of four-co-ordination by a potentially sexidentate ligand, 1405–8

Synthesis and characterization of cobalt(III) and cobalt(IV) complexes of *N*-(2-hydroxyphenyl)salicylamide and homologues, 1613–6

Complexes of technetium, rhenium, and rhodium with sexidentate Schiff-base ligands, 2115–20

SELENIIDE

Chromium chalcogen complexes. Insertion of elemental sulphur into $[\text{Cr}(\text{cp})(\text{CO})_2\text{Se}]$ and $[\text{Cr}(\text{cp})_2(\text{CO})_4\text{Se}_2]$ ($\text{cp} = \eta\text{-C}_5\text{H}_5$), 431–8

Phosphorus-31 and selenium-77 nuclear magnetic resonance spectra of sulphur- or arsenic-substituted derivatives of tetraphosphorus triselenide; fitting of chemical shifts and coupling constants to molecular parameters for cage compounds containing a phosphorus–selenium–phosphorus linkage, 977–84

SELENIUM

Phosphorus-31 and selenium-77 nuclear magnetic resonance spectra of sulphur- or arsenic-substituted derivatives of tetraphosphorus triselenide; fitting of chemical shifts and coupling constants to molecular parameters for cage compounds containing a phosphorus–selenium–phosphorus linkage, 977–84

Metalla(heteroborane) chemistry. Part 5. Synthesis of *closo*-[2,2- $(\text{PR}_3)_2\text{-1,2-SePtB}_{10}\text{H}_{10}]$ ($\text{R}_3 = \text{Et}_3$, $\text{Bu}^n{}_3$, or Me_2Ph) and *arachno*-[9,9-(PM_2Ph) $_2$ -6,9-SePtB $_8\text{H}_{10}]$ and their characterisation by nuclear magnetic resonance techniques: the crystal and molecular structures of *closo*-[2,2-($\text{PEt}_3)_2\text{-1,2-SePtB}_{10}\text{H}_{10}]$ and *arachno*-[9,9-(PM_2Ph) $_2\text{-6,9-SePtB}_8\text{H}_{10}]$, 1657–68

SELENIUM-77

Phosphorus-31 and selenium-77 nuclear magnetic resonance spectra of sulphur- or arsenic-substituted derivatives of tetraphosphorus triselenide; fitting of chemical shifts and coupling constants to molecular parameters for cage compounds containing a phosphorus–selenium–phosphorus linkage, 977–84

SELENOACYL

Insertion of phenylethyne-thiolate or -selenolate into metal–carbene bonds: synths for co-ordinated thioacyl or selenoacyl anions and for anionic thiocarbene or selenocarbene complexes, 1565–78

SELENOETHER

Platinum metal complexes of potentially chelating alkene–thioether and alkene–selenoether ligands: synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{MeE}(\text{CH}_2)_n\text{CH}=\text{CH}_2\}]$ ($\text{M} = Pt$ or Pd ; $X = Cl$, Br , or I ; $E = S$ or Se ; $n = 2$ or 3) and the X -ray structure of *cis*-dibromo(2-thia-6-heptene)platinum(II), $[\text{PtBr}_2\{\text{MeS}(\text{CH}_2)_3\text{CH}=\text{CH}_2\}]$, 985–90

Platinum metal complexes of potentially chelating alkene thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of $[\text{MX}_2\{\text{E}[(\text{CH}_2)_n\text{CR}=\text{CR}_2]\}]$ ($\text{M} = Pt$ or Pd ; $X = Cl$, Br , or I ; $E = S$ or Se ; $n = 2$ or 3 ; $R = H$ or Me) and the X -ray crystal structure of *cis*-di-iodo(5-thianona-1,8-diene)platinum(II), $[\text{PtI}_2\{\text{S}[(\text{CH}_2)_2\text{CH}=\text{CH}_2\]\}]$, 2315–22

SELENOLATE

Insertion of phenylethyne-thiolate or -selenolate into metal–carbene bonds: synths for co-ordinated thioacyl or selenoacyl anions and for anionic thiocarbene or selenocarbene complexes, 1565–78

SILANE

Five-co-ordinated silicon compounds: geometry of formation by intramolecular co-ordination. Crystal structure of 2-(dimethylaminomethyl)phenyl-1-naphthylsilane, 43–52

SILICATE

Oxygen-17 nuclear magnetic resonance spectroscopic studies of aqueous alkaline silicate solutions, 275–82

Thermal and hydrothermal stability of SAPO-5 and SAPO-37 molecular sieves, 1361–2

SILICON

Five-co-ordinated silicon compounds: geometry of formation by intramolecular co-ordination. Crystal structure of 2-(dimethylaminomethyl)phenyl-1-naphthylsilane, 43–52

Oxygen-17 nuclear magnetic resonance spectroscopic studies of aqueous alkaline silicate solutions, 275–82

SILICON (contd)

- Isopropylsilylamine and disilyl-t-butylamine: preparation, spectroscopic properties, and molecular structure in the gas phase, determined by electron diffraction, 779–84
 Thermal and hydrothermal stability of SAPO-5 and SAPO-37 molecular sieves, 1361–2
 Synthesis and properties of trimethyl(trifluorosilyl)stannane, 1879–82
 Aqueous chemistry and thermodynamic modelling of $\text{CaO}\text{-SiO}_2\text{-H}_2\text{O}$ gels, 2371–80

SILOXANE

- Cyclic metallasiloxane compounds of Ti^{IV} and Zr^{IV} . X-Ray crystal structures of a *cis*-bis(pyridine) bis(cyclosiloxy)titanate and a pyridinelithiated tris(cyclosiloxy)zirconate, 2347–52

SILOXIDE

- Trit(*n*-cyclopentadienyl)uranium(IV) silyl and siloxide compounds. Crystal structure of $[\text{U}(\eta^5\text{C}_5\text{H}_5)_3(\text{OSiPh}_3)]$. Insertion of isocyanide into a uranium–silicon bond, 677–82

SILVER

- Kinetics and mechanism of the reaction between 4-t-butylphenolate anion and tetrahydroxoargentate(III) in aqueous alkaline media, 217–22
 Synthesis and structural studies on silver–tin complex salts with bis(diphenylarsino)methane and bis(diphenylphosphino)methane, 247–52
 Oxidation of silver metal and silver(I) cations by molybdenum, tungsten, and uranium hexafluorides and by the nitrosonium cation in acetonitrile. Preparation and properties of silver(III) hexafluorometalates(V), 461–4
 Polyhedral rearrangements in undeca- and dodeca-nuclear Group 1B metal osmium clusters, 505–10
 Reaction of $[\text{Rh}_2\{\mu\text{-1,8-(NH)}_2\text{C}_{10}\text{H}_6\}(\text{CO})_2(\text{PPh}_3)_2]$ with $[\text{M}(\text{PPh}_3)]^+$ ($\text{M} = \text{Au, Ag, or Cu}$): formation of unusual 44-electron Rh_2M clusters. Crystal structure of $[\text{Rh}_2\{\mu\text{-Au}(\text{PPh}_3)\}\{\mu\text{-1,8-(NH)}_2\text{C}_{10}\text{H}_6\}(\text{CO})_2(\text{PPh}_3)_2]\text{ClO}_4\cdot 2\text{CH}_2\text{Cl}_2$, 1249–52
 Synthesis, structure, and reactivity of the heteronuclear clusters $[\text{Os}_8\text{H}(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}]$ and $[\text{Os}_5(\text{CO})_{15}\{\text{M}(\text{PR}_3)\}_2]$ ($\text{M} = \text{Cu, Ag, or Au; R = Ph or Me}$; the X-ray crystal structure of $[\text{Os}_8\text{H}(\text{CO})_{15}\{\text{Au}(\text{PPh}_3)\}]$), 1421–6
 Kinetics of the uncatalysed and silver(I)-catalysed oxidation of hydrogen peroxide by [ethylenebis(biguaniide)]silver(III) ion in acid perchlorate media, 1645–50
 Bi-, tetra-, and poly-nuclear complexes of gold(I) and silver(I) with bis(diphenylphosphino)-methanide or -amide acting as six-electron donors, 1883–6

The heteronuclear cluster chemistry of the Group 1B metals. Part 12. Variable temperature nuclear magnetic resonance spectroscopic studies of the dynamic behaviour of the mixed-metal cluster compounds $[\text{M}_2\text{Ru}_4(\mu_3\text{-H})_2(\mu\text{-L})(\text{CO})_{12}]$ [$\text{M} = \text{Cu or Ag, L} = \text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2$ ($n = 1\text{--}6$) or *cis*- $\text{Ph}_2\text{PCH}=\text{CHPPh}_2$; $\text{M} = \text{Cu, L} = \text{Ph}_2\text{As}(\text{CH}_2)_n\text{PPH}_2$ ($n = 1$ or 2)] in solution, 2039–48

Insertion of CS_2 into Ag-S bonds. Preparations and crystal structures of $[[\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}_3^{\text{i}}\text{-2,4,6})_4\cdot\text{CHCl}_3]_n]$ and its insertion product with CS_2 , $[[\text{Ag}(\text{S}_2\text{CSC}_6\text{H}_2\text{Pr}_3^{\text{i}}\text{-2,4,6})_6]\cdot 8\text{CHCl}_3]$, 2297–302

SILYL

- Trit(*n*-cyclopentadienyl)uranium(IV) silyl and siloxide compounds. Crystal structure of $[\text{U}(\eta^5\text{C}_5\text{H}_5)_3(\text{OSiPh}_3)]$. Insertion of isocyanide into a uranium–silicon bond, 677–82
 Isopropylsilylamine and disilyl-t-butylamine: preparation, spectroscopic properties, and molecular structure in the gas phase, determined by electron diffraction, 779–84

SILYLMETHYL

- Tris(trimethylsilyl)methyl and tris(dimethylphenylsilyl)methyl derivatives of boron. Crystal structures of dihydroxy[tris(trimethylsilyl)methyl]borane and of the lithium–boron complex $[(\text{MeOH})_2\text{Li}(\mu\text{-OMe})_2\text{B}(\text{OMe})_2]$, 447–52

SODIUM

- The equilibrium of strontium ion exchange of sodium titanate, $\text{Na}_4\text{Ti}_9\text{O}_{20}\cdot x\text{H}_2\text{O}$, 101–4
 The nature of a sodium–crown interaction studied by sodium-23 nuclear magnetic resonance and by crystal structure determination of a complex with co-ordinated solvent, 203–10

SODIUM-23

- The nature of a sodium–crown interaction studied by sodium-23 nuclear magnetic resonance and by crystal structure determination of a complex with co-ordinated solvent, 203–10

SOLVENT EFFECT

- A kinetic and mechanistic study of the solvolysis of aryltellurium trihalides in organic solvents, 457–60
 Solvent influence on trifluoromethylation reactions with $\text{Cd}(\text{CF}_3)_2$: one-pot synthesis of trifluoromethyl iron complexes, 739–44
 Reactivity studies on the hydrolysis of iron(II) complexes of substituted phenanthrolines in dioxane–water solvents, 1105–8
 Vibrational study of ionic association in aprotic solvents. Part 12. Isothiocyanate M(NCS) complexes of non-transition-metal ions, 2193–8

SOLVENT EXTRACTION

- Investigations of tin(II) halide aqueous solutions and solvent extracts by ^{119}Sn nuclear magnetic resonance spectroscopy: characterisation of the trihalogenostannate(II) anions $[\text{SnX}_3]^-$, $[\text{SnX}_2\text{Y}]^-$, and $[\text{SnX(Y)Z}]^-$ ($\text{X, Y, or Z} = \text{Cl, Br, or I}$), 2223–8

SOLVOLYSIS

- A kinetic and mechanistic study of the solvolysis of aryltellurium trihalides in organic solvents, 457–60

STABILITY

- Polymeric thiolate complexes in solution. Stability constants of the cadmium(II) complexes of 2-aminopropane-1-thiol, 13–8
 The unusual co-ordination ability of vasopressin-like peptides: potentiometric and spectroscopic studies of some copper(II) and nickel(II) complexes, 173–8
 Stabilisation of trivalent nickel through 1:2 co-ordination by cyclic terdentate ligands $\text{CH}_2\text{CH}_2\text{NH}(\text{CH}_2)_2\text{NH}(\text{CH}_2)_2\text{X}$ ($\text{X} = \text{NH, O, or S}$), 229–32
 The co-ordination chemistry of divalent cobalt, nickel, and copper. Part 13. Characterization and stability constants of copper(II) species isolated from mixed-ligand solutions; crystal structure of the complex (pyridine-2,6-dicarboxylato)(*N,N,N',N'*-tetramethyl-1,3-diaminopropane)copper(II), 253–8
 Thermodynamic and electrochemical behaviour of lanthanide cryptates in *N,N*-dimethylformamide, 367–70
 Possible biological activity in proton, metal, and aminohydroxamic acid equilibria. Protonation and complex formation reactions between *N*-hydroxy-D-asparagine and cobalt(II), nickel(II), copper(II), and hydrogen ions in aqueous solution (1988, 953), 383
 The hydrolysis of metal ions. Part 11. The ionic strength dependence of gallium(III), 399–402
 Interligand interaction in mixed amino acid complexes, 567–8
 Spectrophotometric study of complex formation in copper(II) mono-, di-, and tri-ethanolamine systems, 569–74
 Potentiometric and electron spin resonance investigation on the interactions between adenosine 5'-triphosphate (ATP) and triphosphate ligands with vanadyl ion. Synthesis and characterization of solid vanadyl-ATP compounds from acidic aqueous solution, 575–80
 Spectrophotometric and calorimetric studies on the formation of binary (2,2'-bipyridine)nickel(II) and ternary (2,2'-bipyridine)chloronickel(II) complexes in *N,N*-dimethylformamide, 655–60
 Formation in solution, synthesis, and electrochemical study of oxalato complexes of *N,N'*-ethylenabis(salicylideneiminato)-chromium(III) and -iron(III); crystal structures of piperidinium [*N,N'*-ethylenabis(salicylideneiminato)][oxalato- $O^1\text{O}^2$]-chromate(III) and ferrate(III), 729–38
 Thermodynamics of hydroxy complex formation of dialkyltin(IV) ions in aqueous solution, 773–8
 Complexes of aminophosphonates. Part 3. Copper(II) complexes of several monophosphono and diphasphono dipeptides, 1053–8
 Zinc(II) complexes of 4,7-diazadecane-1,10-diamine, 4,8-diazaundecane-1,11-diamine, and tetraethylenetetrapamine: a carbon-13 nuclear magnetic resonance study, 1171–8
 Formation of mixed-ligand complexes of cadmium(II) halide with tri-n-octylphosphine oxide and pyridine or substituted pyridines in 1,2-dichloroethane, 1259–64
 Composition, stability, and lability of copper(II) dipeptide complexes, 1293–8
 Thermodynamic and spectrophotometric investigation of complex formation between hydrogen, cobalt(II), nickel(II), and copper(II) ions and 2-amino-*N*-hydroxypropanamide in aqueous solution, 1299–310
 Thermal and hydrothermal stability of SAPO-5 and SAPO-37 molecular sieves, 1361–2
 Co-ordination of nickel(II) ions by angiotensin II and its peptide

STABILITY (contd)

- fragments. A potentiometric, proton nuclear magnetic resonance and circular dichroism spectroscopic study, 1471–6
 Mixed-metal complexes in solution. A potentiometric and spectrophotometric study of copper(II)–nickel(II)– δ -hydroxylysine complexes in aqueous solution, 1745–50
 Complex-forming properties of L- α -alaninehydroxamic acid (2-amino-N-hydroxypropanamide), 2247–52
 Copper anti-inflammatory drugs in rheumatoid arthritis. Part 2. A potentiometric and spectroscopic study of copper(II) polyaminodicarboxylate complexes, 2429–34

STACKING

- Chemistry of subvalent main group metals with dithiolates. Part 2. X-Ray structural, thermal, and theoretical investigations on the novel one-dimensional stacking structure $K_2[Pb\{S_2C=C(CN)_2\}_2] \cdot 2H_2O$, 627–30

Stacked planar molecules in antiferromagnetic di-iso thiocyanato(bis(thiourea)chromium(II), a new structural type for chromium(II), 1631–4

STEREOCHEMISTRY

- Synthesis and stereochemical studies of η^3 -allyl palladium(II) complexes containing a chiral chelating ligand, 161–8
 Syntheses and dynamic stereochemistry of methyl- and phenyl-carbonyl heterobinuclear metal–metal bonded complexes, 379–82
 Relation between the torsion angles of acetate chelate rings and the deuteron nuclear magnetic resonance chemical shifts for polyamine-N-carboxylatochromate(III) complexes, 1947–50

STRONTIUM

- The equilibrium of strontium ion exchange of sodium titanate, $Na_4Ti_9O_{20} \cdot xH_2O$, 101–4

SUBSTITUTION

- The reaction of hydroxo-species of palladium(II) with bidentate nucleophiles, 375–8
 A kinetic study of the transmission of ligand effects across the quadruple bond in $[Re_2Cl_8]^{2-}$, 767–72
 Variable-temperature and -pressure proton nuclear magnetic resonance studies of the rates and mechanisms of N,N' -dimethylformamide and acetonitrile exchange with the square-pyramidal complexes mono(solvato){3,7,11-tribenzy-3,7,11,17-tetra-azabicyclo[11.3.1]heptadeca-1(17),13,15-triene}nickel(II), 873–8

Composition, stability, and lability of copper(II) dipeptide complexes, 1293–8

Substitution reactions of the zerovalent iron complexes $[Fe(\eta-C_6H_6)(PMes_2)]$ and $[Fe(\eta-C_6H_6)(dmpe)]$ (dmpe = $Me_2PCH_2CH_2PMe_2$), 1385–92

Substitution reactions on sterically hindered square-planar *trans*- $[NiBr(R)(PR'_3)_2]$ (R = aryl) complexes. Effects of the substituents of the aryl ligand, 1669–74

Kinetics of formation, dissociation, and isomerization reactions in complexes of cyanopyridines and benzonitrile with pentacyanoruthenate(II), 1973–8

SUGAR

- Isotopic multiplets in the carbon-13 nuclear magnetic resonance spectra due to partial deuteration of co-ordinated NH_2 and NH groups of cobalt(III) complexes containing an N-glycoside. Importance of sugar ring oxygen atoms in the long-range isotope shifts, 1241–8

SULPHATE

A reinvestigation of the structures of organotin sulphates and chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74

The structure of 18-molybdodisulphate(vi) (4^-) ion in $(NEt_4)_4S_2Mo_{18}O_{62}CH_3CN$, 1491–8

SULPHIDE

- (Diphenylphosphinomethyl)diphenylphosphine sulphide ($Ph_2PCH_2PPh_2S$) and its methanide anion $Ph_2PCHPPh_2S^-$ as ligands in organogold chemistry. X-Ray crystal structure of *cis*- $[Au(C_6F_5)_2(PPh_2CHPPh_2S)]$, 73–8

Chromium chalcogen complexes. Insertion of elemental sulphur into $[Cr(cp)(CO)_2Se]$ and $[Cr(cp)_2(CO)_4Se_2]$ ($cp = \eta-C_5H_5$), 431–8

Reaction of elemental sulphur with $(5H,14H-6,8,15,17-$ -tetramethyldibenzo[b,i][1,4,8,11]tetra-azacyclotetradecinato)iron(II); synthesis and X-ray structure of a μ -sulphido iron(III) complex, 551–4

Solid-state phosphorus-31 nuclear magnetic resonance spectroscopy of phosphorus sulphides, 809–14

Comparative study on the multielectron reduction of $HOC_2H_4N_3$ catalyzed by $[NBu^+4]_2[MoFe_3S_4(SPh)_3(o-O_2C_6Cl_4)]_2^-$ and $[NBu^+4]_2[Fe_4S_4(SPh)_4]$ -modified glassy carbon electrodes, 1547–54

Reactions of cyclopalladated compounds. Part 21. Various examples of sulphur-assisted intramolecular palladation of aryl and alkyl groups, 1715–20

SULPHIMIDE

- Synthesis of sulphimide complexes of palladium and platinum. Crystal and molecular structure of *cis*-dichloro[S,S-dimethyl- N -(2-pyrimidinyl)sulphimide](triphenylphosphine)palladium(II), 497–504

SULPHINYLAMIDE

Preparation of thionylimide complexes of titanium, zirconium, and hafnium. Crystal structure of $[Zr(cp)(\eta-C_5Me_5)(NSO)_2]$, 1815–8

SULPHITE

Oxidation of a $Pt_2(\mu-SO_2)$ moiety to $\mu-SO_3$ and the crystal structure of $[Pt_2(\mu-SO_3)(\mu-SO_2)(C_8H_{12})_2]$, 557–60

SULPHONAMIDE

Sulphonamide nitrogen-containing, nitrogen protected amino acids interacting with the Cd^{2+} ion. A cadmium-113 nuclear magnetic resonance and potentiometric study. X-Ray diffraction of $[Cd(ts-\beta\text{-alaO})(H_2O)_4]$, 1345–50

SULPHONE

- Metallacyclic complexes. Part 2. Synthesis of metallathietane-3,3-dioxide complexes of platinum(II) and palladium(II); the crystal structures of $[Pt\{CH(COPh)S(O)_2CH(COPh)\}-\langle PPh_3\rangle_2 \cdot 2CH_2Cl_2$ and $[Pd\{CH(COPh)S(O)_2CH(COPh)\}-\langle PMePh_2\rangle_2 \cdot CH_2Cl_2 \cdot H_2O$, 259–66

SULPHOXYLATE

Oxidation of a $Pt_2(\mu-SO_2)$ moiety to $\mu-SO_3$ and the crystal structure of $[Pt_2(\mu-SO_3)(\mu-SO_2)(C_8H_{12})_2]$, 557–60

SULPHUR

Metal complexes (Co^{3+} , Fe^{3+} , Mn^{3+}) of the quinquedentate salen analogue, 2-methyl-4-benzylthio- N,N' -butane-1,2-diybis(salicylideneimine): crystal structures and sulphur coordination effects, 179–84

Chromium chalcogen complexes. Insertion of elemental sulphur into $[Cr(cp)(CO)_2Se]$ and $[Cr(cp)_2(CO)_4Se_2]$ ($cp = \eta-C_5H_5$), 431–8

Reaction of elemental sulphur with $(5H,14H-6,8,15,17-$ -tetramethyldibenzo[b,i][1,4,8,11]tetra-azacyclotetradecinato)iron(II); synthesis and X-ray structure of a μ -sulphido iron(III) complex, 551–4

Solid-state phosphorus-31 nuclear magnetic resonance spectroscopy of phosphorus sulphides, 809–14

Gas-phase molecular structure of NN' -bis(trimethylsilyl)sulphur di-imide, determined by electron diffraction, 859–62

New metal-sulphur–nitrogen compounds from reactions in liquid ammonia. The X-ray structures of *trans*-bis(acetophenone dimethylhydrazone- N^2)dichloropalladium(II) and [di(azathien)-1-yl- $S^1N^2\langle 2-(hydrazonoethoxy)phenyl\rangle$]palladium(II), 1179–86

Contributions to the chemistry of boron. Part 194. Synthesis and characterization of mono-, bi-, and tri-cyclic boron–nitrogen–sulphur ring systems. Crystal structure of 2,7,9,14-tetrakis(diethylamino)-3,5,6,10,12,13-hexathia-1,8-diazab-2,4,7,9,11,14-hexaboratricyclo[9.3.0.0^{4,8}]tetradecane, 1311–20

An *ab initio* self-consistent field molecular-orbital study of novel stereoelectronic effects in linear $R_2S_3N_2$ and cyclic XS_3N_2 systems, 1609–12

Direct insertion of a nitrogen atom into the S–S bond of a 1,2,3,5-dithiadiazole ring in a direct current nitrogen glow discharge, and X-ray crystal structure of 4-methyl-1,2,3,5-dithiadiazole, 1705–14

Preparation and properties of $[M(S_2N_2H)Cl(L)]$ ($M = Pt$, $L = PMe_2Ph$ or C_5H_4 ; $M = Pd$, $L = PMe_2Ph$), $[Pt(S_2N_2)Cl(PMe_2Ph)]$, and corresponding ^{15}N -enriched complexes, 2179–82

SULPHUR DIOXIDE

Oxidation of a $Pt_2(\mu-SO_2)$ moiety to $\mu-SO_3$ and the crystal structure of $[Pt_2(\mu-SO_3)(\mu-SO_2)(C_8H_{12})_2]$, 557–60

TANTALUM

Hydrogen scrambling processes in $[Ta(\eta-C_5Me_5)(\eta-C_5H_5)\langle (\mu-H_2BH_2)\rangle]$: a surprising observation, 2133–8

TECHNETIUM

Complexes of technetium, rhenium, and rhodium with sexidentate Schiff-base ligands, 2115–20

TELLURIUM

Metalla-heteroborane chemistry. Part 3. Synthesis of [2,2-(PR_3)₂-1,2-TePtB₁₀H₁₀] ($\text{R}_3 = \text{Et}_3, \text{Bu}^n_3$, or Me_2Ph), their characterisation by nuclear magnetic resonance spectroscopy, and the crystal and molecular structure of [2,2-(PEt_3)₂-1,2-TePtB₁₀H₁₀] (1988, 2555), 383

A kinetic and mechanistic study of the solvolysis of aryltellurium trihalides in organic solvents, 457-60

Synthesis and reactions of 1,6-bis(2-butyltellurophenyl)-2,5-diazahexa-1,5-diene and related compounds. The crystal and molecular structures of 2-(butyl dichlorotelluro)benzaldehyde and bis[2-(hydroxyiminomethyl)phenyl] ditelluride, 1697-704

TERPYRIDINE

Crystal structure and magnetic properties of 1-aqua- μ -hydroxo-1,2,2-tris(perchlorato)-1,2-bis(2,2'-6,2''-terpyridine)dicopper(II), 237-42

TETRA-AZACYCLOTETRADECANE

6,13-Diamino-6,13-dimethyl-1,4,8,11-tetra-azacyclotetradecane, L⁷, a new, potentially sexidentate polyamine ligand. Variable coordination to cobalt(III) and crystal structure of the complex [Co(L⁷)Cl₂[ClO₄]], 1059-66

TETRAHYDROBORATE

Hydrogen scrambling processes in [Ta(η -C₅Me₅) $(\eta$ -C₅H₅) $\{\langle\mu\rangle\text{H}_2\text{BH}_2\}]:$ a surprising observation, 2133-8

TETRAHYDROFURAN

Effect of axially bound anions on the electroreduction of tin(IV) porphyrins in tetrahydrofuran, 1531-6

Chelate-stabilised tetrahydrofuran complexes of platinum(II). Crystal structure of [PtCl(PPh₂PCH₂C₄H₈O)-(Ph₂PCH₂C₄H₈O)]PF₆], 2069-72

TETRAHYDROTHIOPHENE

Synthesis and reactivity of trinuclear complexes of platinum containing the single bridging ligand SC₄H₈. Molecular structure of [NBu₂]₂[trans-PtCl₂{ $\langle\mu$ -SC₄H₈}Pt(C₆F₅)₃}₂], 1011-6

TETRAZIACYCLOHEXADECANE

Platinum metal thioether macrocyclic complexes: synthesis, electrochemistry, and single-crystal X-ray structures of cis-[RhCl₂L²]PF₆ and trans-[RhCl₂L³]PF₆ ($L^2 = 1,4,8,11$ -tetra-azacyclotetradecane, L³ = 1,5,9,13-tetra-azacyclohexadecane), 1675-80

TETRAZIACYCLOTETRADECANE

Platinum metal thioether macrocyclic complexes: synthesis, electrochemistry, and single-crystal X-ray structures of cis-[RhCl₂L²]PF₆ and trans-[RhCl₂L³]PF₆ ($L^2 = 1,4,8,11$ -tetra-azacyclotetradecane, L³ = 1,5,9,13-tetra-azacyclohexadecane), 1675-80

TETRAZIAFULVALENIUM

Tetra-aziafulvalenium salts of planar Pt^{II}, Pd^{II}, and Cu^{III} 1,2-dithio-oxalato-S,S' anions. Synthesis, chemistry and molecular structures of bis(tetra-aziafulvalenium) bis(1,2-dithio-oxalato-S,S') palladate(II), [ttf]₂[Pd(S₂C₂O₂)₂], and of bis(tetra-aziafulvalenium) tetrathia-fulvalene bis(1,2-dithio-oxalato-S,S') platinum(II), [ttf]₂[Pt(S₂C₂O₂)₂], 719-28

THALLIUM

Monocyclopentadienyl pentafluorothiophenolate complexes of molybdenum and their interactions with thallium(I) ions; dynamic nuclear magnetic resonance studies and crystal and molecular structures of [TlMo(SC₆F₅)₂L₂(cp)] (L = CO or SC₆F₅; cp = η^5 -C₅H₅), 991-1002

Co-ordination compounds of indium. Part 47. Indium(I) and thallium(I) derivatives of alkanedithiols, 1759-64

Synthetic, spectroscopic, and X-ray studies on methylmercury(II) and dimethylthallium(III) complexes with cyclopentanone thiosemicarbazone, 1787-92

Synthesis of arylthallium complexes using organomercury compounds. Crystal and molecular structures of [NMe₄][Tl(C₆H₄N-NPh-2)Cl₃] and [Tl(C₆H₄CH₂NMe₂-2)Cl₂], 2241-6

TERMODYNAMICS

Thermodynamics of the formation of metal complexes of 1,4,10-trioxa-7,13-diazacyclopentadecane-*N,N'*-diacetic acid and of 1,4,10,13-tetraoxa-7,16-diazacyclo-octadecane-*N,N'*-diacetic acid, 133-8

The nickel(II) bromide system in dimethyl sulphoxide: a detailed study of the influences of temperature and mole ratio, 325-30

Thermodynamic and electrochemical behaviour of lanthanide cryptates in *N,N*-dimethylformamide, 367-70

Spectrophotometric and calorimetric studies on the formation of

binary (2,2'-bipyridine)nickel(II) and ternary (2,2'-bipyridine)chloronickel(II) complexes in *N,N*-dimethylformamide, 655-60

Thermodynamics of hydroxy complex formation of dialkyltin(IV) ions in aqueous solution, 773-8

Mixed-metal complexes in solution. A potentiometric and spectrophotometric study of copper(II)-nickel(II)- δ -hydroxylysine complexes in aqueous solution, 1745-50

THERMOLYSIS

Mechanism and energetics of reductive elimination of ethane from some tetramethylplatinum(IV) complexes, 2121-6

THIADIAZOLE

Preparation and crystal structure of a binuclear compound of methylmercury(II) containing bismuthiol I as a bridging ligand, 1409-12

THIOACYL

Insertion of phenylethyne-thiolate or -selenolate into metal carbene bonds: synths for co-ordinated thioacyl or selenoacyl anions and for anionic thiocarbene or selenocarbene complexes, 1565-78

THIOCARBONATE

Insertion of CS₂ into Ag-S bonds. Preparations and crystal structures of [{Ag(SC₆H₂Pr^{1,3})₂,2,4,6}]₄[CHCl₃]_n and its insertion product with CS₂, [{Ag(S₂CSC₆H₂Pr^{1,3}-2,4,6)₂}Ag(SC₆H₂Pr^{1,3}-2,4,6)₆]₈CHCl₃, 2297-302

THIOCYANATE

Crystal structure, spectroscopic and magnetic properties of the complex [Cu(paphy)(NCS)(SCN)] (paphy = pyridine-2-carbaldehyde 2'-pyridylhydrazone). An unusual di- μ -thiocyanato-*N* bridged copper(II) dimer, 53-6

Lewis-base adducts of lead(II) thiocyanate-1,10-phenanthroline(1/2). Crystal structure determinations, 595-600

Charge density in the [Mo(NCS)₆]³⁻ ion in Cs₃Mo(NCS)₆, 1779-86

A barium dithiocyanate complex with an unusual macrocycle containing pendant arms, the amine precursor, and related copper(II) diperchlorate complexes: synthesis and structure, 1993-2002

Unusual chelating properties of the ligand 1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-dithiaoctane (bddo). Crystal structures of Ni(bddo)(NCS)₂, Zn(bddo)(NCS)₂ and Cd₂(bddo)(NCS)₄, 2309-14

Crystal and molecular structure of a lattice adduct, HgPh₂[2(Hg(SCN)₂(PPh₃)₂]: an unusual case of two- and four-co-ordinate mercury in the same unit cell, 2339-42

THIOETHER

Co-ordination chemistry of higher oxidation states. Part 29.

Cobalt(III) complexes of bi- and tetra-thioethers: synthesis, properties, and cobalt-59 nuclear magnetic resonance data, 453-6

A dynamic nuclear magnetic resonance study of trimethylplatinum(IV) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part 1. Low-temperature solution properties, and X-ray crystal structure of [PtIMe₃[(MeS)₂CHCH(SMe)₂]], 701-10

A dynamic nuclear magnetic resonance study of trimethylplatinum(IV) halide complexes of 1,1,2,2-tetrakis(methylthio)ethane. Part 2. High-temperature fluxionality, 711-8

Platinum metal complexes of potentially chelating alkene-thioether and alkene-selenoether ligands: synthesis and dynamic nuclear magnetic resonance study of [MX₂{MeE(CH₂)_nCH=CH₂}]ⁿ (M = Pt or Pd; X = Cl, Br, or I; E = S or Se; n = 2 or 3) and the X-ray structure of cis-dibromo(2-thia-6-heptene)platinum(II), [PtBr₂{MeS(CH₂)₃CH=CH₂}]ⁿ, 985-90

Platinum metal complexes of potentially chelating alkene-thioether and selenoether ligands: the synthesis and dynamic nuclear magnetic resonance study of [MX₂{E[(CH₂)_nCR=CR₂]}]ⁿ (M = Pt or Pd; X = Cl, Br, or I; E = S or Se; n = 2 or 3; R = H or Me) and the X-ray crystal structure of cis-di-iodo(5-thianona-1,8-diene)platinum(II), [PtI₂{S[(CH₂)₂CH=CH₂]}₂]ⁿ, 2315-22

THIOL

Polymeric thiolate complexes in solution. Stability constants of the cadmium(II) complexes of 2-aminopropane-1-thiol, 13-8

Cationic dinuclear rhodium complexes as catalyst precursors for the hydroformylation of alkenes, 1579-82

Co-ordination compounds of indium. Part 47. Indium(I) and thallium(I) derivatives of alkanedithiols, 1759-64

THIOLATE

Rhodium complexes of the binucleating ligands pyridine-2-thiolate

THIOLATE (contd.)

and benzothiazole-2-thiolate. Crystal structures of $[\{\text{Rh}(\mu\text{-SC}_5\text{H}_4\text{N})(\text{CO})_2\}_2]$ and $[\{\text{Rh}(\mu\text{-SC}_5\text{H}_4\text{N})(\text{tfbb})\}_2]\cdot\text{Me}_2\text{CO}$ (tfbb = tetrafluorobenzobarrelene), 25–32

Comparison of the structures and properties of $[\text{N}(\text{C}_2\text{H}_5)_4][\text{Pt}\{\text{S}_2\text{C}_2(\text{CN})_2\}_2]$ and $[\text{H}_3\text{O}]_x[\text{NH}_4]_{1-x}[\text{Pt}\{\text{S}_2\text{C}_2(\text{CN})_2\}_2]\cdot 2 - x\text{H}_2\text{O}$, 61–6

The crystal structures of $\text{K}[\text{Pd}\{\text{S}_2\text{C}_2(\text{CN})_2\}_2]\cdot\text{H}_2\text{O}$ and $[\text{NH}_4]_x[\text{Pd}\{\text{S}_2\text{C}_2(\text{CN})_2\}_2]\cdot\text{H}_2\text{O}$, 67–72

Synthesis and spectroscopic characterisation of $[\text{MoO}(\text{SR})_4]^-$ ($\text{R} = \text{C}_6\text{H}_5, \text{C}_{10}\text{H}_7, 4\text{-CH}_3\text{C}_6\text{H}_4, 4\text{-ClC}_6\text{H}_4, 3\text{-ClC}_6\text{H}_4, 4\text{-FC}_6\text{H}_4, 2,6\text{-Cl}_2\text{C}_6\text{H}_3$, or C_6F_5) complexes and $[\text{MoO}(\text{SC}_6\text{F}_5)_4]^2-$, 413–8

Thiolate complexes of osmium(III) with tertiary phosphines and their dicarbonyl derivatives of osmium(II): X-ray crystal structure of *trans,trans,trans*- $[\text{Os}(\text{SC}_6\text{F}_5)_2(\text{CO})_2(\text{PEt}_3\text{Ph})_2]$, 419–24

Studies on gold(II) complexes with hard and soft donor ligands.

Part 1. Complexes with *o*-aminobenzenethiol (1988, 2607), 555

Chemistry of subvalent main group metals with dithiolates. Part 2. X-Ray structural, thermal, and theoretical investigations on the novel one-dimensional stacking structure

$\text{K}_2[\text{Pb}\{\text{S}_2\text{C}=\text{C}(\text{CN})_2\}_2]\cdot 2\text{H}_2\text{O}$, 627–30

Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten $[\text{M}\{\eta^3\text{-SC}(\text{CF}_3)=\text{C}(\text{CF}_3)\text{H}\}\{\text{CF}_3\text{C}=\text{CCF}_3\}(\eta^3\text{-C}_5\text{H}_5)]$ ($\text{M} = \text{Mo}$ or W). Their relevance to the mechanism of metal-promoted isomerisation of carbon–carbon double bonds, 1027–36

Preparation and crystal structure of a binuclear compound of methylmercury(II) containing bismuthiol I as a bridging ligand, 1409–12

Insertion of phenylethyne-thiolate or -selenolate into metal–carbene bonds: synths for co-ordinated thioacyl or selenoacyl anions and for anionic thiocarbene or selenocarbene complexes, 1565–78

Co-ordination compounds of indium. Part 47. Indium(II) and thallium(I) derivatives of alkanedithiols, 1759–64

Crystal structure and magnetic properties of the heterobinuclear complex $[(\text{CH}_3\text{CN})\text{LMi}(\text{mnt})\text{Cu}(\text{mnt})]\cdot\text{CH}_3\text{CN}$ ($\text{L} = \text{tetrabenzo}[b,f,j,n][1,5,9,13]\text{tetra-azacyclohexadecine}$ and $\text{mnt} = \text{maleonitriledithiolate}$), 2095–8

Spectral intensity and charge distributions in the tetrahedral chromophores, bis(3-amino-2-methylpropane-2-thiolato)- and diacetatobis(ethylenethiourea)-cobalt(II), 2109–14

Interaction of benzo-1,3-thiazoline-2-thione and related ligands with copper(II) salts and the single-crystal X-ray structure of $\mu\text{-}[\text{1,2-bis(diphenylphosphino)ethane]-bis(benzo-1,3-thiazole-2-thiolato-S)[1,2-bis(diphenylphosphino)ethane]copper(II)}$, 2167–72

Iron carbene complexes. Part 3. Aminolysis of di-iron alkoxy carbene complexes with ammonia and dimethylamine. X-Ray crystal structure of $[\text{Fe}_2(\mu\text{-SPh})_2(\text{CO})_5\{\text{C}(\text{NMMe}_2)\text{Ph}\}]$, 2199–204

Insertion of CS_2 into Ag-S bonds. Preparations and crystal structures of $[(\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}_3^{\text{i}})^{\text{2},4,6})_4\cdot\text{CHCl}_3]_n$ and its insertion product with CS_2 , $[\text{Ag}(\text{S}_2\text{CSC}_6\text{H}_2\text{Pr}_3^{\text{i}})^{\text{2},4,6})_2\text{Ag}(\text{SC}_6\text{H}_2\text{Pr}_3^{\text{i}})^{\text{2},4,6}]_6\cdot 8\text{CHCl}_3$, 2297–302

THIONE

The interaction of 1-methylimidazoline-2(*3H*)-thione with copper(II) salts, 39–42

Crystal structure and mass spectrometry of dibromodimethyl[1-methyl-2(*3H*)-imidazolinethione-*S*]tin(IV), 927–30

Blue copper models. Spectroscopic and electrochemical studies of copper(II) complexes with new ligand systems containing sulphur and nitrogen donor atoms, 1161–70

Interaction of benzo-1,3-thiazoline-2-thione and related ligands with copper(II) salts and the single-crystal X-ray structure of $\mu\text{-}[\text{1,2-bis(diphenylphosphino)ethane]-bis(benzo-1,3-thiazole-2-thiolato-S)[1,2-bis(diphenylphosphino)ethane]copper(II)}$, 2167–72

THIOSEMICARBAZONE

Thiosemicarbazones as co-ordinating agents. Part 3. Synthesis, spectroscopic characterisation, and X-ray structure of methyl pyruvate thiosemicarbazone hemihydrate, chloro(ethyl pyruvate thiosemicarbazone)copper(II) (green form), and chloro(pyruvic acid thiosemicarbazone)copper(II) dihydrate (blue form), 361–6

Synthetic, spectroscopic, and X-ray studies on methylmercury(II) and dimethylthallium(III) complexes with cyclopentanone thiosemicarbazone, 1787–92

THIOSULPHATE

Synthesis and properties of the bis(μ -thiosulphato-*S*)-bis(dinitrosylferrate)(2⁻) anion. Crystal and molecular structure

of bis[bis(triphenylphosphoranylidene)ammonium] bis(μ -thiosulphato-*S*)-bis(dinitrosylferrate), 2061–4

THIOUREA

Stacked planar molecules in antiferromagnetic di-isothiocyanatobis(thiourea)chromium(II), a new structural type for chromium(II), 1631–4

Spectral intensity and charge distributions in the tetrahedral chromophores, bis(3-amino-2-methylpropane-2-thiolato)- and diacetatobis(ethylenethiourea)-cobalt(II), 2109–14

TIN

Synthesis and structural studies on silver tin complex salts with bis(diphenylarsino)methane and bis(diphenylphosphino)methane, 247–52

A reinvestigation of the structures of organotin sulphates and chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74

Synthesis, infrared, and tin-119 Mössbauer spectroscopic characterization of seven-co-ordinate diorganotin(IV) adducts with 2,6-diacetylpyridine acylhydrazone, including the X-ray crystal structure of SnEt_3 (dapt) [$\text{H}_2\text{dapt} = 2,6\text{-diacetyl}[\text{pyridine}-\text{bis}(2\text{-thenoylhydrazone})]$], 289–94

One-electron transfer processes in the reaction of tin(II) halides with substituted *o*-quinones; crystal structure of bis(3,5-di-t-butylcatecholate)tin-1,10-phenanthroline-dimethylformamide (1/1/2), 439–46

The binding of trialkyltin(IV) moieties to rat haemoglobin, and the structure of model systems, studied by tin-119 Mössbauer spectroscopy, 519–26

Thermodynamics of hydroxy complex formation of dialkyltin(IV) ions in aqueous solution, 773–8

High-resolution solid-state tin-119 nuclear magnetic resonance spectroscopy of ternary tin oxides, 843–8

Crystal structure and mass spectrometry of dibromodimethyl[1-methyl-2(*3H*)-imidazolinethione-*S*]tin(IV), 927–30

Effect of axially bound anions on the electroreduction of tin(IV) porphyrins in tetrahydrofuran, 1531–6

Phenyltin trichloroacetate complexes: preparations and structures, 1589–98

Synthesis and properties of trimethyl(trifluorosilyl)stannane, 1879–82

Stereochemical role of lone pairs in main-group elements. Part 4.

The crystal structure at 120 K of bis(1,4,7,10,13-pentaoxacyclopentadecane)tin(II) bis(trichlorostannate(II)), confirming the sterically inactive lone pair, 2155–60

Investigations of tin(II) halide aqueous solutions and solvent extracts by ¹¹⁹Sn nuclear magnetic resonance spectroscopy: characterisation of the trihalogenostannate(II) anions $[\text{SnX}_3]^-$, $[\text{Sn}_2\text{X}_2\text{Y}]^-$, and $[\text{SnX(Y)Z}]^-$ (X, Y , or $\text{Z} = \text{Cl}, \text{Br}$, or I), 2223–8

TIN-119

A reinvestigation of the structures of organotin sulphates and chromates, including the crystal and molecular structure of bis(trimethyltin) sulphate dihydrate, 267–74

High-resolution solid-state tin-119 nuclear magnetic resonance spectroscopy of ternary tin oxides, 843–8

Investigations of tin(II) halide aqueous solutions and solvent extracts by ¹¹⁹Sn nuclear magnetic resonance spectroscopy: characterisation of the trihalogenostannate(II) anions $[\text{SnX}_3]^-$, $[\text{Sn}_2\text{X}_2\text{Y}]^-$, and $[\text{SnX(Y)Z}]^-$ (X, Y , or $\text{Z} = \text{Cl}, \text{Br}$, or I), 2223–8

TITANIUM

An investigation of halide abstraction by arsenic(III): reactions of AsCl_3 with MCl_2 ($\text{M} = \text{Mg}$ or Zn) and TiCl_4 in donor solvents (MeCN or POCl_3). Formation and crystal structure of the dimeric titanium(IV) compound $[\text{Ti}_2\text{Cl}_6(\text{POCl}_3)_2(\text{PO}_2\text{Cl}_2)_2]$, 57–60

The equilibrium of strontium ion exchange of sodium titanate, $\text{Na}_4\text{Ti}_4\text{O}_{20}\cdot n\text{H}_2\text{O}$, 101–4

Lamellar inorganic ion exchangers. Hydrogen–lithium ion exchange in γ -titanium phosphate, 829–36

N,N'-Ethylenebis(acetylacetoneiminato)titanium complexes: synthesis, crystal structures, and properties of titanium(III) and titanium(IV) derivatives, 953–8

Hydrazido-complexes of titanium(IV) with tris(pyrazolyl)borates. Crystal structures of $[\text{TiLCl}_2(\text{NMeNMe}_2)]$ and $[\text{TiLCl}_2(\text{NPhNH}_2)]$, $\text{L} = \text{tris(pyrazolyl)borate}$, 1413–6

Phenoxy and diphenoxo complexes of titanium(III) and vanadium(IV): synthesis and X-ray structure, 1793–8

Preparation of thionylimide complexes of titanium, zirconium, and hafnium. Crystal structure of $[\text{Zr}(\text{cp})(\eta\text{-C}_5\text{Me}_5)(\text{NSO})_2]$, 1815–8

TITANIUM (contd)

- Lamellar inorganic ion exchangers. H^+ /Cs⁺ ion exchange in γ -titanium phosphate, 1825–30
- Synthesis and crystal structures of three four-membered ring compounds containing PN₂Ti skeletons, 2173–8
- Cyclic metallasiloxane compounds of Ti^{IV} and Zr^{IV}. X-Ray crystal structures of a *cis*-bis(pyridine) bis(cyclosiloxy)titanate and a pyridinelithiated tris(cyclosiloxy)zirconate, 2347–52
- The chemistry of dinitrogen residues. Part 7. Hydrazido(1–)- and *N,N*-dimethylhydroxylaminato(1–)-complexes of titanium(IV), 2389–96
- On the existence of the anion trichloro(η -cyclopentadienyl)titanate(1–). Crystal structure of the salt [Co(C₅H₅)₂][Ti(C₅H₅)Cl₃], 2453–6

TOLYL

- Synthesis and X-ray crystal structure of tetra(2-methylphenyl)molybdenum(IV), Mo(2-MeC₆H₄)₄. Redox chemistry of M(2-MeC₆H₄)₄ compounds of molybdenum, rhenium, ruthenium, and osmium, 2149–54

TRANSITION METAL

- Thermodynamics of the formation of metal complexes of 1,4,10-trioxa-7,13-diazacyclopentadecane-*N,N'*-diacetic acid and of 1,4,10,13-tetraoxa-7,16-diazacyclo-octadecane-*N,N'*-diacetic acid, 133–8
- Syntheses and dynamic stereochemistry of methyl- and phenyl-carbonyl heterobinuclear metal–metal bonded complexes, 379–82
- Oxidation of silver metal and silver(I) cations by molybdenum, tungsten, and uranium hexafluorides and by the nitrosonium cation in acetonitrile. Preparation and properties of silver(III) hexafluorometalates(V), 461–4
- Bimetallic compounds of *trans*-cyclohexane-1,2-diamine-*NN'N'*-tetra-acetate (cdta). Part 3. Structural and magnetic characterization of the dinuclear [M(OH₂)₅][M'(cdta)]·H₂O (M,M' = Ni,Ni; Mn,Ni; Mn,Cu; Co,Ni; or Co,Cu) and the tetrานuclear [M(OH₂)₄][M'(cdta)(OH₂)₂]·4H₂O (M,M' = Zn,Zn; Zn–Co; Co,Co; or Mn,Co) complexes, 863–72
- Some transition-metal complexes of 1,3,2-dithiarsolane, 885–8
- Preparation and characterisation of linear mixed-metal trinuclear carboxylate complexes of general formula [M₂M'(O₂CR)₆(C₉H₇N)₂]; crystal structure of [Zn₂Ba(O₂CCMe₃)₆(C₉H₇N)₂], 1127–32
- New dialkylamido complexes of transition metals in Groups 4 and 5 stabilised by terdentate ligands, 1253–8
- Complexes of the pyridine-based, tridentate, sulphur-containing ligands 2,6-bis(ethylthiomethyl)pyridine, 2,6-bis(methoxycarbonylethylthiomethyl)pyridine, and 2,6-bis(benzylthiomethyl)pyridine. Crystal structure of [2,6-bis(ethylthiomethyl)pyridine]dichlorocadmium(II) hydrate, 1381–4
- A planar, binucleating, pyrazole derivative ligand. Crystal structure of bis[μ -3,5-bis(2'-pyridyl)pyrazolato-*N*¹*N'*:*N*²*N'*]-bis[dimethanolnickel(II)] dichloride dihydrate, 1401–4
- Complexes of ligands providing endogenous bridges. Part 7. The concurrent formation of 1 + 1 and 2 + 2 oxa-azamacrocycles; the X-ray crystal structure of an exocyclic dinuclear copper(II) complex, 1727–38
- Electronic structure of transition-metal indenyl compounds: a He I and He II photoelectron spectroscopic study of [Mn(η ⁵-C₉H₇)(CO)₃], [Fe(η ⁵-C₉H₇)₂], [Ru(η ⁵-C₉H₇)₂], and [Ru(η ⁵-C₉H₇)(η -C₅Me₅)], 2139–48
- Complex-forming properties of L- α -alaninehydroxamic acid (2-amino-*N*-hydroxypropanamide), 2247–52
- Unusual chelating properties of the ligand 1,8-bis(3,5-dimethyl-1-pyrazolyl)-3,6-dithiaoctane (bddo). Crystal structures of Ni(bddo)-(NCS)₂, Zn(bddo)(NCS)₂ and Cd₂(bddo)(NCS)₄, 2309–14
- Optical absorption and magnetic circular dichroism studies of hydrogen, copper(II), zinc(II), nickel(II), and cobalt(II) crown ether-substituted monomeric and dimeric phthalocyanines, 2397–406
- Tables of bond lengths determined by X-ray and neutron diffraction. Part 2. Organometallic compounds and co-ordination complexes of the *d*- and *f*-block metals, S1–S84

TRIAZACYCLONONANE

- Synthesis and spectroscopic studies of cobalt(III) complexes of the chiral pendant-arm macrocycle *N,N',N''*-tris[(2*S*)-2-hydroxypropyl]-1,4,7-triazacyclononane (H₃L); crystal structure of the hydrogen-bridged dimer [CoLH₃LCO]₂[PF₆]₃, 931–6

TRIAZATRIPHOSPHORINANE

- Reactions of amine hydrochlorides with phosphorus trichloride-oxide; preparation and X-ray crystal structure of 2,4,6-trichloro-

1,3,5-triphenyl-1,3,5,2 λ ⁵,6 λ ⁵-triazatrichosphorinane 2,4,6-trioxide (PhNPOCl)₃, 821–8

TRIAZENIDE

- Synthesis and X-ray crystal structure of [Rh(NO)(PPh₃)₂(*p*-MeC₆H₄NNC₆H₄Me-*p*)][BF₄][–]·CH₂Cl₂, 745–8
- Reduction–oxidation properties of organotransition-metal complexes. Part 30. Triazenido-bridged carbonylrhodium bipyridyl complexes with [Rh₂]²⁺, [Rh₂]³⁺, and [Rh₂]⁴⁺ cores, and the X-ray structures of [Rh₂(CO)₂(bipy)(μ -RNNNR)₂][BF₄][–]·CH₂Cl₂ and [$\{\text{Rh}_2\}(\text{CO})(\text{bipy})(\mu\text{-RNNNR})_2\}_2$][PF₆]₂·2.5CH₂Cl₂ (*R* = *p*-tolyl), 2049–60

TRIAZINE

- Copper(II)-assisted hydrolysis of 2,4,6-tris(2-pyridyl)-1,3,5-triazine. Part 2. Crystal structures of [bis(2-pyridylcarbonyl)amido]·[pyridine-2-carboxamide]copper(II) trifluoromethanesulphonate and [bis(2-pyridylcarbonyl)amido]₂[2,4,6-tris(2-pyridyl)-1,3,5-triazine]copper(II) trifluoromethanesulphonate, 1681–8

TRICHLOROACETATE

- Phenylltin trichloroacetate complexes: preparations and structures, 1589–98

TRICHLOROSTANNYL

- Crystallographic and nuclear magnetic resonance studies of carbonyl(η ³-methylallyl)(trichlorostannyl) complexes of platinum(II) and palladium(II), 615–22

TRIFLUOROMETHYL

- Solvent influence on trifluoromethylation reactions with Cd(CF₃)₂; one-pot synthesis of trifluoromethyl iron complexes, 739–44
- Synthesis of hydridotrifluoromethyl complexes of platinum(II). A spectroscopic investigation of the *trans* and *cis* influence of ligands in hydridoplatinum(II) compounds, 1149–60

TRIFLUOROSILYL

- Synthesis and properties of trimethyl(trifluorosilyl)stannane, 1879–82

TUNGSTEN

- Reactions of the bis(alkyne) complexes [WI₂(CO)(NCMe)(η ²-RC₂R)₂] (*R* = Me or Ph) with mono- and bi-dentate phosphine-donor ligands and the X-ray crystal structure of [WI₂(CO)(Ph₂P(CH₂)PPh₂)₂(n²-MeC₂Me)], 295–302

- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 84. Carbaborane tungsten–platinum complexes having a μ -CC₆H₃Me₂-2,6 ligand; crystal structures of [WP(μ -CC₆H₃Me₂-2,6)(CO)_n(PEt₃)₂(μ - σ : η ⁵-C₂B₉H₈Me₂)_n] (*n* = 2 or 3), 303–12

- Spectroscopic studies in matrix isolated tungsten chlorides and bromides, 313–6

- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 85. Synthesis of chain and ring compounds containing molybdenum, 785–96

- Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten [$M[\eta$ ³-SC(CF₃)=C(CF₃H){CF₃C=CCF₃}(η ⁵-C₅H₅)] (*M* = Mo or W). Their relevance to the mechanism of metal-promoted isomerisation of carbon–carbon double bonds, 1027–36

- Studies on transition metal peroxy complexes. Part 8. The nature of peroroxomolybdates and peroxytungstates in aqueous solution, 1203–8

- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 87. Docosahedral carbaborane(alkylidyne)tungsten complexes as reagents for the synthesis of compounds with heteronuclear metal–metal bonds: crystal structures of [NEt₄]₂[W(\equiv CC₆H₃Me₂-2,6)(CO)₂(η ⁶-C₂B₁₀H₁₀Me₂)] and [NEt₄]₂[WF₆(μ -CC₆H₃Me₂-2,6)(CO)₄(η ⁶-C₂B₁₀H₁₀Me₂)], 1363–74

- Heteronuclear transition metal–alkyne clusters. Part 1. Reactions of tris(alkyne)monocarbonyltungsten complexes with octacarbonyldicobalt. Synthesis, X-ray crystal structure, and two-dimensional nuclear magnetic resonance studies of [WC₂(μ -C₂Et₂)(μ -C₄Et₄)(CO)₃]₂, 1375–80

- Reactions of heteropolyanions in non-polar solvents. Part 3.

- Activation of dioxygen by manganese(II) centres in polytungstate. Oxidation of hindered phenols, 1483–90

- Tertiary phosphine complexes of triply bonded ditungsten bis(t-butylamide) tetrachloride. The X-ray crystal structures of [W₂Cl₄(NHBu^t)₂(PMe₃)₂]₂, [W₂Cl₄(NHBu^t)₂(PEt₃)₂]₂, and *cis*-[W₂Cl₄(NHBu^t)₂(PMe₃Ph)₂]₂, 1537–42

- Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 88. Carbaboranetungsteniridium compounds; crystal structure of the complex [WIr(μ -CC₆H₄Me₄)(CO)₂(PEt₃)₂(η ⁵-C₂B₉H₉Me₂)], 1845–54

TUNGSTEN (contd)

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 89. Tetra- and penta-nuclear tungsten-rhodium complexes: crystal structures of $[W_3Rh_2(\mu\text{-CO})_2(\mu\text{-CMe})(\mu\text{-C(Me)C(O)})](\mu\text{-PPPh}_2)_2(\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3$ and $[W_3Rh_2(\mu\text{-CO})_3(\mu\text{-CMe})](\mu\text{-C(Me)PPPh}_2)_2(\mu_3\text{-CMe})(\text{CO})_2(\eta\text{-C}_5\text{H}_5)_3$, 1855–64

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 90. Synthesis of the complexes $[W_2M_2(\mu\text{-CO})(\mu\text{-CMe})(\mu_3\text{-CMe})(\mu\text{-PPPh}_2)_2(\text{CO})(\mu\text{-C}_5\text{H}_5)_2]$ ($M = \text{Rh}$ or Ir) and related tetra- and penta-nuclear metal compounds; interconversion of ketenyl and λ^5 -phospha-alkyne ligands bridging tungsten-rhodium bonds, 1865–70

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 91. Steric implications in the reactions of the complexes $[M(\equiv\text{CR})(\text{CO})(\eta\text{-C}_5\text{H}_5)]$ ($M = \text{Mo}$ or W ; $R = \text{C}_6\text{H}_4\text{Me}-4$, $\text{C}_6\text{H}_4\text{OMe}-2$, or $\text{C}_6\text{H}_3\text{Me}_2-2,6$) with the compounds $[\text{Mo}(\text{CO})_3(\text{NCMe})_3]$ and $[\text{Cu}(\text{thf})(\eta\text{-C}_5\text{Me}_5)]$ ($\text{thf} = \text{tetrahydrofuran}$); crystal structure of $[\text{Mo}_3(\mu\text{-CC}_6\text{H}_3\text{Me}_2-2,6)_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2]$, 1871–8

Synthetic and structural studies on organomolybdenum and organotungsten complexes containing bismuth. $X\text{-ray}$ crystal structures of $[\text{BiCl}\{\text{Mo}(\text{CO})_3(\eta\text{-C}_5\text{H}_5)\}_2]$ and $[\text{BiCl}\{\text{Mo}(\text{CO})_2(\text{CNBu}')(\eta\text{-C}_5\text{H}_5)\}_2]$, 1988, 2941), 1887

Preparation, characterisation, and reactions with phosphite donor ligands of the new cationic monoalkyne complexes $[\text{WI}(\text{CO})(\text{NCMe})(\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2)(\eta^2\text{-RC}_2\text{R})][\text{BF}_4^-]$ ($n = 1$ –6; $R = \text{Me}$ or Ph). $X\text{-ray}$ crystal structure of $[\text{WI}(\text{CO})-\{\text{Pi}(\text{OPr}')_3\}_2\{\text{Ph}_2\text{P}(\text{CH}_2)\text{PPh}_2\}(\eta^2\text{-MeC}_2\text{Me})][\text{BF}_4^-]_{0.5}[\text{OH}]_{0.5}0.5\text{H}_2\text{O}$, 1903–10

Synthesis and characterization of bis(xanthate) complexes of Mo^{II} and W^{II} . Crystal structure of $[\text{W}(\text{S}_2\text{COPr}'-\text{S})(\text{S}_2\text{COPr}'-\text{SS'})(\text{CO})_2(\text{PMMe}_3)_2]$, 2003–10

Studies of the oxidative addition of some substrates containing carbon-heteroatom bonds to some tungsten and platinum complexes, 2099–108

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 92. Synthesis of alkyanylmetallylidynes-molybdenum and -tungsten complexes and their reactions with $[\text{Co}_2(\text{CO})_4]$ and $[\text{Mo}_2(\text{CO})_6(\eta\text{-C}_5\text{H}_5)_2]$, 2261–8

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 93. Synthesis of heteropolynuclear metal compounds with chains of seven to eleven metal atoms; crystal structure of $[\text{Mo}_2\text{W}_3\text{Pt}_6(\mu_3\text{-CMe})_3(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\text{CO})_{10}(\text{PMMe}_2\text{Ph})_4(\eta\text{-C}_5\text{H}_5)_5]$, 2269–80

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 94. Synthesis of di- and tri-metal compounds *via* the reagents $[\text{M}(\equiv\text{C}-\text{C}\equiv\text{CBu}')(\text{CO})_2(\eta\text{-C}_5\text{H}_5)]$ ($M = \text{Mo}$ or W), 2281–8

Chemistry of polynuclear metal complexes with bridging carbene or carbyne ligands. Part 95. Synthesis and crystal structures of $[\text{W}_2\text{RuPt}(\mu\text{-CC}_6\text{H}_4\text{Me}-4)(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)(\text{CO})_6(\text{PMMe}_2\text{Ph})(\eta\text{-C}_5\text{H}_5)_2]$ and $[\text{W}_2\text{Ru}_2\text{Pt}_2(\mu_3\text{-CC}_6\text{H}_4\text{Me}-4)_2(\mu\text{-CO})_2(\text{CO})_9(\eta\text{-C}_5\text{H}_5)_2]$, 2289–96

URANIUM

Actinide structural studies. Part 17. The crystal and molecular structures of four uranyl(vi) complexes with N,N -dialkylthiocarbamate ligands, 471–80

Tri(η -cyclopentadienyl)uranium(iv) silyl and siloxide compounds. Crystal structure of $[\text{U}(\eta^5\text{-C}_5\text{H}_5)_3(\text{OSiPh}_3)]$. Insertion of isocyanide into a uranium-silicon bond, 677–82

URETHANE

Formation of alkylurethanes from carbon dioxide by regioselective O -alkylation of alkali-metal N,N -diethylcarbamates in the presence of complexing agents, 1007–10

VALEROLACTAMATE

Structural, spectroscopic, and electrochemical characterization of tetrakis- μ -(2-pyrrolidinonato)-dirhodium(ii) and tetrakis- μ -(δ -valerolactamato)-dirhodium(ii), 93–100

VANADIUM

Oxovanadium(iv) and dioxomolybdenum(vi) complexes: synthesis from the corresponding acetylacetone complexes and $X\text{-ray}$ structures, 145–50

Potentiometric and electron spin resonance investigation on the interactions between adenosine 5'-triphosphate (atp) and triphosphate ligands with vanadyl ion. Synthesis and characterization of solid vanadyl-atp compounds from acidic aqueous solution, 575–80

Proton electron nuclear double resonance study of oxovanadium(iv) complexes of α -galacturonic and polygalacturonic acids, 1283–8

Proton electron nuclear double resonance study of oxovanadium(iv) complexes of α -diphenolic ligands, 1289–92

Phenoxy and diphenoxo complexes of titanium(iii) and vanadium(iii): synthesis and $X\text{-ray}$ structure, 1793–8

Monomolybdonavanadate and *cis*- and *trans*-dimolybdo-octavanadate, 1915–24

VANADIUM-51

Monomolybdonavanadate and *cis*- and *trans*-dimolybdo-octavanadate, 1915–24

VASOPRESSIN

The unusual co-ordination ability of vasopressin-like peptides: potentiometric and spectroscopic studies of some copper(ii) and nickel(ii) complexes, 173–8

VINYLDENE

Alkyne insertion into ruthenium carbon bonds: formation and ring-closing reactions of vinyl complexes, 465–70

Synthesis, structures, and reactions of ethenethiolato complexes of molybdenum and tungsten $[\text{M}(\eta^3\text{-SC(CF}_3)=\text{C}(\text{CF}_3)\text{H})_4(\text{CF}_3\text{C}\equiv\text{CCF}_3)(\eta^5\text{-C}_5\text{H}_5)]$ ($M = \text{Mo}$ or W). Their relevance to the mechanism of metal-promoted isomerisation of carbon-carbon double bonds, 1027–36

VINYLDENE

Synthesis, structure, and dynamic behaviour of two isomeric triosmium clusters containing the ethoxyvinylidene (C=CHOEt) and ethoxyethyne ($\text{CH}\equiv\text{COEt}$) ligands respectively, 5–12

Reactions of 1-alkynes with *trans*-

$[\text{ReCl}(\text{N}_2)(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$: preparation of the vinylidene compounds *trans*- $[\text{ReCl}(-\text{C}=\text{CHR})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$ ($R = \text{alkyl or aryl}$) and $X\text{-ray}$ structure of *trans*- $[\text{ReCl}(-\text{C}=\text{CHPh})(\text{Ph}_2\text{PCH}_2\text{CH}_2\text{PPh}_2)_2]$, 2381–8

WATER

Catalytic oxidation of water mediated by rhenium oxides, 849–52

YLIDE

Bi-, tetra-, and poly-nuclear complexes of gold(i) and silver(i) with bis(diphenylphosphino)-methanide or -amide acting as six-electron donors, 1883–6

ZINC

An investigation of halide abstraction by arsenic(iii): reactions of AsCl_3 with MCl_2 ($M = \text{Mg}$ or Zn) and TiCl_4 in donor solvents (MeCN or POCl_3). Formation and crystal structure of the dimeric titanium(iv) compound $[\text{Ti}_2\text{Cl}_6(\text{POCl}_3)_2(\text{PO}_2\text{Cl}_2)_2]$, 57–60

Zinc(ii) complexes of 4,7-diazadecane-1,10-diamine, 4,8-diazaundecane-1,11-diamine, and tetrathylenepentamine: a carbon-13 nuclear magnetic resonance study, 1171–8

ZIRCONIUM

Formation of chromium interlayers in α -zirconium phosphate, 895–900

Zirconium phosphate ester interchange reactions, 1617–24

Preparation of thionylimide complexes of titanium, zirconium, and hafnium. Crystal structure of $[\text{Zr}(\text{cp})(\eta\text{-C}_5\text{Me}_5)(\text{NSO})_2]$, 1815–8

Cyclic metallasiloxane compounds of Ti^{IV} and Zr^{IV} . $X\text{-ray}$ crystal structures of a *cis*-bis(pyridine) bis(cyclosiloxyl)titanate and a pyridinolithiated tris(cyclosiloxyl)zirconate, 2347–55