

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/20447614>

# Synthesis of 5-[1-hydroxy(or methoxy)-2-bromo(or chloro)ethyl]-2'-deoxyuridines and related halohydrin analogs with antiviral and cytotoxic activity

ARTICLE in JOURNAL OF MEDICINAL CHEMISTRY · JUNE 1989

Impact Factor: 5.45 · DOI: 10.1021/jm00125a003 · Source: PubMed

CITATIONS

21

READS

8

8 AUTHORS, INCLUDING:



Leonard I Wiebe

University of Alberta

333 PUBLICATIONS 3,448 CITATIONS

SEE PROFILE



Ed Knaus

University of Alberta

187 PUBLICATIONS 1,832 CITATIONS

SEE PROFILE



Theresa M Allen

University of British Columbia - Vancouver

257 PUBLICATIONS 17,623 CITATIONS

SEE PROFILE

## Articles


# Synthesis of 5-[1-Hydroxy(or methoxy)-2-bromo(or chloro)ethyl]-2'-deoxyuridines and Related Halohydrin Analogues with Antiviral and Cytotoxic Activity

Rakesh Kumar,<sup>†</sup> Leonard I. Wiebe,<sup>†</sup> Tse W. Hall,<sup>†</sup> Edward E. Knaus,<sup>\*,†</sup> Dorothy R. Tovell,<sup>‡</sup> D. Lorne Tyrrell,<sup>‡</sup> Theresa M. Allen,<sup>§</sup> and R. Fathi-Afshar<sup>§</sup>

Faculty of Pharmacy and Pharmaceutical Sciences, University of Alberta, Edmonton, Alberta, Canada T6G 2N8, and Departments of Biochemistry and Pharmacology, Faculty of Medicine, University of Alberta, Edmonton, Alberta, Canada T6G 2H7. Received September 19, 1988


A series of new 5-(1-hydroxy-2-haloethyl)-2'-deoxyuridines (**3**, **6**, **8**) were synthesized in 60–70% yields by addition of HOX (X = Br, Cl, I) to the vinyl substituent of the respective 5-vinyl-2'-deoxyuridines (**2**, **5**, **7**). Treatment of **3a,b** with methanolic sulfuric acid afforded the corresponding 5-(1-methoxy-2-haloethyl)-2'-deoxyuridines (**4a,b**). The 5-(1-hydroxy-2-chloroethyl) (**3b**), 5-(1-methoxy-2-bromoethyl) (**4a**), 5-(1-hydroxy-2-bromo-2-(ethoxycarbonyl)ethyl) (**6a**), and 5-(1-hydroxy-2-iodo-2-(ethoxycarbonyl)ethyl) (**6b**) derivatives exhibited in vitro antiviral activity (ID<sub>50</sub> = 0.1–1 µg/mL range) against herpes simplex virus type 1 (HSV-1). 5-(1-Hydroxy-2-bromo-2-(ethoxycarbonyl)ethyl)-2'-deoxyuridine (**6a**) was the most active cytotoxic agent in the in vitro L1210 screen exhibiting an ED<sub>50</sub> of 11 µg/mL relative to melphalan (ED<sub>50</sub> = 0.15 µg/mL).

(*E*)-5-(2-Bromovinyl)-2'-deoxyuridine (**1a**, BVDU)<sup>1</sup> and 5-(2-chloroethyl)-2'-deoxyuridine (**1b**, CEDU)<sup>2</sup> are two of the most potent and selective antiviral agents from the large number of 5-substituted pyrimidine nucleoside analogues investigated.<sup>3</sup> CEDU was effective in vivo



against systemic herpes simplex virus type 1 (HSV-1) infection and HSV-1 encephalitis in mice at a 5–15-fold lower dose than either BVDU or acyclovir (ACV).<sup>2,4,5</sup> 5-Vinyl-2'-deoxyuridine (**1c**, VDU) is a potent antiviral and antileukemic agent in tissue culture<sup>1,6,7</sup> but was inactive in vivo as an antiviral, antileukemic, or cytotoxic agent due to in vivo degradation by thymidine phosphorylase.<sup>8</sup> Although 5-(2-hydroxyethyl)-2'-deoxyuridine (**1d**, HEDU) is an inactive antiviral agent,<sup>2</sup> 5-ethyl-2'-deoxyuridine (**1e**, EDU) is a potent and selective inhibitor of herpes simplex virus type 1 (HSV-1) and 2 (HSV-2) replication that is being investigated for the topical treatment of HSV-1 and HSV-2 infections in humans. The selectivity of EDU is due to its preferential phosphorylation by the virus-infected cell and its preferential incorporation into viral DNA.<sup>9</sup> It was therefore of interest to investigate the effect of novel 5-halohydrin substituents attached to the C-5 position of 2'-deoxyuridine on biological activity. Halohydrins (**3**, **6**, **8**) could undergo selective phosphorylation,

Scheme I<sup>a</sup>



<sup>a</sup> Reagents: i, *N*-bromosuccinimide (**3a**), *N*-chlorosuccinimide (**3b**), dioxane–water (3:7, v/v), acetic acid, 25 °C; ii, H<sub>2</sub>SO<sub>4</sub>–MeOH, 25 °C.

like BVDU<sup>1</sup> and EDU,<sup>9</sup> by HSV-1 virus-encoded thymidine kinase to exhibit selective antiviral activity and/or


<sup>†</sup> Faculty of Pharmacy and Pharmaceutical Sciences.

<sup>‡</sup> Department of Biochemistry.

<sup>§</sup> Department of Pharmacology.


- (1) De Clercq, E.; Descamps, J.; De Somer, P.; Barr, P.; Jones, A. S.; Walker, R. T. *Proc. Natl. Acad. Sci. U.S.A.* 1979, 76, 2947.
- (2) Griengl, H.; Bodenteich, M.; Hayden, W.; Wanek, E.; Streicher, W.; Stutz, P.; Bachmayer, H.; Ghazzouli, I.; Rosenwirth, B. *J. Med. Chem.* 1985, 28, 1679.
- (3) De Clercq, E. In *Approaches to Antiviral Agents*; Harnden, M. R., Ed.; MacMillan: London, 1985; p 57.
- (4) De Clercq, E.; Descamps, J.; Ogata, M.; Shigeta, S. *Antimicrob. Agents Chemother.* 1982, 21, 33.
- (5) De Clercq, E.; Rosenwirth, B. *Antimicrob. Agents Chemother.* 1985, 28, 246.
- (6) De Clercq, E.; Descamps, J.; Verhelst, G.; Walker, R. T.; Jones, A. S.; Torrence, P. F.; Shugar, D. *J. Infect. Dis.* 1980, 141, 563.
- (7) De Clercq, E.; Balzarini, J.; Torrence, P. F.; Mertes, M. P.; Schmidt, C. L.; Shugar, D.; Barr, P. J.; Jones, A. S.; Verhelst, G.; Walker, R. T. *Mol. Pharmacol.* 1981, 19, 321.



**Table I.** In Vitro Antiviral Activity (HSV-1) and L1210 Cytotoxicity of 5-Substituted 2'-Deoxyuridines


no.	R <sup>2</sup>	antiviral activity ID <sub>50</sub> <sup>a</sup> μg/mL	L1210 cytotoxicity % survival ± SD <sup>b</sup>	
			10 μg/mL	50 μg/mL
3a	CH(OH)CH <sub>2</sub> Br	>10	65.5 ± 8.7	23.7 ± 1.5
3b	CH(OH)CH <sub>2</sub> Cl	0.1-1	74.0 ± 4.5	24.3 ± 2.1
4a	CH(OMe)CH <sub>2</sub> Br	0.1-1	80.8 ± 3.4	11.7 ± 18.6
4b	CH(OMe)CH <sub>2</sub> Cl	>10	75.0 ± 3.2	38.8 ± 1.1
6a	CH(OH)CH(Br)CO <sub>2</sub> Et	1	49.7 ± 2.8	4.9 ± 0.3 <sup>c</sup>
6b	CH(OH)CH(I)CO <sub>2</sub> Et	0.1-1	73.2 ± 3.2	10.4 ± 1.7
8a:8b	CH(OH)CH(Br)I and CH(Br)CH(OH)I <sup>d</sup>	>10	79.2 ± 5.3	40.1 ± 1.3
acyclovir		0.01		
melfalane <sup>e</sup>			2.8 ± 0.68 <sup>f</sup>	0.0

<sup>a</sup>The concentration required to inhibit plaque formation in monolayers of Vero cells by 50%. <sup>b</sup>The result is the mean value ± SD for three experiments. <sup>c</sup>ED<sub>50</sub> = 11 μg/mL. <sup>d</sup>Mixture of regioisomers 8a and 8b in a ratio of 1:1. <sup>e</sup>4-[N-Bis(2-chloroethyl)amino]phenylalanine. <sup>f</sup>Percent survival ± SD determined at 1 μg/mL, ED<sub>50</sub> = 0.15 μg/mL.

**Scheme III<sup>a</sup>**

<sup>a</sup>Reagents: i, N-bromosuccinimide, dioxane-water (3:7, v/v) acetic acid, 25 °C.

pound (3b > 4b, *P* = 0.005). A comparison of the activities for 6a and 3a (6a > 3a, *P* = 0.001) indicated that the introduction of an ethoxycarbonyl group enhanced cytotoxic activity significantly whereas incorporation of an iodine substituent (3a > 8a:8b, *P* = 0.001) decreased activity. The most active cytotoxic agent, 6a (ED<sub>50</sub> = 11 μg/mL), was 73 times less active than the reference compound melfalane (ED<sub>50</sub> = 0.15 μg/mL). Compounds 3, 4, 6, and 8a,b were not evaluated in the P388 leukemia in vivo screen since compounds exhibiting an ED<sub>50</sub> > 5 μg/mL in this in vitro screen are considered to have low cytotoxic activity.

A number of similar correlations were observed in both the antiviral and cytotoxic screens where the relative activities were 4a > 4b, 4a > 3a, 3b > 4b, and 6a > 3a.

**Experimental Section**

Melting points were determined with a Büchi capillary apparatus and are uncorrected. Nuclear magnetic resonance spectra (<sup>1</sup>H NMR, <sup>13</sup>C NMR) were determined for solutions in pyridine-*d*<sub>5</sub>, D<sub>2</sub>O, or Me<sub>2</sub>SO-*d*<sub>6</sub> with Me<sub>4</sub>Si as internal standard (<sup>1</sup>H NMR) with a Bruker AM-300 spectrometer. Chemical-ionization mass spectra (CIMS, NH<sub>3</sub>) were measured on an AEI MS-12 spec-

trometer. Preparative thin-layer chromatography (PTLC) was performed with Whatman PLK5F plates, 1.0 mm in thickness. Silica gel column chromatography was carried out with Fisher S-704 (60-200 mesh) silica gel. Microanalyses were within ±0.4% of theoretical values for all elements listed, unless otherwise indicated. 5-Vinyl-2'-deoxyuridine (2)<sup>14</sup> and (*E*)-5-(2-(ethoxycarbonyl)vinyl)-2'-deoxyuridine (5)<sup>15</sup> were prepared by using the literature procedures.

**5-(1-Hydroxy-2-bromoethyl)-2'-deoxyuridine (3a).** *N*-Bromosuccinimide (0.149 g, 0.844 mmol) was added slowly with stirring to a solution of 2 (0.208 g, 0.82 mmol) in dioxane-water (3:7, v/v, 10 mL) and glacial acetic acid (0.04 mL) during a period of 5 min. The reaction was allowed to proceed for 2 h at 25 °C with stirring and the solvent was removed in vacuo. The product was purified by elution from a silica gel column using chloroform-methanol (94:6, v/v) as eluant to yield 3a (0.20 g, 70%) after recrystallization from methanol: mp 135 °C dec; <sup>1</sup>H NMR (pyridine-*d*<sub>5</sub>) δ 2.7 (complex m, 2 H, H-2'), 3.85 (2 d, *J* = 9.6 and 7.2 Hz, 1 H, CHBr), 4.12-4.32 (complex m, 3 H, H-5', CHBr), 4.58 (m, 1 H, H-4'), 5.05 (m, 1 H, H-3'), 5.25 (br s, 6 H, 3'-OH, 5'-OH, CHOCH<sub>2</sub>Br, <sup>3</sup>/<sub>2</sub>H<sub>2</sub>O, exchange with deuterium oxide), 5.5 (2 d, *J* = 7.2 and 2.4 Hz, 1 H, CHOCH<sub>2</sub>Br), 7.1 (d, *J* = 6 Hz of d, *J* = 6 Hz, 1 H, H-1'), 8.89 (s, 1 H, H-6), 13.54 (s, 1 H, NH, exchanges with deuterium oxide); <sup>13</sup>C NMR (Me<sub>2</sub>SO-*d*<sub>6</sub>) δ 38.18 and 38.48 (C-Br), 39.49 (C-2'), 61.42 (C-5'), 65.62 and 65.98 (CHOCH<sub>2</sub>Br), 69.99 and 70.41 (C-3'), 84.18 (C-1'), 87.31 and 87.36 (C-4'), 114.16 (C-5), 137.47 (C-6), 149.99 (C-2), 162.06 (C-4); MS (CIMS, NH<sub>3</sub>) *m/z* 369 (M<sup>+</sup> + NH<sub>4</sub>). Anal. (C<sub>11</sub>H<sub>15</sub>N<sub>2</sub>O<sub>6</sub>Br·<sup>3</sup>/<sub>2</sub>H<sub>2</sub>O) C, N, H: calcd, 4.35; found, 4.79.

**5-(1-Hydroxy-2-chloroethyl)-2'-deoxyuridine (3b).** *N*-Chlorosuccinimide (0.12 g, 0.90 mmol) was added slowly with stirring to a solution of 2 (0.221 g, 0.87 mmol) in dioxane-water (3:7, v/v, 10 mL) and glacial acetic acid (0.03 mL) during a period of 5 min. The reaction was allowed to proceed for 24 h at 25 °C. The product was isolated and purified by silica gel column chromatography, as described for 3a, to yield 3b as a white solid after recrystallization from methanol (0.16 g, 60%): mp 155 °C dec; <sup>1</sup>H NMR (pyridine-*d*<sub>5</sub>) δ 2.7 (m, 2 H, H-2'), 3.98 (d, *J*<sub>gem</sub> = 10.3 Hz of d, *J*<sub>vic</sub> = 7.16 Hz, 1 H, CHCl), 4.23 (complex m, 2 H, H-5'), 4.38 (d, *J*<sub>gem</sub> = 10.3 Hz of d, *J*<sub>vic</sub> = 2.38 Hz, 1 H, CHCl), 4.58 (m, 1 H, H-4'), 4.9-5.3 (br s, 7 H, 3'-OH, 5'-OH, CHOCH<sub>2</sub>Cl, <sup>3</sup>/<sub>2</sub>H<sub>2</sub>O, H-3', hydroxyls exchange with deuterium oxide), 5.51 (2 d, *J* = 7.16 and 2.38 Hz, 1 H, CHOCH<sub>2</sub>Cl), 7.1 (d, *J* = 6 Hz of

(14) Rahim, S. G.; Duggan, M. J. H.; Walker, R. T.; Jones, A. S.; Dyer, R. L.; Balzarini, J.; De Clercq, E. *Nucleic Acids Res.* 1982, 10, 5285.

(15) Barr, P. J.; Oppenheimer, N. J.; Santi, D. V. *J. Biol. Chem.* 1983, 258, 13627.

d,  $J = 6$  Hz, 1 H, H-1'), 8.85 (s, 1 H, H-6), 13.52 (s, 1 H, NH, exchanges with deuterium oxide). Anal. ( $C_{11}H_{15}N_2O_6Cl \cdot \frac{1}{2}H_2O$ ) C, Calcd: H, 4.75; N, 7.96. Found: H, 5.43; N, 8.39.

**5-(1-Methoxy-2-bromoethyl)-2'-deoxyuridine (4a).** A solution of sulfuric acid in methanol (4 mL, 5 N) was added to a solution of **3a** (0.144 g, 0.41 mmol) in methanol (50 mL). The reaction was allowed to proceed for 48 h at 25 °C with stirring prior to neutralization with methanolic sodium hydroxide. Removal of the solvent in vacuo, dissolution of the residue in methanol (5 mL), adsorption on silica gel (2 g), removal of the solvent in vacuo, and application of this material to the top of a silica gel column followed by elution with chloroform-methanol (96:4, v/v) yielded **4a** (0.14 g, 93%) as a viscous oil. Trituration with cold ether gave **4a** as a white solid: mp 135 °C dec;  $^1H$  NMR ( $Me_2SO-d_6$ ) (mixture of two diastereomers in a ratio of 1:1)  $\delta$  2.15 (m, 2 H, H-2'), 3.25 and 3.28 (2 s, 3 H total, OMe), 3.45 (m, 1 H, CHBr), 3.6 (m, 2 H, H-5'), 3.76 (complex m, 1 H, CHBr), 3.85 (m, 1 H, H-4'), 4.28 (m, 1 H, H-3'), 4.36 (m, 1 H, CHOMe), 5.08 and 5.3 (2 br s, 1 H each, 3'-OH, 5'-OH, exchange with deuterium oxide), 6.2 (2 overlapping d,  $J = 6$  Hz of d,  $J = 6$  Hz, 1 H total, H-1'), 7.92 and 7.95 (2 s, 1 H total, H-6), 11.54 (s, 1 H, NH, exchanges with deuterium oxide). Anal. ( $C_{12}H_{17}N_2O_6Br \cdot \frac{1}{2}H_2O$ ) C, H, N: calcd, 6.94; found, 7.48.

**5-(1-Methoxy-2-chloroethyl)-2'-deoxyuridine (4b).** A solution of sulfuric acid in methanol (5.5 mL, 5 N) was added to a solution of **3b** (0.131 g, 0.43 mmol) in methanol (50 mL). The reaction was completed, and the product was isolated and purified as described for **4a**, to yield **4b** as a white solid (0.135 g, 98%): mp 157 °C dec;  $^1H$  NMR ( $Me_2SO-d_6$ ) (mixture of two diastereomers in a ratio of 1:1)  $\delta$  2.19 (m, 2 H, H-2'), 3.3 and 3.32 (2 s, 3 H total, OMe), 3.4 (m, 1 H, CHCl), 3.64 (m, 2 H, H-5'), 3.72 (m, 1 H, CHCl), 3.88 (m, 1 H, H-4'), 4.3 (m, 1 H, H-3'), 4.36 (m, 1 H, CHOMe), 5.1 and 5.31 (2 m, 1 H each, 3'-OH, 5'-OH, exchange with deuterium oxide), 6.2 (2 overlapping d,  $J = 6$  Hz each, 1 H total, H-1'), 7.92 and 7.95 (2 s, 1 H total, H-6), 11.52 (s, 1 H, NH, exchanges with deuterium oxide). Anal. ( $C_{12}H_{17}N_2O_6Cl \cdot \frac{1}{2}H_2O$ ) C, H, N.

**5-(1-Hydroxy-2-bromo-2-(ethoxycarbonyl)ethyl)-2'-deoxyuridine (6a).** *N*-Bromosuccinimide (0.109 g, 0.61 mmol) was added in small aliquots to a solution of **5** (0.2 g, 0.615 mmol) in water (10 mL) and the reaction was allowed to proceed for 24 h at 25 °C. Removal of the solvent in vacuo gave a viscous oil, dissolution in methanol (5 mL), adsorption on silica gel (2 g), removal of the solvent in vacuo, and application of this material to the top of a silica gel column followed by elution with chloroform-methanol (95:5, v/v) afforded **6a** (0.18 g, 70%) as a white solid: mp 68 °C (sublimes);  $^1H$  NMR ( $Me_2SO-d_6$ ) (mixture of two diastereomers in a ratio of 1:1)  $\delta$  1.22 (t,  $J = 7$  Hz, 3 H,  $CH_2CH_3$ ), 2.12 (m, 2 H, H-2'), 3.6 (m, 2 H, H-5'), 3.8 (m, 1 H, H-4'), 4.2 (q,  $J = 7$  Hz, 2 H,  $CH_2CH_3$ ), 4.26 (m, 1 H, H-3'), 4.82 (2 overlapping m, 2 H, CHOHCHBr), 5.14 and 5.31 (2 m, 1 H each, 3'-OH, 5'-OH, exchange with deuterium oxide), 6.06 (m, 1 H, CHOHCHBr, exchanges with deuterium oxide), 6.21 (2 overlapping d,  $J = 6$  Hz, of d,  $J = 6$  Hz, 1 H total, H-1'), 8.04 and 8.06 (2 s, 1 H total, H-6), 11.54 (s, 1 H, NH, exchanges with deuterium oxide);  $^{13}C$  NMR ( $Me_2SO-d_6$ )  $\delta$  13.71 ( $CH_2CH_3$ ), 39.58 (C-2'), 46.92 and 47.14 (CHBr), 61.24 (C-5'), 61.39 ( $CH_2CH_3$ ), 68.07 and 68.58 (CHOHCHBr), 70.26 and 70.41 (C-3'), 84.16 and 84.34 (C-1'), 87.48 (C-4'), 112.54 (C-5), 139.51 and 139.36 (C-6), 149.84 (C-2), 162.18 and 162.24 (C-4), 168.14 ( $CO_2$ ). Anal. ( $C_{14}H_{19}N_2O_8Br$ ) C, H, N.

**5-(1-Hydroxy-2-iodo-2-(ethoxycarbonyl)ethyl)-2'-deoxyuridine (6b).** A mixture of **5** (49 mg, 0.15 mmol), iodine (19 mg, 0.15 mmol), potassium iodate (6 mg, 0.028 mmol), water (2 mL), and sulfuric acid (15  $\mu$ L, 5 N) were stirred for 12 h at 55 °C. Removal of the solvent in vacuo gave a residue which was purified by PTLC using chloroform-methanol (90:10, v/v) as development solvent. Extraction of the ultraviolet visible spot with chloroform-methanol (88:12, v/v) yielded **6b** as a white solid (42 mg, 60%): mp 110 °C dec;  $^1H$  NMR ( $Me_2SO-d_6$ ) (mixture of two diastereomers in a ratio of 1:1)  $\delta$  1.22 (t,  $J = 7$  Hz, 3 H,  $CH_2CH_3$ ), 2.15 (m, 2 H, H-2'), 3.64 (m, 2 H, H-5'), 3.83 (m, 1 H, H-4'), 4.18 (q,  $J = 7$  Hz, 2 H,  $CH_2CH_3$ ), 4.3 (m, 1 H, H-3'), 4.75-4.9 (2 overlapping m, 2 H, CHOHCHI), 5.15 and 5.3 (2 m, 1 H each, 3'-OH, 5'-OH, exchange with deuterium oxide), 5.9 (m, 1 H, CHOHCHI, exchanges with deuterium oxide), 6.22 (2 overlapping d,  $J = 6$  Hz each, 1 H total, H-1'), 8.0 and 8.02 (2 s, 1 H total,

H-6), 11.52 (s, 1 H, NH, exchanges with deuterium oxide);  $^{13}C$  NMR ( $Me_2SO-d_6$ )  $\delta$  13.53 ( $CH_3$ ), 24.56 and 24.78 (CHI), 39.63 and 39.87 (C-2'), 61.01 ( $CH_2CH_3$ ), 61.25 and 61.39 (C-5'), 68.79 and 69.51 (CHOHCHI), 70.26 and 70.41 (C-3'), 84.09 and 84.31 (C-1'), 87.49 (C-4'), 113.42 and 113.54 (C-5), 139.21 and 139.42 (C-6), 149.84 (C-2), 162.09 and 162.15 (C-4), 170.0 ( $CO_2$ ). Anal. ( $C_{14}H_{19}N_2O_8I$ ) C, H, N.

**5-(1-Hydroxy-2-bromo-2-iodoethyl)-2'-deoxyuridine (8a) and 5-(1-Bromo-2-hydroxy-2-iodoethyl)-2'-deoxyuridine (8b).** A mixture of *N*-bromosuccinimide (91 mg, 0.51 mmol) and **7** (190 mg, 0.5 mmol) in dioxane-water (3:7, v/v, 8 mL) and glacial acetic acid (25  $\mu$ L) was allowed to stir at 25 °C for 2 h. Removal of the solvent in vacuo and purification of the product by elution from a silica gel column using methanol-chloroform (6:94, v/v) as eluant yielded a mixture of regioisomers **8a** and **8b** in a ratio of 1:1, determined by integration of the CHBr resonances at  $\delta$  6.01 and 6.16 respectively, as a white solid (0.152 g, 64%): mp 88 °C (sublimes);  $^1H$  NMR ( $D_2O$ ) (each regioisomer **8a** and **8b** is a mixture of two diastereomers)  $\delta$  2.36 (complex m, 4 H total, H-2'), 3.76 (complex m, 4 H total, H-5'), 4.05 (m, 2 H total, H-4'), 4.45 (m, 2 H total, H-3'), 5.02 (m, 2 H total, CHOHCHBr), 6.01 (m, 1 H, CHBr in one regioisomer), 6.16 (m, 1 H, CHBr in one regioisomer), 6.28 (m, 1 H total, H-1'), 8.05 (4 closely spaced singlets, 2 H total, H-6);  $^{13}C$  NMR ( $Me_2SO-d_6$ )  $\delta$  20.66 and 21.11 (CHBrI of **8a**), 39.65 and 39.95 (C-2'), 51.90 and 52.26 (CHBr of **8b**), 61.85 and 61.75 (C-5'), 70.59 and 70.71 (C-3'), 71.10, 71.43, 71.94 and 72.37 (CHOH of **8a** and **8b**), 84.61 (C-1'), 87.51 and 87.57 (C-4'), 113.13 and 113.63 (C-5), 138.88 and 138.94 (C-6), 149.99 (C-2), 162.21 (C-4). Anal. ( $C_{11}H_{14}N_2O_8BrI \cdot 1H_2O$ ) C, H, N.

**In Vitro Antiviral Assay (HSV-1).** Monolayers of mycoplasma-free Vero cells in 60-mm plates were infected with HSV-1 (strain JLV isolated from a patient with herpes simplex encephalitis). After a 1-h adsorption period at 37 °C, the cells were overlaid with 1% agarose in Eagle minimum essential medium (E-MEM) containing 2% fetal bovine serum and known concentrations of the agents (0.1, 1, and 10  $\mu$ g/mL for **3**, **4**, **6**, and **8**; 0.01, 0.1, and 1  $\mu$ g/mL for acyclovir) being tested. Cells were stained with neutral red after incubation for 4 days at 37 °C in a 5%  $CO_2$  incubator, and the number of plaques were counted. The experiment was performed once. The concentration of test compound required to inhibit plaque formation by 50% ( $ID_{50}$ ) was determined.

**In Vitro L1210 Cytotoxic Assay.** Mouse L1210 leukemia cells were cultivated as a suspension in Fischer's medium supplemented with 10% heat-inactivated horse serum and incubated at 37 °C in a humidified 5%  $CO_2$  atmosphere to prepare a cell stock solution. The number of cells/mL of medium was determined with a Model ZF Coulter Counter 48 h after incubation. The test compound was dissolved in ethanol-water (1:1, v/v, **3a,b**), water (**4a,b**), or ethanol **6a,b**, **8a:8b**) and 10  $\mu$ L of this solution was added to test wells containing 2 mL of suspended L1210 cells ( $10^5$  cells/mL) such that 2 mL of the cell suspension had a test compound concentration of 50, 10, and 1  $\mu$ g/mL of medium, respectively. Control wells were identical, except that the test compound was absent. Compounds for which  $ED_{50}$  values were obtained had the following test compound concentrations ( $\mu$ g/mL of medium): **6a** (50, 25, 10, 5.0, 2.5, and 1) and melphalan (10, 1, 0.5, 0.25, 0.1, and 0.05). All tests and controls were grown in triplicate. The percent cell survival was calculated with the formula: percent survival =  $(T_{48} - T_0) / (C_{48} - C_0) \times 100$ ; where  $T_{48}$  is the number of living cells/mL for each test drug concentration at 48 h,  $T_0$  is the number for test wells at time zero (normally  $10^5$ ),  $C_{48}$  is the number for the control at 48 h, and  $C_0$  is the number for the control at time zero (normally,  $T_0 = C_0 = 10^5$  cells/mL). Compounds exhibiting an  $ED_{50} > 5 \mu$ g/mL are considered to be inactive in this screen.

**Acknowledgment.** We are grateful to the Medical Research Council of Canada (Grant No. MT-5965) (L.I.W. and E.E.K.) and the National Cancer Institute of Canada (T.M.A.) for financial support of this work and to the Alberta Heritage Foundation for Medical Research for a fellowship to one of us (R.K.). We also thank Bruker Spectrospin (Canada) Ltd. for providing the 2-D INADEQUATE spectrum of **8a:8b**.