

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/49802986>

Discovery of New Human A(2A) Adenosine Receptor Agonists: Design, Synthesis, and Binding Mode of Truncated 2-Hexynyl-4'-thioadenosine.

ARTICLE in ACS MEDICINAL CHEMISTRY LETTERS · SEPTEMBER 2010

Impact Factor: 3.12 · DOI: 10.1021/ml1001823 · Source: PubMed

CITATIONS

9

READS

25

11 AUTHORS, INCLUDING:



Mulamoottil Varughese

University of Georgia

21 PUBLICATIONS 225 CITATIONS

SEE PROFILE



Zhan-Guo Gao

The National Institute of Diabetes and Digest...

146 PUBLICATIONS 4,673 CITATIONS

SEE PROFILE



Kenneth A. Jacobson

National Institutes of Health

753 PUBLICATIONS 29,958 CITATIONS

SEE PROFILE



Lak Shin Jeong

Ewha Womans University

225 PUBLICATIONS 2,042 CITATIONS

SEE PROFILE

Published in final edited form as:

ACS Med Chem Lett. 2010 September 1; 2010(9): 516–520. doi:10.1021/ml1001823.

Discovery of New Human A_{2A} Adenosine Receptor Agonists: Design, Synthesis, and Binding Mode of Truncated 2-Hexynyl-4'-thioadenosine

Xiyan Hou¹, Hea Ok Kim¹, Varughese Alexander¹, Kyunglim Kim¹, Sun Choi¹, Seul-gi Park¹, Jin Hee Lee¹, Lena S. Yoo², Zhan-Guo Gao², Kenneth A. Jacobson², and Lak Shin Jeong^{1,3,*}

¹ Laboratory of Medicinal Chemistry, College of Pharmacy, Ewha Womans University, Seoul 120-750, Korea

² Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes, and Digestive and Kidney Disease, National Institutes of Health, Bethesda, Maryland 20892, USA

³ Department of Bioinspired Science, Ewha Womans University, Seoul 120-750, Korea

Abstract

The truncated C2- and C8-substituted-4'-thioadenosine derivatives **4a-d** were synthesized from D-mannose, using palladium-catalyzed cross coupling reactions as key steps. In this study, an A₃ adenosine receptor (AR) antagonist, truncated 4'-thioadenosine derivative **3** was successfully converted into a potent A_{2A}AR agonist **4a** ($K_i = 7.19 \pm 0.6$ nM) by appending a 2-hexynyl group at the C2-position of a derivative of **3** that was N⁶-substituted. However, C8-substitution greatly reduced binding affinity at the human A_{2A}AR. All synthesized compounds **4a-d** maintained their affinity at the human A₃AR, but **4a** was found to be a competitive A₃AR antagonist/A_{2A}AR agonist in cyclic AMP assays. This study indicates that the truncated C2-substituted-4'-thioadenosine derivatives **4a** and **4b** can serve as a novel template for the development of new A_{2A}AR ligands.

Keywords

A_{2A} adenosine receptor agonists; truncated 2-hexynyl-4'-thioadenosine; palladium-catalyzed cross coupling reactions; binding mode

The endogenous cytoprotective modulator adenosine (**1**) exerts its pharmacological effects against hypoxia, ischemia, and inflammation through binding to four subtypes (A₁, A_{2A}, A_{2B}, and A₃) of adenosine receptors (ARs), members of the G protein-coupled receptor (GPCR) family.¹ Among these, activation of A_{2A}AR has been known to play a role in the suppression of immune and inflammatory responses and in vascular responses to adenosine.² It is also highly localized within the central nervous system (CNS), and selective antagonists have become an attractive target for the treatment of Parkinson's disease.³ The

*To whom correspondence should be addressed: Lak Shin Jeong, Ph.D., Laboratory of Medicinal Chemistry, College of Pharmacy and Department of Bioinspired Science, Ewha Womans University, Seoul 120-750, Korea, Tel) 82-2-3277-3466, Fax) 82-2-3277-2851, lakjeong@ewha.ac.kr.

SUPPORTING INFORMATION AVAILABLE: Complete experimental procedures and characterization data and ¹H and ¹³C NMR copies of **4a-d**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

A₃AR is the most recently identified subtype and is found in the cardiovascular system, CNS, immune cells, lung, and liver.^{4,5} The activation of A₃AR is beneficial in models of myocardial and cerebral ischemia and cancer, while its antagonism is of interest for treating asthma, inflammation, and glaucoma.⁶

Thio-Cl-IB-MECA (**2**)⁷ which is bioisosteric with 2-chloro-*N*⁶-(3-iodobenzyl)adenosine-5'-*N*-methyluronamide (Cl-IB-MECA)⁸ was discovered as a potent and selective A₃AR agonist ($K_i = 0.38$ nM). This compound showed a potent anticancer activity by inhibiting the Wnt signaling pathway.⁹ On the basis of rational design, truncated 4'-thioadenosine derivative **3** lacking the 5'-uronamide of **2** essential for the receptor activation was discovered as a potent, selective, and species-independent A₃AR antagonist ($K_i = 1.66$ nM).¹⁰ This compound and other related A₃AR antagonists with nucleoside skeletons are expected to be suitable for evaluation in small animal models and for further development as drugs (Figure 1).

Based on the observation that truncation resulting in the 4'-thioadenosine antagonist derivative **3** preserved AR affinity and selectivity, we designed and synthesized the truncated C2- and C8-substituted 4'-thioadenosine derivatives **4a-d** as potential new ligands for the A_{2A}AR. This expectation was supported by reports that C2- or C8-substitution sometimes leads to substantial enhancement in the binding affinity or selectivity at the A_{2A}AR or other AR subtype.¹¹ C2- or C8-substitution was readily achieved through Sonogashira¹² and Suzuki¹³ cross coupling reactions. From this study, a C2-alkynyl derivative was found to be potent, mixed A_{2A}AR agonist and A₃AR antagonist, which is an excellent combination for the anti-asthmatic activity. Herein, we describe the synthesis and pharmacological activity of novel C2- and C8-substituted 4'-thioadenosine derivatives **4a-d** from D-mannose.

D-Mannose was converted to the glycosyl donor **5** according to our previously published procedure.¹⁰ The glycosyl donor **5** was condensed with 2-amino-6-chloropurine in the presence of TMSOTf as a Lewis acid to give the β -anomer **6** (30%) as a single stereoisomer (Scheme 1). The anomeric assignment was easily accomplished in a ¹H NMR experiment that showed a NOE between H-8 and 3'-H. Treatment of 2-amino-6-chloro derivative **6** with isoamyl nitrite, iodine, and methylene iodide in the presence of CuI afforded the 2-iodo-6-chloro derivative **7**, which was converted to the 2-iodo-6-amino derivative **8** upon treatment with methanolic ammonia. Sonogashira¹² coupling reaction of **8** with 1-hexyne in the presence of bis(triphenylphosphine)palladium dichloride yielded the 2-hexynyl derivative **9**. Finally, removal of the isopropylidene of **9** with 1 *N* HCl produced the final 2-hexynyl-4'-thioadenosine derivative **4a**. Suzuki¹³ coupling reaction of the 2-iodo derivative **8** with (*E*)-1-catecholboranylhexene¹⁴, prepared by treating with 1-hexyne and catecholborane, in the presence of tetrakis(triphenylphosphine)palladium(0) afforded the 2-hexenyl derivative **10**. Removal of the acetonide of **10** with 1 *N* HCl gave the 2-hexenyl-4'-thioadenosine derivative **4b**.

Using a strategy similar to Scheme 1, 8-substituted adenosine derivatives **4c** and **4d** were synthesized from the glycosyl donor **5** (Scheme 2). Condensation of **5** with 8-bromoadenine¹⁵ under Lewis acid conditions, afforded the 8-bromo derivative **11**. Coupling of **11** with 1-hexyne under Sonogashira conditions gave 8-hexynyl derivative **12**, which was treated with 1 *N* HCl to yield the final 8-hexynyl-4'-thioadenosine derivative **4c**.

The 8-bromo derivative **11** was condensed with (*E*)-1-catecholboranylhexene¹⁴ under Suzuki conditions¹³ to give the 2-hexenyl derivative **13**. Removal of the isopropylidene group of **13** under acidic conditions afforded the final 8-hexenyl-4'-thio adenosine derivative **4d**.

Binding assays were carried out using standard radioligands and membrane preparations from Chinese hamster ovary (CHO) cells stably expressing the human (h) A₁ or A₃AR or human embryonic kidney cells (HEK-293) expressing the hA_{2A}AR.¹⁶ Unlike the parent *N*⁶-substituted compound **3** that only weakly bound to the A_{2A}AR, C2-substituted variations of compound **3** led to a dramatic increase in the binding affinity ($K_i = 7.19 \pm 0.6$ nM for **4a** and 72.0 ± 19.1 nM for **4b**) at the hA_{2A}AR, while maintaining high binding affinity ($K_i = 11.8 \pm 1.3$ nM for **4a** and 13.2 ± 0.8 nM for **4b**) at the hA₃AR (Table 1).

These results indicate that bulky hydrophobic pockets exist in the binding sites of A_{2A}AR and A₃AR, allowing the C2-substituent to form favorable hydrophobic interactions. The 2-alkynyl derivative **4a** showed a better binding affinity than the 2-alkenyl derivative **4b**. Interestingly, C8-substitution on **3** abolished the binding affinity at the hA_{2A}AR, but the binding affinity at the hA₃AR was maintained although decreased. These findings suggest that a bulky hydrophobic group at position 8 could be tolerated at the binding site of the hA₃, but not hA_{2A}AR. The ability to enhance affinity at the A_{2A}AR in the truncated series by extending an unsaturated carbon chain at the 2 position implies a mode of receptor binding in common with the riboside series.¹¹ All compounds showed very weak binding affinity at the hA₁AR.

Compounds **4a** and **4b** were found to be potent and full antagonists in a cyclic AMP functional assay at the hA₃AR. In this assay, **4a** dose-dependently shifted the concentration-response curve for agonist Cl-IB-MECA to the right as an antagonist, corresponding to a K_B value of 1.69 nM calculated by Schild analysis (Figure 2). This is consistent with previous studies in which truncated *N*⁶-substituted 4'-thioadenosine derivatives have generally displayed A₃ AR antagonist activity.¹⁰ However, in a cyclic AMP functional assay at the hA_{2A}AR expressed in CHO cells, compound **4a** behaved as a full agonist compared to the standard 2-[p-(2-carboxyethyl)phenyl-ethylamino]-5'-*N*-ethylcarboxamidoadenosine (CGS21680) and displayed an EC₅₀ of 12 nM. At the hA_{2B}AR expressed in CHO cells, **4a** was a weak partial agonist in cyclic AMP accumulation (EC₅₀ ~10 μM). The finding that compound **4a** is both a potent and selective agonist at the hA_{2A}AR and a competitive antagonist at the hA₃AR is similar to the pharmacological profile of a more heavily 2,5'-substituted adenosine derivative¹⁷ which inhibited both formation of ROS and eosinophils degranulation for the anti-asthmatic activity.

To investigate the binding mode, we performed a study of docking the C2- and C8-substituted-4'-thioadenosine derivatives **4a-d** in the hA_{2A}AR X-ray crystallographic structure (PDB code: 3EML)¹⁸ using the GOLD software¹⁹, considering the flexibility of the binding site residues. As shown in Figure 3, the C2-substituted-4'-thioadenosine derivatives **4a** and **4b**, whose binding affinities are in the nM range, occupied the binding site very well. Their bulky and rigid C2-substituents oriented toward the extracellular region, forming hydrophobic interactions. The adenine moieties appeared to form three H-bonds with Glu169 and Asn253, and the ring systems were in π - π stacking with Phe168. Also, the thio-sugar rings were located deep inside the binding pocket, and the 3'-OH groups were able to donate a H-bond to Ser277. Based on this result, the C8-substituents were expected to be oriented in a hydrophobic pocket deep inside the binding site, which was not occupied by **4a** and **4b** in Figure 3. However, the C8-substituted derivatives **4c** and **4d** showed various binding modes (data not shown). In addition to the expected binding mode, the C8-substituents alternately pointed toward the extracellular region through a rotation of the bond between the adenine and thio-sugar rings. It might be due to spatial restriction of the long and rigid C8-substituents in the hydrophobic pocket inside the binding site, which would cause the nucleosides to lose some H-bonding and/or π - π stacking interactions that were shown for the C2-substituted derivatives. These results might explain why the A_{2A}AR affinities of **4c** and **4d** were reduced.

In conclusion, we synthesized the truncated C2- and C8-substituted-4'-thioadenosine derivatives **4a-d**, starting from D-mannose, using palladium-catalyzed cross coupling reactions as key steps. Although the antagonist activity of various truncated 4'-thionucleosides at the A₃AR was well explored previously, this is the first characterization of the functional activity of such derivatives at the A_{2A}AR. From this study, we successfully identified potent and sterically compact A_{2A}AR agonists, **4a** and **4b** by placing extended hydrophobic 2-hexynyl or 2-hexenyl groups on truncated and N⁶-unsubstituted 4'-thioadenosine derivatives. This observation was supported by molecular modeling that placed the chain at the 2 position in a hydrophobic region of the A_{2A}AR. However, C8-substitution greatly reduced binding affinity at the hA_{2A}AR. Thus, the absence of a 5'-uronamide of typical A_{2A}AR agonists² or the native -CH₂OH of adenosine did not preclude potent binding and full activation of the hA_{2A}AR. This suggested a major difference between A_{2A}AR and the A₃AR in the pathway of receptor activation. All synthesized compounds **4a-d** maintained their binding affinity at the human A₃AR, and as for the 2-H or 2-Cl analogues that were N⁶-substituted, competitive A₃AR antagonism was demonstrated. This study establishes that the truncated C2-substituted-4'-thioadenosine derivatives **4a** and **4b** can serve as a novel template for the development of new A_{2A}AR ligands, although they still may interact at the A₃AR. This mixed activity as A_{2A}AR agonist/A₃AR antagonist might also be advantageous in disease models such as asthma.¹⁷ Thorough elucidation of the structure-activity relationship of this series is in progress in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the Basic Science Research grant (NRF-2008-314-E00304), the National Core Research Center grant (R15-2006-020), the World Class University grant (R31-2008-000-10010-0) from National Research Foundation (NRF), Korea, and the Intramural Research Program of NIDDK, NIH, Bethesda, MD, USA.

References

1. Olah ME, Stiles GL. The role of receptor structure in determining adenosine receptor activity. *Pharmacol Ther* 2000;85:55–75. [PubMed: 10722120]
2. (a) Fredholm BB, Cunha RA, Svenningsson P. Pharmacology of adenosine receptors and therapeutic applications. *Curr Top Med Chem* 2002;3:413–426. [PubMed: 12570759] (b) Sitkovsky MV, Lukashev D, Apasov S, Kojima H, Koshiba M, Cladwell C, Ohta A, Thiel M. Physiological control of immune response and inflammatory tissue damage by hypoxia-inducible factors and adenosine A_{2A} receptors. *Ann Rev Immunol* 2004;22:657–682. [PubMed: 15032592] (c) Lappas CM, Sullivan GW, Linden J. Adenosine A_{2A} agonists in development for the treatment of inflammation. *Expert Opin Invest Drugs* 2005;14:797–806.
3. Svenningsson P, Hall H, Sedvall G, Fredholm BB. Distribution of adenosine receptors in the postmortem human brain: an extended autoradiographic study. *Synapse* 1997;27:322–335. [PubMed: 9372555]
4. Linden J. Cloned adenosine A₃ receptors: pharmacological properties, species differences and receptor functions. *Trends Pharmacol Sci* 1994;15:298–306. [PubMed: 7940998]
5. Baraldi PG, Cacciari B, Romagnoli R, Merighi S, Varani K, Borea PA, Spalluto G. A₃ adenosine receptor ligands: history and perspectives. *Med Res Rev* 2000;20:103–128. [PubMed: 10723024]
6. Jacobson KA, Gao ZG. Adenosine receptors as therapeutic targets. *Nature Rev Drug Disc* 2006;5:247–264.
7. (a) Jeong LS, Jin DZ, Kim HO, Shin DH, Moon HR, Gunaga P, Chun MW, Kim Y-C, Melman N, Gao Z-G, Jacobson KA. N⁶-Substituted D-4'-thioadenosine-5'-methyluronamides: Potent and selective agonists at the human A₃ adenosine receptor. *J Med Chem* 2003;46:3775–3777. [PubMed:

- 12930138] (b) Jeong LS, Lee HW, Jacobson KA, Kim HO, Shin DH, Lee JA, Gao ZG, Lu C, Duong HT, Gunaga P, Lee SK, Jin DZ, Chun MW, Moon HR. Structure-activity relationships of 2-chloro- N^6 -substituted-4'-thioadenosine-5'-uronamides as highly potent and selective agonists at the human A_3 adenosine receptor. *J Med Chem* 2006;49:273–281. [PubMed: 16392812]
8. Kim HO, Ji X-d, Siddiqi SM, Olah ME, Stiles GL, Jacobson KA. 2-Substitution of N^6 -benzyladenosine-5'-uronamides enhances selectivity for A_3 adenosine receptors. *J Med Chem* 1994;37:3614–3621. [PubMed: 7932588]
9. Lee EJ, Min HY, Chung HJ, Park EJ, Shin DH, Jeong LS, Lee SK. A novel adenosine analog, thio-Cl-IB-MECA, induces G_0/G_1 cell cycle arrest and apoptosis in human promyelocytic leukemia HL-60 cells. *Biochem Pharmacol* 2005;70:918–924. [PubMed: 16051194]
10. (a) Jeong LS, Choe SA, Gunaga P, Kim HO, Lee HW, Lee SK, Tosh DK, Patel A, Palaniappan KK, Gao ZG, Jacobson KA, Moon HR. Discovery of a new nucleoside template for human A_3 adenosine receptor ligands: D-4'-thioadenosine derivatives without 4'-hydroxymethyl group as highly potent and selective antagonists. *J Med Chem* 2007;50:3159–3162. [PubMed: 17555308] (b) Jeong LS, Pal S, Choe SA, Choi WJ, Jacobson KA, Gao Z-G, Klutz AM, Hou X, Kim HO, Lee HW, Tosh DK, Moon HR. Structure-activity relationships of truncated D- and L-4'-thioadenosine derivatives as species-independent A_3 adenosine receptor antagonists. *J Med Chem* 2008;51:6609–6613. [PubMed: 18811138]
11. (a) Matsuda A, Shinozaki M, Yamaguchi T, Homma H, Nomoto R, Miyasaka T, Watanabe Y, Abiru T. Nucleosides and Nucleotides. 103. 2-Alkynyladenosines: a novel class of selective adenosine A_2 receptor agonists with potent antihypertensive effects. *J Med Chem* 1992;35:241–252. [PubMed: 1732541] (b) Cristalli G, Volpini R, Vittori S, Camaioni E, Monopoli A, Conti A, Dionisotti S, Zocchi C, Ongini E. 2-Alkynyl derivatives of adenosine-5'-N-ethyluronamide (NECA): selective A_2 adenosine receptor agonists with potent inhibitory activity on platelet aggregation. *J Med Chem* 1994;37:1720–1726. [PubMed: 8201607] (c) Lambertucci C, Costanzi S, Vittori S, Volpini R, Cristalli G. Synthesis and adenosine receptor affinity and potency of 8-alkynyl derivatives of adenosine. *Nucleosides, Nucleotides & Nucleic Acids* 2001;20:1153–1157.
12. Chinchilla R, Nájera C. The Sonogashira reaction: A booming methodology in synthetic organic chemistry. *Chem Rev* 2007;107:874–922. [PubMed: 17305399]
13. Suzuki A. Carbon-carbon bonding made easy. *Chem Commun* 2005;38:4759–4763.
14. Miyaura N, Suzuki A. Palladium-catalyzed Reaction of 1-alkenylboronates with vinylic halides: (1Z,3E)-1-phenyl-1,3-octadiene. *Org Syn Coll Vol* 1993;8:532–534.
15. Laxer A, Major DT, Gottlieb HE, Fischer B. ($^{15}N_5$)-Labeled Adenine Derivatives: Synthesis and Studies of Tautomerism by ^{15}N NMR Spectroscopy and Theoretical Calculations. *J Org Chem* 2001;66:5463–5481. [PubMed: 11485471]
16. (a) Perreira M, Jiang JK, Klutz AM, Gao ZG, Shainberg A, Lu C, Thomas CJ, Jacobson KA. *J Med Chem* 2005;48:4910–4918. [PubMed: 16033270] (b) Jarvis MF, Schutz R, Hutchison AJ, Do E, Sills MA, Williams M. *J Pharmacol Exp Ther* 1989;251:888–893. [PubMed: 2600819] (c) Olah ME, Gallo-Rodriguez C, Jacobson KA, Stiles GL. *Mol Pharmacol* 1994;45:978–982. [PubMed: 8190112] (d) Nordstedt C, Fredholm BB. *Anal Biochem* 1990;189:231–234. [PubMed: 2177960]
17. Bevan N, Butchers PR, Cousins R, Coates J, Edgar EV, Morrison V, Sheehan MJ, Reeves J, Wilson DJ. Pharmacological characterisation and inhibitory effects of (2R,3R,4S,5R)-2-(6-amino-2-[[[(1S)-2-hydroxy-1-(phenylmethyl)ethyl]amino]-9H-purin-9-yl]-5-(2-ethyl-2H-tetrazol-5-yl)tetra hydro-3,4-furandiol, a novel ligand that demonstrates both adenosine A_{2A} receptor agonist and adenosine A_3 receptor antagonist activity. *Eur J Pharmacol* 2007;564:219–225. [PubMed: 17382926]
18. Jaakola VP, Griffith MT, Hanson MA, Cherezov V, Chien EYT, Lane JR, Ijzerman AP, Stevens RC. The 2.6 angstrom crystal structure of a human A_{2A} adenosine receptor bound to an antagonist. *Science* 2008;322:1211–1217. [PubMed: 18832607]
19. GOLD, version 4.1.2. Cambridge Crystallographic Data Centre; Cambridge, UK: 2009.




Figure 1.
The rationale for the design of A_{2A} adenosine receptor agonists




Figure 2. Parallel right shifts induced by compound **4a** on the concentration-response curve of a full agonist in the inhibition of cyclic AMP production at the hA_3AR expressed in CHO cells (A), the corresponding Schild plot (B) and the activity of **4a** as a full agonist at the $hA_{2A}AR$ expressed in CHO cells, compared to CGS21680 (C).






Figure 3.

Predicted binding modes of (A) **4a** and (B) **4b** docked in the hA_{2A}AR crystal structure. The key interacting residues are marked and displayed in capped-stick, except Phe168 in ball-and-stick, with carbon atoms in white. The ligands are depicted as ball-and-stick with carbon atoms in magenta (**4a**) and purple (**4b**). Hydrogen bonds are shown in yellow dashed lines. The Van der Waals surfaces of the ligands were generated by MOLCAD and colored by hydrogen bonding property (red: H-bond donating regions; blue: H-bond accepting regions). The Fast Connolly surface of the protein is Z-clipped and non-polar hydrogens are undisplayed for clarity.



Scheme 1. Synthesis of the 2-substituted derivatives 4a and 4b

Reagents and conditions: a) silylated 2-amino-6-chloropurine, TMSOTf, DCE, rt to 80 °C, 3 h; b) CuI, isoamyl nitrite, I₂, CH₂I₂, THF, 110 °C, 45 min; c) NH₃/MeOH, 80 °C, 2 h; d) 1-hexyne, CuI, TEA, DMF, bis(triphenylphosphine)palladium dichloride, rt, 3 h; e) 1 *N* HCl, THF, rt, 15 h; f) (*E*)-1-catecholboranylhexene, then tetrakis(triphenylphosphine)palladium(0), Na₂CO₃, DMF, H₂O, 90 °C, 15 h.



Scheme 2. Synthesis of the 8-substituted derivatives 4c and 4d

Reagents and conditions: a) silylated 8-bromoadenine, TMSOTf, DCE, rt to 90 °C, 2 h; b) 1-hexyne, CuI, TEA, DMF, bis(triphenylphosphine) palladium dichloride, rt, 3 h; c) 1 *N* HCl, THF, rt, 15 h; d) (*E*)-1-catecholboranylhexene, then tetrakis(triphenylphosphine) palladium(0), Na₂CO₃, DMF, H₂O, 90 °C, 15 h.

Table 1

Binding affinities of known A₃AR antagonist **3** and truncated 2- and 8-substituted-4'-thioadenosine derivatives **4a-d** at three subtypes of hARs.

Compounds	Affinity ^a		
	hA ₁ (% inhibition)	hA _{2A} (% inhibition)	hA ₃ (K _i , nM)
3	37.9%	17.7%	1.66 ± 0.90
4a	38.9 ± 9.9%	97.2 ± 4.1%	11.8 ± 1.3
4b	16.2 ± 8.4%	95.9 ± 8.7%	13.2 ± 0.8
4c	49.3 ± 4.9%	46.5 ± 4.3%	20.0 ± 4.0
4d	3.7 ± 2.9%	22.8 ± 6.4%	259 ± 10

^a All binding experiments were performed using adherent mammalian cells stably transfected with cDNA encoding the appropriate hAR (A₁AR and A₃AR in CHO cells and A_{2A}AR in HEK-293 cells). Binding was carried out using 1 nM [³H]CCPA, 10 nM [³H]CGS21680, or 0.5 nM [¹²⁵I]-AB-MECA as radioligands for A₁, A_{2A}, and A₃ARs, respectively. Values are expressed as mean ± sem, n = 3–4 (outliers eliminated) and normalized against a non-specific binder, 5'-N-ethylcarboxamidoadenosine (NECA, 10 μM). Values expressed as a percentage refer to percent inhibition of specific radioligand binding at 10 μM, with nonspecific binding defined using 10 μM NECA.