

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/231526719>

Mechanistic Studies of Palladium(II)-Catalyzed Hydrosilation and Dehydrogenative Silation Reactions

ARTICLE *in* JOURNAL OF THE AMERICAN CHEMICAL SOCIETY · FEBRUARY 1997

Impact Factor: 12.11 · DOI: 10.1021/ja962979n

CITATIONS

139

READS

33

3 AUTHORS, INCLUDING:



Anne M. LaPointe

Cornell University

26 PUBLICATIONS 890 CITATIONS

SEE PROFILE

Mechanistic Studies of Palladium(II)-Catalyzed Hydrosilation and Dehydrogenative Silation Reactions


Anne M. LaPointe, Francis C. Rix, and Maurice Brookhart*

Contribution from the Department of Chemistry, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599-3290

Received August 26, 1996[⊗]

Abstract: The cationic Pd(II) complexes, $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{L})]^+[\text{BAR}'_4]^-$ phen = 1,10-phenanthroline; L = Et₂O, Me₃SiC≡CSiMe₃; Ar' = 3,5-(CF₃)₂C₆H₃) catalyze the hydrosilation and dehydrogenative silation of olefins. Hydrosilation of ethylene, *tert*-butylethylene, 1-hexene, and cyclohexene by HSiR₃ (R = CH₂CH₃, C₆H₅) occurs in the presence of 1 mol % $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{L})]^+[\text{BAR}'_4]^-$. The reaction of *tert*-butylethylene with HSi(*i*-Pr)₃ in the presence of $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{L})]^+[\text{BAR}'_4]^-$ yields neohexane and *t*-BuCH=CHSi(*i*-Pr)₃. Low-temperature NMR experiments revealed that the catalyst resting state for the silations of ethylene and alkyl-substituted olefins is $[(\text{phen})\text{Pd}(\text{SiR}_3)(\eta^2\text{-H}_2\text{C=CHR}')^+][\text{BAR}'_4]^-$. Evidence for rapid, reversible silyl migration at -70 °C was observed by ¹H NMR spectroscopy. Deuterium labeling studies show that the intermediate Pd(II) alkyl complexes can isomerize via a series of β -hydride eliminations followed by reinsertions of olefin prior to reaction with DSiEt₃. Styrene undergoes both hydrosilation and dehydrogenative silation in the presence of $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{L})]^+[\text{BAR}'_4]^-$ or $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_3)\text{C}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ yielding ethylbenzene, R₃SiCH₂CH₂C₆H₅ and *trans*-R₃SiCH=CHPh (R = CH₂-CH₃, CH(CH₃)₂). ¹H NMR spectroscopy revealed that the π -benzyl complexes $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_2\text{SiR}_3)\text{C}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ and $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_3)\text{C}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ are the catalyst resting states for the silation reactions of styrene.


Hydrosilation of olefins (eq 1) can be achieved either through



a free radical process or with the use of a transition metal catalyst, typically an electron-rich complex of a late transition metal such as Co, Rh, Pd, or Pt.^{1,2} More recently, zirconocene derivatives,^{3,4} lanthanide complexes,^{5–7} and electrophilic late transition metal complexes⁸ have been reported to catalyze the hydrosilation of olefins.

A variety of mechanisms have been proposed for this process. A widely accepted mechanism was first proposed by Chalk and Harrod (Scheme 1).^{9,10} The key feature in the Chalk–Harrod mechanism is migratory insertion of an olefin hydride complex followed by reductive coupling of the alkyl and silyl fragments.^{9,10} If the intermediate alkyl complex undergoes reversible β -hydride elimination and reinsertion with opposite regiochemistry, then the Chalk–Harrod mechanism provides an explanation for the olefin isomerization and deuterium scrambling observed in many hydrosilation reactions.¹

However, many catalysts form vinylsilanes and alkanes in addition to the hydrosilation product (eq 2).^{1,11–14} In some cases, dehydrogenative silation occurs more readily than



hydrosilation.^{13–15} Because of the utility of vinylsilanes in organic synthesis,¹⁶ the dehydrogenative silation reaction is of interest. The Chalk–Harrod mechanism does not account for dehydrogenative silation, and a number of groups have proposed mechanisms based on silyl (rather than hydride) migration to olefin.^{8,13,15,17–19}

One such possible mechanism is shown in Scheme 2. Silyl migration to olefin results in a β -silylalkyl intermediate, which can then undergo two possible reactions. The first possibility is β -hydride elimination to form an olefin–hydride complex. If reinsertion of the olefin with opposite regiochemistry occurs, isomerization of the alkyl complex results. Displacement of the olefin from the metal results in dehydrogenative silation. The second possibility is cleavage of the β -silylalkyl fragment by reaction by HSiR₃. This reaction may proceed via oxidative addition of silane followed by reductive elimination of the hydrosilated product or via a σ -bond metathesis reaction.

Efforts to detect intermediate species in many transition metal catalyzed hydrosilation reactions have been hindered by extremely high reaction rates¹ and heterogeneous systems.²⁰ In an earlier study from these laboratories, the cationic cobalt

[⊗] Abstract published in *Advance ACS Abstracts*, January 1, 1997.

(1) Ojima, I. In *The Chemistry of Organic Silicon Compounds*; Patai, S., Rappaport, S., Eds.; Wiley: New York, 1989.

(2) Collman, J. P.; Hegedus, L. S.; Norton, J. R.; Finke, R. G. *Principles and Applications of Organotransition Metal Complexes*; University Science Books: Mill Valley, CA, 1987.

(3) Takahashi, T.; Hasegawa, M.; Suzuki, M.; Saburi, M.; Rousset, C. J.; Fanwick, P. E.; Negishi, E.-I. *J. Am. Chem. Soc.* **1991**, *113*, 8564.

(4) Kesti, M. R.; Waymouth, R. M. *Organometallics* **1992**, *11*, 1095.

(5) Fu, P.-F.; Brard, L.; Li, Y.; Marks, T. J. *J. Am. Chem. Soc.* **1995**, *117*, 7157.

(6) Molander, G. A.; Julius, M. *J. Org. Chem.* **1992**, *57*, 6347.

(7) Sakakura, T.; Lautenschlager, H.-J.; Tanaka, M. *J. Chem. Soc., Chem. Commun.* **1991**, 40.

(8) Brookhart, M.; Grant, B. E. *J. Am. Chem. Soc.* **1993**, *115*, 2151.

(9) Chalk, A. J.; Harrod, J. F. *J. Am. Chem. Soc.* **1965**, *87*, 16.

(10) Harrod, J. F.; Chalk, A. J. *J. Am. Chem. Soc.* **1965**, *87*, 1133.

(11) Ojima, I.; Fuchikami, T.; Yatabe, M. *J. Organomet. Chem.* **1984**, *260*, 335.

(12) Onopchenko, A.; Sabourin, E. T.; Beach, D. L. *J. Org. Chem.* **1983**, *48*, 5101.

(13) Takeuchi, R.; Yasue, H. *Organometallics* **1996**, *15*, 2098.

(14) Christ, M. L.; Sabo-Etienne, S.; Chaudret, B. *Organometallics* **1995**, *14*, 1082.

(15) Tanke, R. S.; Crabtree, R. H. *Organometallics* **1991**, *10*, 415.



(16) Fleming, I.; Donogues, J.; Smithers, R. *Org. React.* **1989**, *37*, 57.

(17) Randolph, C. L.; Wrighton, M. S. *J. Am. Chem. Soc.* **1986**, *108*, 2266.


(18) Seitz, F.; Wrighton, M. S. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 289.

(19) Bergens, S. H.; Noheda, P.; Whelan, J.; Bosnich, B. *J. Am. Chem. Soc.* **1992**, *114*, 2128.

(20) Lewis, L. N. *J. Am. Chem. Soc.* **1990**, *112*, 5998.


Scheme 1. Chalk–Harrod Mechanism for the Hydrosilation of Olefins**Scheme 2.** Silyl Migration Mechanism for Hydrosilation and Dehydrogenative Silylation of Olefins

complex $[\text{Cp}^*\text{Co}(\text{P}(\text{OMe})_3)(\text{CH}_2\text{CH}_2-\mu\text{-H})][\text{BAR}'_4]^-$ was found to be an efficient precatalyst for the hydrosilation of 1-hexene. Deuterium labeling studies and spectroscopic observation of intermediate species in this system suggested that the reaction proceeds by silyl migration to olefin followed by a series of β -hydrogen elimination and reinsertion reactions. The terminal cobalt alkyl complex $[\text{Cp}^*\text{Co}(\text{CH}_2\text{CH}_2-\mu\text{-H})(\text{CH}_2)_4\text{SiEt}_3)(\text{P}(\text{OMe})_3)][\text{BAR}'_4]^-$ then reacts with HSiEt_3 to form $\text{Et}_3\text{Si}(\text{CH}_2)_5\text{CH}_3$ (eq 3). Thus, silane adds with overall 1,6 regiochemistry.⁸



Cationic square-planar Pd(II) complexes have been found to be efficient catalysts for the formation of carbon–carbon bonds, including the dimerization^{21,22} and polymerization of α -olefins²³ and the alternating co-polymerization of α -olefins with carbon monoxide.^{24–28} We report here that the cationic Pd(II) com-

plexes, $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{L})][\text{BAR}'_4]^-$ (phen = 1,10-phenanthroline; L = Et_2O (**1a**), $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$ (**1b**); $\text{Ar}' = 3,5\text{-(CF}_3)_2\text{C}_6\text{H}_3$) are efficient precatalysts for the formation of silicon–carbon bonds.




1a: L = Et_2O
1b: L = $\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$

Hydrosilation and dehydrogenative silylation of olefins are achieved under mild conditions (25–85 °C, 0.1–2 mol % **1**). The mechanisms of the hydrosilation and dehydrogenative silylation reactions were studied by a variety of low-temperature NMR experiments and deuterium labeling experiments. The results of these studies are presented below.

Results and Discussion

I. Survey of Silylation Reactions. A. Silylation of Ethylene and Alkyl-Substituted Olefins. In the presence of a catalytic amount of $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{L})][\text{BAR}'_4]^-$, hydrosilation of olefins proceeds at 25 °C in CH_2Cl_2 (eq 4). The addition of HSiR_3 to



α -olefins proceeds with a 1,2 regiochemistry. **1b** is a more convenient catalyst than the thermally unstable ether adduct **1a** since **1b** is thermally stable, moderately air- and water-stable, and stable at 25 °C in CH_2Cl_2 . The results are summarized in Table 1.

When the reaction was conducted in Et_2O , lower yields were obtained. Likewise, only 5% yield is obtained when $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{NCCH}_3)(\text{NCCH}_3)][\text{BAR}'_4]^-$ (**1c**) is used as the catalyst. The low yields are due to inhibition of the silylation reaction rather than catalyst decomposition; NMR studies (see below) suggest that coordinating ligands such as acetonitrile or ether can compete with olefin or silane for the vacant coordination site and thus inhibit the hydrosilation reaction.

No dehydrogenative silylation is observed in the reactions of ethylene, 1-hexene, or cyclohexene with HSiR_3 (R = Et, Ph). Some dehydrogenative silylation (5%) is observed in the reaction of *tert*-butylethylene with HSiEt_3 in the presence of **1a** or **1b**. Efforts to increase the yield of *t*-BuCH=CHSiEt₃ by increasing the *t*-BuCH=CH₂/HSiEt₃ ratio were not successful.

The reaction of ethylene with $\text{HSi}(i\text{-Pr})_3$ in the presence of 1 mol % **1a** yields the hydrosilation product, *i*-Pr₃SiEt (74% yield). In contrast, the reaction of *tert*-butylethylene with 0.5 equiv of $\text{HSi}(i\text{-Pr})_3$ in the presence of 1% **1a** yields *trans*-*t*-BuCH=CHSi-

(21) Rix, F. C.; Brookhart, M. *J. Am. Chem. Soc.* **1995**, *117*, 1137.

(22) DiRenzo, G. M.; White, P. S.; Brookhart, M. *J. Am. Chem. Soc.* **1996**, *118*, 6225.

(23) Johnson, L. K.; Killian, C. M.; Brookhart, M. *J. Am. Chem. Soc.* **1995**, *117*, 6414.

(24) Sen, A. *Acc. Chem. Res.* **1993**, *26*, 303.

(25) Rix, F. C.; Brookhart, M.; White, P. S. *J. Am. Chem. Soc.* **1996**, *118*, 4746.

(26) Brookhart, M.; Rix, F. C.; DeSimone, J. M.; Barborak, J. C. *J. Am. Chem. Soc.* **1992**, *114*, 5894.

(27) Drent, E.; van Broekhoven, J. A. M.; Doyle, M. *J. Organomet. Chem.* **1991**, *417*, 235.


(28) van Asselt, R.; Gielens, E. E. C. G.; Rülke, R. E.; Vrieze, K.; Elsevier, C. *J. Am. Chem. Soc.* **1994**, *116*, 997.

Table 1. Summary of Reaction Conditions^{a,c} and Yields^b for the Reaction of Ethylene and Alkyl-Substituted Olefins with HSiR₃ in the Presence of 1 mol % [(phen)Pd(CH₃)(L)]⁺[BAR'₄]⁻ (**1a**, L = Et₂O; **1b**, L = Me₃CCSiMe₃; **1c**, L = CH₃CN)

entry no.	HSiR ₃ R =	olefin	catalyst	substrate ratio, HSiR ₃ /olefin	solvent	product	yield, ^b %
1	Et	ethylene	1a	<i>c</i>	CH ₂ Cl ₂	SiEt ₄	85
2	Et	1-hexene	1b	1.1	CH ₂ Cl ₂	Et ₃ Si(CH ₂) ₅ CH ₃	88
3	Et	cyclohexene	1b	1.1	CH ₂ Cl ₂	Et ₃ Si(Cy)	88
4	Et	<i>t</i> -BuCH=CH ₂	1b	1.1	CH ₂ Cl ₂	Et ₃ SiCH ₂ CH ₂ - <i>t</i> -Bu ^d	79
5	Et	<i>t</i> -BuCH=CH ₂	1b	1.0	CH ₂ Cl ₂	Et ₃ SiCH ₂ CH ₂ - <i>t</i> -Bu ^d	80
6	Et	<i>t</i> -BuCH=CH ₂	1b	1.1	Et ₂ O	Et ₃ SiCH ₂ CH ₂ - <i>t</i> -Bu ^d	14
7	Et	<i>t</i> -BuCH=CH ₂	1c	1.1	CH ₂ Cl ₂	Et ₃ SiCH ₂ CH ₂ - <i>t</i> -Bu ^d	5
8	Ph	cyclohexene	1b	1.1	CH ₂ Cl ₂	Ph ₃ Si(Cy)	78
9	Ph	<i>t</i> -BuCH=CH ₂	1b	1.1	CH ₂ Cl ₂	Ph ₃ SiCH ₂ CH ₂ - <i>t</i> -Bu	20 ^e
10	<i>i</i> -Pr	ethylene	1a	<i>c</i>	CH ₂ Cl ₂	Si(<i>i</i> -Pr) ₃ Et	75
11	<i>i</i> -Pr	<i>t</i> -BuCH=CH ₂	1a	0.5	CH ₂ Cl ₂	(<i>i</i> -Pr) ₃ SiCH=CH- <i>t</i> -Bu	82
12	<i>i</i> -Pr	<i>t</i> -BuCH=CH ₂	1a	1.1	CH ₂ Cl ₂	(<i>i</i> -Pr) ₃ SiCH=CH- <i>t</i> -Bu	94 ^f


^a 24 h, 25 °C. ^b Yields were calculated based on the limiting reagent. ^c C₂H₄ was bubbled through the solution until C₂H₄ uptake ceased. ^d Et₃SiCH=CH-*t*-Bu (5%) was also formed. ^e Recrystallized. ^f An equivalent amount of *t*-BuCH₂CH₃ was also produced.

(*i*-Pr)₃ (82% yield) (eq 5). The hydrosilation product, *t*-BuCH₂-



CH₂Si(*i*-Pr)₃, is not observed. When a slight excess of HSi(*i*-Pr)₃ (1.1 equiv) was used, no hydrosilation is observed.

B. Silation Reactions of Styrene. The reaction of styrene with HSiEt₃ in the presence of **1b** or [(phen)Pd(η³-CH(CH₃)-C₆H₅)]⁺ (**7**)²⁹ was investigated. The results are summarized in Table 2. Both hydrosilation and dehydrogenative silation of styrene occur, yielding a mixture of ethylbenzene, Et₃SiCH₂-CH₂Ph, and *trans*-Et₃SiCH=CHC₆H₅ (eq 6). The product ratio Et₃SiCH₂CH₂Ph/*trans*-Et₃SiCH=CHC₆H₅ is dependent on the styrene/HSiEt₃ ratio. Hydrosilation predominates when styrene/HSiEt₃ < 1 (entry nos. 1 and 5). Dehydrogenative silation predominates when styrene/HSiEt₃ ≥ 1. At 85 °C, yields are higher.




The reaction of HSi(*i*-Pr)₃ with styrene in the presence of 1 mol % **7** was investigated. Higher temperatures (85 °C) are required. Under identical conditions, the proportion of dehydrogenative silation observed is higher than in analogous reactions of HSiEt₃.

II. Silation of Ethylene and Alkyl-Substituted Olefins.

A. NMR Studies. The hydrosilation and dehydrogenative silation reactions were investigated by low-temperature NMR spectroscopic methods in order to gain mechanistic insight. Complexes containing the [BAR'₄]⁻ anion frequently exhibit enhanced stability and solubility and are well-suited for low-temperature NMR studies.^{8,22,25,29,30}

The reaction of the precatalyst **1a** with 2 equiv of HSiR₃ (R = Et, Ph) at -78 °C in CD₂Cl₂ proceeds with loss of methane and formation of a new complex with the general formation [(phen)Pd(SiR₃)₂(H)]⁺[BAR'₄]⁻ (**2a**, R = Et; **2b**, R = Ph) (eq 7).




(29) Rix, F. C.; Brookhart, M.; White, P. S. J. Am. Chem. Soc. **1996**, *118*, 2436.

Table 2. Reaction Conditions^{a,c} and Product Ratios^d for the Reaction of Styrene with HSiR₃ (R = Et, *i*-Pr) in the Presence of 1 mol % [(phen)Pd(η³-CH(CH₃)-C₆H₅)]⁺[BAR'₄]⁻ (**7**)

entry no.	HSiR ₃ R =	substrate ratio, styrene/HSiR ₃	T, °C	solvent	yield, ^{b,d} %	product ratio, PhCH ₂ CH ₂ SiR ₃ / PhCH=CHSiR ₃
1	Et	1:4	35	CH ₂ Cl ₂	27	100:0
2	Et	1:1	35	CH ₂ Cl ₂	50	45:55
3	Et	2:1	35	CH ₂ Cl ₂	58	25:75
4	Et	4:1	35	CH ₂ Cl ₂	77	13:87
5	Et	1:4	85	none ^c	85	96:4
6	Et	1:1	85	none ^c	65	43:57
7	Et	2:1	85	none ^c	84	32:68
8	Et	4:1	85	none ^c	88	21:79
9	<i>i</i> -Pr	1:4	85	none ^c	58	41:59
10	<i>i</i> -Pr	1:1	85	none ^c	52	13:87
11	<i>i</i> -Pr	2:1	85	none ^c	82	9:91
12	<i>i</i> -Pr	4:1	85	none ^c	97	0:100

^a 24 hours. ^b Yield = (PhCH₂CH₂SiR₃ + PhCH=CHSiR₃). The yield is calculated based on the limiting reagent. For the reactions of equimolar amounts of HSiR₃ and styrene, the isolated yield will be <100% due to the conversion of styrene into ethylbenzene. ^c At 85 °C, **7** is soluble in neat HSiR₃/styrene and no additional solvent was used. ^d Ethylbenzene was removed *in vacuo* during product isolation. For this reason, the yield of ethylbenzene was not calculated but was assumed to be equal to the yield of PhCH=CHSiR₃.

Scheme 3. Possible Structures for **2**




In the presence of only 1 equiv of silane, a 1:1 mixture of **2** and **1a** is obtained. In **2a** the two SiEt₃ fragments are equivalent by ¹H and ¹³C NMR spectroscopy at temperatures down to -95 °C. A singlet is observed at δ = -9.99 (**2a**) or -12.13 (**2b**) and is attributed to Si-H. At -95 °C no ²⁹Si satellites are observed, but the peak is quite broad and ²⁹Si-H coupling constants less than 40 Hz would not be detected. Based on the spectroscopic evidence **2a** and **2b** are either Pd(IV) hydrides with the general formula [(phen)Pd(SiR₃)₂(H)]⁺[BAR'₄]⁻ or a rapidly fluxional Pd(II) species which contains an η¹-silyl group and a η²-silane (Scheme 3). A less likely possibility is that the hydride is symmetrically bridged between the silyl groups, as

(30) Brookhart, M.; Grant, B.; Volpe, A. F. Organometallics **1992**, *11*, 3920.


shown in Scheme 3. The reaction of **1a** with 2 equiv of $\text{HSi}(i\text{-Pr})_3$ in CD_2Cl_2 at -78°C does not yield any well-defined species, although CH_4 evolution is observed. It is possible that the larger steric requirements of $\text{Si}(i\text{-Pr})_3$ prevent formation of a species with the general formula $[(\text{phen})\text{Pd}(\text{Si}(i\text{-Pr})_3)_2(\text{H})]^+[\text{BAR}'_4]^-$.

HSiEt_3 is readily displaced from **2a** by a variety of ligands, allowing the formation of a variety of cationic Pd(II) silyl complexes. Addition of triphenylphosphine to **2a** results in formation and isolation of $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\text{PPh}_3)]^+[\text{BAR}'_4]^-$ (**3a**) (eq 8). When allyldiphenylphosphine is added to **2a** at



-78°C , orange $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\text{Ph}_2\text{PC}_3\text{H}_5)]^+[\text{BAR}'_4]^-$ (**3b**) is isolated. ^1H and ^{31}P NMR spectroscopy suggest that in **3b** allyldiphenylphosphine is coordinated to palladium via phosphorus rather than through the olefin. The isolable Pd silyl complexes **3a,b** are not suitable hydrosilation precatalysts; no reaction of cyclohexene with HSiEt_3 was observed in the presence of 1 mol % **3a** or **3b** (CH_2Cl_2 , 25°C).

Addition of ethylene to a solution of **2a** at -78°C results in displacement of HSiEt_3 and formation of $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\eta^2\text{-C}_2\text{H}_4)]^+[\text{BAR}'_4]^-$ (**4a**) (eq 9). No hydrosilation is observed at -78°C . In the presence of excess ethylene, exchange between free and bound ethylene is observed at -78°C and only a single averaged signal can be observed ($\delta = 5.0$ ppm). In the absence of excess ethylene, the coordinated ethylene ligand displays an AA'BB' pattern centered at 4.9 ppm. The olefinic doublets collapse to a singlet as the temperature is raised. At the coalescence temperature, $T_c = -70(1)^\circ\text{C}$, $k = 3.9 \times 10^2 \text{ s}^{-1}$, $\Delta G^\ddagger = 9.2 \text{ kcal/mol}$. The fluxional behavior of **4a** is similar to that observed for the cationic alkyl complexes $[(\text{phen})\text{Pd}(\text{R})(\text{C}_2\text{H}_4)]^+[\text{BAR}'_4]^-$ ($\text{R} = \text{CH}_3$, $\Delta G^\ddagger = 9.2(0.2) \text{ kcal/mol}$; $\text{R} = \text{C}_2\text{H}_5$, $\Delta G^\ddagger = 9.2(0.2) \text{ kcal/mol}$).^{21,25}



Addition of *tert*-butylethylene to a CD_2Cl_2 solution of **2a** at -78°C results in displacement of HSiEt_3 and formation of $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\eta^2\text{-CH}_2\text{CH}(t\text{-Bu}))]^+[\text{BAR}'_4]^-$ (**4b**). At -78°C , no exchange is observed between free and bound *t*-butylethylene.

The reaction of **1a** with 1 equiv of HSiEt_3 followed by addition of $\text{H}_2\text{C}=\text{CHR}'$ ($\text{R}' = \text{H}, t\text{-Bu}$) at -78°C yields **4a** and **4b**, respectively. This method permits the generation of silane-free solutions of **4a,b** as well as the generation of tri-(isopropyl)silyl analogues of **4**; the reaction of **1a** with 1 equiv of $\text{HSi}(i\text{-Pr})_3$ at -78°C followed by addition of *tert*-butylethylene results in formation of $[(\text{phen})\text{Pd}(\text{Si}(i\text{-Pr})_3)(\eta^2\text{-CH}_2\text{CH}(t\text{-Bu}))]^+[\text{BAR}'_4]^-$ (**4c**).

The rapid reaction of the precatalyst **1a** with 2 equiv of HSiR_3 to form **2** and the subsequent displacement of HSiR_3 by olefin to form **4** demonstrate how catalyst generation occurs in the silation reactions. Additionally, the reaction of **1a** with HSiR_3 suggests that alkyl complexes of the general formula $[(\text{phen})\text{PdR}''(\text{L})]^+$ ($\text{R}'' = \text{alkyl}$) react readily with silane to form alkane





Figure 1. A plot for k_{obs} vs $[\text{HSiEt}_3]$ for the reaction of $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\text{H}_2\text{C}=\text{CH}-t\text{-Bu})]^+$ (**4b**) with HSiEt_3 to form **2a** and $t\text{-BuCH}_2\text{-CH}_2\text{SiEt}_3$ (-38°C ; $[\text{4b}] = 0.014 \text{ M}$; $k_{\text{obs}} = 1.7 \times 10^{-3} \text{ s}^{-1} [\text{HSiEt}_3]$).


and regenerate a Pd(II) silyl complex; this reaction is a key step in the catalytic hydrosilation reaction (see below).

In the presence of **4a** and excess HSiEt_3 and ethylene (2–10 equiv), hydrosilation proceeds readily at -40°C . During catalysis, **4a** is the only organometallic species observed by ^1H NMR spectroscopy. Upon complete consumption of ethylene, **2a** is regenerated. Similar results were obtained in the reaction of **4b** with excess *tert*-butylethylene and HSiEt_3 . These results establish that the silyl-olefin complexes $[(\text{phen})\text{Pd}(\text{SiR}_3)(\eta^2\text{-CH}_2\text{CHR}')^+[\text{BAR}'_4]^-$ are the catalyst resting state for the silation of olefins. This led us to propose a mechanism consisting of silyl migration to olefin followed by reaction of the β -silylethyl complex with HSiR_3 to form the hydrosilation product and regenerate $[\text{Pd}]\text{-SiR}_3$ (eq 10).




Based on the rapid reaction of **1a** with HSiEt_3 and the rapid formation of **4** from **2**, it was anticipated that the rate-determining step would be the silyl migration to olefin, and the hydrosilation reaction would be zero-order in $[\text{HSiEt}_3]$. CD_2Cl_2 solutions of **4a** and **4b** were monitored by ^1H NMR spectroscopy in the temperature range -60 to -20°C . It was found that in the absence of HSiEt_3 , complexes **4a,b** are stable in CD_2Cl_2 at temperatures up to ca. -20°C . Neither the β -silylalkyl complexes $[(\text{phen})\text{Pd}(\text{CHR}'\text{CH}_2\text{SiEt}_3)]^+$ ($\text{R}' = \text{H}, t\text{-Bu}$) nor any other species attributable to β -silyl migration was observed. This result suggested that β -silyl migration ($\text{4} \rightleftharpoons [\text{Pd}]\text{-R}''$) may be rapid and reversible, favoring **4** in the catalytic cycle. To examine this possibility, the kinetics of the reaction of **4b** with HSiEt_3 under pseudo-first-order conditions (10–30 equiv of HSiEt_3) was monitored at -38°C . A first-order rate dependence on $[\text{HSiEt}_3]$ was observed at all silane concentrations. ($k_{\text{obs}} = 1.7 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1} [\text{HSiEt}_3]$) (Figure 1).

A crossover experiment was performed to determine whether the silyl and olefin fragments in complexes **4** are coupled in the hydrosilation product or if the bound olefin fragment is in some way attacked by free HSiR_3 . **4b** was generated in CD_2Cl_2 at -78°C ; care was taken not to introduce excess HSiEt_3

Scheme 4. Nondegenerate Silyl Scrambling and Olefin Displacement Reactions in $[(\text{phen})\text{Pd}(\text{SiR}_3)(\eta^2\text{-CH}_2=\text{CHCH}_2\text{SiR}'_3)]^+$ 

or *tert*-butylethylene. HSiPh_3 was added and the solution was allowed to warm to 25 °C. The hydrosilation product was analyzed by ^1H and ^{13}C NMR spectroscopy and found to be exclusively *t*- $\text{BuCH}_2\text{CH}_2\text{SiEt}_3$ (eq 11). The results of this




experiment suggest that intramolecular silyl migration to olefin occurs rather than an intermolecular reaction of bound *tert*-butylethylene with HSiPh_3 .

The results of the kinetics and crossover experiments strongly support rapid, reversible silyl migration. To test the reversibility of the silyl migration, a silyl scrambling experiment was performed (Scheme 4). $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\eta^2\text{-H}_2\text{C}=\text{CHCH}_2\text{-SiPh}_3)]^+[\text{BAR}'_4]^-$ (**5a**) was generated in CD_2Cl_2 at -78 °C by addition of allyltriphenylsilane to a solution of **2a**. If silyl migration to olefin occurs with a 2,1 regiochemistry (as is suggested by the regiochemistry of the hydrosilated products), the Pd(II) alkyl intermediate would have two nonequivalent β -silyl substituents. If β -silyl elimination occurs it would be nondegenerate and would result in formation of a mixture of **5a** and a new complex, $[(\text{phen})\text{Pd}(\text{SiPh}_3)(\eta^2\text{-H}_2\text{C}=\text{CHCH}_2\text{-SiEt}_3)]^+[\text{BAR}'_4]^-$.

5a was generated at -78 °C and the NMR sample was inserted into a precooled (-78 °C) NMR probe. Initially, $\approx 95\%$ **5a** was observed, but as the sample was allowed to warm to -60 °C, a mixture of **5a** (87%) and a new species **5b** (13%) whose ^1H NMR spectrum closely resembled that of **5a** was formed. $[(\text{phen})\text{Pd}(\text{SiPh}_3)(\eta^2\text{-H}_2\text{C}=\text{CHCH}_2\text{SiEt}_3)]^+[\text{BAR}'_4]^-$ was independently generated at -78 °C and its NMR spectrum was identical with that of complex **5b**. At -60 °C, **5b** rapidly equilibrated to a mixture of **5a** (87%) and **5b** (13%). For the equilibrium $\text{5a} \rightleftharpoons \text{5b}$, $K(\text{eq}) = 0.15$ at -60 °C. These observations clearly establish that rapid and reversible silyl migration is occurring at -60 °C.

After an hour at -60 °C, two new species were also observed and identified as $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\eta^2\text{-H}_2\text{C}=\text{CHCH}_2\text{SiEt}_3)]^+[\text{BAR}'_4]^-$ (**5c**) and $[(\text{phen})\text{Pd}(\text{SiPh}_3)(\eta^2\text{-H}_2\text{C}=\text{CHCH}_2\text{SiPh}_3)]^+[\text{BAR}'_4]^-$ (**5d**) (based on independent generation of these complexes).

The observed preference for **5a** may be due to preferential binding of the more electron-rich silyl fragment SiEt_3 to the electrophilic Pd(II) cation. The eventual formation of **5c** and **5d** results from nondegenerate exchange of allylsilane. The exchange may be explained by either an associative or dissociative mechanism in the presence of traces of free allylsilane. Mechanistic studies in related cationic Pd(II) systems suggest that olefin displacement proceeds by an associative mechanism.^{23,29}


Scheme 5. Reactions of α -Olefins with DSiEt_3 in the Presence of **1b**

At -70 °C, the equilibration of **5a** and **5b** occurred competitively with the displacement of HSiR_3 from **2**. Accurate rate constants for the interconversion of **5a** and **5b** could not be obtained.

B. Deuterium Labeling Studies. Deuterium labeling studies were conducted to determine whether β -hydride elimination and reinsertion reactions occur in the β -silylalkyl intermediates prior to cleavage by silane. The reactions of cyclohexene, *tert*-butylethylene, and 1-hexene with DSiEt_3 in the presence of **1b** were investigated. The catalyst and substrate concentrations were similar to those used in the bulk hydrosilations. The deuteriosilated products were isolated and analyzed by ^{13}C NMR spectroscopy. The results are shown in Scheme 5.


For cyclohexene, deuteriosilation occurs, with a 1,2 regiochemistry and $<5\%$ deuterium incorporation is observed at any other position. These results are consistent with migratory insertion of cyclohexene followed by reaction of the Pd cyclohexyl intermediate with DSiEt_3 , as shown in Scheme 5.

In contrast, the deuteriosilation reaction of *tert*-butylethylene proceeds with a net 1,1 regiochemistry, and *t*- $\text{BuCH}_2\text{CHDSiEt}_3$ is the only product observed by ^{13}C NMR spectroscopy. A plausible mechanism is proposed consisting of migratory insertion of *tert*-butylethylene, followed by β -hydride elimina-



styrene, CD_2Cl_2 solutions of **6** are stable for several days at 25 °C; however, when solutions of **6** were concentrated, decomposition to palladium black and *trans*- $\text{Et}_3\text{SiCH}=\text{CHC}_6\text{H}_5$ occurred.


The η^2 -styrene complex, $[(\text{phen})\text{Pd}(\text{SiEt}_3)(\eta^2\text{-H}_2\text{C}=\text{CHC}_6\text{H}_5)]^+[\text{BAR}'_4]^-$, was not observed in the reaction of **2a** with styrene at -78°C . Thus, silyl migration is quite rapid even at low temperatures. This result is in agreement with the results from the silyl scrambling experiments described above. Following silyl migration to styrene, isomerization of the initially formed σ -benzyl intermediate to the more stable π -benzyl complex **6** occurs (eq 15). Qualitatively, the silyl migration to styrene is



faster than the analogous methyl migration to styrene in $[(\text{phen})\text{Pd}(\text{CH}_3)(\eta^2\text{-H}_2\text{C}=\text{CHC}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ ($\Delta H^\ddagger = 16.6 \pm 1.3$ kcal/mol, $\Delta S^\ddagger = -6.6 \pm 5.4$ eu).²⁹

The reaction of **6** with excess HSiEt_3 results in formation of $\text{PhCH}_2\text{CH}_2\text{SiEt}_3$ and regeneration of **2a**. The rate was measured under pseudo-first-order conditions (10–30 equiv of HSiEt_3) at 13 °C and found to be first order in HSiEt_3 ($k_{\text{obs}} = 1.1 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1} [\text{HSiEt}_3]$) (Figure 2). The rate is much slower than that observed for the reaction of **4b** with HSiEt_3 (-38°C ; $k_{\text{obs}} = 1.7 \times 10^{-3} \text{ M}^{-1} \text{ s}^{-1} [\text{HSiEt}_3]$). Since the steric demands of a *tert*-butyl group are actually greater than that of a phenyl group, the low reactivity of **6** with HSiEt_3 must be attributed to the equilibrium between the σ - and π -benzyl isomers of **6**, which strongly favors the π -benzyl isomer.

In the presence of excess styrene at 25 °C, **6** reacts with styrene to produce *trans*- $\text{Et}_3\text{SiCH}=\text{CHC}_6\text{H}_5$ and $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_3)\text{C}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ (**7**), (eq 16). The displacement of



trans- $\text{Et}_3\text{SiCH}=\text{CHC}_6\text{H}_5$ is reversible, but the equilibrium lies in favor of the less-crowded π -benzyl complex **7** (25 °C; $K_{\text{eq}} = 26$). In a related reaction, $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_2\text{CH}_3)\text{C}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ reacts with excess styrene to produce **7** (25 °C; $K_{\text{eq}} = 249$) and *trans*-methylstyrene.²⁹ Mechanistic studies for this process suggest that the benzyl transfer reaction proceeds via β -hydride elimination followed by associative displacement of *trans*- $\text{CH}_3\text{CH}=\text{CHPh}$ by styrene. β -Hydride migration and rapid isomerization produces the new π -benzyl complex **7**. Based on the kinetics experiments (see below) and the similarities between the two systems, it is likely that the interconversion of **6** and **7** proceeds through a similar series of reactions (Scheme 7).





Figure 2. Plots of k_{obs} vs $[\text{HSiEt}_3]$ for the reaction of $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_2\text{SiEt}_3)\text{C}_6\text{H}_5)]^+$ (**6**) and $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_3)\text{C}_6\text{H}_5)]^+$ (**7**) ($[\text{6 or 7}] = 0.027 \text{ M}$) with HSiEt_3 at 13 °C. (**6** + $\text{HSiEt}_3 \rightarrow 2a + \text{PhCH}_2\text{CH}_2\text{SiEt}_3$; $k_{\text{obs}} = 1.1 \times 10^{-4} \text{ s}^{-1} [\text{HSiEt}_3]$; $R^2 = 0.995$.) (**7** + $\text{HSiEt}_3 \rightarrow 2a + \text{PhCH}_2\text{CH}_3$; $k_{\text{obs}} = 7.1 \times 10^{-4} \text{ s}^{-1} [\text{HSiEt}_3]$; $R^2 = 0.998$.)

Scheme 7. Proposed Associative Benzyl Exchange Mechanism for the Reaction of $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_2\text{SiR}_3)\text{C}_6\text{H}_5)]^+$ (**6**) and Styrene ($[\text{Pd}] = (\text{phen})\text{Pd}^+$)




The kinetics of the reaction of **6** with excess styrene were measured at 13 °C. The results are shown in Figure 3. The system approached saturation at higher styrene concentrations. The saturation rate constant was obtained as the reciprocal of the intercept of the plot of $1/k_{\text{obs}}$ vs $1/[\text{styrene}]$ ($k_{\text{sat}} = 1.1 \times 10^{-4} \text{ s}^{-1}$). The rate dependence on $[\text{styrene}]$ suggests that displacement of *trans*- $\text{Et}_3\text{SiCH}=\text{CHPh}$ occurs via an associative mechanism. Similar kinetic behavior was observed in the reaction of $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_2\text{CH}_3)\text{C}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ with styrene.²⁹ In a typical catalytic reaction (Table 2), $[\text{styrene}]_{\text{initial}} > 2 \text{ M}$, so saturation behavior is assumed.

The π -benzyl complex **7** reacts with HSiEt_3 to produce ethylbenzene and **2a**. The reaction of **7** with HSiEt_3 was monitored under pseudo-first-order conditions (0.027 M **7**, 10–30 equiv of HSiEt_3) at 13 °C. The reaction of **7** with HSiEt_3 was found to be first order in both **7** and HSiEt_3 ($k_{\text{obs}} = 7.1 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1} [\text{HSiEt}_3]$) (Figure 2).

¹H NMR spectroscopy of a working catalyst solution revealed the presence of both **6** and **7** in the presence of excess styrene and HSiEt_3 . Thus, the π -benzyl complexes **6** and **7** are the catalyst resting state for the silylation reactions of styrene. **7** can be recovered in >90% yield (based on **1b**) from the reaction mixture by precipitation with hexane upon completion of the catalytic reaction.

B. Deuterium Labeling Studies. The hydrosilylation and dehydrogenative silylation reactions of styrene with DSiEt_3 were

Scheme 8. Reactions of Styrene with DSiEt₃ in the Presence of **1b**

investigated (Scheme 8). The reaction of a 3-fold molar excess of DSiEt₃ with styrene in the presence of 1% **1b** yielded C₆H₅-CHDCH₂SiEt₃. In contrast, the reaction of excess styrene with DSiEt₃ in the presence of 1% **1b** yielded C₆H₅CH₂CH₂D as the only deuterated product. No evidence for deuterium incorporation into *trans*-C₆H₅CHCHSiEt₃ was observed. The reaction of **7** with 4 equiv of DSiEt₃ in CD₂Cl₂ yielded C₆H₅CH₂CH₂D. Once again, steric factors appear to govern the regiochemistry of cleavage by silane.

C. Mechanism of Hydrosilation and Dehydrogenative Silation of Styrene. A mechanistic scheme can be proposed based on the results of the NMR experiments and the deuterium labeling studies. This mechanism is shown in Scheme 9. As was the case with the hydrosilation, silyl migration to coordinated olefin is a key feature in the silation reactions of styrene. However, rapid isomerization to the π -benzyl complex **6** occurs, so that the complexes [(phen)Pd(SiEt₃)(η^2 -CH₂CHPh)]⁺ and [(phen)Pd(η^1 -CH(CH₂SiEt₃)(C₆H₅))], which are plausible intermediates in the conversion of **2a** to **6**, are not observed.

The concentration-dependent rate constants for the conversion of **6** to **7** and the reaction of **6** and **7** with HSiEt₃ at 13 °C can be compared directly (Scheme 10). The reaction of the π -benzyl complexes **6** and **7** with excess HSiEt₃ show a first-order dependence on [HSiEt₃] and no evidence of saturation kinetics under the reaction conditions (Figure 2). In contrast, saturation






Figure 3. (A) A plot of k_{obs} vs [styrene] for the reaction of [(phen)Pd(η^3 -CH(CH₂SiEt₃)C₆H₅)]⁺ (**6**) ([**6**] = 0.027 M) and styrene yielding [(phen)Pd(η^3 -CH(CH₃)C₆H₅)]⁺ (**7**) and *trans*-PhCH=CHSiEt₃. (B) A plot of $1/k_{\text{obs}}$ vs $1/[\text{styrene}]$ for the conversion of **6** to **7**. The k_{sat} was determined from the inverse of the intercept of the inverse plot ($k_{\text{sat}} = 1.1 \times 10^{-4} \text{ s}^{-1}$). See ref 29 for the derivation of the kinetic expression for the benzyl exchange reaction.

behavior is observed in the conversion of **6** to **7**. For the conversion of **6** to **7**, $k_{\text{sat}} = 1.1 \times 10^{-4} \text{ s}^{-1}$. In a typical catalytic reaction (Table 2), the initial styrene concentration > 2 M and saturation is assumed.

Scheme 9. Proposed Mechanism for the [(phen)Pd]⁺-Catalyzed Hydrosilation and Dehydrogenative Silation of Styrene ([Pd] = (phen)Pd⁺)

Scheme 10. Comparison of the Concentration-Dependent Rate Constants for the Rate-Determining Steps of the Hydrosilation and Dehydrogenative Silation Reactions of Styrene with HSiEt_3 at 13 °C



As shown in Scheme 10, the displacement of *trans*- $\text{C}_6\text{H}_5\text{CH}=\text{CHSiEt}_3$ from **6** results in dehydrogenative silation of styrene, and the reaction of **6** with HSiEt_3 results in hydrosilation. The similarities in the rates of the two processes and the fact that the rate of displacement of *trans*- $\text{C}_6\text{H}_5\text{CH}=\text{CHSiEt}_3$ saturates at high styrene concentration suggest that the distribution of hydrosilation and dehydrogenative silation in the catalytic reactions might be significantly altered by changing the relative ratios of styrene and HSiEt_3 . This was observed in the catalytic silitations of styrene (Table 2), in agreement with the proposed mechanism. No quantitative predictions of product distributions were possible since the rates were determined at 13 °C in CD_2Cl_2 and the preparative scale silitations were conducted at higher temperature (35–85 °C).

The difference in the rate constants for the reactivity of the π -benzyl complexes **6** and **7** with HSiEt_3 may be due to steric factors. If this is the case, then it suggests that the reaction of the π -benzyl complexes **6** or **7** with a bulkier silane (such as $\text{HSi}(i\text{-Pr})_3$) would be even slower. This was confirmed qualitatively by noting that no reaction was observed between **7** and excess $\text{HSi}(i\text{-Pr})_3$ in CD_2Cl_2 at 25 °C. Since the rate of associative displacement of styrene would be unlikely to change as much, the use of bulkier silanes should favor the dehydrogenative silation reaction rather than the hydrosilation reaction. This was observed in the reactions of styrene with $\text{HSi}(i\text{-Pr})_3$ (Table 2). Similar results were recently observed in a Rh-catalyzed dehydrogenative silation of styrene.¹³

Even when steric factors are taken into account, the silation reactions of styrene proceed much more slowly than the silation reactions of alkyl-substituted olefins. It is likely that this difference results from the high stability of the π -benzyl intermediates **6** and **7**. The relatively low reactivity of **6** toward HSiR_3 allows an alternate reaction, the displacement of *trans*- $\text{R}_3\text{SiCH}=\text{CHPh}$ by styrene, to occur at a similar rate. Thus, dehydrogenative silation is competitive with hydrosilation in this system.

Conclusions

Several conclusions can be drawn from the results of the mechanistic studies and deuterium labeling studies.

(1) In the silation reactions of alkyl-substituted α -olefins and ethylene, the catalyst resting state is the η^2 -olefin complex, $[(\text{phen})\text{Pd}(\text{SiR}_3)(\eta^2\text{-H}_2\text{C}=\text{CHR}')^+]$. This is true for both hydrosilation and dehydrogenative silation.

(2) *Silyl migration to olefin is rapid and reversible at -60 °C.* The ease of β -silyl elimination reactions in this system

suggests that β -silyl elimination may be a facile reaction of other transition metal β -silyl(alkyl) complexes. The silyl migration and elimination reactions are similar to hydride and alkyl migration reactions and β -hydrogen and β -methyl abstraction reactions, respectively.²

(3) The deuterium labeling studies and formation of vinyl silanes suggest that in addition to the rapid, the degenerate β -silyl migrations, β -hydrogen elimination and reinsertion reactions are operative in this system. Both the β -silyl migrations and the β -hydride migrations are operative at low temperatures (< -40 °C).

(4) The eventual reaction of the Pd(II) alkyl intermediate $[(\text{Pd})-\text{R}']$ with HSiR_3 appears to be governed by steric factors, with the most accessible isomer reacting the fastest. If intermediate $[(\text{Pd})-\text{R}']$ and/or the silane HSiR_3 are too bulky, the cleavage of $[(\text{Pd})-\text{R}']$ by HSiR_3 is disfavored and associative displacement of vinylsilane by alkene can occur, resulting in dehydrogenative silation. In this system, *dehydrogenative silation results when the reaction with HSiR_3 is inhibited*, rather than when associative displacement is promoted (e.g., by addition of a coordinating ligands such as PPh_3).

(5) The proposed mechanism of styrene hydrosilation and dehydrogenative silation is similar to that proposed for the silation reactions of ethylene and alkyl-substituted olefins. The main difference is that in the silation reactions of styrene, the catalyst resting states are the π -benzyl complexes **6** and **7** rather than the η^2 -olefin complexes, $[(\text{phen})\text{Pd}(\text{SiR}_3)(\eta^2\text{-H}_2\text{C}=\text{CHR}')^+]$. The stability of the π -benzyl complexes results in a slow reaction with HSiR_3 . As a result of the low reactivity of **6** with HSiR_3 , the associative displacement of vinylsilane by styrene becomes competitive with cleavage by HSiR_3 , and significant amounts of dehydrogenative silation are observed, despite the relatively low steric demands of SiEt_3 and phenyl groups.

(6) It should be emphasized that the catalysts used in this study are highly electrophilic, cationic Pd(II) complexes. The Pd(II)-catalyzed hydrosilation and dehydrogenative silation reactions reported here may proceed by a different mechanism than Pd(0)-catalyzed hydrosilation of olefins, in which dehydrogenative silation is not observed and hydrosilation of styrene occurs with silation of the benzylic carbon.

The results reported here contribute to an understanding of the factors that govern hydrosilation and dehydrogenative silation reactions. Using the mechanistic information obtained in this study, we are currently investigating the development of new catalysts that are highly selective for dehydrogenative silation.

Experimental Section

General. Unless otherwise noted, all reactions were conducted under an atmosphere of dry, deoxygenated argon or nitrogen using standard Schlenk techniques or in a Vacuum Atmospheres glovebox. Pentane, hexane, ether, toluene, and tetrahydrofuran were distilled from sodium benzophenone ketyl under a nitrogen atmosphere prior to use. Dichloromethane was distilled from P_4O_{10} under a nitrogen atmosphere. CD_2Cl_2 (CIL) was dried over CaH_2 under argon and was degassed and vacuum transferred. CDCl_3 was used as received. $(\text{phen})\text{PdMe}_2$,²⁹ $\text{H}(\text{OEt})_2\text{BAR}'_4$,³⁰ $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{OEt}_2)]^+[\text{BAR}'_4]^-$ (**1a**),²⁹ $[(\text{phen})\text{Pd}(\text{CH}_3)(\text{NCCCH}_3)]^+[\text{BAR}'_4]^-$ (**1c**),²⁶ and $[(\text{phen})\text{Pd}(\eta^3\text{-CH}(\text{CH}_3)\text{C}_6\text{H}_5)]^+[\text{BAR}'_4]^-$ (**7**)²⁹ were prepared according to reported procedures. Ethylene (polymer grade) was purchased from Matheson Gases. Unless otherwise noted, silanes, chlorosilanes, and olefins were purchased from Aldrich and used without further purification. Et_3SiD was prepared from Et_3SiCl and LiAlD_4 in Et_2O .⁸ ^1H NMR of the product revealed <5% H incorporation. Allyltriethylsilane was prepared from allylmagnesium bromide and Et_3SiCl in THF.

^1H and ^{13}C chemical shifts were referenced to residual protio solvent peaks and solvent ^{13}C peaks, respectively. Coupling constants are

reported in hertz; routine coupling constants are not listed. Elemental analyses were performed by Oneida Laboratories.

Atom labeling schemes for the phenanthroline and $((\text{CF}_3)_2\text{C}_6\text{H}_3)_4\text{B}^-$ counterion resonances are as follows:



The ^1H NMR resonances were assigned into groups of a, b, c, d or a, a', b, b', c, c', d, d' according to their characteristic coupling patterns. The ^{13}C NMR resonances were assigned in pairs such as C_a or $\text{C}_{a'}$ and $\text{C}_{a'}$ or C_a , based on their chemical shifts and J_{CH} . The relation between the ^1H and ^{13}C assignments and the stereochemistry with respect to the ligands X and Y has not been ascertained.

^1H and ^{13}C data attributed to the counterion BAR'_4^- ($\text{Ar}' = 3,5\text{-(CF}_3)_2\text{C}_6\text{H}_3$) follow. These are consistent for all examined cationic complexes and are not included in each compound characterized below. ^1H NMR (CD_2Cl_2) δ 7.72 (s, 8, H_o), 7.56 (s, 4, H_p). ^1H chemical shifts are accurate to within ± 0.02 ppm. ^{13}C NMR (CD_2Cl_2) δ 162.1 (q, $J_{\text{C-B}} = 50$ Hz, C_i), 135.2 (C_o), 129.3 (q, $^2J_{\text{C-F}} = 31$ Hz, C_m), 125.0 (q, $J_{\text{C-F}} = 272$ Hz, CF_3), 117.8 (C_p). ^{13}C NMR chemical shifts and coupling constants are consistent to within ± 1 ppm and ± 2 Hz, respectively.

[(phen)Pd(CH_3)($\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$)] $^+[\text{BAR}'_4]^-$ (1b**). Solid (phen)-Pd(CH_3)₂ (116 mg, 0.37 mmol) and $[\text{H}(\text{OEt}_2)_2]^+[\text{BAR}'_4]^-$ (365 mg, 0.37 mmol) were combined. The reaction flask was cooled to -30°C and Et_2O (10 mL) and CH_2Cl_2 (5 mL) were added. The resulting slurry was allowed to warm to 25°C to dissolve solid (phen)Pd(CH_3)₂, and then the solution was cooled to -30°C . Bis(trimethylsilyl)acetylene (85 μL , 0.37 mmol) was added and colorless microcrystals formed. The mixture was allowed to warm to room temperature, and the solid dissolved. The mixture was stirred for 1 h and then the volume was reduced to 5 mL *in vacuo* and cooled slowly to -78°C . Colorless needles formed and were washed with 10 mL of cold Et_2O , collected, and dried (yield 285 mg, 59%). ^1H NMR (CD_2Cl_2 , 20°C) δ 8.95 (d, 1, phen H_a), 8.71 (d, 1, phen H_a), 8.59 (dd, 1, phen H_c), 8.54 (dd, 1, phen H_c), 8.05 (s, 2, phen H_d), 8.00 (dd, 1, phen H_b), 7.92 (dd, 1, phen H_b), 1.06 (s, 3, Pd CH_3), 0.32 (s, 18, Si(CH_3)₃). ^{13}C NMR (CD_2Cl_2 , 20°C) δ 147.2, 145.2 (C_a , $\text{C}_{a'}$), 146.2, 143.2 (C_e , $\text{C}_{e'}$), 139.7, 138.4 (C_c , $\text{C}_{c'}$), 129.9, 129.3 (C_f , $\text{C}_{f'}$), 126.8, 126.5 (C_d , $\text{C}_{d'}$), 124.6, 124.4 (C_b , $\text{C}_{b'}$), 103.6 ($\text{Me}_3\text{SiC}\equiv\text{CSiMe}_3$), 7.9 (Pd CH_3), -1.7 (SiMe_3). Anal. Calcd for $\text{C}_{53}\text{H}_{41}\text{N}_3\text{BF}_4\text{PdSi}_2$: C, 47.67; H, 3.09; N, 2.10. Found: C, 47.83; H, 3.04; N, 1.94.**

Hydrosilation with HSiEt_3 . General Procedure. In a glovebox, solid [(phen)Pd(CH_3)($\text{Me}_3\text{SiCCSiMe}_3$)] $^+[\text{BAR}'_4]^-$ (150 mg, 0.10 mmol) was loaded into a Schlenk flask. The flask was removed from the glovebox and 5 mL of CH_2Cl_2 was added. Olefin (10.0 mmol) was added to the solution and HSiEt_3 (1.80 mL, 11.2 mmol) was then added. The yellow solution was stirred overnight. CH_2Cl_2 , HSiEt_3 , and olefin were removed *in vacuo*, leaving a clear oil and a precipitate of Pd black and deactivated catalyst residue. The oil was dissolved in hexane (5 mL) and passed through alumina (4×1 cm). Hexane was removed *in vacuo*, leaving a colorless oil which was pure by ^1H NMR. Yields and NMR data for the hydrosilation products are given below.

SiEt_3Cy . Yield = 88%. ^1H NMR (CDCl_3 , 20°C) δ 1.70 (m, 6), 1.17 (m, 4, $\text{CH}_2\text{CHSiEt}_3$), 0.92 (t, 9, SiCH_2CH_3), 0.7 (t, 1, CHSiEt_3), 0.49 (q, 6, SiCH_2CH_3). ^{13}C NMR (CDCl_3 , 20°C) δ 28.4, 27.9, 27.2, 23.6 (C_6H_{11}), 7.7 (SiCH_2CH_3), 2.0 (SiCH_2CH_3).

$\text{SiEt}_3(\text{CH}_2\text{CH}_2\text{-}t\text{-Bu})$. Yield = 79%. ^1H NMR (CDCl_3 , 20°C) δ 1.1 (m, 2, $\text{CH}_2\text{-}t\text{-Bu}$), 0.90 (t, 9, SiCH_2CH_3), 0.82 (s, 9, $t\text{-Bu}$) 0.5 (m, 2, CH_2SiEt_3), 0.47 (q, 6, SiCH_2CH_3). ^{13}C NMR (CDCl_3 , 20°C) δ 37.8 ($\text{CH}_2\text{-}t\text{-Bu}$), 31.1 (CMe_3), 28.8 (CMe_3), 7.47 (SiCH_2CH_3), 5.05 ($\text{CH}_2\text{-SiEt}_3$), 3.20 (SiCH_2CH_3).

$\text{SiEt}_3([\text{CH}_2]_5\text{CH}_3)$. Yield = 88%. NMR data matched reported data.⁸

SiEt_4 . In a modification of the above procedure, **1a** (0.1 mmol) was generated from (phen)PdMe₂ and $[\text{H}(\text{OEt}_2)_2][\text{BAR}'_4]$ at -78°C in 10 mL of CH_2Cl_2 . HSiEt_3 (1.55 mL, 9.8 mmol) was added. Ethylene

was bubbled through the solution and the mixture was allowed to warm to room temperature. After 20 min, a precipitate of Pd black had formed and ethylene addition was stopped. Solvent was removed *in vacuo*, yielding a colorless oil that was pure SiEt_4 by ^1H NMR (1.20 g, 85%).

$\text{Si}(i\text{-Pr})_3(\text{Et})$. $\text{Si}(i\text{-Pr})_3(\text{Et})$ was prepared in a procedure similar to that used for SiEt_4 . The resulting colorless oil was pure by ^1H and ^{13}C NMR (yield 75%). ^1H NMR (CDCl_3 , 20°C) δ 1.0 (m, 24, $i\text{-Pr} + \text{SiCH}_2\text{CH}_3$), 0.82 (q, 2, SiCH_2CH_3). ^{13}C NMR (CDCl_3 , 20°C) δ 18.0 (CHMe_2), 10.0 (CHMe_2), 7.33, 0.0 (SiEt). Anal. Calcd for $\text{C}_{11}\text{H}_{26}\text{Si}$: C, 70.87; H, 14.06. Found: C, 70.66; H, 14.63.

SiPh_3Cy . In a drybox, solid $\text{HSi}(\text{C}_6\text{H}_5)_3$ (1.37 g, 5.27 mmol) and **1b** (76 mg, 0.05 mmol) were loaded into a Schlenk flask. The solids were then dissolved in 10 mL of CH_2Cl_2 at -78°C and cyclohexene (500 μL , 5.0 mmol) was added. The solution was warmed to room temperature and allowed to stir overnight. Solvent was removed *in vacuo*, yielding a gray solid. The solid was dissolved in 50 mL of hot hexane and filtered. The volume of the filtrate was reduced to 20 mL, at which point crystals began to form. Colorless crystals were collected and dried (1.37 g, 78%) ^1H NMR (CDCl_3 , 20°C) δ 7.3–7.5 (m, 15, SiPh_3), 1.94 (br d, 2, CH_2), 1.7 (br s, 2, CH_2), 1.6 (t, 1, CHSiPh_3), 1.4–1.1 (br m, 6, CH_2). ^{13}C NMR (CDCl_3 , 20°C) δ 136.0, 134.6, 129.2, 122.7 (Ph), 28.2, 28.1, 26.8 (CH_2), 24.1 (CHSiPh_3).

$\text{SiPh}_3(\text{CH}_2\text{CH}_2\text{-}t\text{-Bu})$. $\text{Si}(\text{C}_6\text{H}_5)_3(\text{CH}_2\text{CH}_2\text{-}t\text{-Bu})$ was prepared using the same procedure described for $\text{Si}(\text{C}_6\text{H}_5)_3(\text{C}_6\text{H}_{11})$. A colorless oil was obtained which crystallized from hexane at -30°C after 2 months (recrystallized yield 20%). ^1H NMR (CDCl_3 , 20°C) δ 7.3–7.5 (m, 15, SiPh_3), 1.55 (br s, 2, $\text{CH}_2\text{-}t\text{-Bu}$), 1.32 (br s, 2, CH_2SiPh_3), 0.85 (s, 9, $t\text{-Bu}$). ^{13}C NMR (CDCl_3 , 20°C) δ 135.7, 135.4, 129.3, 127.8 (Ph), 37.6 ($\text{CH}_2\text{-}t\text{-Bu}$), 31.3 (CMe_3), 28.8 (CMe_3), 7.5 (CH_2SiPh_3). Anal. Calcd for $\text{C}_{24}\text{H}_{28}\text{Si}$: C, 83.66; H, 8.19. Found: C, 83.07; H, 7.92.

$t\text{-BuCH=CHSi}(i\text{-Pr})_3$. **1a** was generated *in situ* at -78°C from phenPdMe₂ (31 mg, 0.1 mmol) and $[\text{H}(\text{OEt}_2)_2]^+[\text{BAR}'_4]^-$ (101 mg, 0.1 mmol) in CH_2Cl_2 . $\text{HSi}(i\text{-Pr})_3$ (2.0 mL, 10 mmol) and $t\text{-BuCH=CH}_2$ (2.6 mL, 20 mmol) were added via syringe. The mixture was allowed to warm to room temperature and stir for 24 h. CH_2Cl_2 and $t\text{-BuCH}_2\text{-CH}_3$ were removed *in vacuo*. The resulting oil was dissolved in hexane (10 mL) and filtered through a pipet of alumina (4×1 cm). Hexane was removed *in vacuo*, leaving a colorless oil that was pure $t\text{-BuCH=CHSi}(i\text{-Pr})_3$ by ^{13}C NMR spectroscopy (1.97 g, 82%). ^1H NMR (CDCl_3) δ 6.07 (d, 1, $J_{\text{H-H}} = 19$ Hz, $\text{CH-}t\text{-Bu}$), 5.37 (d, 1, $J_{\text{H-H}} = 19$ Hz, $\text{CHSi}(i\text{-Pr})_3$), 1.0 (br m, 30 total, $\text{Si}(i\text{-Pr})_3 + t\text{-Bu}$). ^{13}C NMR (CDCl_3) δ 159.9 ($J_{\text{C-H}} = 145$ Hz, $\text{CH-}t\text{-Bu}$), 115.7 ($J_{\text{C-H}} = 134$ Hz, $\text{CHSi}(i\text{-Pr})_3$), 35.4 (CMe_3), 29.1 (CMe_3), 18.6 ($\text{CH}(\text{Me})_2$), 10.9 ($\text{CH}(\text{Me})_2$). Anal. Calcd for $\text{C}_{15}\text{H}_{32}\text{Si}$: C, 74.91; H, 13.41. Found: C, 74.16; H, 13.24.

Reactions of Styrene. In a typical procedure, **1b** or **7** (0.05 mmol) was loaded into a flask that was fitted with a Teflon plug. For the reactions conducted at 35°C , CH_2Cl_2 (1 mL) was added to dissolve the catalyst. For the reactions conducted at 85°C , the catalyst was soluble in styrene/silane and no solvent was added. Styrene and HSiR_3 ($\text{R} = \text{C}_2\text{H}_5$, $\text{CH}(\text{CH}_3)_2$) were then added and the reaction was allowed to stir for 24 h. CH_2Cl_2 , ethylbenzene, and unreacted styrene and HSiR_3 were then removed *in vacuo*, leaving a pale yellow oil and a yellow solid. The oil was dissolved in hexane and passed through a pipet full of alumina (4×1 cm) to remove any remaining **7**. Hexane was then removed *in vacuo*, leaving a colorless oil that was a mixture of $\text{C}_6\text{H}_5\text{-CH}_2\text{CH}_2\text{SiR}_3$ and *trans*- $\text{C}_6\text{H}_5\text{CH=CHSiR}_3$. ^1H and ^{13}C NMR data matched reported values.¹³ Yields and product distribution are given in Table 1. The hexane-insoluble yellow solid isolated from the reaction mixture was washed with hexane and dried. ^1H NMR data for the solid matched that of **7**.

[(phen)Pd(SiEt_3)(HSiEt_3)] $^+[\text{BAR}'_4]^-$ (2a**). [(phen)Pd(CH_3)(OEt_2)] $^+[\text{BAR}'_4]^-$ (53 mg, 0.042 mmol) was loaded into a NMR tube. CD_2Cl_2 (700 μL) was added at -78°C , the mixture was shaken to dissolve the solid, and HSiEt_3 (14 μL , 0.11 mmol) was added. The sample was inserted into a precooled NMR probe. ^1H NMR (CD_2Cl_2 , -60°C) δ 9.05 (d, 2, phen), 8.54 (d, 2, phen), 7.96 (br. s, 4, phen), 1.00 (m, 30, $\text{SiCH}_2\text{CH}_3 + \text{SiCH}_2\text{CH}_3$), -9.99 (s, 1, Si-*H*-Si). ^{13}C NMR (CD_2Cl_2 , -60°C) δ 151.8, 144.4, 140.3, 130.0, 127.6, 125.8 (phen) 9.2 (SiCH_2CH_3), 8.5 (SiCH_2CH_3). ^1H and ^{13}C NMR revealed that the two silyl groups were equivalent at -80°C .**

[(phen)Pd(SiPh₃)(HSiPh₃)]⁺[BAR'₄][−] (**2b**). Solid [(phen)Pd(CH₃)-(OEt₂)⁺][BAR'₄][−] (30 mg, 0.024 mmol) and HSiPh₃ (18 mg, 0.069 mmol) were combined in an NMR tube. CD₂Cl₂ (700 μL) was added at −78 °C and the tube was shaken briefly to dissolve the solids. The sample was inserted into a precooled NMR probe. ¹H NMR (CD₂Cl₂, −80 °C) δ 10.2 (br s, 2, phen H_{a,a'}) 8.2–7 (36, phen + SiPh₃), −12.13 (s, 1, Si-H).

[(phen)Pd(SiEt₃)(PPh₃)]⁺[BAR'₄][−] (**3a**). Solid (phen)Pd(CH₃)₂ (64 mg, 0.20 mmol) and [H(OEt₂)₂]⁺[BAR'₄][−] (210 mg, 0.21 mmol) were combined. CH₂Cl₂ (10 mL) was added at −78 °C and the mixture was stirred until the solids had dissolved and methane evolution ceased. HSiEt₃ (65 μL, 0.40 mmol) was added to generate **2a**. A solution of PPh₃ (62 mg, 0.20 mmol) in CH₂Cl₂ (3 mL) was added to the solution of **2a** at −78 °C. The mixture became orange and was stirred for 20 min. The volume was reduced to 3 mL and pentane (5 mL) was added. An orange microcrystalline solid formed. The solid was collected, washed with 5 mL of pentane and dried (145 mg, 48%). ¹H NMR (CD₂Cl₂) δ 7.2–9.0 (m, 23 total, phen + phenyl), 0.82 (t, 9, SiCH₂CH₃), 0.65 (q, 6, SiCH₂CH₃). ³¹P NMR (CD₂Cl₂) δ 37.8. Anal. Calcd for C₆₈H₅₀N₂BF₂₄PPdSi: C, 53.47; H, 3.30; N, 1.83. Found: C, 53.03; H, 2.89; N, 0.75.

[(phen)Pd(SiEt₃)(Ph₂PCH₂CH=CH₂)]⁺[BAR'₄][−] (**3b**). **3b** was prepared in a manner similar to that described for **3a** (yield 74%). ¹H NMR (CD₂Cl₂) δ 8.52, 8.50 (br s, 4 tot., phen + Ph), 7.99 (s, 2, phen), 7.1–7.8 (m, 12, phen + Ph), 5.93 (m, 1, CH₂=CHCH₂PPh₂), 5.20 (dd, 1, CH_aH_b=CHCH₂PPh₂), 5.07 (dd, 1, CH_aH_b=CHCH₂PPh₂), 3.42 (m, 2, CH₂PPh₂), 0.95 (m, 15, SiEt₃). ³¹P NMR (CD₂Cl₂) δ 29.3. Anal. Calcd for C₆₅H₅₀N₂BF₂₄PPdSi: C, 52.35; H, 3.38; N, 1.87. Found: C, 52.12; H, 3.20; N, 0.98.

[(phen)Pd(SiEt₃)(η²-C₂H₄)]⁺[BAR'₄][−] (**4a**). [(phen)Pd(CH₃)(OEt₂)⁺][BAR'₄][−] (10.2 mg, 0.08 mmol) was loaded into a NMR tube. CD₂Cl₂ (700 μL) was added at −78 °C, the mixture was shaken to dissolve the solid, and HSiEt₃ (2.0 μL, 0.015 mmol) was added. Ethylene (0.50 mL, 0.022 mmol) was added via syringe. The sample was inserted into a precooled (−90 °C) NMR probe. ¹H NMR (CD₂Cl₂, −90 °C) δ 8.74, 8.37 (d, 1 each, phen H_{a,a'}), 8.65, 8.51 (d, 1 each, phen H_{c,c'}), 8.05, 7.88 (dd, 1 each, phen H_{b,b'}), 7.99 (s, 2, phen H_{d,d'}), 5.12, 4.70 (br d, 2 each, C₂H₄), 0.88 (t, 9, SiCH₂CH₃), 0.55 (q, 6, SiCH₂CH₃). In the presence of excess ethylene, exchange between free and bound ethylene occurs and only a single averaged resonance is observed at 5.0 ppm.

[(phen)Pd(SiEt₃)(η²-CH₂=CH-*t*-Bu)]⁺[BAR'₄][−] (**4b**). [(phen)Pd-(SiEt₃)(η²-CH₂=CH-*t*-Bu)]⁺[BAR'₄][−] was generated at −78 °C using a procedure similar to that described for [(phen)Pd(SiEt₃)(C₂H₄)]⁺[BAR'₄][−]. ¹H NMR (CD₂Cl₂, −80 °C) δ 9.33, 9.25 (d, 1 each, phen H_{a,a'}), 8.50 (m, phen H_{c,c'}), 9.93 (phen H_{b,b'} and H_{d,d'}), 4.17 (dd, 1, H₂C=CH-*t*-Bu), 3.63 (d, 1, H_aH_bC=CH-*t*-Bu), 3.12 (d, 1, H_aH_bC=CH-*t*-Bu), 1.02 (s, 9, *t*-Bu), 0.78 (t, 9, SiCH₂CH₃), 0.61 (q, 6, SiCH₂CH₃). ¹³C NMR (CD₂Cl₂, −60 °C) δ 152.2, 151.5 (phen C_{a,a'}), 144.1, 143.2 (phen C_{c,c'}), 139.8, 139.5 (phen C_{c,c'}), 130.0, 129.7 (phen C_{f,f'}), 127.7, 127.5 (phen C_{d,d'}), 125.9, 125.85 (phen C_{b,b'}), 94.2 (H₂C=CH-*t*-Bu), 55.0 (H₂C=CH-*t*-Bu), 35.8 (CMe₃), 30.1 (CMe₃), 15.5, 9.9 (SiEt₃).

[(phen)Pd(Si(*i*-Pr)₃)(η²-CH₂=CH-*t*-Bu)]⁺[BAR'₄][−] (**4c**). In a 5-mm NMR tube, **1a** (52 mg, 0.042 mmol) was dissolved in CD₂Cl₂ (700 μL) at −78 °C. HSi(*i*-Pr)₃ (8.5 μL, 0.042 mmol) was added and the mixture was shaken once to mix. *tert*-Butylethylene (6 μL, 0.046 mmol) was added, yielding a yellow solution. The sample was inserted into a precooled (−65 °C) NMR probe. ¹H NMR (CD₂Cl₂, −65 °C) δ 9.28, 9.00 (d, 1 each, phen H_{a,a'}) 8.55 (overlapping doublets, 2 total, phen H_{c,c'}), 9.95 (s, 2, phen H_{d,d'}), 7.86 (m, 2, phen H_{b,b'}), 3.68 (d, 1, H_aH_b=CH-*t*-Bu), 1.2 (br s, 30 total, *i*-Pr + *t*-Bu). Note: the peaks associated with H_aH_b=CH-*t*-Bu and H₂C=CH-*t*-Bu were not located and were assumed to be obscured by the free ether peak at 3.4 ppm. ¹³C NMR (CD₂Cl₂, −65 °C) δ 154.3, 152.4 (C_{a,a'}) 146.2, 144.3 (C_{c,c'}), 139.5, 139.1 (C_{c,c'}), 130.2, 129.8 (C_{f,f'}), 127.4, 127.3 (C_{d,d'}), 126.1, 125.5 (C_{b,b'}), 72.0 (H₂C=CH-*t*-Bu), 37.5 (H₂C=CH-*t*-Bu), 30.2 (CMe₃), 29.4 (CMe₃), 18.3, 18.2 (CHMe₂), 11.3 (CHMe₂). Free *t*-BuCH=CHSi(*i*-Pr)₃ was observed when a solution of **4c** was warmed to −20 °C.

Kinetic Study of the Reaction of 4b with HSiEt₃. **1a** (12 mg, 0.01 mmol) was loaded into a 5-mm NMR tube and dissolved in 700 μL of CD₂Cl₂ at −78 °C. HSiEt₃ was added via syringe to generate **2a**; *t*-BuCH=CH₂ (1.2 μL, 0.009 mmol) was added and the NMR tube was shaken briefly and inserted into a precooled (−38 ± 1 °C) NMR

Table 3

[4a], M	[HSiEt ₃], M	vol of HSiEt ₃ , μL	10 ⁴ <i>k</i> _{obs} , s ^{−1}
0.014	0.14	15.6	2.25
0.014	0.21	24	3.43
0.014	0.30	35	5.46
0.013	0.40	48	6.54

probe. The concentration of **4b** was monitored as a function of time; the Ar' H_{para} resonance at 7.25 ppm was used as an internal standard. Rate constants were obtained from a plot of ln [**4b**] vs time. Reaction conditions and observed rate constants are shown in Table 3. The rate dependence on [HSiEt₃] was determined from the slope of the line obtained from a plot of *k*_{obs} vs [HSiEt₃] (Figure 1) (*k*_{obs} = 1.7 × 10^{−3} M^{−1} s^{−1} [HSiEt₃], R² = 0.979).

Reaction of 4b with HSiPh₃. A CD₂Cl₂ solution (700 μL) of **4b** was generated from **1a** (52 mg, 0.042 mmol), HSiEt₃ (7 μL, 0.043 mmol), and *tert*-butylethylene (6 μL, 0.046 mmol) at −78 °C. A solution of HSiPh₃ (24 mg, 0.09 mmol) in 300 μL of CD₂Cl₂ was then added and the solution was allowed to warm to room temperature. *t*-BuCH₂CH₂SiEt₃ was observed by ¹H and ¹³C NMR spectroscopy; *t*-BuCH₂CH₂SiPh₃ was not detected.

Observation of [(phen)Pd(SiEt₃)(η²-H₂C=CHCH₂SiPh₃)]⁺[BAR'₄][−] (5a**).** **1a** (33 mg, 0.027 mmol) was loaded into a 5-mm NMR tube and dissolved in CD₂Cl₂ (400 μL) at −78 °C. HSiEt₃ (8 μL, 0.05 mmol) was added and the mixture was shaken briefly to generate **2a**. Allyltriphenylsilane (10 mg, 0.033 mmol) was dissolved in 300 μL of CD₂Cl₂ and added to the solution of **2a** at −78 °C. The sample was inserted into a precooled (−78 °C) NMR probe. ¹H NMR (CD₂Cl₂, −60 °C) δ 5.85 (br m, 1, H₂C=CHCH₂SiPh₃), 4.22, 3.43 (d, 1 each, H₂C=CHCH₂SiPh₃), 2.20 (dd, CH_aH_bSiPh₃), 1.62 (t, 1, CH_aH_bSiPh₃). Because of the equilibrium between **5a** and **5b**, the resonances associated with SiPh₃, phen, and SiEt₃ are quite complicated and are not listed here.

Observation of [(phen)Pd(SiPh₃)(η²-H₂C=CHCH₂SiEt₃)]⁺[BAR'₄][−] (5b**).** A solution of **5b** in CD₂Cl₂ was prepared in a manner similar to that described for **5a**. ¹H NMR (CD₂Cl₂, −60 °C) δ 4.8 (br m, 1, H₂C=CHCH₂SiEt₃), 4.05, 3.67 (d, 1 each, H₂C=CHCH₂SiEt₃), 2.32 (dd, CH_aH_bSiEt₃), 1.78 (t, 1, CH_aH_bSiEt₃). Because of the equilibrium between **5a** and **5b**, the resonances associated with SiPh₃, phen, and SiEt₃ are quite complicated and are not listed here.

[(phen)Pd(SiEt₃)(η²-H₂C=CHCH₂SiEt₃)]⁺[BAR'₄][−] (**5c**). **1a** (33 mg, 0.027 mmol) was loaded into a NMR tube. CD₂Cl₂ (700 μL) was added at −78 °C, the mixture was shaken to dissolve the solid, and HSiEt₃ (4.2 μL, 0.036 mmol) was added. The mixture was shaken briefly and allyltrisethylsilane (5.4 μL, 0.27 mmol) was added via syringe. The NMR sample was then inserted into a precooled (−78 °C) NMR probe and the spectra were acquired. ¹H NMR (CD₂Cl₂, −78 °C) δ 9.17 (d, 2, phen), 8.49 (d, 2, phen) 7.92 (s, 2, phen) 7.90 (m, 2, phen), 4.8 (br m, 1, H₂C=CH₂CH₂SiEt₃), 3.64, 3.36 (d, 1 each, H₂C=CH₂CH₂SiEt₃), 1.74 (d, 1, H₂C=CH₂CH_aH_bSiEt₃), 1.20 (t, 1, H₂C=CH₂CH_aH_bSiEt₃), 0.86 (t, 18, SiCH₂CH₃; note that the two SiEt₃ resonances are apparently coincident), 0.63, 0.55 (q, 6 each, SiCH₂CH₃); ¹³C NMR (CD₂Cl₂) δ 150.7 (phen C_{a,a'}), 136.3 (phen C_{c,c'}), 139.4 (phen C_{c,c'}), 129.6 (phen C_{f,f'}), 125.4 (phen C_{b,b'}), 90.0 (H₂C=CHCH₂-SiEt₃), 61.8 (H₂C=CHCH₂SiEt₃), 21.0 (H₂C=CHCH₂SiEt₃), 8.3, 8.2 (SiCH₂CH₃), 6.9 (SiCH₂CH₃); note that the two SiEt₃ resonances are apparently coincident).

[(phen)Pd(SiPh₃)(η²-H₂C=CHCH₂SiPh₃)]⁺[BAR'₄][−] (**5d**). Solid HSiPh₃ (15 mg, 0.05 mmol) and **1a** (33 mg, 0.027 mmol) were loaded into a NMR tube. The solids were dissolved in 400 μL of CD₂Cl₂ at −78 °C. Allyltriphenylsilane (20 mg, 0.067 mmol) was dissolved in 400 μL of CD₂Cl₂ and added via syringe to the cold solution of [(phen)Pd(SiPh₃)(HSiPh₃)]⁺[BAR'₄][−]. The NMR tube was shaken briefly to mix the solution and then inserted into a precooled (−78 °C) NMR probe. ¹H NMR (CD₂Cl₂, −60 °C) δ 4.08, 3.42 (d, 1 each, H₂C=CHCH₂SiPh₃), 4.0 (br m, 1, H₂C=CHCH₂SiPh₃), 3.08 (d, 1, CH_aH_bSiPh₃), 2.35 (t, 1, CH_aH_bSiPh₃). The peaks associated with phenyl, Ar', and phen are not listed.

[(phen)Pd(η³-CH(CH₂SiEt₃)C₆H₅)]⁺[BAR'₄][−] (**6**). [(phen)Pd(CH₃)-(OEt₂)⁺][BAR'₄][−] (60 mg, 0.049 mmol) was loaded into a NMR tube. CD₂Cl₂ (700 μL) was added at −78 °C. HSiEt₃ (13 μL, 0.081 mmol)

Table 4

[6], M	[HSiEt ₃], M	vol of HSiEt ₃ , μ L	$10^4 k_{\text{obs}}$, s ⁻¹
0.0263	0.49	60	0.53
0.0286	0.87	98	1.05
0.0273	1.18	141	1.30

was added via syringe to the cold solution. The mixture was shaken briefly to dissolve the solid and generate [(phen)Pd(SiEt₃)(HSiEt₃)]⁺[BAR'₄]⁻. Styrene (6 μ L, 0.0525 mmol) was added. ¹H NMR (CD₂Cl₂, -60 °C) δ 9.02, 8.56, 8.45, 8.41, 7.38, 7.22, 6.38 (d, 1 each, phen + C₆H₅), 7.97 (dd, 1, phen), 7.93 (d, 2, phen), 7.56 (m, 2 phen), 7.29 (t, 1, phen), 4.11 (dd, 1, CHCH₂SiEt₃), 1.32 (m, 2, CH₂SiEt₃), 0.93 (t, 9, SiCH₂CH₃), 0.60 (q, 6, SiCH₂CH₃). ¹³C NMR (CD₂Cl₂, -60 °C) δ 150.0, 149.9, 148.9, 147.7, 147.6, 146.5, 145.4, 144.0, 140.9, 140.8, 139.5, 139.2, 135.3, (phen + C₆H₅), 64.2 (CHCH₂SiEt₃), 14.6 (CHCH₂SiEt₃), 7.13 (SiCH₂CH₃), 2.64 (SiCH₂CH₃). Note: [(phen)Pd(η^3 -CH(CH₂SiEt₃)(C₆H₅)]⁺[BAR'₄]⁻ is stable at 25 °C in CD₂Cl₂ solution, but in the solid state it decomposes to form Pd black and (2-triethylsilyl)styrene. For this reason it could not be isolated in analytically pure form.

Kinetic Study of the Reaction of 6 with HSiEt₃. 6 (0.02 mmol) was generated *in situ* as described above. When the total volume of HSiEt₃ to be added was <100 μ L, 700 μ L of CD₂Cl₂ was added; when the volume of HSiEt₃ was >100 μ L, 600 μ L of CD₂Cl₂ was added. Mesitylene (0.5 μ L) was added as an internal standard and the solution was cooled to -78 °C. HSiEt₃ was added via a gas-tight syringe and the sample was inserted into a precooled (13 \pm 1 °C) NMR probe. The concentration of 6 was monitored as a function of time. Rate constants were obtained as the slope of the line -ln [6] vs time. Two runs were performed at each HSiEt₃ concentration and the rates were averaged. The results are shown in Table 4 and in Figure 2. The rate dependence on [HSiEt₃] was determined from the slope of the line obtained from a plot of k_{obs} vs [HSiEt₃] ($k_{\text{obs}} = 1.1 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$ [HSiEt₃]).

Kinetic Study of the Reaction of 7 with HSiEt₃. 7 (25 mg, 0.02 mmol) was loaded into a 5-mm NMR tube. Mesitylene (0.5 μ L) was added as an internal standard. When the total volume of HSiEt₃ to be added was <100 μ L, 700 μ L of CD₂Cl₂ was added; when the volume of HSiEt₃ was >100 μ L, 600 μ L of CD₂Cl₂ was added. The solution was cooled to -78 °C and HSiEt₃ was added via gas-tight syringe. The sample was inserted into a precooled (13 \pm 1 °C) NMR probe. The concentration of 7 was monitored as a function of time. Rate constants were obtained as the slope of the line -ln [7] vs time. Two runs were performed at each HSiEt₃ concentration and the rates were averaged. The results are shown in Table 5 and in Figure 2. The rate dependence on [HSiEt₃] was determined from the slope of the line obtained from a plot of k_{obs} vs [HSiEt₃] ($k_{\text{obs}} = 7.1 \times 10^{-4} \text{ M}^{-1} \text{ s}^{-1}$ [HSiEt₃]).

Kinetic Study of the Reaction of 6 with Styrene. 6 (0.02 mmol) was generated *in situ* from 1a (25 mg, 0.02 mmol), HSiEt₃ (3.5 μ L, 0.022 mmol), and styrene (2.5 μ L, 0.022 mmol) in CD₂Cl₂ (600 μ L). Mesitylene (0.5 μ L) was added as an internal standard and the solution was cooled to -78 °C. Styrene was added via gas-tight syringe and

Table 5

[7], M	[HSiEt ₃], M	vol of HSiEt ₃ , μ L	$10^4 k_{\text{obs}}$, s ⁻¹
0.0272	0.287	34	2.12
0.0267	0.414	50	3.07
0.026	0.560	70	4.12
0.0252	0.742	95	5.18

Table 6

[6], M	[styrene], M	vol of styrene, μ L	$10^5 k_{\text{obs}}$, s ⁻¹
0.0286	1.25	100	7.14
0.0266	1.75	150	8.33
0.025	2.18	200	8.62

the sample was inserted into a precooled (13 \pm 1 °C) NMR probe. The methine signals of 6 and 7 were monitored. Rate constants were obtained as the slope of the line of -ln ([6]/[6]+[7]) vs time. Two runs were performed at each styrene concentration and the rates were averaged. The results are shown in Table 6 and in Figure 3. The limiting rate constant (k_{sat}) was obtained as the inverse of the intercept of the plot of $1/k_{\text{obs}}$ vs $1/[\text{styrene}]$. $k_{\text{sat}} = 1.1 \times 10^{-4} \text{ s}^{-1}$.

Determination of the Equilibrium Constant for 6 + Styrene \rightleftharpoons 7 + *trans*- β -Triethylsilylstyrene (20 °C). A 5-mm NMR tube was charged with 7 (21.5 mg, 0.0171 mmol), *trans*- β -triethylsilylstyrene (45 mg, 0.204 mmol), and CD₂Cl₂ (700 μ L). The tube was subjected to three freeze-pump-thaw cycles and then flame-sealed under vacuum. The ¹H NMR spectrum was monitored until equilibrium was reached (ca. 7 days). The equilibrium constant was calculated from $K_{\text{eq}} = [\text{trans-}\beta\text{-triethylsilylstyrene}][7]/[\text{styrene}][6]$; the concentrations were determined by the relative integrals. $K_{\text{eq}} = 26$.

Reactions of RCH=CH₂ with DSiEt₃: General Procedure. 1b (15 mg, 0.01 mmol) was dissolved in 1 mL of CH₂Cl₂. Olefin (1 mmol) and DSiEt₃ (1.2 mmol) were added and the reaction was allowed to stir overnight at 25 °C (1-hexene, cyclohexene, and *tert*-butylethylene). When styrene was used as the substrate, the stoichiometry was varied (styrene/DSiEt₃ = 0.33 and 2) and the reaction was stirred at 35 °C overnight. CH₂Cl₂ was removed in vacuo and the resulting oil was analyzed by ¹³C NMR spectroscopy. The results are shown below (only the peaks showing D incorporation are listed).

Cyclohexene/DSiEt₃. ¹³C NMR (CDCl₃) δ 27.8 (t, $J_{\text{C-D}} = 19$, CHDCHSiEt₃).

***t*-BuCH=CH₂/DSiEt₃.** ¹³C NMR (CDCl₃) δ 4.78 (t, $J_{\text{C-D}} = 17$, *t*-BuCH₂CHDSiEt₃).

1-Hexene/DSiEt₃. ¹³C NMR (C₆D₆) δ 34.6, 32.6, 24.9, 15.0 (t).

Styrene (2 equiv)/DSiEt₃. ¹³C NMR (CDCl₃) δ 26.5 (t, $J_{\text{C-D}} = 19$ Hz, PhCH₂CH₂D).

Styrene (1 equiv) DSiEt₃ (3 equiv). ¹³C NMR (CDCl₃) δ 11.9 (t, $J_{\text{C-D}} = 19$, PhCHDCH₂SiEt₃).

Acknowledgment. We thank the National Institutes of Health for funding (GM 28938). A.M.L. thanks the National Science Foundation for a postdoctoral fellowship.

JA962979N