

N-Containing Compounds of Macromycetes


Ji-Kai Liu*

Kunming Institute of Botany, the Chinese Academy of Sciences, Kunming 650204, P. R. China

Received June 1, 2004

Contents

1. Introduction	2723
2. Nitrogen Heterocycles	2724
3. Nucleosides	2729
4. Nonprotein Amino Acids	2730
5. Cyclic Peptides and Peptide Antibiotics	2733
6. Sphingolipids	2734
7. Cytochalasins	2738
8. Miscellaneous	2738
9. Acknowledgment	2742
10. References	2742



1. Introduction

Macromycetes are characterized by the production of macroscopic fruiting bodies which generate and distribute their spores. This review surveys the chemical, biological, and mycological literature dealing with the isolation, structure elucidation, biological activities, and synthesis of nitrogen-containing compounds from the fruiting bodies of macromycetes, concentrating on work that has appeared in the literature up to December 2003. In addition, this paper examines the research into some nitrogen-containing compounds produced by macromycetes grown in mycelial culture.

Macromycetes, among the many diverse organisms, are a major source of biologically active natural products. They have often been found to contain biologically active compounds, and they provide a rich variety of active secondary metabolites. There are potentially many compounds still to be discovered in macromycetes since until now only a relatively small number of macromycetes have been chemically investigated, and many of the remaining species are involved in interesting biological phenomena. These

Ji-Kai Liu was born in Anhui province of P. R. China in 1962, studied pharmacy from 1978 to 1982 at Anhui College of Traditional Chinese Medicine in Hefei and obtained his M.S. in 1985 at the China Pharmaceutical University in Nanjing. In 1988 he acquired his doctorate at Lanzhou University (organic chemistry). After that he worked at the Department of Chemistry, Zhongshan University in Guangzhou. During 1993-1994 he worked as a research fellow of Alexander von Humboldt at the Institute of Pharmacognosy and Analytical Phytochemistry, University of the Saarland in Germany. Between 1996-1997 he worked as a chemical researcher at the Pharma Research Center of Bayer AG in Wuppertal/Germany. Since 1998 he has been professor of natural products chemistry at the Kunming Institute of Botany, the Chinese Academy of Sciences. His field of work concerns natural bioactive compounds from higher fungi. He has published over 80 scientific papers in international journals. He is the author of the book *Mycochemistry* and also one of the inventors for 8 patents.

as yet unstudied species hold the promise of providing new natural products, many of which will no doubt be nitrogen-containing compounds. That these fungi are often involved in interesting biological processes indicates not only that the new metabolites involved will be chemically interesting but also that the new metabolites may be biologically interesting and significant.

Nitrogen-containing compounds of macromycetes represent unique secondary metabolites with respect to their chemodiversity. The large biodiversity of macromycetes provides a huge resource for extending

* E-mail: jkliu@mail.kib.ac.cn.


the chemodiversity of nitrogen-containing compounds and for finding new lead structures for medicinal chemistry.

The isolation, characterization, and chemistry of pigments from fungi belonging to the macromycetes have been reviewed comprehensively.^{1–5} Turner and Aldridge's monograph presented a comprehensive fungus secondary metabolite list including N-containing compounds that was classified according to biosynthetic origin.⁶ While these reviews deal primarily with the chemistry of coloring substances, or more generally, of all types of metabolites manufactured by macromycetes, they also touch upon the subject of nitrogen-containing compounds from macromycetes.

2. Nitrogen Heterocycles


The genus *Cortinarius* represents a complex group of fungi found throughout the world. The abundance of species, together with the high potency of some of their toxins, has led to regular reports of serious intoxications. For example, the toxicity of *C. orellanus* and *C. speciosissimus* is well-known and well-documented.⁷ The structure of the toxin was elucidated as 3,3',4,4'-tetrahydroxy-2,2'-bipyridine-1,1'-dioxide (orellanine) **1** by Antkowiak and Gessner.^{8,9} These authors were also the first to mention that this compound undergoes chemical reduction and decomposition by heat or light to the orellanine **2** and finally to a nontoxic compound, the orelline **3** (Scheme 1).¹⁰

Scheme 1




Orellanine is a nephrotoxic bipyridine *N*-oxide found in some *Cortinarius* species. Its mechanism of action is not fully understood, but it has been shown to inhibit protein synthesis and to generate free oxygen radicals. It was reported that generation of oxygen radicals from an iron complex of orellanine may play a major role in some of the known toxic effects of orellanine, and the mechanism of toxicity was correlated with a depletion of glutathione and ascorbate, which are implicated in the defense against oxidative damage.¹¹ Early symptoms of poisoning by this compound are often lacking or vague, and so initially the possibility of poisoning may be overlooked or the symptoms may be misinterpreted. Unfortunately, this means that affected patients usually appear at the clinic with established renal damage for which supportive care including kidney transplant is the only therapeutic option.¹² Recently, the use of antioxidant therapy in two patients with acute failure caused by *Cortinarius speciosissimus* intoxication was reported from Munich, Germany.¹³ The authors presented the case reports of a 30-year-old man and his 29-year-old wife who ingested a mushroom meal containing *C. speciosissimus*.

Scheme 2



As mentioned above, the deadly poisonous toadstool *Cortinarius orellanus* causes an irreversible loss of kidney function within 1–2 weeks of ingestion. The responsible toxin, orellanine, was assigned the structure **1**, which has been confirmed by several syntheses and a crystal structure analysis. During the investigation on the biosynthesis of orellanine, it was discovered that the toxin occurs mainly as the water-soluble 4,4'-diglucopyranoside **4** (Scheme 2).¹⁴ On the basis of the results obtained from biosynthetic experiments, a hypothetical pathway depicted in Scheme 3 was proposed even though the order of some steps


Scheme 3. Hypothetical Biosynthesis of Orellanine **1** from Anthranilic Acid **5**^a



^a Intermediates in brackets have not been identified in the toadstool.

remains arbitrary. The biosynthesis starts with anthranilic acid **5**, which is converted via **6** into **7**. After oxidative ring opening and a new ring closure, **8** could be generated and then converted by oxidative decarboxylation into **9**, the precursor of **10**. Finally, orellanine (**1**) itself is generated by hydrolysis of the diglucoside **4**.¹⁴

Orelline **13** is another substance commonly present in the poisonous mushroom *C. orellanus*.⁸ It was synthesized by Dehmlow and Schulz^{15,16} (eight steps, 1.4% yield), Tiecco et al.¹⁷ (eight steps, 4.4%


Scheme 4^a

^a Reagents and conditions: (i) 2.2 PhLi/THF/0 °C/1 h; B(OMe)₃/–70 °C; CH₃CO₃H (ii) CH₂N₂; (iii) 2.2 *n*-BuLi/THF/–70 or 0 °C/1 h; I₂ (iv) NiCl₂/PPh₃/Zn; DMF/50 °C/1.5 h; (v) HBr/AcOH.

yield), Hasseberg and Gerlach¹⁸ (five steps, 4.8% yield), and Trecoart et al.¹⁹ (five steps, 16.6% yield, Scheme 4).

Three disulfide metabolites were isolated from the fruiting bodies of a basidiomycete *Cortinarius* sp., collected in the Catlins, New Zealand. The structures of these compounds were determined as the unsymmetric disulfide cortamidine oxide **19**, as 2,2'-dithio-bis(pyridine *N*-oxide) **20**, and as the symmetric disulfide **21** (Scheme 5). Both **19** and **20** showed

Scheme 5





significant antimicrobial activity and cytotoxicity.²⁰

The illudins are a group of sesquiterpene antibiotics from *Clitocybe illudens* that have been widely reported as antibacterial and antitumor agents. From the culture of the same species (*C. illudens*), illudinine **22**, a sesquiterpenoid alkaloid, was isolated.²¹

"Magic mushrooms" is the name most commonly given to hallucinogenic fungi containing the psychoactive constituents psilocin **23** and psilocybin **24**, the principal active constituents of *Psilocybe* mushrooms.^{22,23} These compounds closely resemble the neurotransmitter serotonin, and the hallucinogenic effect of magic mushrooms is probably caused by their interference with the normal actions of brain serotonin.²⁴ The use of magic mushrooms has become popular among young people because it is relatively inexpensive and because there is lower awareness of guilt than with other drugs. The concise large-scale syntheses of **23** and **24** were achieved without chromatographic purification (Scheme 7).²⁵


Muscimol **25**, a structural analogue of γ -aminobutyric acid (GABA), exhibits central nervous system

Scheme 6

Scheme 7^a

^a Reagents and conditions: (i) Ac₂O, pyridine, CH₂Cl₂, 0 °C to room temperature; (ii) (COCl)₂, ether, 0 °C, *n*-hexane, then –20 °C; (iii) (CH₃)₂NH, THF; (iv) LiAlH₄, THF, Δ ; (v) [(BnO)₂PO]₂O, *n*-BuLi, THF, –78 °C to 0 °C; (vi) H₂, Pd/C, MeOH, rt.


activity and is a constituent of the psychotropic mushroom *Amanita muscaria*.²⁶ It has been the subject of considerable pharmacological interest and synthetic effort (Scheme 8).^{27–29}

Scheme 8^a

^a Reagents and conditions: (i) *N*-dichloroacetylpropargylamine, K₂CO₃/AcOEt; (ii) 36% HBr (H₂O), 120 °C, 4 h; (iii) KOH, MeOH, reflux 48 h; (iv) 33% HBr (CH₃COOH), reflux 10 min.

Phenoxazone **26** and α -amino-phenoxazone **27** have been detected in cultured mycelium of *Calocybe gambosa*.³⁰ Culture fluids of the wood-rotting basidiomycete *Pycnoporus cinnabarinus* produce cinnabarinic acid **28** by the laccase-induced oxidation of


Scheme 9



3-hydroxyanthranilic acid.³¹ Cinnabaric acid **28** is active against several Gram-positive bacteria of the *Streptococcus* group. Cinnabarin **29** from *Pycnoporus cinnabarinus* and *Pycnoporus sanguineus* shows antibacterial, antifungal, and antiviral activities.³² Cinnabarin has a basic ring similar to that of actinomycin D, which is an antibiotic used routinely to treat certain forms of cancer. However, actinomycin D is very toxic to humans. By contrast, cinnabarin at a concentration of 0.31 mg/mL had no effect on mouse neuroblastoma cells, and at doses of 1000 mg/kg it did not cause toxic effects in mice. In addition, at concentrations of 0.31 mg/mL, it produced a 4-fold reduction in the titers of the rabies virus.


The unusual pale-blue complex amavadin, which contains a vanadium ion and two molecules of *N*-(1-carboxyethyl)-*N*-hydroxyalanine, was first isolated from aqueous extracts of *Amanita muscaria* and was originally assigned the structure **30**. Further investigation with the *N*-(1-hydroxyethyl)-*N*-hydroxyalanine ligand and various cations has revealed exceptionally high selectivity for, and strong binding of, vanadium(IV) and has led to the revised structure **31** for amavadin (Scheme 10).³³

Scheme 10



The simple indolone **32** was isolated from the edible mushroom *Pleurotus salmoneostramineus* as a glycoprotein conjugate by aqueous extraction.³⁴ Indolone **32** plays a role in the photochemical generation of oxygen from water, suggesting the possible involvement of indolone in photosynthesis. Another indole-based pigment, haematopodin **33**, has been isolated from the red-violet methanolic extracts of the wood-rotting fungus *Mycena haematopus*³⁵ and was the first pyrroloquinone alkaloid found in a fungus (Scheme 11). Indoles **34–36** were isolated from the fruiting bodies of *Tricholoma sciodes* and *T. virgatum*.³⁶ The indole derivatives found in basidiomycetes are shikimate metabolites whose biosyntheses involve tryptophan. A possible precursor of the indoles **34–36** is lascivol **37**, isolated as the bitter principle of the fruiting bodies of *T. lascivum*.³⁷ Upon treatment with a strong acid, lascivol **37** yields the 2,4-dimethyl-5-methoxyindole derivative **38**. The dimeric indoles **39–43** are obtained from the bitter-tasting


Scheme 11



fruiting bodies of *Tricholoma sciodes* and *T. virgatum* (Scheme 11).^{38,39}

In the course of screening for free-radical scavenging substances from microorganisms, two simple indoles, **44** and **45**, have been found in the fruiting bodies of *Agrocybe cylindracea* (Scheme 12).⁴⁰ **44** and


Scheme 12



45 inhibit lipid peroxidation in rat liver microsomes, with IC₅₀ values of 4.1 and 3.9 μg/mL, respectively. Extracts of another toadstool, *Lactarius necator*, exhibit considerable mutagenic activity according to the Ames test.^{41–43} Later, Suortti and co-workers isolated a highly mutagenic compound, necatorin, from this fungus and proposed a coumarocinnoline formula for it.⁴⁴ Independent work on the pigments of *L. necator* has led to the isolation of an alkaloid, necatorone, for which structure **46** was deduced from spectroscopic data (Scheme 12).⁴⁵ Necatorone has the same molecular formula as necatorin, and direct comparison of the two compounds has established their identity. Because of the scarcity of necatorone and its interesting biological activity, Steglich and co-workers⁴⁶ finished its synthesis.

Condensation of 2-(3,4-dimethoxyphenyl)ethylamine with 5-methoxy-2-nitrobenzoyl chloride yielded the amide **47**, which on Bischler-Napieralski cyclization (POCl₃, acetonitrile, reflux, 4.5 h) was converted into the dihydroisoquinoline derivative **48**. Dehydrogenation was achieved most satisfactorily with MnO₂ in benzene under azeotropic water removal; this produced a yield of 90–98% of the isoquinoline derivative **49**. Demethylation of **49** with 48% HBr (11-h reflux, argon atmosphere) and hydrogenation of the


Scheme 13



resulting phenol **50** with H₂/Pd–C in methanol/water/acetic acid (2:2:1) gave amine **51**, which could be oxidatively cyclized to necatorone by stirring with 5% aqueous sodium hydroxide (Scheme 13).⁴⁶

Necatorone **46**, 4,4'-binecatorone **52**, and the 10-deoxy-4,4'-binecatorone **53** are the principal pigments of the green-brown cap skin of the toadstool *Lactarius necator* (Scheme 14).⁴⁷ In young specimens,

Scheme 14




necatorone **46** and its dehydrodimer **52** are found in almost equal proportions whereas in aged fruiting bodies the dimer predominates. Necatorone is responsible for the characteristic purple color that develops when the cap of *L. necator* is exposed to alkali. The American species *L. atroviridis* contains 10,10'-dideoxy-4,4'-binecatorone **54** as the main alkaloid; this is responsible for the dark green appearance of this toadstool.⁴⁷


The highly fluorescent β -carboline derivatives infractin **55** and 6-hydroxyinfractin **56** and the pentacyclic infractopicrin **57** have been isolated from *Cortinarius infractus* (Scheme 14).⁴⁸ The latter alkaloid is responsible for the bitter taste of this toadstool. Recently, we also identified a highly fluorescent compound, flazin **58**, from the fruiting bodies of *Suillus granulatus*. This compound was previously isolated from Japanese soy sauce.⁴⁹ Another N-containing compound with the canthin-6-one skeleton **59** and several S-containing derivatives **60–65** are present in *Boletus curtisii* (Scheme 15).⁵⁰

Chalciporus piperatus owes its peppery taste to the 2*H*-azepines chalciporone **66** and chalciporyl propionate **67**, which co-occur with the 3*H*-azepines isochalciporone **68** and dehydroisochalciporone **69**.⁵¹ Because of the unique structure of chalciporone **66**, its

Scheme 15




Scheme 16



biosynthesis was investigated. It appeared reasonable to speculate that **66** could be formed by cyclization of a linear polyketide chain. Young fruiting bodies of *Chalciporus piperatus*, while growing in their natural forest habitat, were fed with a mixture of [U-¹³C]-labeled fats. This experiment revealed the degradation of the fatty acids to doubly labeled acetate and the incorporation of seven acetate units into chalciporone, leading to ¹³C-enrichment of carbons 3–16 (see Scheme 17). In contrast, the methyl

Scheme 17. Incorporation of Seven Acetate Units into Chalciporone




group and the neighboring ring carbon atom C-2 exhibited no detectable ¹³C-enrichment.⁵² Labeled L-alanine is also incorporated into chalciporone. It provides the methyl group, C-2, and probably the nitrogen atom with concomitant loss of the carboxyl group.


König et al. reported the isolation of ophiocordin **70** from the fungus *Cordyceps ophioglossoides*.⁵³ A later study was designed to determine if ophiocordin **70** and balanol **71**, a potent protein kinase C inhibitor from *Verticillium balanoides*, are the same compound. The results indicated that the two fungi produced the same compound, the structure being that assigned to balanol **71** (Scheme 18).⁵⁴

In a screening program, new aldose reductase inhibitors, designated as salfredins A₃ **72**, A₄ **73**, A₇ **74**, C₁ **75**, C₂ **76**, and C₃ **77**, were obtained from the fermentation broth of *Crucibulum* sp. RF-3817 (Scheme 19).⁵⁵


Scheme 18



Scheme 19




Scheme 20



A pyrazine derivative, 3,6-dibenzyl-2-hydroxy-5-methoxypyridine **78**, and emeheterone **79** were isolated from the mushroom *Albatrellus confluens* (Scheme 19).⁵⁶ The compound **78** was reported to promote melanin synthesis by B16 melanoma cells.⁵⁷ Recently, guided by acute toxicity against mice, a toxic principle, 2-butyl-1-azacyclohexane iminium salt **80**, was obtained from a *Tylopilus* species (Boletaceae) (Scheme 19).⁵⁸ 2-Butyl-1-azacyclohexane iminium salt **80** exhibited moderate acute toxicity against ddY mice.

A green fluorescent substance extracted with cold water from the luminous mushroom *Lampteromyces japonicus* was identified as riboflavin **81**.⁵⁹ Riboflavin is widely distributed in macromycetes.¹ 3-N-Methyl-riboflavin **82** and riboflavin are responsible for the color of *Panellus serotinus*.⁶⁰ Lampteroflavin **83**, which was the first riboflavinyl α -ribofuranoside acting as a light emitter in the bioluminescence of moon night mushroom *Lampteromyces japonicus*, was reported (Scheme 21).⁶¹ Lampteroflavin was


Scheme 21



synthesized through coupling between 5-trityl-2,3-(*p*-methoxy)benzylidene-1- β -D-ribofuranosyl trichloroacetimidate and the 2',4'-(*p*-methoxy)benzylidene derivative of riboflavin to produce the crucial alpha riboside.^{62,63}

4-Hydroxymethyl-quinoline **84** was isolated from the wood-rotting fungi *Trametes versicolor* and


Scheme 22



Pycnoporus sanguineus. This was the first detection in fungi of an unoxidized quinoline nucleus, and it is active against malaria.⁶⁴ Hericerin **85** from *Hericiaceae* inhibits pollen growth.⁶⁵ The fruiting bodies of *Laccaria vinaceoavellanea* contain laccarin **86** which shows phosphodiesterase inhibitory activity.⁶⁶ The first reports of compounds related to plant hormones concerned the tryptophol esters **87–89** which were formed in large amounts by *Craterellus cornucopioides*.⁶⁷ The tryptophol esters detected would be suitable for transport through lipid membranes and storage in lipophilic cell compartments.

In an effort to identify neuroprotective compounds effective against the excitatory neurotoxins in edible and medicinal mushrooms, dictyoquinazols A **90**, B **91**, and C **92** have been isolated from the methanolic extract of the mushroom *Dictyophora indusiata* (Scheme 23). Dictyoquinazols protected primary cul-


Scheme 23




tured mouse cortical neurons from glutamate- and NMDA-induced excitotoxicities in a dose-dependent manner.⁶⁸

During screening for antifungal and antibacterial fungal metabolites, the fungus *Aporpium caryae* was investigated. The isolation and structural elucidation of two indole alkaloids **93** and **94**, which possess antifungal activity against the phytopathogenic fungus *Cladosporium cucumerinum*, are reported from a culture of *Aporpium caryae* (Scheme 24).⁶⁹ A culture of *Serpula himantoides* (Coniophoraceae) also pro-

Scheme 24




Scheme 25



duced antibiotic metabolites, himanimides A–D **95**–**98** (Scheme 25).⁷⁰ Each of these four compounds is either a succinimide or a maleimide derivative, and two of them are N-hydroxylated. The himanimide C **97** exhibited fungicidal effects, especially against *Alternaria porri*, *Aspergillus ochraceus*, and *Pythium irregulare*. It also showed moderate antibacterial activity against Gram-positive bacteria and yeast. Furthermore, **97** also showed cytotoxic activity on HL 60 (IC₅₀ 10 µg/mL) and L1210 (IC₅₀ 25 µg/mL).

Agaricus xanthoderma (Agaricales) is a poisonous mushroom that has relatively large caps with prominent rings on the stalks. This mushroom can be easily distinguished because it stains bright yellow when bruised or cut. Careful treatment and chromatographic separation of extracts of *A. xanthoderma* at 0–3 °C isolates agaricone **99**.⁷¹ It was found that *Agrocybe cylindracea* produces a new type of alkaloid, agrocybenine **100** (Scheme 26).⁷² Two unusual ni-

Scheme 26




trogenous metabolites **101** and **102** with a *p*-terphenyl core were isolated from the fruiting bodies of the basidiomycetes *Sarcodon leucopus* and *S. scabrosus*, respectively.^{73,74}


3. Nucleosides

Scheme 28 depicts three *N*(9)-alkylated adenines isolated from *Lentinula edodes*.^{75,76} Eritadenine **103** in particular demonstrates a significant hypocholesteremic effect by improving the cholesterol transport

Scheme 27




Scheme 28




along the blood vessels.^{77,78} Therefore, in Asia this mushroom is looked upon as a life elixir. Many eritadenine derivatives (e.g., **104** and **105**) have been synthesized in the past. Synthetic derivative **106** demonstrates the highest hypocholesteremic effect (Scheme 28).⁷⁹ Chemical investigation of insect pest control agents produced by mushrooms has led to the isolation of cliticine **107**. This compound showed strong insecticidal activity against the pink bollworm *Pectinophora gossypiella* (Scheme 29).⁸⁰ 1-β-D-Arabinofuranosyl-cytosine (ara-C) **108**, a synthetic anticancer drug, has been reported for the first time as

Scheme 29




Scheme 30



occurring naturally in the mushroom *Xerocomus nigromaculatus* (Scheme 29).⁸¹

Unsubstituted purine does not exist in nature: the simplest form is the β -D-ribose nucleoside nebularine **109**, a nucleoside antibiotic isolated from the mushroom *Lepista nebularis* (Scheme 30). Nebularine **109** is a highly toxic nucleoside. It exerts its toxicity by being a very potent inhibitor of adenosine deaminase.^{82,83} Cordycepin **110** and 3'-amino-3'-deoxyadenosine **111**, isolated from the mushroom *Cordyceps militaris*, act as feedback inhibitors of purine nucleotide biosynthesis by inhibiting the phosphoribosylpyrophosphate amidotransferase.^{84,85} Cordycepin **110** is also a chain terminator for the 3'-end of the growing RNA chain.^{86,87} Antibiotics **112** and **113** from the same fungus inhibit aminoacyl-*t*-RNA synthetases.^{88,89} Successive purification of a crude extract of cultured *Mi Huan Jun* (*Armillariella mellea*) mycelia, followed by an assay of the effect on complete ischemia in mice, led to the isolation of AMG-1 **114**, an N⁶-substituted adenosine with cerebral protecting activity.⁹⁰ The first C-alkylated purine, 6-methylpurine **115**, and its riboside **116** and also **117** were isolated from cultures of the mushroom *Collybia maculata* (Scheme 30). Since then, it has been found that these compounds exhibit significant antiviral activity, however, they demonstrate no antibacterial properties.⁹¹ 6-Methylpurine **115** had already been synthesized by Taylor and Martin before its isolation from mushrooms.⁹² The reaction involves alkylation of tetrahydropyran-yl-protected 6-chloropurine **118** by the Wittig reagent and subsequent hydrolysis of intermediate **119** (Scheme 31).


Scheme 31



4. Nonprotein Amino Acids

It is known that the genus *Amanita* contains various nonprotein amino acids of specific structure, and this area was concisely reviewed by Hatanaka.⁹³ The fly agaric, *Amanita muscaria*, and *A. pantherina* are the species mainly involved in the pantherina-muscaria poisoning syndrome which is characterized by central nervous system dysfunction. Ibotenic acid **120** and muscimol **25** are the most likely to be the biologically active components, but other active principles are suspected. Ibotenic acid **120** is an isoxazole derivative (Scheme 32).⁹⁴ Muscazone **121** would


Scheme 32



result from photorearrangement of muscimol **25** and is produced during isolation.⁹⁵ This amino acid has always been isolated as a racemate. (-)-(*R*)-4-hydroxy-pyrrolidone-(2) **122**, whose structure is closely related to ibotenic acid and muscimol, was found in *Amanita muscaria* (Scheme 32).⁹⁶ Tricholomic acid **123**, a dihydro derivative of ibotenic acid, has been identified in *Tricholoma muscarium*.⁹⁷

A naturally occurring allenic amino acid **124**, *trans*-2-amino-5-chloro-4-hexenoic acid **125**, and *trans*-2-amino-5-chloro-6-hydroxy-4-hexenoic acid **126** have been found in *Amanita solitaria* (Scheme 33).^{98,99}



Scheme 33



2-Amino-4-chloro-4-pentenoic acid **127**, a closely related derivative, has been reported from *A. pseudoporphyria*.¹⁰⁰

(2*S*)-Amino-3-methylenehexanoic acid **128** has been found in *Amanita vaginata*.¹⁰¹ Two compounds reported in *A. pantherina* are (1'*R*,2*R*)-[(2-amino-2-carboxyethyl)thio]butanedioic acid **129** and (1'*S*,2*R*)-[(2-amino-2-carboxyethyl)thio]butanedioic acid **130** (Scheme 34).¹⁰² Stizolobic **131** and stizolobinic acids **132**, which are known to occur in *Stizolobium hasjoo*, have also been found in *A. pantherina* and in low levels in *A. gemmata*, but these compounds have not


Scheme 34

Scheme 35^a

^a Reagents and conditions: (i) BnBr/K₂CO₃/DMF/rt (89%); (ii) Ac₂O/3-(CF₃)C₆H₄C(O)NHCH₂COOH/NaOAc/100 °C (72%); (iii) MeOH/Na₂CO₃/Δ (95%); (iv) 10% Pd(C)/MeOH/H₂/rt (100%); (v) AcOH/AcOOH/Fe(OAc)₂OH/rt (45%); (vi) c.HCl/100 °C (65%).

been detected in *A. muscaria*.¹⁰³ **131** and **132** exhibit an excitatory action in isolated rat spinal cord.¹⁰⁴ A biomimetic synthesis of stizolobinic acid (**±**)**132** was achieved from an analogue of L-DOPA via an oxidative cleavage reaction (Scheme 35).¹⁰⁵


Scheme 36




Scheme 36 depicts (2*R*)-2-amino-6-hydroxy-4-hexynoic acid **133**, (2*S*)-2-amino-4-hexynoic acid **134**, and 2-amino-5-chloro-5-hexenoic acid **135**, which were all isolated from the fruiting bodies of *Amanita miculifera*.¹⁰⁶ **134** had been found first in the fruiting bodies of *Tricholomopsis rutilans*.¹⁰⁷ **133** exists in pure D-form and its possible direct precursor, **134**, occurs predominantly in the L-form.

Muscarine was long considered to be the active principle in *A. muscaria*, because of a presumed action on the central nervous system. This has since been discredited. Muscarine **136**, *epi*-muscarine **137**, and *allo*-muscarine **138** have been detected in the mycelium of *A. muscaria* (Scheme 37).¹⁰⁸ The occur-

Scheme 37



rence of *epiallo*-muscarine **139** was not demonstrated unambiguously.¹⁰⁹ Synthesis of muscarine has been accomplished several times from different precursors.^{110–113} Moreover, a chemoenzymatic syn-


Scheme 38^a

^a Reagents: (i) Br₂, BaCO₃, H₂O; (ii) CF₃CO₂H, H₂O; (iii) MeSO₂Cl, DMAP, pyridine; (iv) (CF₃CO)₂O, Et₃N, THF; then AcOH, MeOH; (v) 10% Pd-C, EtOAc; (vi) LiBH₄, THF; then NaOAc, MeCN; (vii) TsCl, THF, pyridine; (viii) Me₃N, MeOH.

thesis of eight stereoisomers of muscarine has been reported.¹¹⁴ The renewed interest in this molecule is due, in part, to the suggestion that various subtypes of muscarinic receptors may be implicated in Alzheimer's Disease. Synthesis of (+)-muscarine from L-rhamnose was finished later. This synthesis does not require the use of any protecting group (Scheme 38).¹¹⁵

Several compounds, the muscaaurins I–VII **151**, **152**, **153**, have been proved to be the pigments responsible for the characteristic red-orange color of the caps of several *Amanita* species (Scheme 39).¹¹⁶ Mus-


Scheme 39




caflavin **154**, a yellow pigment, muscapurpurin **155**, a red-violet pigment, and 1,2,3,4-tetrahydro-1-methyl-β-carboline-3-carboxylic acid **156** have also been found in *A. muscaria* (Scheme 40).^{116,117} The structure of **154** has been confirmed by synthesis.^{118,119} It has also been detected together with L-Dopa in the blackening fruiting bodies of *Hygrocybe conica*.¹²⁰

The isolation and concise syntheses of acromelic acid A **157** and acromelic acid B **158**, neuroexcitatory compounds from the mushroom *Clitocybe acromelalgae*, have been variously reported.^{121–124} The synthesis of **157** involves an oxidative cleavage–recyclization strategy to construct the C-4 pyridone from an

Scheme 40




Scheme 41




intermediate catechol. Acromelic acids C **159**, D **160**, and E **161** have also been found in *C. acromelalgae*.^{125,126} **162** from the same fungus is a possible intermediate in the biogenesis of acromelic acids. The structure of **162** is confirmed by synthesis.¹²⁷ The first synthesis of lycoperdic acid **163**, which occurs in the mushroom *Lycoperdon perlatum*, was reported.¹²⁸ Other nonprotein amino acids isolated from *C. acromelalgae* include **164–168** (Scheme 41) and **169–174** (Scheme 42).^{129–136}

The toxic mushroom *Russula subnigricans* is very similar to an edible mushroom, *R. nigricans*, and

Scheme 42




Scheme 43



therefore poses a considerable risk to humans. The amino acid (2S,3R)-(-)-3-hydroxy-baikiaian **175**, together with baikiaian **176** and pipecolic acid **177**, was isolated from *R. subnigricans*.¹³⁷ N-γ-Glutamyl boletine **178**, together with a toxin (as discussed earlier), has been found in the mushroom *Tylophila* sp.⁵⁸ The absolute stereostructure of **178** was clarified on the basis of spectroscopic analysis, acidic hydrolysis, and total synthesis. It exhibited moderate antibacterial activity. Two homoserine lipids **179** and **180** from *Xerocomus langbianensis* have been reported (Scheme 43).¹³⁸


Particularly interesting examples of flavor generation by “allium-like” pathways in very different species are illustrated by the mushrooms *Marasmius alliaceus* and *Lentinula edodes* (shiitake mushroom). The flavor precursor in the former mushroom, which, as suggested by its Latin name, develops “a strong, garlic like odor – especially when wet or when crushed with water”, is γ-glutamyl-marasmin **181**.^{139,140} It is suggested that **181** is sequentially cleaved by a γ-glutamyl transpeptidase and by a C-S lyase giving a highly reactive sulfenic acid intermediate.^{141,142} Of particular interest is the occurrence in *Lentinula edodes* of lentinic acid **182**, a “tetrathio” flavor precursor with a structure curiously reminiscent of that of the polyketides (Scheme 44).¹⁴³

Scheme 44



(2S,4S)-4-Hydroxy-2-pyrrolidinecarboxylic acid **183** and (2S,3R,4R)-3,4-dihydroxy-2-pyrrolidinecarboxylic acid **184** were isolated from *Amanita phalloides* and *A. verna*, respectively (Scheme 45).^{144,145} Scl-

Scheme 45




rothionine (STH) **185** is a sulfur-containing imidazole compound that has been isolated from sclerotia of *Sclerotinia libertiana* cultured on bran.¹⁴⁶

Morel mushroom, a common edible fungi, especially *Morchella esculenta*, contains *cis*-3-amino-L-proline **186**,¹⁴⁷ which is also present in a free state in the fruiting bodies of *M. esculenta*, *M. conica*, and *M. crassipes*, as well as in their cultured mycelia. Two polyene pigments, boletocrocin A **187** and B **188**, were isolated (Scheme 45)¹⁴⁸ from the fruiting bodies of the Japanese mushroom *Boletus laetissimus*. These compounds are diamides of hexadecaheptaenedioic acid with isoleucine and either aspartic acid or asparagine.


Extracts of the mushroom *Lactarius helvus* have been reported to contain several unusual amino acids.¹⁴⁹ For one of these, the structure **189** has been tentatively proposed, although the position of one double bond has not been determined.¹⁵⁰ The structure β -methylene-L-(+)-norvaline **190** was proposed¹⁵¹ for an unusual amino acid also isolated from the fruiting bodies of this mushroom. Confirmation of this structure has been provided by a Strecker synthesis of the desired racemate from α -ethylacrolein. Another species of the same genus, *Lactarius piperatus*, was found to contain (2*S*,3'*S*)-1-(3-amino-3-carboxypropyl)-5-oxo-2-pyrrolidinecarboxylic acid **191** (Scheme 46).¹⁵²

Scheme 46



Cortinarius violaceus is a spectacular mushroom well-known for its dark blue-violet color and its cedar woodlike smell. The experimental evidence indicated that the pigment from *C. violaceus* is an ion–ligand 1:2 complex; however, it cannot be determined if it is mono- or binuclear (**192** or **193**).¹⁵³ The ligand is (*R*)- β -dopa. Steglich and co-workers confirmed by a successful feeding experiment with *rac*-3-fluoro- β -tyrosine that the biosynthesis of (*R*)- β -dopa proceeds through β -tyrosine, which is then hydroxylated (Scheme 47).¹⁵⁴


Scheme 47. Biosynthesis of (*R*)- β -Dopa in *C. Violaceus*: (i) Tyrosine-2,3-aminomutase; (ii) Monooxygenase



5. Cyclic Peptides and Peptide Antibiotics

Most of the reported cases of fatal mushroom poisoning in the world occur after the ingestion of


Scheme 48



Amanita species, primarily of *Amanita phalloides*. They contain toxins that are lethal to humans even in extremely small doses. The toxin molecules are cyclopeptides. They come in three varieties, known as amatoxins, which contain eight amino acid molecules, phallotoxins, which contain seven amino acid molecules, and virotoxins, which are monocyclic heptapeptides and contain D-serine instead of L-cysteine. Amatoxins and phallotoxins are bicyclic, cross-linked by the 2'-bound sulfur or sulfoxide group. When injected into mice, the phallotoxins are 10 times more lethal than cyanide: their LD₅₀ is 2 mg/kg. However, when taken by mouth, they have no effect. In contrast, the much more deadly amatoxins are actively toxic when eaten (LD₅₀ = 0.1 mg/kg). The structures of all groups of toxic peptides are shown: the amatoxins **197–205** in Scheme 48, the phallotoxins **206–213** in Scheme 49, and the vivotoxins **214–219** in Scheme 50.^{155,156} Another group of cyclopeptides, antamanide **220** and cycloamanides A–D **221–224**, are also produced by *Amanita phalloides* (Scheme 51).¹⁵⁷ The chemistry, biochemistry, and molecular biological aspects of the toxic cyclic peptides in *Amanita* were concisely reviewed by Wieland.¹⁵⁸ The chemical structures of these toxins, including their stereochemistry and conformational analysis of their hydroxyamino acids, were reported mainly during the 1960s and 1970s from Wieland's laboratory.


Hypelcins A I–IX **225–233** isolated from *Hypocrea peltata* are peptide antibiotics containing unusual amino acids, α -aminoisobutyric acid (Aib) and isovaline (Iva), and having the N- and C-terminal residues protected by an acetyl group (Ac) and an amino alcohol (Lol), respectively (Scheme 52).¹⁵⁹ These hypelcins belong to the class of membrane-modifying peptides named peptaibols. Hypelcin B is a mixture of five antibiotic peptides (hypelcins B-I, B-II, B-III, B-IV, and B-V, **234–238**) produced by *Hypocrea peltata* (Scheme 52).¹⁶⁰

Scheme 49



	Compounds	R ₁	R ₂	R ₃	R ₄
206	Phallacidin	COOH	CH(CH ₃) ₂	CH ₃ C(OH)CH ₂ OH	OH
207	Phallacin	COOH	CH(CH ₃) ₂	CH ₃ CHCH ₂ OH	OH
208	Phallisin	COOH	CH(CH ₃) ₂	CH ₂ OHC(OH)CH ₂ OH	OH
209	Phallisin	CH ₃	CH ₃	CH ₂ OHC(OH)CH ₂ OH	OH
210	Phalloidin	CH ₃	CH ₃	CH ₃ C(OH)CH ₂ OH	OH
211	Phallin B	CH ₃	CH ₂ Ph	C(OH)(CH ₃) ₂	H
212	Phalloin	CH ₃	CH ₃	C(OH)(CH ₃) ₂	OH
213	Prophallin	CH ₃	CH ₃	C(OH)(CH ₃) ₂	H


Scheme 50



	Compounds	X	R ₁	R ₂
214	Viroidin	SO ₂	CH ₃	CH(CH ₃) ₂
215	Deoxiviroidin	SO	CH ₃	CH(CH ₃) ₂
216	/Ala ¹ /viroidin	SO ₂	CH ₃	CH ₃
217	/Ala ¹ /deoxiviroidin	SO	CH ₃	CH ₃
218	Viroisin	SO ₂	CH ₂ OH	CH(CH ₃) ₂
219	Deoxiviroisin	SO	CH ₂ OH	CH(CH ₃) ₂

From a submerge culture of the ascomycetous fungus *Hypocrea muroiana*, two major groups of peptides belonging to the peptaibol family, designated hypomurocin (HM) A 1–5a **239–244** and B 1–6 **245–250**, were characterized by amino acid analysis and sequence determination using fast atom bombardment and electrospray tandem MS (Scheme 53).¹⁶¹ Both groups showed antibiotic activity (*Bacil-*

Scheme 51



lus subtilis) and caused hemolysis of rat erythrocytes, with HM A being less active than HM B.

The natural product omphalotin A **251** belongs to a family of cyclic dodecapeptides from the basidiomycete *Omphalotus olearius*^{162–164} and shows a selective activity against phytopathogenic nematodes such as *Meloidogyne incognita* (Scheme 54).¹⁶⁴ Under in vitro conditions, omphalotin A outperforms known nematocides such as ivermectin in terms of potency and selectivity.¹⁶⁵ The high specificity and the structure of **251**, both of which are unusual for a nematocide, lead to the assumption that a hitherto unknown biological target is responsible for the activity of **251**. To elucidate this target, larger amounts of the cyclopeptide are required than can be produced by fermentation alone.¹⁶⁵

Jung et al. reported on the synthesis of **251** using Fmoc amino acids on a polystyrene support with a trityl linker (TCP resin). Starting from TCP resin preloaded with Fmoc-sarcosine, and using a combination of BTC (triposgene), DIC/HOAt (diisopropylcarbodiimide/hydroxyaza-benzotriazole), and HATU-couplings, the linear dodecapeptide with C-terminal Sar [omphalotin A (7–6)] was obtained. Diastereomerically pure omphalotin A was obtained in a cyclization yield of 37% and in an overall yield of 31% with respect to the first loading of the resin with Fmoc-sarcosine.¹⁶⁶

Beauvericin **252** is produced by the bright yellow polypore *Laetiporus sulfureus*, commonly known as “Chicken-of-the-Woods” (Scheme 55).¹⁶⁷ Beauvericin is a mycotoxin produced by hypocrealean ascomycetes in grain.¹⁶⁸

6. Sphingolipids

Sphingolipids, for example, glycosphingolipid (GSL), sphingomyelin, and sphingosine derivatives or analogues are important building blocks of the plasma membranes of eukaryotic cells. In recent years, however, a great deal of attention has been devoted to studies of the biological processes regulated by sphingolipids, and evidence of newly discovered roles of these compounds in cell functions is continually being demonstrated. Their functions include anchoring lipid-bound carbohydrates to cell surfaces to create an epidermal water permeability barrier and participating in both antigen–antibody reactions and biological information transmission.^{169,170}

We investigated the chemical constituents of the mushrooms *Russula cyanoxantha* and *Albatrellus ellisii*. The new ceramide **253** from *R. cyanoxantha* and the new glycosphingolipid **254** from *A. ellisii* contain unusual sphingoid bases (Scheme 56).^{171,172}

Scheme 52. Primary Structures of Hypelcin A's and B's

Hypelcin

225	A-I	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Aib-Gln-Gln-Lol
226	A-II	Ac-Aib-Pro-Aib-Ala-Ala-Aib-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
227	A-III	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
228	A-IV	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Ile-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
229	A-V	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
230	A-VI	Ac-Aib-Pro-Aib-Ala-Aib-Ala-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
231	A-VII	Ac-Aib-Pro-Aib-Ala-Aib-Ala-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
232	A-VIII	Ac-Aib-Pro-Aib-Ala-Aib-Ala-Gln-Aib-Ile-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
233	A-IX	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Ile-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Gln-Gln-Lol
234	B-I	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Aib-Glu-Gln-Lol
235	B-II	Ac-Aib-Pro-Aib-Ala-Aib-Ala-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Aib-Glu-Gln-Lol
236	B-III	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Iva-Glu-Gln-Lol
237	B-IV	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Ile-Aib-Gly-Aib-Aib-Pro-Val-Aib-Aib-Glu-Gln-Lol
238	B-V	Ac-Aib-Pro-Aib-Ala-Aib-Aib-Gln-Aib-Leu-Aib-Gly-Aib-Aib-Pro-Val-Aib-Aib-Glu-Gln-Lol

Scheme 53. Primary Structures of Hypomurocin A's and B's

HM


239	A-1	Ac-Aib-Gln-Val-Val-Aib-Pro-Leu-Leu-Aib-Pro-Lol
240	A-2	Ac-Iva-Gln-Val-Val-Aib-Pro-Leu-Leu-Aib-Pro-Lol
241	A-3	Ac-Aib-Gln-Val-Leu-Aib-Pro-Leu-Ile-Aib-Pro-Lol
242	A-4	Ac-Aib-Gln-Ile-Val-Aib-Pro-Leu-Leu-Aib-Pro-Lol
243	A-5	Ac-Aib-Gln-Ile-Ile-Aib-Pro-Leu-Leu-Aib-Pro-Lol
244	A-5a	Ac-Aib-Gln-Ile-Leu-Aib-Pro-Leu-Ile-Aib-Pro-Lol
245	B-1	Ac-Aib-Ser-Ala-Leu-Aib-Gln-Aib-Val-Aib-Gly-Aib-Aib-Pro-Leu-Aib-Aib-Gln-Vol
246	B-2	Ac-Aib-Ser-Ala-Leu-Aib-Gln-Aib-Val-Aib-Gly-Aib-Aib-Pro-Leu-Aib-Aib-Gln-Lol
247	B-3	Ac-Aib-Ala-Ala-Leu-Aib-Gln-Aib-Val-Aib-Gly-Aib-Aib-Pro-Leu-Aib-Aib-Gln-Vol
248	B-4	Ac-Aib-Ser-Ala-Leu-Aib-Gln-Iva-Val-Aib-Gly-Aib-Aib-Pro-Leu-Aib-Aib-Gln-Vol
249	B-5	Ac-Aib-Ser-Ala-Leu-Aib-Gln-Aib-Val-Aib-Gly-Iva-Aib-Pro-Leu-Aib-Aib-Gln-Vol
250	B-6	Ac-Aib-Ser-Ala-Leu-Aib-Gln-Aib-Val-Aib-Gly-Iva-Aib-Pro-Leu-Aib-Aib-Gln-Lol

Truffles, also known as “black diamonds”, are underground mushrooms which grow in symbiosis with certain trees. They are thought to be a “miracle of nature” and have long been regarded as the ultimate in gastronomic experience. There are more than a hundred different kinds of truffles, but some are not edible and only a few are of interest to world cuisine. The Chinese truffle *Tuber indicum*, a generally hypogeous fungus belonging to the family Tu-


beraceae, is found in the provinces of Yunnan and Sichuan in southwest China. This truffle closely resembles the European black winter truffle *Tuber melanosporum*. *T. indicum*'s skin is reddish before becoming brown/black, and the flesh is white veins and has a rubbery consistency.

The glycosingolipid and ceramide composition of the fresh fruiting bodies of the mushroom *Cortinarius umidicola* were investigated resulting in the isolation

Scheme 54



Scheme 55




of a new glycosphingolipid **259** containing rare C_{17} -phytosphingosine and cerebroside B and D. In addition, three ceramides **260**–**262** derived from a 4-hydroxy-6-methyl-sphingosine were isolated (Scheme 57).¹⁷⁴


In Scheme 58, the structures of cortenuamide A **263**, cortenuamide B **264**, and cortenuamide C **265** from the fruiting bodies of the basidiomycetes *Corrinarius tenuipes* together with (4*E*,8*E*)-*N*-D-2-hydroxyoctasanoyl-1-*O*-β-D-glycopyranosyl-9-methyl-4,8-sphinga-dienine **266** and (4*E*,8*E*)-*N*-D-2-hydroxypalmitoyl-1-*O*-β-D-glycopyranosyl-9-methyl-4,8-sphinga-dienine **267** are shown. The structures of these compounds were elucidated by spectroscopic and chemical methods.¹⁷⁵ Hygrophamide **268** was also obtained from the fruiting bodies of the basidiomycete *Hygrophorus eburneus*.¹⁷⁶ The white-rot fungus, *Phellinus pini*, contained two ceramides **269** and **270**.¹⁷⁷

Fractionation of the extract of an edible Chinese mushroom, *Termitomyces albuminosus*, afforded six novel neuritogenic cerebroside, termitomycesphins A–F **271**–**276**, which induced neuronal differentiation in rat PC12 cells (Scheme 59). The absolute stereostructures of the six metabolites were elucidated by spectroscopic analysis and chemical transformation.^{178,179} Moreover, termitomycesphin E **275** was also obtained from *Russula ochroleuca*.¹⁸⁰ Cerebroside A **277**, B **278**, and C **279**, along with glucoceramide **280**, were isolated from *Schizophyllum commune* (Scheme 60).¹⁸² The study shows that

Scheme 56



Scheme 57




sphingolipids have elicitor activity in plants. Treatment of rice leaves with cerebroside A induced the accumulation of antimicrobial compounds (phytoalexin), cell death, and increased resistance to subsequent infection by compatible pathogens.¹⁸¹


Novel glycoinositolphosphingolipids (GIPSL), such as the basidiolipids **281**–**284**, have been obtained from the edible mushrooms *Agaricus bisporus* and *A. campestris* (Schemes 61 and 62).¹⁸³

Five mushrooms, *Panellus serotinus*, *Lyophyllum connatum*, *Amanita pantherina*, *Sarcodon asperatus*, and *Lepista nuda*, have been investigated chemically. Two new ceramides, **285** and **286**, have been isolated from *Panellus serotinus*. Compound **286** was also found in *Lyophyllum connatum*. Two more new ceramides, **288** and **289**, have been isolated from *Amanita pantherina* along with **287**, a known syn-

Scheme 58




Scheme 59



thetic compound. Additionally, compounds **287** and **288** have been identified from *Sarcodon asperatus* and *Lepista nuda*, respectively.¹⁸⁴ The fruiting bodies of *Grifola frondosa* (Polyporaceae) contain the new ceramides **290–293** (Scheme 63).¹⁸⁵

Three glycosphingolipids with a *cis*- Δ^{17} -fatty acyl moiety, namely, catacerebrosides A–C **294–296**, were obtained from the fungus *Catathelasma ventricosum* (Scheme 64).¹⁸⁶ Lactariamides A **297** and B **298** were obtained from *Lactarius volemus*, and com-


Scheme 60



pound **299** was obtained from *Engleromyces goetzei*. (Scheme 65).^{187,188}

A recent publication described glycoinositolphosphosphingolipids (basidiolipids) from the six basidiomycetes *Amanita virosa*, *Calvatia excipuliformis*, *Cantharellus cibarius*, *Leccinum scabrum*, *Lentinula edodes*, and *Pleurotus ostreatus*.¹⁸⁹ The immune properties of the basidiolipids have been also investigated.¹⁹⁰ It was determined that sera of normal adult human subjects contained IgG2 and IgM heterophile antibodies (hetAbs) that immunoreacted with these lipids. However, this immune recognition was not shared by the glycolipids of all mushroom species. The basidiolipids of *Amanita virosa* and *Cantharellus cibarius* did not bind antibodies of normal human sera. Only certain basidiolipids of *Agaricus bisporus*, *Leccinum scabrum*, *Lentinula edodes*, and *Pleurotus ostreatus* immunoreacted with human hetAbs. The basidiolipids that were recognized by the human hetAbs had either terminal Galalpha1-6Gal or Galbeta1-6Man epitopes. Enzymatic destruction of the respective carbohydrate epitopes negated the previous immune reactivity. It is assumed that contact with nonhuman antigens causes generation of the antibasidiolipid antibodies.

Scheme 61




7. Cytochalasins


The cytochalasins are a group of toxic fungal metabolites that show marked cytostatic effects on mammalian cells in tissue culture and have a wide range of biological activities. They were first isolated and characterized by Aldridge and Turner¹⁹¹ and independently by Tamm.¹⁹² Since their initial isolation, these compounds have been identified as metabolites of many fungi. Cytochalasin E **300** is a metabolite of *Rosellinia necatrix*,^{193,194} and engleromycin **301** is produced by *Engleromyces goetzi*.^{195,196} In Scheme 66, the structures of cytochalasins **302**–**307** from the culture medium and mycelium of fungus *Hypoxylon terricola* are shown.¹⁹⁷

L-696,474 **308**, an inhibitor of the HIV-1 protease, was discovered in extracts of fungal culture of *Hypoxylon fragiforme* (Scheme 67). L-696,474 inhibited HIV-1 protease activity with an IC₅₀ of 3 μM, and the mode of inhibition was competitive with respect to substrate. Furthermore, it was not a low-binding inhibitor. The inhibition due to L-696,474 was also independent of the HIV-1 protease concentration.¹⁹⁸ Cytochalasins **309** and **310** from *Xylaria obovata* were lethal to brine shrimp (LC₅₀ 2.5 μg/mL) and cytotoxic to HL-60 cells at 1 μg/mL (Scheme 67). These cytochalasins also inhibited mammalian cell growth with high potency as demonstrated in the Vero monkey cell growth inhibition assay for cytotoxicity (IC₅₀ for **309**, 0.46 μg/mL; for **310**, 1.9 μg/mL).¹⁹⁹ Additionally, 17 new cytochalasins **311**–**327** have been isolated from an unidentified *Daldinia* sp. of fungus.^{200–202} Their structures are shown in Schemes 67 and 68.

Scheme 62



Scheme 63




8. Miscellaneous


Neoengleromycin **328**, which contains the rare N-substituted hydroxamic acid structural moiety, was isolated from the fruiting bodies of the ascomycete *Engleromyces goetzii*.²⁰³ Another compound, vibratilicin **329**, was isolated from the fruiting bodies of the basidiomycete *Cortinarius vibratilis*.²⁰⁴ Vibratilicin, which contains a glycerol and a hydroxamic acid moiety, has a very similar structure to neoengleromycin with the only difference being that the N-substituted ethyl group of neoengleromycin is replaced by a proton in vibratilicin.

Piptamine **330**, an antibiotic produced by *Piptoporus betulinus*, showed antimicrobial activity against

Scheme 64



Scheme 65




a series of Gram-positive bacteria, yeasts, and fungi, with particularly notable MIC values for *Staphylococcus aureus* SG 511 (0.78 $\mu\text{g/mL}$) and *Enterococcus faecalis* 1528, (1.56 $\mu\text{g/mL}$).²⁰⁵ Piptamine exhibited hemolytic activity at 10–50 $\mu\text{g/mL}$ using heparinized blood from beagle dogs.


The characteristic violet color reaction with aqueous ferric chloride exhibited by fruiting bodies of *Lyophyllum connatum* is due to the hydroxamic acid derivatives connatin **331** and *N*-hydroxy-*N'*,*N'*-dimethylurea **332**.²⁰⁶

A third metabolite of *L. connatum* is the unusual hydroxycarboxamide lyophyllin **333**.^{206,207} The biosynthesis of lyophyllin depicted in Scheme 70 was confirmed by experiments in which the formation of the azoxy compound lyophyllin **333** by oxidative condensation of *N*-hydroxy-*N'*,*N'*-dimethylurea **334** with *N*-methylhydroxylamine **335** was observed in fruiting bodies of the toadstool *Lyophyllum connatum*.²⁰⁶ The condensing enzyme is remarkably un-

Scheme 66




Scheme 67




specific and transforms a variety of hydroxyureas and *N*-alkyl-hydroxylamines into the corresponding lyophyllin analogues.


Scheme 68



Scheme 69




Scheme 70




A nitrogen-containing aristolane sesquiterpenoid compound, lepidamine **336**, was isolated from the fruiting bodies of the basidiomycete *Russula lepida* (Scheme 71). It is the first aristolane-type sesquiterpene alkaloid isolated from nature.²⁰⁸ **337**, an aromatic aminoaldehyde, was isolated from the fruiting bodies of *Cortinarius umidicola*.²⁰⁹ Fruiting bodies of the mushroom *Hebeloma sacchariolens* reduce anthranilic acid and some of its analogues to the corresponding aromatic aminoaldehydes with high efficiency (Scheme 71).²¹⁰

Arcaine **342** was identified from *Panus tigrinus*.²¹¹ Basidalin **343** occurs in some basidiomycetes such as *Leucoagaricus carneifolia*.²¹² Muscaridine **344** (2R, 3R-form) is another compound present in *Amanita*

Scheme 71




Scheme 72



muscaria.²¹³ The stereochemistry of this natural product could not be confirmed. It was claimed to have the erythro-configuration, but its properties do not coincide with those of the well-characterized synthetic stereoisomers later obtained. 10-Hydroxyundeca-2,4,6,8-tetraynamide **345**, produced by *Myceina viridimarginata*, is highly antibiotic and cytotoxic.²¹⁴ Xylaramide **346**, possessing potent antifungal activity toward *Nematospora coryli* and *Saccharomyces cerevisiae*, was isolated from the culture fluids of the wood-inhabiting ascomycete *Xylaria longipes*.²¹⁵ Gyromitrin **347**, acetaldehyde *N*-methyl-*N*-formylhydrazone, is a toxin present in the edible wild mushroom *Gyromitra esculenta* (Scheme 72).²¹⁶ Under pH and temperature conditions mimicking the milieu of the human stomach, gyromitrin is converted through the intermediate *N*-methyl-*N*-formylhydrazine to methylhydrazine, a known tumor inducer in mice and hamsters. In addition, methylhydrazine is formed in the mouse stomach after oral administration of gyromitrin.²¹⁷ These findings imply that

Scheme 73



consumption of *G. esculenta* could be a carcinogenic, as well as an acutely toxic, health hazard.


Lilacinone **348**, a red aminobenzoquinone pigment from *Lactarius lilacinus*, and blennione **349**, a green aminobenzoquinone derivative from *Lactarius blennius*, were extracted and purified (Scheme 73). They represent a novel type of fungal pigment and may be formed biosynthetically from anthranilic acid units.^{218,219} A steroidal derivative **350** with the urea junction at the C-3 position was isolated from the fruiting bodies of *Chlorophyllum molybdites* (Agaricaceae) and was found to exhibit cytotoxicity against Kato III cells.²²⁰

In the initial separation of *Agaricus xanthoderma* on Sephadex, the last fractions contain yellow solid sodium 4-hydroxybenzenediazosulfonate **351**.⁷¹ **351** is an artifact that is formed from the 4-diazo-2,5-cyclo-hexadien-1-one **352** present in the fruiting bodies of the aforementioned fungus. **352** can be detected in the methanolic extract of the fungus by azo coupling with resorcinol or β -naphthol. 4-(Hydroxymethyl)benzene-diazonium chloride **353** was reported from other *Agaricus* species.²²¹ Interestingly, neither phenol nor 4,4'-dihydroxyazobenzene **354** are detectable during the workup of the fungi with methanol containing SO_2 , so these compounds isolated by Gill et al. are probably formed from the precursors during the workup.²²²


The carrot truffle, *Stephanospora caroticolor*, is a rare Gasteromycete. The fruiting bodies are tubers of 1–3 cm in diameter that mature underground. They may be recognized by their bright orange appearance in places where the soil is partially removed. The interior of the mushroom (gleba) is also of an intense orange color. Extraction of the fruiting bodies of *S. caroticolor* with methanol gives a yellow solution in which two pigments, **355** and **356** (Scheme 74), can be detected by HPLC.²²³ Structure **355** was confirmed by synthesis.

The orange pigment rubroflavin **357** from the dried fruiting bodies of *Calvatia rubro-flava* (Lycoperdaceae) owes its high optical rotation ($[\alpha]_{\text{D}}^{22}$ 2180, MeOH) to a methanesulfinyl group directly attached to a 1,4-benzoquinone semicarbazone chromophore

Scheme 74



Scheme 75



(Scheme 74). Rubroflavin is present in fresh fungi in its leuco form **358**, which is easily oxidized to **357**.²²⁴

Only a few nitroaromatic metabolites have so far been reported from basidiomycetes; *para*-nitroanisole **359** and *para*-nitrobenzaldehyde **360** occur in low concentrations in cultures of *Lepista diemii*,²²⁵ while 3,5,6-trichloro-1,4-dimethoxy-2-nitrobenzene **361** and 6-nitro-*iso*-vanillic acid **362** have been isolated from the fruiting bodies of *Fomes robiniae* and *Cortinarius anomalus*, respectively.^{226,227}

Agaricales species synthesize compounds in which the glutamic acid is linked to phenylhydrazine (agaritine **363** and related compounds **364**–**366**, Scheme 75).^{228–232} *p*-Hydrazinobenzoic acid **367** as a biosyn-

thetic intermediate has been characterized in *Agaricus bisporus*.²³³ Another interesting biologically active metabolite, calvatic acid **368**, was reported from fermentation of a mushroom, *Calvatia craniformis*. Total synthesis of **368** was completed in four steps starting from *p*-hydrazinobenzoic acid hydrochloride.²³⁴ In Scheme 75, the structure of pistillarine **369** from *Clavariadelphus pistillari* is shown.²³⁵ Pistillarine is responsible for the dark green color reaction with FeCl₃ that is of significance in the chemotaxonomy of coral fungi, for example, the genus *Ramaria*.

9. Acknowledgment

I wish to acknowledge the National Natural Science Foundation of China for financial support (30470027 and 30225048).

10. References

- Gill, M.; Steglich, M. *Prog. Chem. Org. Nat. Prod.* **1987**, *51*, 1.
- Gill, M. *Nat. Prod. Rep.* **1994**, *11*, 67.
- Gill, M. *Nat. Prod. Rep.* **1996**, *13*, 513.
- Gill, M. *Nat. Prod. Rep.* **1999**, *16*, 301.
- Gill, M. *Nat. Prod. Rep.* **2003**, *20*, 535.
- Turner, W. B.; Aldridge, D. C. *Fungal Metabolites*; Academic Press: 1971; Vol. 1; 1983; Vol. 2.
- Schumacher, T.; Hoiland, K. *Arch. Toxiol.* **1983**, *53*, 87.
- Antkowiak, W. Z.; Gessner, W. P. *Tetrahedron Lett.* **1979**, *20*, 1931.
- Antkowiak, W. Z.; Gessner, W. P. *Tetrahedron Lett.* **1984**, *25*, 4045.
- Antkowiak, W. Z.; Gessner, W. P. *Experientia* **1985**, *41*, 769.
- Oubrahim, H.; Richard, J. M.; Cantin-Esnault, D. *Free Radical Res.* **1998**, *28*, 497.
- Karlson-Stiber, C.; Persson, H. *Toxicon* **2003**, *42*, 339.
- Wornle, M.; Angstwurm, M. W.; Sitter, T. *Am. J. Kidney Dis.* **2004**, *43*, e3.
- Spiteller, P.; Spiteller, M.; Steglich, W. *Angew. Chem., Int. Ed.* **2003**, *42*, 2864.
- Dehmlow, E. V.; Schulz, H. J. *Tetrahedron Lett.* **1985**, *26*, 4903.
- Dehmlow, E. V.; Schulz, H. J. *Liebigs Ann. Chem.* **1987**, 857.
- Tiecco, M.; Tingoli, M.; Testaferri, L.; Chianelli, D.; Wenkert, E. *Tetrahedron* **1986**, *42*, 1475.
- Hasseberg, H. A.; Gerlach, H. *Helv. Chim. Acta* **1988**, *71*, 957.
- Trecourt, F.; Mallet, M.; Mongin, O.; Gervais, B.; Queguiner, G. *Tetrahedron* **1993**, *49*, 8373.
- Nicholas, G. M.; Blunt, J. W.; Munro, M. H. G. *J. Nat. Prod.* **2001**, *64*, 341.
- Nair, M. S. R.; Takeshita, H.; McMorris T. C.; Anchel, M. *J. Org. Chem.* **1969**, *34*, 240.
- Hofmann, A.; Heim R.; Kobel, H. *Experientia* **1958**, *14*, 107.
- Hofmann, A.; Heim, R.; Brack, A.; Kobel, H.; Frey, A.; Ott, H.; Petzilk, T.; Troxler, F. *Helv. Chim. Acta* **1959**, *42*, 1557.
- Vollenweider, F. X.; Vollenweider-Scherpenhuyzen, M. F.; Babler, A.; Vogel, H.; Hell, D. *Neuroreport* **1998**, *9*, 3897.
- Shirota, O.; Hakamata W.; Goda, Y. *J. Nat. Prod.* **2003**, *66*, 885.
- Boden, K.; Crank, K.; Ross, W. J. *J. Chem. Soc.* **1968**, 172.
- Krogsgaard-Larsen, P.; Falch, E.; Christensen, V. *Drugs Future* **1984**, *9*, 577.
- Chiarino, D.; Napoletano, M.; Sala, A. *Tetrahedron Lett.* **1986**, *27*, 3181.
- Pevarello, P.; Varasi, M. *Synth. Commun.* **1992**, *22*, 1939.
- Schlunegger, U. P.; Kuchen, A.; Clemencon, H. *Helv. Chim. Acta* **1976**, *59*, 1383.
- Eggert, C. *Microbiol. Res.* **1997**, *152*, 315.
- Smania, A., Jr.; Marques, C. J. S.; Smania, E. F. A.; Zanetti, C. R.; Carobrez, S. G.; Tramonte, R.; Loguercio-Leite, C. *Phytother. Res.* **2003**, *17*, 1069.
- Bayer, E.; Koch, E.; Anderegg, G. *Angew. Chem., Int. Ed. Engl.* **1987**, *26*, 545.
- Takekuma, S.; Takekuma, H.; Matsubara, Y.; Inaba, K.; Yosshida, Z. *J. Am. Chem. Soc.* **1994**, *116*, 8849.
- Baumann, C.; Bröckelmann, M.; Fugmann, B.; Steffan, B.; Steglich, W.; Sheldrick, W. *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 1087.
- Garlaschelli, L.; Pang, Z.; Sterner, O.; Vidari, G. *Tetrahedron* **1994**, *50*, 3571.
- Eizenhöfer, T.; Fugmann, B.; Sheldrick, W. S.; Steffan, B.; Steglich, W. *Liebigs Ann. Chem.* **1990**, 1115.
- Pang, Z.; Sterner, O. *J. Nat. Prod.* **1994**, *57*, 852.
- Pang, Z.; Sterner, O. *Acta Chem. Scand.* **1996**, *50*, 303.
- Kim, W. G.; Lee, I. K.; Kim, J. P.; Ryoo, I. J.; Koshino, H.; Yoo, I. D. *J. Nat. Prod.* **1997**, *60*, 721.
- Sterner, O.; Bergmann, E.; Kesler, E.; Magnusson, G.; Nilsson, L.; Wickberg, B.; Zimerson, E.; Zetterberg, G. *Mutat. Res.* **1982**, *101*, 269.
- Sterner, O.; Bergmann, R.; Franzen, C.; Kesler, E.; Nilsson, L. *Mutat. Res.* **1982**, *104*, 233.
- Wright, A. v.; Suortti, T. *Mutat. Res.* **1983**, *121*, 103.
- Suortti, T.; Wright, A. v.; Koskinen, A. *Phytochemistry* **1983**, *22*, 2873.
- Fugmann, B.; Steffan, B.; Steglich, W. *Tetrahedron Lett.* **1984**, *25*, 3575.
- Hilger, C. S.; Fugmann, B.; Steglich, W. *Tetrahedron Lett.* **1985**, *26*, 5975.
- Klamann, J. D.; Fugmann, B.; Steglich, W. *Phytochemistry* **1989**, *28*, 3519.
- Steglich, W.; Kopanski, L.; Wolf, M.; Moser, M.; Tegtmeyer, G. *Tetrahedron Lett.* **1984**, *25*, 2341.
- Nakatsuka, S.; Feng, B.; Goto, T.; Kihara, K. *Tetrahedron Lett.* **1986**, *27*, 3399.
- Bröckelmann, M. G.; Dasenbrock, B.; Steffan, B.; Steglich, W.; Wang Y.; Raabe, G.; Fleischhauer, J. *Eur. J. Org. Chem.* **2004**, *23*, 4856.
- Sterner, O.; Steffan, B.; Steglich, W. *Tetrahedron* **1987**, *43*, 1075.
- Spiteller, P.; Hamprecht, D.; Steglich, W. *J. Am. Chem. Soc.* **2001**, *123*, 4837.
- König, W. A.; Sinnwell, V.; Witt, S.; Kneifel, H. *Chem. Ber.* **1980**, *113*, 2221.
- Boros, C.; Hamilton, S. M.; Katz, B.; Kulanthaivel, P. *J. Antibiot.* **1994**, *47*, 1010.
- Matsumoto, K.; Nagashima, K.; Kamigauchi, T.; Kawamura, Y.; Yasuda, Y.; Ishii, K.; Uotani, N.; Sato, T.; Nakai, H.; Terui, Y.; Kikuchi, J.; Ikenisi, Y.; Yoshida, T.; Kato, T.; Itazaki, H. *J. Antibiot.* **1995**, *48*, 439.
- Ding, Z. H.; Dong, Z. J.; Liu, J. K. *Helv. Chim. Acta* **2001**, *84*, 259.
- Kawagishi, H.; Tanaka, A.; Sugiyama, K.; Mori, H.; Sakamoto, H.; Ishiguro, Y.; Kobayashi, K.; Uramato, M. *Phytochemistry* **1996**, *42*, 547.
- Watanabe, R.; Kita, M.; Uemura, D. *Tetrahedron Lett.* **2002**, *43*, 6501.
- Isobe, M.; Uyakul, D.; Goto, T. *J. Biolumin. Chemilumin.* **1987**, *1*, 181.
- Steglich, W.; Zechlin, L. Z. *Naturforsch.* **1977**, *32c*, 520.
- Uyakul, D.; Isobe, M.; Goto, T. *Tetrahedron* **1990**, *46*, 1367.
- Isobe, M.; Takahashi, H.; Goto, T. *Tetrahedron Lett.* **1990**, *31*, 717.
- Takahashi, H.; Isobe, M.; Goto, T. *Tetrahedron* **1991**, *47*, 6215.
- Abraham, W. R.; Spassov, G. *Phytochemistry* **1991**, *30*, 371.
- Kimura, Y.; Nishibe, M.; Nakajima, H.; Hamasaki, T.; Shimada, A.; Tsuneda, A.; Shigematsu, N. *Agric. Biol. Chem.* **1991**, *55*, 2679.
- Matsuda, M.; Kobayashi, T.; Nagao, S.; Ohta, T.; Nozoe, S. *Heterocycles* **1996**, *43*, 685.
- Magnus, V.; Lacan, G.; Iskrac, S.; Lewer, P.; Aplin, R. T.; Thaller, V. *Phytochemistry* **1989**, *28*, 2949.
- Lee, I. K.; Yun, B. S.; Han, G.; Cho, D. H.; Kim, Y. H.; Yoo, I. D. *J. Nat. Prod.* **2002**, *65*, 1769.
- Levy, L. M.; Cabrera, G. M.; Wright, J. E.; Seldes A. M. *Phytochemistry* **2000**, *54*, 941.
- Aqueveque, P.; Anke, T.; Sterner, O. *Z. Naturforsch.* **2002**, *57c*, 257.
- Hilbig, S.; Andries, T.; Steglich, W. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 1063.
- Koshino, H.; Lee, I. K.; Kim, J. P.; Kim, W. G.; Uzawa, J.; Yoo, I. D. *Tetrahedron Lett.* **1996**, *37*, 4549.
- Geraci, C.; Neri, P.; Paterno, C.; Rocco, C.; Tringali, C. *J. Nat. Prod.* **2000**, *63*, 347.
- Ma, B. G.; Liu, J. K. *Z. Naturforsch.* **2005**, *60b*, 565.
- Shibukawa, N. S.; Tokuda, S.; Kaneda, T. *J. Jpn. Soc. Food Nutr.* **1967**, *19*, 373.
- Chibata, I.; Okumura, K.; Takeyama, S.; Kotera, K. *Experientia* **1969**, *25*, 1237.
- Tokuda, S.; Kaneda, T. *J. Jpn. Soc. Food Nutr.* **1966**, *17*, 297.
- Tokuda, S.; Kano, E.; Kaneda, T. *J. Jpn. Soc. Food Nutr.* **1972**, *25*, 609.
- DiMenna, W. S.; Piantadosi, C.; Lamb, G. R. *J. Med. Chem.* **1978**, *21*, 1073.
- Kubo, I.; Kim, M.; Wood, W. F.; Naoki, H. *Tetrahedron Lett.* **1986**, *27*, 4277.
- Takahashi, A.; Nunozawa, T.; Endo, T.; Nozoe, S. *Chem. Pharm. Bull.* **1992**, *40*, 1313.
- Suhadolnik, R. J. *Nucleosides as Biological Probes*; John Wiley & Sons: New York, 1979; pp 244–245.
- Evans, B.; Wolfenden, R. *J. Am. Chem. Soc.* **1970**, *92*, 4751.
- Cunningham, K. G.; Hutchinson, S. A.; Manson, W.; Spring, F. S. *J. Chem. Soc.* **1951**, 2299.
- Rottman, F.; Guarino, A. J. *Biochim. Biophys. Acta* **1964**, *89*, 465.

- (86) Rich, M. A.; Meyers, P.; Weinbaum, G.; Cory, J. G.; Suhadolnik, R. J. *Biochim. Biophys. Acta* **1965**, *95*, 194.
- (87) Suhadolnik, R. J. *Nucleosides as Biological Probes*; John Wiley & Sons: New York, 1979; pp 118–135.
- (88) Guarino, A. J.; Ibershof, M. L.; Swain, R. *Biochim. Biophys. Acta* **1963**, *72*, 62.
- (89) Guarino, A. J.; Kredich, N. M. *Fed. Proc.* **1964**, *23*, 371.
- (90) Watanabe, N.; Obuchi, T.; Tamai, M.; Araki, H.; Omura, S.; Yang, J. S.; Yu, D. Q.; Liang, X. T. *Planta Med.* **1990**, *56*, 48.
- (91) Leonhardt, K.; Anke, T.; Hillen-Maske, E.; Steglich, W. Z. *Naturforsch.* **1987**, *42c*, 420.
- (92) Taylor, E. C.; Martin, S. F. *J. Am. Chem. Soc.* **1974**, *96*, 8095.
- (93) Hatanaka, S. I. *Prog. Chem. Org. Nat. Prod.* **1992**, *59*, 1.
- (94) Eugster, C. H.; Müller, G. F. R.; Good, R. *Tetrahedron Lett.* **1965**, 1813.
- (95) Göth, H.; Gagneux, A. R.; Eugster, C. H.; Schmid, H. *Helv. Chim. Acta* **1967**, *50*, 137.
- (96) Matsumoto, T.; Trueb, W.; Gwinner, R.; Eugster, C. H. *Helv. Chim. Acta* **1969**, *52*, 716.
- (97) Takemoto, T.; Nakajima, T. *J. Pharm. Soc. Jpn.* **1964**, *84*, 1230.
- (98) Chilton, W. S.; Tsou, G.; Kirk, L.; Benedict, R. G. *Tetrahedron Lett.* **1968**, 6283.
- (99) Chilton, W. S.; Tsou, G. *Phytochemistry* **1972**, *11*, 2853.
- (100) Hatanaka, S. I.; Kaneko, S.; Niimura, Y.; Kinoshita, F.; Soma, G. I. *Tetrahedron Lett.* **1974**, 3931.
- (101) Vervier, R.; Casimir, J. *Phytochemistry* **1970**, *9*, 2059.
- (102) Fushiya, S.; Gu, Q. Q.; Ishikawa, K.; Funayama, S.; Nozoe, S. *Chem. Pharm. Bull.* **1993**, *41*, 484.
- (103) Chilton, W. S.; Hsu, C. R.; Zdybak, W. T. *Phytochemistry* **1974**, *13*, 1179.
- (104) Ishida, M.; Shinozaki, H. *Brain Res.* **1988**, *473*, 193.
- (105) Baldwin, J. E.; Spyvee, M. R.; Whitehead, R. C. *Tetrahedron Lett.* **1997**, *38*, 2771.
- (106) Hatanaka, S. I.; Niimura, Y.; Takishima, K.; Sugiyama, J. *Phytochemistry* **1998**, *49*, 573.
- (107) Hatanaka, S. I.; Niimura, Y. *Phytochemistry* **1972**, *11*, 3327.
- (108) Stadelmann, R. J.; Müller, E.; Eugster, C. H. *Sydowia* **1976**, *29*, 15.
- (109) Michelot, D.; Melendez-Howell, L. M. *Mycol. Res.* **2003**, *107*, 131.
- (110) Pirrung, M. C.; DeAmicis, C. V. *Tetrahedron Lett.* **1988**, *29*, 159.
- (111) Adams, J.; Poupart, M. A.; Grenier, L. *Tetrahedron Lett.* **1989**, *30*, 1753.
- (112) Takanaka, S.; Iwabuchi, Y.; Ogasawara, J. *J. Chem. Soc., Chem. Commun.* **1989**, 1371.
- (113) Chan, T. H.; Li, C. J. *Can. J. Chem.* **1992**, *70*, 2726.
- (114) De Amici, M.; De Micheli, C.; Molteni, G.; Pitre, D.; Carrea, G.; Riva, S.; Spezia, S.; Zetta, L. *J. Org. Chem.* **1991**, *56*, 67.
- (115) Mantell, S. J.; Fleet, G. W. J.; Brown, D. J. *Chem. Soc., Perkin Trans. 1* **1992**, 3023.
- (116) Döpp, H.; Musso, H. *Chem. Ber.* **1973**, *106*, 3473.
- (117) Ardenne, R.; Döpp, H.; Musso, H.; Steglich, W. Z. *Naturforsch.* **1974**, *29c*, 637.
- (118) Barth, H.; Kobayashi, M.; Musso, H. *Helv. Chim. Acta* **1979**, *62*, 1231.
- (119) Burger, G.; Döpp, H.; Kobayashi, M.; Musso, H. *Liebigs Ann. Chem.* **1981**, 2164.
- (120) Steglich, W.; Preuss, R. *Phytochemistry* **1975**, *14*, 1119.
- (121) Konno, K.; Shirahama, H.; Matsumoto, T. *Tetrahedron Lett.* **1983**, *24*, 939.
- (122) Konno, K.; Hashimoto, K.; Ohfune, Y.; Shirahama, H.; Matsumoto, T. *J. Am. Chem. Soc.* **1988**, *110*, 4807.
- (123) Baldwin, J. E.; Fryer, A. M.; Pritchard, G. J.; Spyvee, M. R.; Whitehead, R. C.; Wood, M. E. *Tetrahedron Lett.* **1998**, *39*, 707.
- (124) Takano, S.; Tomita, S.; Iwabuchi, Y.; Ogasawara, K. *Heterocycles* **1989**, *29*, 1473.
- (125) Fushiya, S.; Sato, S.; Kanazawa, T.; Kusano, G.; Nozoe, S. *Tetrahedron Lett.* **1990**, *31*, 3901.
- (126) Fushiya, S.; Sato, S.; Kera, Y.; Nozoe, S. *Heterocycles* **1992**, *34*, 1277.
- (127) Yamano, K.; Shirahama, H. *Chem. Lett.* **1993**, 21.
- (128) Kaname, M.; Yoshifuji, S. *Tetrahedron Lett.* **1992**, *33*, 8103.
- (129) Yamano, K.; Hashimoto, K.; Shirahama, H. *Heterocycles* **1992**, *34*, 445.
- (130) Yamano, K.; Shirahama, H. *Tetrahedron* **1993**, *49*, 2427.
- (131) Yamano, K.; Shirahama, H. Z. *Naturforsch.* **1994**, *49*, 157.
- (132) Yamano, K.; Shirahama, H. *Tetrahedron* **1992**, *48*, 1457.
- (133) Hirayama, F.; Konno, K.; Shirahama, H. *Phytochemistry* **1989**, *28*, 1133.
- (134) Kanno, K.; Hayano, K.; Shirahama, H.; Saito, H.; Matsumoto, T. *Tetrahedron* **1982**, *38*, 3281.
- (135) Yamano, K.; Shirahama, H. *Phytochemistry* **1994**, *35*, 897.
- (136) Kanno, K.; Shirahama, H.; Matsumoto, T. *Phytochemistry* **1984**, *23*, 1003.
- (137) Kusano, G.; Ogawa, H.; Takahashi, A.; Nozoe, S.; Yokoyama, K. *Chem. Pharm. Bull.* **1987**, *35*, 3482.
- (138) Kiet, T. T.; Schlegel, B.; Gräfe, U. *J. Basic Microbiol.* **2002**, *42*, 133.
- (139) Gmelin, R.; Luxa, H. H.; Roth, K.; Höfle, G. *Phytochemistry* **1976**, *15*, 1717.
- (140) Gmelin, R.; N'Galamulume-Treves, M.; Höfle, G. *Phytochemistry* **1980**, *19*, 553.
- (141) Block, E.; Deorazio, R. *Pure Appl. Chem.* **1994**, *66*, 2205.
- (142) Sneed, E. Y.; Harris, H. H.; Pickering, I. J.; Prince, R. C.; Johnson, S.; Li, X. J.; Block, E.; George, G. N. *J. Am. Chem. Soc.* **2004**, *126*, 458.
- (143) Höfle, G.; Gmelin, R.; Luxa, H. H.; N'Galamulume-Treves, M. Hatanaka, S. I. *Tetrahedron Lett.* **1976**, 3129.
- (144) Jones, G. P.; Naidu, B. P.; Paleg, L. G.; Tiekink, E. R. T.; Snow, M. R. *Phytochemistry* **1987**, *26*, 3343.
- (145) Baku, A.; Faulstich, H.; Wieland, T.; Dabrowski, J. *Proc. Natl. Acad. Sci. U.S.A.* **1980**, *77*, 2370.
- (146) Matsuo, M.; Sato, K.; Kitani, S.; Satomura, Y. *Agric. Biol. Chem.* **1967**, *31*, 356.
- (147) Hatanaka, S. I. *Phytochemistry* **1969**, *8*, 1305.
- (148) Kahner, L.; Dasenbrock, J.; Spiteller, P.; Steglich, W.; Marumoto, R.; Spiteller, M. *Phytochemistry* **1998**, *49*, 1693.
- (149) Casimir, J.; Virtanen, A. I. *Acta Chem. Scand.* **1959**, *13*, 2139.
- (150) Honkanen, E.; Moisio, T.; Virtanen, A. I.; Melera, A. *Acta Chem. Scand.* **1964**, *18*, 1319.
- (151) Levenberg, B. J. *Biol. Chem.* **1968**, *243*, 6009.
- (152) Fushiya, S.; Watari, F.; Tashiro, T.; Kusano, G.; Nozoe, S. *Chem. Pharm. Bull.* **1988**, *36*, 1366.
- (153) Nussbaum, F.; Spiteller, P.; Rütth, M.; Steglich, W.; Wanner, G.; Gambelin, B.; Stieven, L.; Wagner, F. E. *Angew. Chem., Int. Ed.* **1998**, *37*, 3292.
- (154) Spiteller, P.; Rütth, M.; Nussbaum, F.; Steglich, W. *Angew. Chem., Int. Ed.* **2000**, *39*, 2754.
- (155) Wieland, Th.; Faulstich, H. *Experientia* **1991**, *47*, 1186.
- (156) Vetter, J. *Toxicon* **1998**, *36*, 13.
- (157) Gauhe, A. *Annalen* **1977**, 859.
- (158) Wieland, Th. *Peptides of Poidonora Amanita Mushrooms*; Springer: Berlin, 1986.
- (159) Matsuura, K.; Yesilada, A.; Iida, A.; Takaishi, Y.; Kanai, M.; Fujita, T. *J. Chem. Soc., Perkin Trans. 1* **1993**, 381.
- (160) Matsuura, K.; Shima, O.; Takeda, Y.; Takaishi, Y.; Nagaoka, Y.; Fujita, T. *Chem. Pharm. Bull.* **1994**, *42*, 1063.
- (161) Becker, D.; Kiess, K.; Brückner, H. *Liebigs Ann.* **1997**, 767.
- (162) Mayer, A.; Anke, H.; Sterner, O. *Nat. Prod. Lett.* **1997**, *10*, 25.
- (163) Sterner, O.; Etzel, W.; Mayer, A.; Anke, H. *Nat. Prod. Lett.* **1997**, *10*, 33.
- (164) Mayer, A.; Anke, H.; Sterner, O.; Kilian, M.; Hain, R.; Berg, D.; Etzel, W.; Gau, H. (Bayer AG), WO 97/20857, 1996.
- (165) Mayer, A.; Kilian, M.; Hoster, B.; Sterner, O.; Anke, H. *Pestic. Sci.* **1999**, *55*, 27.
- (166) Thern, B.; Rudolph, J.; Jung, G. *Angew. Chem., Int. Ed.* **2002**, *41*, 2307.
- (167) Deol, B. S.; Ridley, D. D.; Singh, P. *Aust. J. Chem.* **1978**, *31*, 1397.
- (168) Morrison, E.; Kosiak, B.; Ritieni, A.; Aastveit, A. H.; Uhlig, S.; Bernhoft, A. J. *Agric. Food Chem.* **2002**, *50*, 3070.
- (169) Kolter, T.; Sandhoff, K. *Angew. Chem., Int. Ed.* **1999**, *38*, 1532.
- (170) Hannun, Y. A.; Bell, R. M. *Science* **1989**, *243*, 500.
- (171) Gao, J. M.; Dong, Z. J.; Liu, J. K. *Lipids* **2001**, *36*, 175.
- (172) Gao, J. M.; Hu, L.; Dong, Z. J.; Liu, J. K. *Lipids* **2001**, *36*, 521.
- (173) Gao, J. M.; Wang, C. Y.; Zhang, A. L.; Liu, J. K. *Lipids* **2001**, *36*, 1365.
- (174) Liu, J. K.; Hu, L.; Dong, Z. J. *Lipids* **2003**, *38*, 669.
- (175) Tan, J. W.; Dong, Z. J.; Liu, J. K. *Lipids* **2003**, *38*, 81.
- (176) Qu, Y.; Zhang, H. B.; Liu, J. K. Z. *Naturforsch.* **2004**, *59b*, 241.
- (177) Lourenco, A.; Lobo, A. M.; Rodriguez, B.; Jimeno, M. L. *Phytochemistry* **1996**, *43*, 617.
- (178) Qi, J.; Ojika, M.; Sakagami, Y. *Tetrahedron* **2000**, *56*, 5835.
- (179) Qi, J.; Ojika, M.; Sakagami, Y. *Bioorg. Med. Chem.* **2001**, *9*, 2171.
- (180) Gao, J. M.; Shen, J.; Yang, X.; Liu, J. K. *Acta Bot. Yunnanica* **2001**, *23*, 385.
- (181) Koga, J.; Yamauchi, T.; Shimura, M.; Ogawa, N.; Oshima, K.; Umemura, K.; Kikuchi, M.; Ogasawara, N. *J. Biol. Chem.* **1998**, *273*, 31985.
- (182) Kawai, G.; Ikeda, Y.; Tubaki, K. *Agric. Biol. Chem.* **1985**, *49*, 2137.
- (183) Jennemann, R.; Bauer, B. L.; Bertalanffy, H.; Geyer, R.; Gschwind, R. M.; Selmer, T.; Wiegandt, H. *Eur. J. Biochem.* **1999**, *259*, 331.
- (184) Yaoita, Y.; Kohata, R.; Kakuda, R.; Machida, K.; Kikuchi, M. *Chem. Pharm. Bull.* **2002**, *50*, 681.
- (185) Yaoita, Y.; Ishizuka, T.; Kakuda, R.; Machida, K.; Kikuchi, M. *Chem. Pharm. Bull.* **2000**, *48*, 1356.
- (186) Zhan, Z. J.; Yue, J. M. J. *Nat. Prod.* **2003**, *66*, 1013.
- (187) Yue, J. M.; Fan, C. Q.; Xu, J.; Sun, H. D. *J. Nat. Prod.* **2001**, *64*, 1246.
- (188) Zhan, Z. J.; Sun, H. D.; Wu, H. M.; Yue, J. M. *Acta Bot. Sin.* **2003**, *45*, 248.
- (189) Jennemann, R.; Geyer, R.; Sandhoff, R.; Gschwind, R. M.; Levery, S. B.; Gröne, H. J.; Wiegandt, H. *Eur. J. Biochem.* **2001**, *268*, 1190.
- (190) Jennemann, R.; Sandhoff, R.; Gröne, H. J.; Wiegandt, H. *Immunol. Invest.* **2001**, *30*, 115.
- (191) Aldridge, D. C.; Armstrong, J. J.; Speake, R. N.; Turner, W. B. *J. Chem. Soc. C* **1967**, 1667.

- (192) Rothweiler, W.; Tamm, C. *Experientia* **1966**, *22*, 750.
- (193) Aldridge, D. C.; Burrows, B. F.; Turner, W. B. *J. Chem. Soc., Chem. Commun.* **1972**, 148.
- (194) Aldridge, D. C.; Burrows, B. F.; Turner, W. B. *J. Chem. Soc., Chem. Commun.* **1972**, 551.
- (195) Pederson, E. J.; Boll, P.; Boll, P. M. *Tetrahedron Lett.* **1980**, *21*, 5079.
- (196) Pendse, G. *Experientia* **1974**, *30*, 107.
- (197) Edwards, R. L.; Maitland, D. J. *J. Chem. Soc., Perkin Trans. 1* **1989**, 57.
- (198) Lingham, R. B.; Hsu, A.; Silverman, K. C.; Bills, G. F.; Domrowsky, A.; Goldman, M. E.; Darke, P. L.; Huang, L.; Koch, G. *J. Antibiot.* **1992**, *45*, 686.
- (199) Dagne, E.; Gunatilaka, A. A. L.; Asmellash, S.; Abate, D.; Kinston, D. G. I.; Hofmann, G. A.; Johnson, R. K. *Tetrahedron* **1994**, *50*, 5615.
- (200) Buchanan, M.; Hashimoto, T.; Asakawa, Y. *Phytochemistry* **1995**, *40*, 135.
- (201) Buchanan, M.; Hashimoto, T.; Asakawa, Y. *Phytochemistry* **1996**, *41*, 821.
- (202) Buchanan, M.; Hashimoto, T.; Takaoka, S.; Kan, Y.; Asakawa, Y. *Phytochemistry* **1996**, *42*, 173.
- (203) Liu, J. K.; Tan, J. W.; Dong, Z. J.; Ding, Z. H.; Wang, X. H.; Liu, P. G. *Helv. Chim. Acta* **2002**, *85*, 1439.
- (204) Wang, F.; Tan, J. W.; Liu, J. K. *Helv. Chim. Acta* **2004**, *87*, 1912.
- (205) Schlegel, B.; Luhmann, U.; Härtl, A.; Gräfe, U. *J. Antibiot.* **2000**, *53*, 973.
- (206) Fugmann, B.; Steglich, W. *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 72.
- (207) Ye, Y.; Aulinger, K.; Arnold, N.; Spahl, W.; Steglich, W. *Tetrahedron Lett.* **1997**, *38*, 8013.
- (208) Tan, J. W.; Dong, Z. J.; Liu, J. K. *Helv. Chim. Acta* **2003**, *86*, 307.
- (209) Hu, L.; Liu, J. K. *Z. Naturforsch.* **2003**, *58c*, 659.
- (210) Nussbaum, F.; Spahl, W.; Steglich, W. *Phytochemistry* **1997**, *46*, 261.
- (211) Thailambal, V. G. *Acta Crystallogr. C* **1985**, *41*, 105.
- (212) Huff, T.; Kuball, H. G.; Anke, T. *Z. Naturforsch.* **1994**, *49c*, 407.
- (213) Saleminck, C. A. *Recl. Trav. Chim. Pays-Bas* **1963**, *82*, 21.
- (214) Berle, J.; Anke, T.; Jente, R.; Bosold, F. *Arch. Microbiol.* **1982**, *132*, 194.
- (215) Schneider, G.; Anke, H.; Sterner, O. *Z. Naturforsch.* **1996**, *51c*, 802.
- (216) Pyysalo, H.; Niskanen, A. *J. Agric. Food Chem.* **1977**, *25*, 644.
- (217) Nagel, D.; Wallcave, L.; Toth, B.; Kupper, R. *Cancer Res.* **1977**, *37*, 3458.
- (218) Spiteller, P.; Arnold, N.; Spiteller, M.; Steglich, W. *J. Nat. Prod.* **2003**, *66*, 1402.
- (219) Spiteller, P.; Steglich, W. *J. Nat. Prod.* **2002**, *65*, 725.
- (220) Yoshikawa, K.; Ikuta, M.; Arihara, S.; Matsumura, E.; Katayama, S. *Chem. Pharm. Bull.* **2001**, *49*, 1030.
- (221) Levenberg, B. *Biochim. Biophys. Acta* **1962**, *63*, 212.
- (222) Gill, M.; Strauch, R. *J. Z. Naturforsch.* **1984**, *39c*, 1027.
- (223) Lang, M.; Spiteller, P.; Hellwig, V.; Steglich, W. *Angew. Chem., Int. Ed.* **2001**, *40*, 1704.
- (224) Fugmann, B.; Arnold, S.; Steglich, W.; Fleischhauer, J.; Repges, C.; Koslowski, A.; Raabe, G. *Eur. J. Org. Chem.* **2001**, 3097.
- (225) Thaller, V.; Turner, J. L. *J. Chem. Soc., Perkin Trans. 1* **1972**, 2032.
- (226) Butruille, D.; Dominguez, X. A. *Tetrahedron Lett.* **1972**, 211.
- (227) Gill, M.; Gimenez, A.; Strauch, R. *J. Phytochemistry* **1987**, *26*, 2815.
- (228) Levenberg, B. *J. Am. Chem. Soc.* **1961**, *83*, 503.
- (229) Kelly, R. B.; Daniels, E. G.; Hinman, J. W. *J. Org. Chem.* **1962**, *27*, 3229.
- (230) Levenberg, B. *J. Biol. Chem.* **1964**, *239*, 2267.
- (231) Hilbig, S.; Andries, T.; Steglich, W.; Anke, T. *Angew. Chem., Int. Ed. Engl.* **1985**, *24*, 1063.
- (232) Chulia, A. J.; Bernillon, J.; Favre-Bonvin, J.; Kaouadji, M.; Arpin, N. *Phytochemistry* **1988**, *27*, 929.
- (233) Chauhan, Y.; Nagel, D.; Issenberg, P.; Toth, B. *J. Agric. Food Chem.* **1984**, *32*, 1067.
- (234) Umezawa, H.; Takeuchi, T.; Iinuma, H.; Ito, M.; Ishizuka, M.; Kurakata, Y.; Nakamura, T.; Obayashi, A.; Tanabe, O. *J. Antibiot.* **1975**, *28*, 87.
- (235) Steglich, W.; Steffan, B.; Strähle, K.; Wolf, M. *Z. Naturforsch.* **1984**, *39c*, 10.

CR0400818