

# IL-1RN VNTR polymorphism and genetic susceptibility to cervical cancer in Portugal

Hugo Sousa · Alexandra M. Santos · Raquel Catarino ·  
Daniela Pinto · José Moutinho · Paulo Canedo ·  
José Carlos Machado · Rui Medeiros

Received: 24 November 2011 / Accepted: 1 October 2012 / Published online: 10 October 2012  
© Springer Science+Business Media Dordrecht 2012

**Abstract** Human Papillomavirus infection is considered as the main etiological factor of cervical cancer (ICC), although, the role of host genetic factors in ICC susceptibility has been increasing. Immunological response is crucial for the prevention of viral associated diseases. Interleukin 1 receptor antagonist (IL-1RN) is considered to be an important regulator of host immunity and several studies have shown a potential role of a 86 bp VNTR

polymorphism within intron 2 of the IL-1RN gene in host immune response variability. We investigated the role of this polymorphism in cervical cancer development in Portugal with a case–control study developed with peripheral blood samples from 196 healthy women and 340 women with cervical lesions from the Northern Region of Portugal. We observed that IL-1RN Allele 2 homozygosity was significantly higher in cases than in controls. In fact, IL-1RN A2\*A2 homozygous revealed to be associated with an increased risk of HSIL + ICC (OR = 1.90; 95 % IC 1.13–3.21;  $p = 0.015$ ). Furthermore, we also observed that median age of onset of HSIL + ICC was significantly different (46.0 vs 52.0) in IL-1RN A2\*A2 homozygous comparing to non-A2\*A2 ( $p = 0.028$ ). Our results indicated that IL-1RN A2 allele is associated with an increased susceptibility to cervical cancer development, probably by increasing predisposition to shorter immune responses.

H. Sousa (✉) · A. M. Santos · R. Catarino · D. Pinto ·  
R. Medeiros  
Grupo de Oncologia Molecular—CI, Laboratórios 4º Piso,  
Instituto Português de Oncologia do Porto FG,  
EPE, Rua Dr. António Bernardino Almeida,  
4200-072 Porto, Portugal  
e-mail: hugomls@gmail.com

H. Sousa · R. Medeiros  
Serviço de Virologia, Laboratórios 4º Piso, Instituto Português  
de Oncologia do Porto FG, EPE, Rua Dr. António Bernardino  
Almeida, 4200-072 Porto, Portugal

J. Moutinho  
Serviço de Ginecologia, Instituto Português de Oncologia  
do Porto FG, EPE, Rua Dr. António Bernardino Almeida,  
4200-072 Porto, Portugal

P. Canedo · J. C. Machado  
IPATIMUP - Institute of Molecular Pathology and Immunology  
of the University of Porto, Rua Dr. Roberto Frias s/n, 4200-465  
Porto, Portugal

J. C. Machado  
Faculty of Medicine of University of Porto, Al. Prof. Hernâni  
Monteiro, 4200-319 Porto, Portugal

R. Medeiros  
ICBAS—Abel Salazar Institute for the Biomedical Sciences of  
University of Porto, Rua de Jorge Viterbo Ferreira n.º 228,  
4050-313 Porto, Portugal

**Keywords** Cervical cancer · Human Papillomavirus (HPV) · Genetic polymorphism · Interleukin 1 · IL-1RN

## Introduction

Cervical cancer is the second most common cancer in women with almost 500,000 new cases each year and over 260,000 deaths [1, 2]. Moreover, it is estimated that over 1 million women worldwide have cervical cancer and the majority of them have not yet been diagnosed [2].

In the early 90's persistent infection by the oncogenic types of the Human Papillomavirus (HPV) was established as the etiological factor for the development of cervical cancer [3–8]. Over time, several studies have been focusing on the role of risk factors that influence either the acquisition of persistent HPV infection or by contributing for the

progression from pre-invasive lesions to cervical cancer. Literature has demonstrated that despite its oncogenic potential, HPV is not sufficient for cervical carcinogenesis [9–11]. Epidemiological studies have revealed a number of co-factors for cervical cancer development (number of sexual partners, age at first sexual intercourse, parity, tobacco and alcohol consumption, co-infection with other sexually transmitted agents and host genetic factors) [9–11]. Despite the role of some genetic factors is being emphasized in scientific community, there is still a great discussion on their potential inclusion clinically.

Immunologic factors are especially important in the prevention of pathogenic infections, and therefore they are expected to have great impact on viral associated diseases where they act as markers of individual susceptibility [12–14]. The interleukin 1 receptor antagonist (IL-1RN) is an important immunologic regulator that competes with other interleukin 1 family members (IL-1 $\alpha$  and IL-1 $\beta$ ) for the IL-1 receptor in target cells, acting as its negative regulator with anti-inflammatory effect [15–18]. Its expression blocks pro-inflammatory signals from unwanted cellular damages during extensive immunologic responses [15–22].

Literature refers the existence of several polymorphism on the *IL-1RN* gene sequence, and despite the majority are single nucleotide polymorphisms [23, 24], the majority of studies have focused on a common variable number of tandem repeats (VNTR) polymorphism characterized by a 86 bp variation within intron 2 [25–28]. Genetic studies have shown that the distribution of IL-1RN alleles is associated with ancestral distribution with increased distribution of alleles 1 and 2 and repression of all other alleles [29, 30]. The role of IL-1RN VNTR has being studied for many years in the development of inflammatory disorders [19, 22, 31], nevertheless great discussion has being held on the potential role of this polymorphism in cancer development [28, 30]. In fact, several studies showed different associations of IL-1RN VNTR alleles with the development of several cancers, such as of gastric [26–28, 32–39], oesophageal [40], bladder [41–43], breast [44], colorectal [45, 46], lung [47–49], brain [50, 51], nasopharyngeal [52] and gallbladder [41]. Moreover, and despite differences in ethnic distribution [29, 30], some authors have discussed a possible role of this polymorphism in cervical carcinoma [53–56].

Since literature is suggesting that this polymorphism may be correlated with increased susceptibility of viral diseases, including viral-associated neoplasias, we have developed a hospital-based case control study to characterize the role IL1RN VNTR polymorphism in cervical carcinoma development in the Northern region of Portugal.

## Materials and methods

### Cases and specimens

The group of women with cervical lesions consisted in a total of 340 women with median age of 46 years old (Table 1) attended at the Gynaecologic Oncology Department from the Portuguese Institute of Oncology of Porto. Patients were further classified according to histological data: 28 patients with histological confirmed low grade squamous intraepithelial lesions (LSIL); 84 with high grade SIL (HSIL); and 228 patients with invasive cervical cancer (ICC).

The study included a control group constituted by 196 randomly selected healthy females without evidence of neoplastic disease with median age of 44 years old (Table 1) collected from a Healthy Donor's database at the *Institute of Molecular Pathology and Immunology of University of Porto* (IPATIMUP).

Informed consent according to the Declaration of Helsinki was obtained from each individual. Individuals from control group were submitted to a clinical evaluation and have no individual history of cancer.

### Sample processing and genotyping

Freshly collected peripheral blood samples were collected following standard venipuncture technique in EDTA-containing tubes, and the DNA was extracted from the white blood cell fraction using a salting out protocol.

The *IL-1RN* VNTR polymorphism was genotyped by polymerase chain reaction (PCR) with 5'-CCCCTCAGCAACTCC-3' forward primer and 5'-GGTCAGAAGG GCAGAG-3' reverse primer as previously described [26]. PCR products were analysed by electrophoresis in a 3 % Agarose gel (Cambrex Bio Science Rockland Inc, USA) stained with 5 % ethidium bromide for identification of the following fragments: 410 bp (allele 1), 240 bp (allele 2), 325 bp (allele 3), 500 bp (allele 4) and 595 bp (allele 5). All possible genotypes were considered: A1\*A1, A1\*A2, A1\*A3, A1\*A4, A1\*A5, A2\*A2, A2\*A3, A2\*A4, A2\*A5, A3\*A3, A3\*A4, A3\*A5, A4\*A4, A4\*A5 and A5\*A5.

### Statistical analysis

For genetic comparison alleles were combined according to the number of repeats: A2, the short allele that corresponds to two repeats; and L, the long alleles that corresponds to three or more repeats. Hardy-Weinberg equilibrium (HWE) was calculated considering the stratification of IL-1RN genotypes as A2\*A2, A2\*L and L\*L with the public software available at <http://www.oege.org/software/hwe-mr-calc.shtml> [57, 58].

Genotypes were analysed with Statistical Package for Social Sciences for Windows (SPSS version 16.0) using Chi-square ( $\chi^2$ ) analysis to compare the categorical variables with a 5 % significance level. Statistical analysis was performed by considering the status of IL1-RN Allele homozygosity in a dichotomic variable (A2\*A2 homozygous vs non-A2\*A2) with odds ratio (OR) and its 95 % confidence interval (CI) to be used as a measure of the association between the genotype and the risk of developing disease. Logistic regression analysis was also performed by adjusting data for median age of onset. Kaplan–Meier (Log-rank and Breslow test) was used for calculation of

cumulative hazard considering genotype-specific distributions according to age of disease onset.

The sample size and power of analysis was calculated with the OpenEpi: Open Source Epidemiologic Statistics for Public Health, Version 2.3. [www.OpenEpi.com](http://www.OpenEpi.com), updated 2009/20/05, accessed 2012/01/14.

## Results

In this study, it was not possible to identify the following genotypes: A2\*A4, A2\*A5, A3\*A3, A3\*A4, A3\*A5, A4\*A4, A4\*A5 and A5\*A5 (Table 2). HWE was tested and results showed that genotypes were according to the expected in the control group ( $p = 0.363$ ) while patients with cervical lesions did not respect the HWE ( $p < 0.001$ ). The genotype distribution showed that 19.7 % of the cases with cervical lesion/cancer were A2\*A2 homozygous in comparison with 11.7 % in controls, which turn to be statistically different ( $p = 0.017$ ).

When stratifying patients according to the histological data for A2\*A2 genotype, we observed that there were no statistical significant differences between controls and patients with LSIL or HSIL independently ( $p > 0.050$ ). Nevertheless, results have shown an almost twofold increased risk for the development of ICC (OR = 1.85;

**Table 1** Characteristics of the study population

Population description	Mean age ( $\pm$ SD), y.o.	Median age (min–max), y.o.
Controls ( $n = 196$ )	43.7 $\pm$ 15.6	44.0 (18–81)
Cases ( $n = 340$ )	46.6 $\pm$ 12.6	46.0 (20–83)
LSIL ( $n = 28$ )	38.3 $\pm$ 12.2	34.5 (21–65)
HSIL ( $n = 84$ )	41.3 $\pm$ 12.9	38.5 (21–81)
ICC ( $n = 228$ )	49.6 $\pm$ 11.5	48.0 (20–83)

y.o. years old, LSIL low-grade squamous intraepithelial lesion, HSIL high-grade squamous intraepithelial lesion, ICC invasive cervical cancer

**Table 2** Distribution of IL-1RN VNTR genotypes

	A1*A1	A1*A2	A1*A3	A1*A4	A1*A5	A2*A2	A2*A3
Controls ( $n = 196$ )	96 (49.0)	69 (35.2)	6 (3.1)	–	1 (0.5)	23 (11.7)	1 (0.5)
Cervical lesions ( $n = 340$ )	159 (46.8)	106 (31.2)	6 (1.8)	1 (0.3)	–	67 (19.7)	1 (0.3)
LSIL ( $n = 28$ )	14 (50.0)	8 (28.6)	1 (3.6)	–	–	5 (17.9)	–
HSIL ( $n = 84$ )	36 (42.9)	30 (35.7)	1 (1.2)	–	–	17 (20.2)	–
ICC ( $n = 228$ )	109 (47.8)	68 (29.8)	4 (1.8)	1 (0.4)	–	45 (19.7)	1 (0.4)

LSIL low-grade squamous intraepithelial lesion, HSIL high-grade squamous intraepithelial lesion, ICC invasive cervical cancer

Values in parenthesis indicate percentage

**Table 3** Association of IL-1RN VNTR genotypes and cervical cancer development

	Non A2*A2 n (%)	A2*A2 n (%)	$p$	OR	95 % CI	$p^a$	OR <sup>a</sup>	95 % CI <sup>a</sup>
Controls ( $n = 196$ )	173 (88.3)	23 (11.7)		1.00	Reference		1.00	Reference
Cervical lesions ( $n = 340$ )	273 (80.3)	67 (19.7)	<b>0.017</b>	<b>1.85</b>	1.11–3.08	<b>0.017</b>	<b>1.87</b>	1.12–3.11
LSIL ( $n = 28$ )	23 (82.1)	5 (17.9)	0.359	1.64	0.57–4.72	0.275	1.82	0.62–5.36
HSIL ( $n = 84$ )	67 (79.8)	17 (20.2)	0.062	1.91	0.96–3.80	0.067	1.91	0.96–3.80
ICC ( $n = 228$ )	183 (80.3)	45 (19.7)	<b>0.025</b>	<b>1.85</b>	1.07–3.19	<b>0.030</b>	<b>1.85</b>	1.06–3.21
HSIL + ICC ( $n = 312$ )	250 (80.1)	62 (19.9)	<b>0.017</b>	<b>1.87</b>	1.11–3.13	<b>0.015</b>	<b>1.91</b>	1.13–3.21

LSIL low-grade squamous intraepithelial lesion, HSIL high-grade squamous intraepithelial lesion, ICC invasive cervical cancer,  $\chi^2$  Chi-square,  $P$  Pearson  $\chi^2$ , OR odds ratio, CI confidence interval

<sup>a</sup> Logistic regression adjusted for age

Bold values denote the statistical significance

95 % IC 1.07–3.19;  $p = 0.025$ ). Statistical analysis also revealed an almost twofold increased risk for IL1-RN A2\*A2 homozygous of developing lesions more severe than HSIL (OR = 1.87; 95 % IC 1.11–3.13;  $p = 0.017$ ). Data from multivariate logistic regression (Table 3), adjusting the OR for age, revealed no significant differences from the predicted ORs either for ICC only (OR = 1.85; 95 % IC 1.06–3.21;  $p = 0.030$ ) or for HSI-L + ICC (OR = 1.90; 95 % IC 1.13–3.21;  $p = 0.015$ ).

Kaplan–Meier revealed that the median age of onset of cases with cervical lesion/cancer with *IL1RN* A2\*A2 homozygosity was of 46.0 years old comparing to 52.0 in non-A2\*A2 ( $p = 0.042$ )—Fig. 1a. Although we have not found statistical significant data considered the different types of lesion isolated (data not shown), results were similar if considering HSIL and ICC together (48.0 vs 52.0;  $p = 0.028$ )—Fig. 1b.

## Discussion


For decades, cervical cancer has been one of the most studied cancer with thousands of publications from all over the world. Actually, the infection by oncogenic HPVs is established as the necessary cause for cervical cancer development, nevertheless, not all infected women will develop cancer. The establishment of a persistent infection by oncogenic HPVs is the key point for the progression from normal epithelium to pre-invasive lesions and ultimately to ICC.

In the past decade, our group has been developing several studies regarding the role of genetic factors in the

development of several cancers, including cervical cancer [59–62]. Literature has been revealing that HPV infection is influenced by environmental and host conditions, such as diet and host immune status, which can lead to immunosuppressed conditions [63, 64]. Hence, the study of genetic factors involved in immune response regulation is expected to reveal important data on HPV-associated carcinogenesis [61, 65].

The interleukin 1 superfamily combines some of the most important cytokines involved in host-pathogen immune response [20, 66, 67]. While IL-1 $\alpha$  and IL-1 $\beta$  have potent pro-inflammatory activity, the antagonist of the receptor for IL-1 (IL-1RN) avoids the transmission of pro-inflammatory signals abolishing the immune response [14, 17, 18]. Many studies have reported that IL-1RN levels are useful predictors of host immune response, since its levels increase during the final steps of inflammatory response [22]. Moreover, several studies have shown that this molecule is also altered in cancer development [30, 68] either by promoting the suppression of immune response to pathogens [69–73] that are able to promote cancer development or by promoting chronic inflammation states that induce cell lesions and tissue differentiation [33, 35, 74].

Up to date, there are controversial data regarding the potential role of this polymorphism in the development of cervical cancer. While Qian et al., showed no association for cervical cancer development, Tamandani et al., showed protection and Singh et al., and Mustea et al., have proved that the allele 2 is associated with increased risk [53–56]. Hence, in this study we attempted to characterize the role of the VNTR polymorphism within the IL-1RN gene in the development of cervical cancer in Portugal.



**Fig. 1** Association between IL-1RN A2\*A2 genotypes and the age of onset by Kaplan–Meier methodology with log-rank test for: **a** cases with cervical lesion/cancer; and **b** HSIL + ICC

Despite our study may reveal a small sample size, nevertheless, the power of the analysis was of 69.2 % (the sample size required to a power of analysis of 80 % with 95 % IC is of 529 cases and 265 controls). Our study suggests that IL-1RN Allele 2 homozygosis was significantly associated with lesions severe than HSIL with an almost twofold increased risk. This data is in accordance with literature suggestions, since (1) it has been established that in LSILs HPV is frequently transient as some of these lesions regress spontaneously and therefore immune surveillance is active; and (2) in lesions severe than HSIL HPV is supposed to be integrated more often, and with a shorter immune response, it will favour the establishment of persistent infections and the development of high-grade lesions that can progress to cancer [18].

This study characterizes the role of *IL-1RN* VNTR polymorphism in cervical cancer susceptibility in Portugal. IL-1RN Allele 2 homozygosis may be determinant for reduced immune responses in our population contributing for increased susceptibility for HPV infection and development of cervical cancer. Therefore, *IL-1RN* VNTR polymorphism may be a useful marker of host immune response and viral associated cancers susceptibility. Although the evidences pointed in our study, we consider that it would be of extreme importance to obtain data regarding the HPV status and IL-1Ra levels to corroborate our hypothesis, and therefore further studies are required.

**Acknowledgments** We would like to thank to the Portuguese League Against Cancer (*Liga Portuguesa Contra o Cancro-Núcleo Regional do Norte*) for the support on the research group and to the Ministry of Health of Portugal (CFICS-32/2007) for the financial support of this project. First author was supported by an individual grant for Doctoral degree from the Portuguese Ministry of Science, Technology and Superior Education—FCT (*Fundação para a Ciência e Tecnologia*; SFRH/BD/40718/2007).

## References

1. Ferlay J et al (2008) GLOBOCAN 2008, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10 2008
2. Sankaranarayanan R, Ferlay J (2006) Worldwide burden of gynaecological cancer: the size of the problem. *Best Pract Res Clin Obstet Gynaecol* 20(2):207–225
3. zur Hausen H (1991) Human papillomaviruses in the pathogenesis of anogenital cancer. *Virology* 184(1):9–13
4. zur Hausen H, Gissmann L, Schlehofer JR (1984) Viruses in the etiology of human genital cancer. *Prog Med Virol* 30:170–186
5. Franco EL, Rohan TE, Villa LL (1999) Epidemiologic evidence and human papillomavirus infection as a necessary cause of cervical cancer. *J Natl Cancer Inst* 91(6):506–511
6. Bosch FX et al (1993) Human papillomavirus and cervical intraepithelial neoplasia grade III/carcinoma in situ: a case-control study in Spain and Colombia. *Cancer Epidemiol Biomark Prev* 2(5):415–422
7. Munoz N et al (1993) Risk factors for cervical intraepithelial neoplasia grade III/carcinoma in situ in Spain and Colombia. *Cancer Epidemiol Biomark Prev* 2(5):423–431
8. Munoz N et al (1993) Human papilloma virus in the etiology of cervicouterine cancer. *Bol Oficina Sanit Panam* 115(4):301–309
9. Castellsague X, Bosch FX, Munoz N (2002) Environmental cofactors in HPV carcinogenesis. *Virus Res* 89(2):191–199
10. Trottier H, Franco EL (2006) The epidemiology of genital human papillomavirus infection. *Vaccine* 24(Suppl 1):S1–S15
11. zur Hausen H (2009) Human papillomavirus & cervical cancer. *Indian J Med Res* 130(3):209
12. Cavallo MG et al (1992) Viral infection induces cytokine release by beta islet cells. *Immunology* 75(4):664–668
13. Cavallo R, Cavallo F, Cavallo G (1994) The role of cytokines in viral infections. *G Batteriol Virol Immunol* 86(1–12):65–78
14. Hurme M (1998) Is susceptibility to infections in the genes? *Duodecim* 114(8):733–734
15. Granowitz EV et al (1991) Interleukin-1 receptor antagonist competitively inhibits the binding of interleukin-1 to the type II interleukin-1 receptor. *J Biol Chem* 266(22):14147–14150
16. Arend WP et al (1998) Interleukin-1 receptor antagonist: role in biology. *Annu Rev Immunol* 16:27–55
17. Hurme M, Santtila S (1998) IL-1 receptor antagonist (IL-1Ra) plasma levels are co-ordinately regulated by both IL-1Ra and IL-1beta genes. *Eur J Immunol* 28(8):2598–2602
18. Witkin SS, Gerber S, Ledger WJ (2002) Influence of interleukin-1 receptor antagonist gene polymorphism on disease. *Clin Infect Dis* 34(2):204–209
19. Fischer E et al (1992) Interleukin-1 receptor antagonist circulates in experimental inflammation and in human disease. *Blood* 79(9):2196–2200
20. Dinarello CA (1996) Biologic basis for interleukin-1 in disease. *Blood* 87(6):2095–2147
21. Dinarello CA (1991) Interleukin-1 and interleukin-1 antagonism. *Blood* 77(8):1627–1652
22. McIntyre KW et al (1991) Inhibition of interleukin 1 (IL-1) binding and bioactivity in vitro and modulation of acute inflammation in vivo by IL-1 receptor antagonist and anti-IL-1 receptor monoclonal antibody. *J Exp Med* 173(4):931–939
23. Graziano F et al (2009) Variations in the interleukin-1 receptor antagonist gene impact on survival of patients with advanced colorectal cancer. *Pharmacogenomics J* 9(1):78–84
24. Lindmark F et al (2005) Interleukin-1 receptor antagonist haplotype associated with prostate cancer risk. *Br J Cancer* 93(4):493–497
25. Tarlow JK et al (1993) Polymorphism in human IL-1 receptor antagonist gene intron 2 is caused by variable numbers of an 86-bp tandem repeat. *Hum Genet* 91(4):403–404
26. El-Omar EM et al (2000) Interleukin-1 polymorphisms associated with increased risk of gastric cancer. *Nature* 404(6776):398–402
27. El-Omar EM et al (2001) The role of interleukin-1 polymorphisms in the pathogenesis of gastric cancer. *Nature* 412(6842):99
28. Sehoul J et al (2002) Polymorphism of IL-1 receptor antagonist gene: role in cancer. *Anticancer Res* 22(6A):3421–3424
29. Xu DP et al (2011) VNTR polymorphism of human IL1RN in Chinese Han and She ethnic populations. *Int J Immunogenet* 38(1):13–16
30. Maruta Y et al (2008) Determination of ancestral allele for possible human cancer-associated polymorphisms. *Cancer Genet Cytogenet* 180(1):24–29
31. Xue J et al (2010) No association of interleukin-1 receptor antagonist VNTR polymorphism and rheumatoid arthritis susceptibility: a meta analysis. *Clin Exp Rheumatol* 28(5):654–660
32. Persson C et al (2011) Polymorphisms in inflammatory response genes and their association with gastric cancer: a HuGE systematic review and meta-analyses. *Am J Epidemiol* 173(3):259–270
33. Peleteiro B et al (2010) Association between cytokine gene polymorphisms and gastric precancerous lesions: systematic



- review and meta-analysis. *Cancer Epidemiol Biomark Prev* 19(3):762–776
34. Machado JC et al (2001) Interleukin 1B and interleukin 1RN polymorphisms are associated with increased risk of gastric carcinoma. *Gastroenterology* 121(4):823–829
  35. Machado JC et al (2003) A proinflammatory genetic profile increases the risk for chronic atrophic gastritis and gastric carcinoma. *Gastroenterology* 125(2):364–371
  36. Figueiredo C et al (2002) *Helicobacter pylori* and interleukin 1 genotyping: an opportunity to identify high-risk individuals for gastric carcinoma. *J Natl Cancer Inst* 94(22):1680–1687
  37. Perri F et al (2010) Role of interleukin polymorphisms in gastric cancer: “pros and cons”. *World J Gastrointest Oncol* 2(6):265–271
  38. Xue H et al (2010) Interleukin-1B and interleukin-1 RN polymorphisms and gastric carcinoma risk: a meta-analysis. *J Gastroenterol Hepatol* 25(10):1604–1617
  39. He BS et al (2011) Polymorphisms in interleukin-1B (IL-1B) and interleukin 1 receptor antagonist (IL-1RN) genes associate with gastric cancer risk in the Chinese population. *Dig Dis Sci* 56(7):2017–2023
  40. Upadhyay R et al (2008) Potential influence of interleukin-1 haplotype IL-1 beta-511\*T-IL-1RN\*1 in conferring low risk to middle third location of esophageal cancer: a case-control study. *Hum Immunol* 69(3):179–186
  41. Vishnoi M et al (2008) IL-1 gene polymorphisms and genetic susceptibility of gallbladder cancer in a north Indian population. *Cancer Genet Cytogenet* 186(2):63–68
  42. Ahirwar DK et al (2009) Cytokine gene polymorphisms are associated with risk of urinary bladder cancer and recurrence after BCG immunotherapy. *Biomarkers* 14(4):213–218
  43. Bid HK, Manchanda PK, Mittal RD (2006) Association of interleukin-1Ra gene polymorphism in patients with bladder cancer: case control study from North India. *Urology* 67(5):1099–1104
  44. Lee KM et al (2006) Genetic polymorphisms of interleukin-1 beta (IL-1B) and IL-1 receptor antagonist (IL-1RN) and breast cancer risk in Korean women. *Breast Cancer Res Treat* 96(2):197–202
  45. Viet HT et al (2005) Interleukin-1 receptor antagonist gene polymorphism in human colorectal cancer. *Oncol Rep* 14(4):915–918
  46. Lurje G et al (2009) Polymorphisms in interleukin 1 beta and interleukin 1 receptor antagonist associated with tumor recurrence in stage II colon cancer. *Pharmacogenet Genomics* 19(2):95–102
  47. Lind H et al (2005) Interleukin 1 receptor antagonist gene polymorphism and risk of lung cancer: a possible interaction with polymorphisms in the interleukin 1 beta gene. *Lung Cancer* 50(3):285–290
  48. Lim WY et al (2011) Polymorphisms in inflammatory pathway genes, host factors and lung cancer risk in Chinese female never-smokers. *Carcinogenesis* 32(4):522–529
  49. Hu Z et al (2006) Allele 2 of the interleukin-1 receptor antagonist gene (IL1RN\*2) is associated with a decreased risk of primary lung cancer. *Cancer Lett* 236(2):269–275
  50. Settin A, Ali N, Salem FK (2008) Cytokine gene polymorphisms in Egyptian cases with brain tumors. *Egypt J Immunol* 15(2):15–23
  51. Badr El-Din NK et al (2009) Cytokine gene polymorphisms in Egyptian cases with brain tumors. *J Egypt Natl Cancer Inst* 21(2):101–106
  52. Yang ZH et al (2011) Association of IL-1 polymorphisms and IL-1 serum levels with susceptibility to nasopharyngeal carcinoma. *Mol Carcinog* 50(3):208–214
  53. Tamandani DM et al (2008) Impact of polymorphism in IL-1RA gene on the risk of cervical cancer. *Arch Gynecol Obstet* 277(6):527–533
  54. Singh H et al (2008) Genetic variants of interleukin-1RN and interleukin-1beta genes and risk of cervical cancer. *BJOG* 115(5):633–638
  55. Mustea A et al (2003) Interleukin 1 receptor antagonist (IL-1RA) polymorphism in women with cervical cancer. *Anticancer Res* 23(2A):1099–1102
  56. Qian N et al (2010) Circulating IL-1beta levels, polymorphisms of IL-1B, and risk of cervical cancer in Chinese women. *J Cancer Res Clin Oncol* 136(5):709–716
  57. Rodriguez S, Gaunt TR, Day IN (2009) Hardy–Weinberg equilibrium testing of biological ascertainment for Mendelian randomization studies. *Am J Epidemiol* 169(4):505–514
  58. Lauretto MS et al (2009) A straightforward multiallelic significance test for the Hardy–Weinberg equilibrium law. *Genet Mol Biol* 32(3):619–625
  59. Catarino R et al (2005) Increased risk of cervical cancer associated with cyclin D1 gene A870G polymorphism. *Cancer Genet Cytogenet* 160(1):49–54
  60. Costa S et al (2002) A slow acetylator genotype associated with an increased risk of advanced cervical cancer. *J Cancer Res Clin Oncol* 128(12):678–682
  61. Duarte I et al (2005) G-308A TNF-alpha polymorphism is associated with an increased risk of invasive cervical cancer. *Biochem Biophys Res Commun* 334(2):588–592
  62. Craveiro R et al (2004) TP73 alterations in cervical carcinoma. *Cancer Genet Cytogenet* 150(2):116–121
  63. Farrell PJ (2002) Tumour viruses—could they be an Achilles’ heel of cancer? *Eur J Cancer* 38(14):1815–1816
  64. Vousden KH, Farrell PJ (1994) Viruses and human cancer. *Br Med Bull* 50(3):560–581
  65. Coelho A et al (2007) The influence of chemokine receptor CCR2 genotypes in the route to cervical carcinogenesis. *Gynecol Obstet Invest* 64(4):208–212
  66. Dinarello CA (1994) The interleukin-1 family: 10 years of discovery. *FASEB J* 8(15):1314–1325
  67. Dinarello CA, Wolff SM (1993) The role of interleukin-1 in disease. *N Engl J Med* 328(2):106–113
  68. Kapoor S (2009) Role of polymorphisms of interleukin-1 receptor antagonist gene in systemic oncogenesis: significant etiopathologic connections besides in benign prostatic hyperplasia. *Urology* 73(1):215
  69. Roberge CJ et al (1996) The IL-1 and IL-1 receptor antagonist (IL-1Ra) response of human neutrophils to EBV stimulation. Preponderance of IL-1Ra detection. *J Immunol* 156(12):4884–4891
  70. Hurme M, Helminen M (1998) Polymorphism of the IL-1 gene complex in Epstein-Barr virus seronegative and seropositive adult blood donors. *Scand J Immunol* 48(3):219–222
  71. Rocha GA et al (2005) IL1RN polymorphic gene and cagA-positive status independently increase the risk of noncardia gastric carcinoma. *Int J Cancer* 115(5):678–683
  72. Queiroz DM et al (2004) IL1B and IL1RN polymorphic genes and *Helicobacter pylori* cagA strains decrease the risk of reflux esophagitis. *Gastroenterology* 127(1):73–79
  73. Wang Y et al (2003) Interleukin-1beta gene polymorphisms associated with hepatocellular carcinoma in hepatitis C virus infection. *Hepatology* 37(1):65–71
  74. Canedo P et al (2008) The interferon gamma receptor 1 (IFNGR1) -56C/T gene polymorphism is associated with increased risk of early gastric carcinoma. *Gut* 57(11):1504–1508