

Inhibition of CYP 1A2-Dependent MROD Activity in Rat Liver Microsomes: An Explanation of the Hepatic Sequestration of a Limited Subset of Halogenated Aromatic Hydrocarbons

Jin Jun Chen,* Guo Sheng Chen, Nigel J. Bunce*

Toxicology Program, University of Guelph, Guelph, Ontario, N1G 2W1, Canada

Received 18 July 2002; revised 15 November 2002; accepted 25 November 2002

ABSTRACT: Many classes of halogenated aromatic compounds (HACs) are highly lipophilic environmental contaminants that exert toxic effects via the Ah receptor signal transduction pathway and whose metabolism generally involves monooxygenase enzymes of the CYP 1A family. Despite their lipophilicity, a high proportion of the body burden of certain polychlorinated dibenzo-*p*-dioxins and coplanar polychlorinated biphenyls is sequestered in liver, a process believed to involve CYP 1A2. In this work we examined HAC-induced inhibition of the demethylation of 7-methoxyresorufin, a process that is selectively catalyzed by CYP 1A2. 2,3,7,8-Tetrachlorodibenzo-*p*-dioxin, 3,3',4,4'-tetrachlorobiphenyl (PCB 77) and 3,3',4,4',5-pentachlorobiphenyl (PCB 126) were found to be strong competitive inhibitors of methoxyresorufin-O-demethylase activity, consistent with the high ability of hepatic tissue to sequester these compounds selectively. © 2003 Wiley Periodicals, Inc. *Environ Toxicol* 18: 115–119, 2003.

Keywords: MROD assay; CYP 1A2; enzyme kinetics; halogenated aromatic compounds; liver sequestration

INTRODUCTION

Halogenated aromatic compounds (HACs), including polychlorinated dibenzo-*p*-dioxins (PCDDs), dibenzofurans (PCDFs), biphenyls (PCBs), and polybrominated diphenyl ethers (PBDEs), are highly lipophilic environmental contaminants that arise through incineration (PCDDs and PC-

DFs), improper disposal practices (PCBs), and open use as flame retardants (PBDEs). They are found in the environment as complex mixtures, with food usually the principal route of human exposure (Diliberto et al., 1999). The related compounds, polycyclic aromatic hydrocarbons (PAHs), are released into the environment as a result of incomplete combustion.

The toxicity of the planar or near-planar members of these classes, of which 2,3,7,8-tetrachlorodibenzo-*p*-dioxin (TCDD) is a highly toxic prototype, is mediated through the Ah receptor signal transduction pathway (Wang et al., 1997). One end point of the latter is the induction of monooxygenase enzymes of the cytochrome P-450 (CYP) 1A family (Sutter et al., 1992; Tritscher et al., 1992; Van den Berg et al., 1994; Drahuskuk et al., 1999). CYP 1A1 and 1A2 catalyze the metabolism of many xenobiotics,

Correspondence to: Nigel J. Bunce, Department of Chemistry and Biochemistry, University of Guelph, Guelph, Ontario, N1G 2W1, Canada; e-mail: bunce@chembio.uoguelph.ca.

*Present address: College of Animal Sciences and Veterinary Medicine, Northwest Sci-Tech University of Agriculture and Forestry, Yangling, Shaanxi, 712100 China.

Contract grant sponsor: Natural Sciences and Engineering Council and Canadian Toxic Substances Research Institute.

Published online in Wiley InterScience (www.interscience.wiley.com). DOI 10.1002/tox.10107

© 2003 Wiley Periodicals, Inc.




Fig. 1. Deethylation of 7-ethoxyresorufin.

including drugs, environmental pollutants, and chemical carcinogens (Guengerich, 1988; Nebert, 1989). The CYP-dependent monooxygenase system, which comprises CYP, NADPH-CYP reductase, and phospholipids, plays a major role in the metabolism of HACs and PAHs, sometimes activating these unreactive contaminants into more reactive, toxic metabolites (Burke et al., 1995).

CYP 1A1 and 1A2 are monooxygenase enzymes with broad and overlapping substrate specificities. In both humans and rodents CYP 1A2 is constitutively expressed in the liver in much greater amounts than CYP 1A1, although the latter is highly inducible by planar xenobiotics such as TCDD and by certain PAHs such as 3-methylcholanthrene (Hu and Bunce, 1999a). Administration of TCDD to male C57BL/6J mice increased both CYP 1A1 and CYP 1A2 hepatic protein levels, whereas the nonplanar PCB 153 (2,2',4,4',5,5'-hexachlorobiphenyl) increased the levels of CYP 1A2 but not CYP 1A1. Moreover, CYP 1A2 activities (but not protein levels) in mice cotreated with TCDD and PCB 153 were higher than those of mice treated with TCDD alone (De Jongh et al., 1995). CYP 1A1 and 1A2 may be distinguished by immunoblotting or by exploiting their selectivity toward particular substrates. CYP 1A1, for example, preferentially deethylates 7-ethoxyresorufin, whereas CYP 1A2 is selective for demethylation of 7-methoxyresorufin (Sohn et al., 1994)—EROD (ethoxyresorufin-*O*-deethylase) and MROD (methoxyresorufin-*O*-demethylase) activities, respectively. Both EROD and MROD involve oxidation of the substrate at the aliphatic carbon atom α to the ether linkage, affording a hemiacetal that spontaneously dissociates to give resorufin, which can be quantitated by fluorescence spectroscopy (Fig. 1, shown for EROD). Activation of the Ah receptor by a given HAC correlates with the synthesis of the CYP 1A protein and with increased enzyme activity (Petrulis et al., 2000; Chen and Bunce, 2001). Other possible biomarkers for CYP 1A2 induction include induction of estradiol-2-hydroxylase activity (Graham et al., 1988) and acetanilide-4-hydroxylase (ACOH; Liu et al., 1991). Nerurkar et al. (1993) showed that both ACOH and MROD activities correlated well with CYP 1A2 protein levels as determined by Western blot analysis, although MROD activity is preferred because of speed and convenience (DeVito et al., 1997; Nerurkar et al., 1993).

Most HACs are highly lipophilic, as reflected in octanol-water partition coefficients ($K_{ow} > 10^5$) and therefore tend to accumulate in biota. Indeed, the release of these compounds from adipose tissue may be the rate-limiting step in their elimination (Van den Berg, 1994; Hu and Bunce,

1999b). Sequestration in adipose tissue may also explain the long whole-body half-lives of substances like TCDD in large mammals, for example, adult humans (5–9 years) versus rodents (weeks). The specific problem addressed in this article is the tendency of a small group of HACs, typified by TCDD and the coplanar PCBs, to be sequestered in liver (DeVito et al., 1998; Van Birgelen et al., 1994) and, in PCB 77 (3,3',4,4'-tetrachlorobiphenyl), to be mainly metabolized by CYP 1A2 (Kramer et al., 2000). Sequestration was hypothesized to be the result of a specific interaction between TCDD—and presumably the coplanar PCBs—and CYP 1A2. In this study we measured the competitive inhibition constants (K_i) of several HACs in the MROD assay, using microsomes prepared from the livers of immature male Sprague-Dawley rats, as a measure of their affinities for CYP 1A2.

MATERIALS AND METHODS

Chemicals

2,3,7,8-Tetrachlorodibenzo-*p*-dioxin (TCDD) was a generous gift from Wellington Laboratories (Guelph, Ontario, Canada); 2,2',4,4',5,5'-hexachlorobiphenyl (PCB 153), 3,3',4,4'-tetrachlorobiphenyl (PCB 77), 2,2',4,4'-tetrabromodiphenyl ether (PBDE 47), 3,3',4,4'-tetrabromodiphenyl ether (PBDE 77), 2,2',3,4,4'-pentabromodiphenyl ether (PBDE 85), 2,3',4,4',6-pentabromodiphenyl ether (PBDE 119), and 3,3',4,4',5-pentabromodiphenyl ether (PBDE 126) were synthesized in this laboratory; 1,2,3,6,7,8-hexachlorodibenzofuran (1,2,3,6,7,8-HexaCDF), 2,3,4,6,7,8-hexachlorodibenzofuran (2,3,4,6,7,8-HexaCDF), 2,2',5,5'-tetrachlorobiphenyl (PCB 52), 3,3',4,4',5-pentachlorobiphenyl (PCB 126), 2,2',4,4',6,6'-hexachlorobiphenyl (PCB 155), 2,3,3',4,4',5-hexachlorobiphenyl (PCB 156), 3,3',4,4',5,5'-hexachlorobiphenyl (PCB 169) were purchased from Accustandard Inc. (New Haven, CT); 1-methylphenanthrene (1-MPA), benzo[*a*]pyrene (BaP), benzo[*e*]pyrene (BeP) were obtained from Supelco Canada (Mississauga, Ontario); dimethyl sulfoxide (DMSO), MgSO₄, and tris-(hydroxymethyl)-aminomethane (TRIS) were purchased from Fisher Scientific Co. (Toronto, Ontario, Canada); β -NADP⁺ was purchased from CALBIOCHEM (La Jolla, CA); Resorufin was purchased from ICN Biomedicals Inc. (Aurora, OH); sodium pentobarbital (Somnitel®) was obtained from MTC Pharmaceuticals (Cambridge, Ontario, Canada); 3-methylcholanthrene (3-MC), β -naphthoflavone (β -NF), Percoll, Hank's balanced salt solution, Dulbecco's phosphate buffered saline (PBS), DL-isocitric acid, isocitric dehydrogenase, NADH, BSA, and 7-MR were purchased from Sigma Chemical Co. (St. Louis, MO).




Fig. 2. Comparison of competitive inhibition of CYP 1A2 catalytic MROD activity by TCDD, PCB 77, and PCB 126, respectively, in rat hepatic microsomes.

Animal Treatment and Microsome Preparation

Male Sprague-Dawley rats (100 g) obtained from Charles River Breeding Laboratories (Canada) were administered 100 μmol 3-MC in corn oil per kilogram by intraperitoneal injection on 4 consecutive days. On the fifth day, the animals were sacrificed by CO_2 /cervical dislocation. The livers were perfused *in situ* with ice-cold buffer [10 mM HEPES, 1 mM EDTA, 10% (v/v) glycerol, pH 7.6]. The livers were excised, rinsed with buffer, minced with scissors and rinsed again. The tissue was homogenized and spun at $9000 \times g$ for 20 min at 4°C . Surface lipid was removed by aspiration, and the supernatant was spun at $100\,000 \times g$ for 68 min in a Beckman L7-65 ultracentrifuge at 4°C . The pellet was rinsed, resuspended in buffer, and spun a second time. The microsomal pellet was suspended in buffer and stored in 1-mL aliquots at -70°C . Microsomal protein concentrations were determined according to the method of Bradford (1976).

Kinetics of MROD Inhibition in Rat Hepatic Microsomes

Rat hepatic microsomes (stored with EROD activity of 85 nmol resorufin $\text{min}^{-1} \text{mL}^{-1}$) were diluted 100-fold with 10 mM TRIS buffer (pH 7.4). Ten microliters was then added to each well of a 48-well microplate containing 450 μL of NADPH-generating system (5 mM MgSO_4 , 7.5 mM DL-isocitric acid, 5 mM $\beta\text{-NADP}^+$, isocitrate dehydrogenase at 0.5 units/mL, NADH at 0.5 mg/mL, and BSA at 0.5 mg/mL in 50 mM TRIS-HCl, pH 7.4). To this, HAC in 2.5 μL of DMSO was added, followed by incubation for 5 min at 23°C . Inhibition curves were generated by adding various concentrations of the substrate 7-MR, incubating for a further 15 min and assaying for the fluorescent product resorufin by monitoring fluorescence at 590 nm using a Bio-Rad FLUOstar fluorescent plate reader (BMG Lab Technologies, Germany).

Data Analysis

Through the use of a standard resorufin curve, fluorescence intensities were converted to reaction rates (V , nmol resorufin $\text{min}^{-1} \text{mL}^{-1}$). Spreadsheet software (Microsoft Excel '97) was used to obtain linear regressions between $1/V$ and $1/[7\text{-MR}]$ for reactions carried out in the absence and in the presence of at least two concentrations of the inhibitor. The slope of the regression line between $1/V$ and $1/[7\text{-MR}]$ is $(K_M/V_{\text{max}}) \times ([\text{HAC}]/K_i)$, from which K_i may be determined.

RESULTS AND DISCUSSION

Our principal objective was to determine K_i for a wide range of Ah receptor ligands that might interact with CYP 1A2. TCDD and other HACs showed dose-dependent decreases in MROD activity with an increasing concentration of HAC. The slopes of the Lineweaver-Burk plots were calculated using linear regression, and a comparison of the slope in the presence of inhibitor with the slope in the absence of inhibitor was used to calculate the inhibition constant (K_i) for each compound. $K_i = [\text{Inhibitor}]/(\alpha - 1)$, where α is the ratio of the slopes in the presence and absence of inhibitor. Figure 2 shows typical examples of the data.

Each experiment was replicated twice; values in Table I are the means of the replicates; overall variation in all cases was $< 5\%$. TCDD and PCB 126 bound strongly to CYP 1A2, with $K_i \sim 3 \times 10^{-8} \text{ M}$, with PCB 169, $K_i \sim 1 \times 10^{-7} \text{ M}$, consistent with their sequestration in the liver, where CYP 1A2 is particularly abundant (DeVito et al., 1997; Chen CY et al., 2001). PCB 77 also bound strongly to CYP 1A2, with $K_i \sim 3 \times 10^{-8} \text{ M}$, but it has not been reported to sequester in the liver because of rapid metabolism (Chen CY et al., 2001). 2,3,4,7,8-Pentachlorodibenzofuran has also been reported to be sequestered in hepatic tissue (DeVito et al., 1997), but we did not have a sample of this compound available.

TABLE I. Competitive inhibition constants (K_i) of aromatic compounds to MROD activity (CYP 1A2)

Congeners	K_i (nM)	Congeners	K_i (nM)
TCDD	33	1-MPA	320
1,2,3,6,7,8-HexaCDF	620	3-MC	92
2,3,4,6,7,8-HexaCDF	610	β -NF	50
PCB 52	80	BaP	46
PCB 77	31	BeP	65
PCB 126	26	PBDE 47	800
PCB 153	80	PBDE 77	400
PCB 155	1200	PBDE 85	1200
PCB 156	220	PBDE 119	2000
PCB 169	110	PBDE 126	400

Among the other compounds studied, attention may be drawn to β -naphthoflavone, B[a]P, B[e]P, and 3-MC, all with K_i values in the range of $5\text{--}9 \times 10^{-8}$ M; despite being Ah receptor agonists, they do not sequester in the liver because they are readily metabolized (Gonzalez et al., 1994; Zhang et al., 1997; Petrulis et al., 2000). Hepatic sequestration by CYP 1A2 is thus limited to Ah receptor agonists that resist Phase I metabolism. We also studied several PBDE congeners, which are flame retardants that are increasingly found in the environment and in biota (Darnerud et al., 2001). These HACs are only weak Ah receptor agonists (Chen G et al., 2001); none of them was a significant inhibitor of MROD activity.

Unlike CYP 1A1, CYP 1A2 is constitutively expressed in hepatic tissues, and its induction occurs through tissue-specific mechanisms (Goldstein and Linko, 1983; Tukey and Nebert, 1984). That CYP 1A2 is the binding protein for dioxin in the liver is shown by the lack of sequestration of TCDD in the livers of the CYP 1A2 knockout mice (Diliberto et al., 1997). Also in contrast to CYP 1A1, CYP 1A2 is not highly inducible by PAHs (Hu and Bunce, 1999a). The tendency of TCDD, PCB 77, and PCB 126 to bind CYP 1A2 is thus of potential toxicological significance, in that such an association may deplete the pool of CYP 1A2 available for metabolizing other xenobiotics. Whether this is positive or negative from the perspective of toxicity depends on the substrate. Some xenobiotic substrates for CYP 1A, such as PAHs, aromatic amines, and aflatoxins, are bioactivated into carcinogenic species (Crespi et al., 1989, 1991; Shimada et al., 1989; Gonzalez et al., 1994; Eaton et al., 1995; Rendic et al., 1997). In these cases the inhibition of CYP 1A2 by HACs might be expected to exert a protective effect.

Recent work indicates widening toxicological implications for CYP 1A2—sequestering HACs such as TCDD and PCBs 77 and 126. PCB 126 was already known as the most toxic PCB congener in the context of Yusho poisoning, an incident of human poisoning in Japan resulting from consumption of PCB-contaminated rice oil (Yoshimura et al., 1979). PCB 126 is also known to depress the hepatic concentrations of class 1 alcohol dehydrogenase (Ishii et al., 2001), aldolase B, carbonic anhydrase III, glucose-regulated protein 78 (GRP78), GRP94, calreticulin, and calnexin (Ishii et al., 1997; Tasaki et al., 1999; Ikeda et al., 2000; Yoshioka et al., 2001).

We thank the Chinese government for support of a study leave by J.J.C.

REFERENCES

- Bradford MM. 1976. A rapid and sensitive method for quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254.
- Burke MD, Thompson S, Elcombe CR, Halpert J, Haaparanta T, Mayer RT. 1985. Ethoxy-, pentoxy-, and benzyloxyphenoxas-ones and homologues: A series of substrates to distinguish between different induced cytochromes P450. *Biochem Pharmacol* 34:3337–3345.
- Chen CY, Hamm JT, Hass JR, Birnbaum LS. 2001. Disposition of polychlorinated dibenzo-*p*-dioxins, dibenzofurans, and non-ortho polychlorinated biphenyls in pregnant Long Evans rats and the transfer to offspring. *Toxicol Appl Pharmacol* 173:65–88.
- Chen G, Bunce NJ. 2000. Polybrominated diphenyl ether congeners as Ah receptor agonists and antagonists. *Organohalogen Comp* 53:353–356.
- Chen G, Konstantinov AD, Chittim BG, Joyce EM, Bols NC, Bunce NJ. 2001. Synthesis of polybrominated diphenyl ethers and their capacity to induce CYP 1A by the Ah receptor mediated pathway. *Environ Sci Technol* 35:3749–3756.
- Crespi CL, Langenbach R, Rudo K, Chen YT, Davies RL. 1989. Transfection of a human cytochrome P-450 gene into the human lymphoblastoid cell line, AHH-1, and use of the recombinant cell line in gene mutation assays. *Carcinogenesis* 10:295–301.
- Crespi CL, Penman BW, Steimel DT, Gelboin HV, Gonzalez FJ. 1991. The development of a human cell line stably expressing human CYP 3A4: Role in the metabolic activation of aflatoxin B1 and comparison to CYP1A2 and CYP2A3. *Carcinogenesis* 12:355–359.
- Darnerud PO, Eriksen GS, Johannesson T, Larsen PB, Viluksela M. 2001. Polybrominated diphenyl ethers: Occurrence, dietary exposure, and toxicology. *Environ Health Perspect* 109:49–68.
- De Jongh J, DeVito M, Nieboer R, Birnbaum L, Van den Berg M. 1995. Induction of cytochrome P450 isoenzymes after toxicokinetic interactions between 2,3,7,8-tetrachlorodibenzo-*p*-dioxin and 2,2',4,4',5,5'-hexachlorobiphenyl in the liver of the mouse. *Fund Appl Toxicol* 25:264–270.
- DeVito MJ, Diliberto JJ, Ross DG, Menache MG, Birnbaum LS. 1997. Dose-response relationship for polyhalogenated dioxins and dibenzofurans following subchronic treatment in mice. 1: CYP 1A1 and CYP1A2 enzyme activities in liver, lung, and skin. *Toxicol Applied Pharmacol* 147:267–280.
- DeVito MJ, Ross DG, Dupuy AE Jr, Ferrario J, McDaniel D, Birnbaum LS. 1998. Dose-response relationships for disposition and hepatic sequestration of polyhalogenated dibenzo-*p*-dioxins, dibenzofurans, and biphenyls following subchronic treatment in mice. *Toxicol Sci* 46:223–234.
- Diliberto JJ, Burgin D, Birnbaum LS. 1997. Role of CYP1A2 in hepatic sequestration of dioxin: Studies using CYP1A2 knockout mice. *Biochem Biophys Res Commun* 236:431–433.
- Diliberto JJ, Burgin DE, Birnbaum LS. 1999. Effects of CYP 1A2 on disposition of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin, 2,3,4,7,8-pentachlorodibenzofuran, and 2,2',4,4',5,5'-hexachlorobiphenyl in CYP 1A2 knockout and parental (C57BL/6N and 129/sv) strains of mice. *Toxicol Appl Pharmacol* 159:52–64.
- Drahushuk AT, McGarrigle BP, Slezak BP, Stegeman JJ, Olson JR. 1999. Time- and concentration-dependent induction of CYP1A1 and CYP1A2 in precision-cut rat liver slices incubated in dynamic organ culture in the presence of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin. *Toxicol Applied Pharmacol* 155:127–138.
- Eaton DL, Gallagher EP, Bammler TK, Kunze KL. 1995. Role of cytochrome P4501A2 in chemical carcinogenesis: Implications

- for human variability in expression and enzyme activity. *Pharmacogenetics* 5:259–274.
- Goldstein JA, Linko P. 1984. Differential induction of two 2,3,7,8-tetrachlorodibenzo-*p*-dioxin-inducible forms of cytochrome P-450 in extrahepatic versus hepatic tissues. *Mol Pharmacol* 25:185–191.
- Gonzalez FJ, Gelboin HV. 1994. Role of human cytochromes P450 in the metabolic activation of chemical carcinogenesis and toxins. *Drug Metab Rev* 26:165–183.
- Graham MJ, Lucier GW, Linko P, Maronpot RR, Goldstein JA. 1988. Increases in cytochrome P-450 mediated 17- β -estradiol 2-hydroxylase activity in rat liver microsomes after both acute administration and subchronic administration of 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in a two-stage hepatocarcinogenesis model. *Carcinogenesis* 9:1935–1941.
- Guengerich FP. 1988. Roles of cytochrome P-450 enzymes in chemical carcinogenesis and cancer chemotherapy. *Cancer Res* 48:2946–2954.
- Hu K, Bunce NJ. 1999a. Metabolism of polychlorinated dibenzo-*p*-dioxins by rat liver microsomes. *J Mol Biochem Toxicol* 13:307–315.
- Hu K, Bunce NJ. 1999b. Metabolism of polychlorinated dibenzo-*p*-dioxins and related dioxin-like compounds. *J Toxicol Environ Health B* 2:183–210.
- Ikeda M, Ishii Y, Kato H, Akazawa D, Hatsumura M, Ishida T, Matsusue K, Yamada H, Oguri K. 2000. Suppression of carbonic anhydrase III in rat liver by a dioxin-related toxic compound, coplanar polychlorinated biphenyl, 3,3',4,4',5-pentachlorobiphenyl. *Arch Biochem Biophys* 380:159–164.
- Ishii Y, Kato H, Hatsumura M, Ishida T, Ariyoshi N, Oguri K. 1997. Significant suppression of rat liver aldolase B by a toxic coplanar polychlorinated biphenyl, 3,3',4,4',5-pentachlorobiphenyl. *Toxicology* 116:193–199.
- Ishii Y, Kato H, Hatsumura M, Ishida T, Ariyoshi N, Yamada H, Oguri K. 2001. Role of the dioxin-like toxic compound coplanar polychlorinated biphenyl, 3,3',4,4',5-pentachlorobiphenyl in reducing hepatic alcohol dehydrogenase levels in rats *in vivo*. *J Health Sci* 47:575–578.
- Kramer HJ, Drenth H-J, Olinga P, Groothuis G, Van Holstein I, Maas R, DeJongh J, Van den Berg M. 2000. Identification of cytochrome P450 enzymes involved in the biotransformation of PCB 77, PCB 136 and Ugilec 141 isomers using human hepatic microsomes. *Organohalogen Comp* 49:299–302.
- Liu G, Gelboin HV, Myers MJ. 1991. Role of cytochrome P450 1A2 in acetanilide 4-hydroxylation as determined with cDNA expression and monoclonal antibodies. *Arch Biochem Biophys* 284:400–406.
- Nebert DW. 1989. The Ah locus; genetic differences in toxicity, cancer, mutation, and birth defects. *Crit Rev Toxicol* 20:153–174.
- Nerurkar PV, Park SS, Thomas PE, Nims RW, Lubet RA. 1993. Methoxyresorufin and benzyloxyresorufin: Substrates preferentially metabolized by cytochromes P4501A2 and 2B, respectively, in the rat and mouse. *Biochem Pharmacol* 46:933–943.
- Petrulis JR, Chen G, Benn S, LaMarre J, Bunce NJ. 2000. Application of the ethoxyresorufin-*O*-deethylase (EROD) assay to mixtures of halogenated aromatic compounds. *Environ Toxicol* 11:177–184.
- Rendic S, Di Carlo FJ. 1997. Human cytochrome P450 enzymes: A status report summarizing their reactions, substrates, inducers, and inhibitors. *Drug Metab Rev* 29:413–580.
- Safe SH. 1993. Toxicology, structure–function relationship, and human environmental health impacts of polychlorinated biphenyls: Progress and problems. *Environ Health Perspect* 100:259–268.
- Safe SH. 1994. Polychlorinated biphenyls (PCBs): Environmental impact, biochemical and toxic responses, and implications for risk assessment. *Crit Rev Toxicol* 24:87–149.
- Shimada T, Iwasaki M, Martin MV, Guengerich FP. 1989. Human liver microsomal cytochrome P-450 enzymes involved in the bioactivation of procarcinogens detected by Umu gene response in *Salmonella typhimurium* TA 1535/pSK1002. *Cancer Res* 49:3218–3228.
- Sohn OS, Surace A, Fiala ES, Richie JP Jr, Colosimo S, Zang E, Weisburger JH. 1994. Effects of green and black tea on hepatic xenobiotic metabolizing systems in the male F344 rat. *Xenobiotica* 24:119–127.
- Sutter TR, Greenlee WF. 1992. Classification of members of the Ah gene battery. *Chemosphere* 25:223–226.
- Tasaki K, Ishii Y, Ishida T, Oguri K. 1999. Suppression of stress proteins in endoplasmic reticulum in liver cytosol of rats treated with a highly toxic coplanar PCB. *Fukuoka Igaku Zasshi* 90: 251–258 (in Japanese).
- Tritscher AM, Goldstein JA, Portier CJ, McCoy Z, Clark GC, Lucier GW. 1992. Dose–response relationships for chronic exposure to 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in a rat liver tumor promotion model: Quantitation and immunolocalization of CYP1A1 and CYP1A2 in the liver. *Cancer Res* 52:3436–3442.
- Tukey RH, Nebert DW. 1984. Regulation of mouse cytochrome P3-450 by the Ah receptor. Studies with a P3-450 cDNA clone. *Biochemistry* 23:6003–6008.
- Van Birgelen AP, Van der Kolk J, Fase KM, Bol I, Poiger H, Brouwer A, Van den Berg M. 1994. Toxic potency of 3,3',4,4',5-pentachlorobiphenyl relative to and in combination with 2,3,7,8-tetrachlorodibenzo-*p*-dioxin in a subchronic feeding study in the rat. *Toxicol Appl Pharmacol* 127:209–221.
- Van den Berg M, De Jongh J, Poiger H, Olson JR. 1994. The toxicokinetics and metabolism of polychlorinated dibenzo-*p*-dioxins (PCDDs) and dibenzofurans (PCDFs) and their relevance for toxicity. *Crit Rev Toxicol* 24:1–74.
- Wang X, Santostefano MJ, Evans MV, Richardson VM, Diliberto JJ, Birnbaum LS. 1997. Determination of parameters responsible for pharmacokinetic behavior of TCDD in female Sprague–Dawley rats. *Toxicol Applied Pharmacol* 147:151–168.
- Yoshimura H, Yoshihara S, Ozawa N, Miki M. 1979. Possible correlation between induction modes of hepatic enzymes by PCBs and their toxicity in rats. *Ann NY Acad Sci* 320:179–192.
- Yoshioka Y, Ishii Y, Ishida T, Yamada H, Oguri K, Motojima K. 2001. Suppression of stress proteins, GRP78, GRP94, calreticulin, calnexin in liver endoplasmic reticulum of rat treated with a highly toxic coplanar PCB. *Fukuoka Igaku Zasshi* 92:201–216 (in Japanese).
- Zhang QY, Wikoff J, Dunbar D, Fasco M, Kaminsky L. 1997. Regulation of cytochrome P4501A1 expression in rat small intestine. *Drug Metab Dispos* 25:21–26.